

Winter Meeting, 4–5 December 2018, Optimal diet and lifestyle strategies for the management of cardio-metabolic risk

Nutritional factors and cardiovascular disease risk in Black African and Black Caribbean women: a cross-sectional study

C. Ewlierhoma¹, A.P. Moore², L.M. Goff², A. Aghili¹, S. Comegna³, G. Begum³ and A.R.A. Adegboye⁴

¹London Metropolitan University, 166-120 Holloway Road, London N7 8DB,

²King's College London, Stamford Street, London SE1 9NH,

³University of Westminster, 115 New Cavendish Street, London W1W 6UW and

⁴University of Greenwich, Avery Hill Campus, Eltham, SE9 2UG.

People of Black African (BA) and Caribbean (BC) heritage form the third largest ethnic group in England and Wales⁽¹⁾. Evidence shows they experience higher rates of overweight/obesity, stroke and type 2 diabetes compared to the general population but lower risk of heart disease, which may be explained by the favourable lipid profile they exhibit^(2,3). There are limited UK studies on their dietary habits and health. The aim of the current study was to assess nutritional intake and cardiovascular disease (CVD) risk factors in UK BA and BC women.

A convenience sample of self-ascribed BA and BC women, aged 19-64 years, were recruited (n = 44) from the ATTITUDE study. Cholesterol was measured using a portable CardioChek Blood Analyser, blood pressure using a digital blood pressure monitor and dietary intake via triple pass 24hr recall. Ethical approval was obtained from London Metropolitan University, King's College London and Westminster University. Percentage energy, total fat, saturated fat, carbohydrate, and fibre and salt intake were calculated and under-reporting was assessed using the Goldberg equation⁽⁵⁾. A sensitivity analysis conducted on nutrient intakes with under-reporters removed.

Nutritional intake and CVD risk factors are shown in Table 1. Sixty three percent of participants were overweight or obese. Analysis of the dietary data revealed higher intakes salt, free sugars, fat and saturated fat than recommendations and lower intake of carbohydrate⁽⁴⁾. Sensitivity analysis was conducted to investigate the impact of under-reporting (n = 22). Reported data remained unchanged except for fibre intake, which was lower in the under-reporters (14.9g compared to 21.0g) (P = 0.004). Blood lipid profiles and blood pressure data were within recommendations.

Table 1. CVD risk factors and nutritional intake data (n = 44)

	Mean or median (%)	SD or [IQR (25 th ,75 th)]	% of participants not within recommended limits
Age	34.2	11.3	
Weight (kg)	78.6	20.4	
BMI (kg/m ²)	26.5	[23.7, 32.6]	Overweight 27 %, Obese 25 %, Morbidly obese 11 %
Waist Circum. (cm)	89.9	16.0	68 % above recommendation of <80cm
Waist:Hip Ratio	0.8	0.1	
Total Cholesterol (mmol/L)	4.1	0.9	
HDL-cholesterol (mmol/L)	1.5	0.4	
TotalCholesterol/HDL-cholesterol	2.8	0.7	
Blood pressure (systol.)	114.6	9.3	
Blood pressure (diastol.)	72.6	9.3	
Dietary Intake			
CHO (% total energy)	45 %	8	75 % below recommendation of 50 %
Free sugar (% total energy)	8 %	5	64 % above recommendation of 5 %
Fibre (g)	17.0	7.4	93 % below recommendation of 30g
Saturated Fat (% total energy)	12 %	5	55 % above recommendation of 11 %
Fat (% total energy)	36 %	6	45 % above recommendation of 35 %
Salt (g)	5.4	3.2	30 % above recommendation of 6g

In conclusion, the anthropometric and certain dietary measures would indicate increased risk for developing CVD in BA and BC women, however, blood lipid profile and blood pressure measures were within healthy ranges.

- Office for National Statistics. (2012). *Ethnicity and National Identity in England and Wales* 2011.
- Lip G, Barnett A, Bradbury A *et al.* (2007) *J Hum Hypertens*, **21**:183–211.
- Goff LM, Griffin BA, Lovegrove JA *et al.* (2013) *Diab Vasc Dis Res*, **10**(4):315–323.
- British Nutrition Foundation (2017) *Nutrition Requirements*. www.nutrition.org.uk
- Goldberg GR, Black AE, Jebb SA *et al.* (1991) *Eur J Clin Nutr*, **45**:569–581.