

QUASI-CONVEXITY AND HERMITE-HADAMARD TYPE INEQUALITIES

(Kekuasicembungan dan Ketaksamaan Jenis Hermite-Hadamard)

ALAWIAH IBRAHIM

ABSTRACT

In this paper, some inequalities connected with the left and right hand side of Hermite-Hadamard type inequalities are established for functions whose the first and second derivatives absolute values are quasi-convex.

Keywords: Hermite-Hadamard inequality; convex function; quasi-convex function

ABSTRAK

Dalam makalah ini, beberapa ketaksamaan bersabit dengan ketaksamaan jenis Hermite-Hadamard sebelah kiri dan kanan diperoleh untuk fungsi yang nilai mutlak terbitan peringkat pertama dan keduanya adalah fungsi kuasi-cembung.

Kata kunci: ketaksamaan Hermite-Hadamard; fungsi-cembung; fungsi kuasi-cembung

1. Introduction

Let I be an interval of real numbers. A function $f : I \rightarrow \mathbb{R}$ is said to be convex on I if the inequality

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y) \quad (1)$$

holds for all $x, y \in I$ and $t \in [0, 1]$. If the inequality (1) is reversed, then f is said to be concave.

The following double inequality,

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2} \quad (2)$$

is well known in the literature as the Hermite-Hadamard integral inequality (see Dragomir & Agarwal 1998), which provides the estimates of the mean value of a continuous convex function defined on an real interval $[a, b]$ with $a < b$. Inequality (2) has attracted many researchers over the years to demonstrate new proofs, refinements, extensions and generalization, see for instance Niculescu and Persson (2003), Ciurdariu (2012), Iscan (2013), González *et al.* (2015), Khan *et al.* (2018) and the references cited therein.

Definition 1.1. A function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ is said to be quasi-convex on I if

$$f(tx + (1-t)y) \leq \max\{f(x), f(y)\} \quad (3)$$

holds for all $x, y \in I$ and $t \in [0, 1]$. If f is quasi-concave, then inequality (3) is reversed.

Dragomir and Argawal (1998), Pearce and Pécarić (2000), Sarikaya and Aktan (2011) provided the inequalities connected with the right part of (3) for functions whose first and second derivatives are convex. Similar results also have been established by Ion (2007) and Alomari *et al.* (2010) for the class of functions whose derivatives absolute values are quasi-convex, where some of the results are pointed out as follows.

Theorem 1.1. (Ion 2007) Assume $a, b \in \mathbb{R}$ with $a < b$ and $f : [a, b] \rightarrow \mathbb{R}$ is a differentiable function on (a, b) , then we have the inequalities

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \begin{cases} \frac{(b-a)}{4} \sup\{|f'(a)|, |f'(b)|\} & (4) \\ \text{for } |f'| \text{ is quasi-convex on } [a, b], \\ \frac{(b-a)}{2(p+1)^{\frac{1}{p}}} \left(\sup\{|f'(a)|^{\frac{p}{p-1}}, |f'(b)|^{\frac{p}{p-1}}\} \right)^{\frac{p-1}{p}}, p > 1 & (5) \\ \text{for } |f'|^{\frac{p}{p-1}} \text{ is quasi-convex on } [a, b]. \end{cases}$$

Theorem 1.2. (Alomari *et al.* 2010) Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a twice differentiable function on the interior points of I , $a, b \in I$ with $a < b$ and f'' be integrable on $[a, b]$. Then we have

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \begin{cases} \frac{(b-a)^2}{12} \sup\{|f''(a)|, |f''(b)|\} & (6) \\ \text{for } |f''| \text{ is quasi-convex on } [a, b], \\ \frac{(b-a)^2}{8} \left(\frac{\sqrt{\pi}}{2} \right)^{\frac{1}{p}} \left(\frac{\Gamma(1+p)}{\Gamma(\frac{3}{2}+p)} \right)^{\frac{1}{p}} & (7) \\ \times \left(\sup\{|f'(a)|^{\frac{p}{p-1}}, |f'(b)|^{\frac{p}{p-1}}\} \right)^{\frac{p-1}{p}}, p > 1 \\ \text{for } |f'|^{\frac{p}{p-1}} \text{ is quasi-convex on } [a, b]. \end{cases}$$

Theorem 1.3. (Alomari *et al.* 2010) Let $L[a, b]$ be class of continuous functions defined on $[a, b]$. Let $f : I \subseteq [0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. If $|f'|^{\frac{p}{p-1}}$, $p > 1$ is quasi-convex on $[a, b]$. Then we have

$$\left| \frac{f(a)+f(b)}{2} - \frac{1}{b-a} \int_a^b f(x)dx \right| \leq \frac{(b-a)}{4(p+1)^{\frac{1}{p}}} \left[\left(\max \left\{ \left| f' \left(\frac{a+b}{2} \right) \right|^{\frac{p}{p-1}}, |f'(b)|^{\frac{p}{p-1}} \right\} \right)^{\frac{p-1}{p}} + \left(\max \left\{ \left| f' \left(\frac{a+b}{2} \right) \right|^{\frac{p}{p-1}}, |f'(a)|^{\frac{p}{p-1}} \right\} \right)^{\frac{p-1}{p}} \right]. \quad (8)$$

The main purpose of this study is to provide some more inequalities related with the Hermite-Hadamard type inequality for twice differentiable functions whose the absolute values of the first and second derivatives are quasi-convex. In particular, we generalise the result of Ion (2007).

2. Some Inequalities of Hermite-Hadamard Type for Quasi-Convex Functions

In order to prove our main results, we need the following integral equalities.

Lemma 2.1. (Iqbal et al. 2012) Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be differentiable function on the interior points of I , where $a, b \in I$ with $a < b$. If $f' \in L[a, b]$ and $\mu, \lambda \in [0, \infty)$ with $\mu + \lambda \neq 0$, then

$$\begin{aligned} & \frac{\mu f(a) + \lambda f(b)}{\mu + \lambda} - \frac{1}{b-a} \int_a^b f(x)dx \\ &= \frac{b-a}{(\mu + \lambda)^2} \left[\int_0^\mu (-t) f' \left(\frac{\lambda + t}{\mu + \lambda} a + \frac{\mu - t}{\mu + \lambda} b \right) dt + \int_0^\lambda t f' \left(\frac{\lambda - t}{\mu + \lambda} a + \frac{\mu + t}{\mu + \lambda} b \right) dt \right] \end{aligned} \quad (9)$$

Lemma 2.2. (Xi & Qi 2013) Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be a twice differentiable function on the interior points of I and $\lambda \in \mathbb{R}$. If $f'' \in L[a, b]$ where $a, b \in I$ with $a < b$, then

$$\begin{aligned} & \lambda \frac{f(a) + f(b)}{2} + (1 - \lambda) f \left(\frac{a+b}{2} \right) - \frac{1}{b-a} \int_a^b f(x)dx \\ &= \frac{(b-a)^2}{16} \int_0^1 t(2\lambda - t) \left[f'' \left((1-t)a + t \frac{a+b}{2} \right) + f'' \left(t \frac{a+b}{2} + (1-t)b \right) \right] dt. \end{aligned} \quad (10)$$

Lemma 2.3. (Xi & Qi 2013) Let $0 \leq \lambda \leq 1$, $t \in [0, 1]$ and $r > -1$. Then,

$$\int_0^1 t |2\lambda - t|^r dt = \begin{cases} \frac{(2\lambda)^{r+2} + (2\lambda + r + 1)(1 - 2\lambda)^{r+1}}{(r+1)(r+2)}, & 0 \leq \lambda \leq \frac{1}{2}, \\ \frac{(2\lambda)^{r+2} - (2\lambda + r + 1)(2\lambda - 1)^{r+1}}{(r+1)(r+2)}, & \frac{1}{2} \leq \lambda \leq 1. \end{cases} \quad (11)$$

We are now in a position to establish some new integral inequalities of Hermite-Hadamard type for twice differentiable and quasi-convex functions.

Theorem 2.1. Assume $a, b \in \mathbb{R}$ with $a < b$ and $f : [a, b] \rightarrow \mathbb{R}$ is a differentiable function on (a, b) . Then for $\mu, \lambda \in [0, \infty)$ with $\mu + \lambda \neq 0$, the following inequalities hold true

$$\left| \frac{\mu f(a) + \lambda f(b)}{\mu + \lambda} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \begin{cases} \frac{(b-a)(\mu^2 + \lambda^2)}{2(\mu + \lambda)^2} \max\{|f'(a)|, |f'(b)|\} \\ \text{for } |f'| \text{ is quasi-convex on } [a, b], \end{cases} \quad (12)$$

$$\left. \begin{cases} \frac{(b-a)(\mu^2 + \lambda^2)}{(\mu + \lambda)^2 (p+1)^{\frac{1}{p}}} \left(\max\{|f'(a)|^{\frac{p}{p-1}}, |f'(b)|^{\frac{p}{p-1}}\} \right)^{\frac{p-1}{p}}, p > 1 \\ \text{for } |f'|^{\frac{p}{p-1}} \text{ is quasi-convex on } [a, b]. \end{cases} \right\} \quad (13)$$

Proof. By Lemma 2.1 with the fact that $|f'|$ is a quasi-convex function on $[a, b]$ and some elementary integration calculus we have

$$\begin{aligned} & \left| \frac{\mu f(a) + \lambda f(b)}{\mu + \lambda} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \frac{b-a}{(\mu + \lambda)^2} \left[\int_0^\mu \left| f' \left(\frac{\lambda+t}{\mu+\lambda} a + \frac{\mu-t}{\mu+\lambda} b \right) \right| dt + \int_0^\lambda \left| f' \left(\frac{\lambda-t}{\mu+\lambda} a + \frac{\mu+t}{\mu+\lambda} b \right) \right| dt \right] \\ & \leq \frac{b-a}{(\mu + \lambda)^2} \max\{|f'(a)|, |f'(b)|\} \left[\int_0^\mu t dt + \int_0^\lambda t dt \right] \\ & = \frac{b-a}{(\mu + \lambda)^2} \left(\frac{\mu^2 + \lambda^2}{2} \right) \max\{|f'(a)|, |f'(b)|\}. \end{aligned}$$

Also, from Lemma 2.1 and applying the weighted Hölder integral inequality, one obtains

$$\begin{aligned} & \left| \frac{\mu f(a) + \lambda f(b)}{\mu + \lambda} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \frac{b-a}{(\mu + \lambda)^2} \left[\left(\int_0^\mu t^p dt \right)^{\frac{1}{p}} \left(\int_0^\mu \left| f' \left(\frac{\lambda+t}{\lambda+\mu} a + \frac{\mu-t}{\lambda+\mu} b \right) \right|^q dt \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\int_0^\lambda t^p dt \right)^{\frac{1}{p}} \left(\int_0^\lambda \left| f' \left(\frac{\lambda-t}{\lambda+\mu} a + \frac{\mu+t}{\lambda+\mu} b \right) \right|^q dt \right)^{\frac{1}{q}} \right], \end{aligned}$$

where $p > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$. Using the fact that $|f'|^q$ is quasi convex function on $[a, b]$, then we have

$$\begin{aligned} & \left| \frac{\mu f(a) + \lambda f(b)}{\mu + \lambda} - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \frac{b-a}{(\mu + \lambda)^2} \left(\max \{ |f'(a)|^q, |f'(b)|^q \} \right)^{\frac{1}{q}} \left[\left(\int_0^{\mu} t^p dt \right)^{\frac{1}{p}} \left(\int_0^{\mu} dt \right)^{\frac{1}{q}} + \left(\int_0^{\lambda} t^p dt \right)^{\frac{1}{p}} \left(\int_0^{\lambda} dt \right)^{\frac{1}{q}} \right] \\ & = \frac{b-a}{(\mu + \lambda)^2} \left(\max \{ |f'(a)|^q, |f'(b)|^q \} \right)^{\frac{1}{q}} \left[\left(\frac{\mu^{p+1}}{p+1} \right)^{\frac{1}{p}} \mu^{\frac{1}{q}} + \left(\frac{\lambda^{p+1}}{p+1} \right)^{\frac{1}{p}} \lambda^{\frac{1}{q}} \right] \\ & = \frac{(b-a)(\mu^2 + \lambda^2)}{(\mu + \lambda)^2 (p+1)^{\frac{1}{p}}} \left(\max \{ |f'(a)|^q, |f'(b)|^q \} \right)^{\frac{1}{q}}. \end{aligned}$$

The proof of Theorem 2.1 is completed. \square

Remark 2.1. If in Theorem 2.1 we choose $\mu = \lambda = 1$, then we recapture the Ion's results (4) and (5).

Theorem 2.2. Assume $a, b \in \mathbb{R}$ with $a < b$ and $f : [a, b] \rightarrow \mathbb{R}$ is a twice differentiable function on $[a, b]$ such that $f''(x)$ is integrable. If $|f''|$ and $|f''|^{\frac{p}{p-1}}$ ($p > 1$) are quasi-convex functions on $[a, b]$, then for $0 \leq \lambda \leq 1$, the following inequalities hold true:

$$\left| \lambda \frac{f(a) + f(b)}{2} + (1-\lambda) f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(x) dx \right|$$

$$\left[\begin{aligned} & \frac{(b-a)^2}{16} H_1(\lambda) \left[\max \left\{ |f''(a)|, \left| f''\left(\frac{a+b}{2}\right) \right| \right\} + \max \left\{ \left| f''\left(\frac{a+b}{2}\right) \right|, |f''(b)| \right\} \right] \quad (14) \\ & \text{for } |f''| \text{ is quasi-convex on } [a, b], \\ & \leq \frac{(b-a)^2}{16} \left(H_p(\lambda) \right)^{\frac{1}{p}} \left(\frac{1}{2} \right)^{\frac{p-1}{p}} \times \left[\left(\max \left\{ |f''(a)|^{\frac{p}{p-1}}, \left| f''\left(\frac{a+b}{2}\right) \right|^{\frac{p}{p-1}} \right\} \right)^{\frac{p-1}{p}} \right. \\ & \quad \left. + \left(\max \left\{ \left| f''\left(\frac{a+b}{2}\right) \right|^{\frac{p}{p-1}}, |f''(b)|^{\frac{p}{p-1}} \right\} \right)^{\frac{p-1}{p}} \right], p > 1 \quad (15) \\ & \text{for } |f''|^{\frac{p}{p-1}} \text{ is quasi-convex on } [a, b], \end{aligned} \right.$$

where

$$H_1(\lambda) = \begin{cases} \frac{8\lambda^3 - 3\lambda + 1}{3}, & 0 \leq \lambda \leq \frac{1}{2}, \\ \frac{3\lambda - 1}{3}, & \frac{1}{2} \leq \lambda \leq 1, \end{cases} \quad (16)$$

and

$$H_p(\lambda) = \begin{cases} \frac{(2\lambda)^{p+2} + (2\lambda + p + 1)(1 - 2\lambda)^{p+1}}{(p + 1)(p + 2)}, & 0 \leq \lambda \leq \frac{1}{2}, \\ \frac{(2\lambda)^{p+2} - (2\lambda + p + 1)(2\lambda - 1)^{p+1}}{(p + 1)(p + 2)}, & \frac{1}{2} \leq \lambda \leq 1. \end{cases} \quad (17)$$

Proof: From Lemma 2.2 we have

$$\begin{aligned} & \left| \lambda \frac{f(a) + f(b)}{2} + (1 - \lambda)f\left(\frac{a + b}{2}\right) - \frac{1}{b - a} \int_a^b f(x) dx \right| \\ & \leq \frac{(b - a)^2}{16} \int_0^1 |t(2\lambda - t)| \left(\left| f''\left((1 - t)a + t\frac{a + b}{2}\right) \right| + \left| f''\left(t\frac{a + b}{2} + (1 - t)b\right) \right| \right) dt \end{aligned} \quad (18)$$

Since $|f''|$ is a quasi convex function on $[a, b]$, thus from (18) we get

$$\begin{aligned} & \left| \lambda \frac{f(a) + f(b)}{2} + (1 - \lambda)f\left(\frac{a + b}{2}\right) - \frac{1}{b - a} \int_a^b f(x) dx \right| \\ & \leq \frac{(b - a)^2}{16} \left(\max \left\{ |f''(a)|, \left| f''\left(\frac{a + b}{2}\right) \right| \right\} + \max \left\{ \left| f''\left(\frac{a + b}{2}\right) \right|, |f''(b)| \right\} \right) \int_0^1 |t(2\lambda - t)| dt \\ & \leq \frac{(b - a)^2}{16} H_1(\lambda) \left(\max \left\{ |f''(a)|, \left| f''\left(\frac{a + b}{2}\right) \right| \right\} + \max \left\{ \left| f''\left(\frac{a + b}{2}\right) \right|, |f''(b)| \right\} \right), \end{aligned}$$

where from Lemma 2.3 we have that

$$H_1(\lambda) = \begin{cases} \frac{8\lambda^3 - 3\lambda + 1}{3}, & 0 \leq \lambda \leq \frac{1}{2}, \\ \frac{3\lambda - 1}{3}, & \frac{1}{2} \leq \lambda \leq 1. \end{cases}$$

Hence, we obtain the desired result (14).

Again, from Lemma 2.2 and applying the weighted Hölder's inequality for integrals, we have

$$\begin{aligned} & \left| \lambda \frac{f(a)+f(b)}{2} + (1-\lambda)f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(x)dx \right| \\ & \leq \frac{(b-a)^2}{16} \left[\left(\int_0^1 t|2\lambda-t|^p dt \right)^{\frac{1}{p}} \left(\int_0^1 t|f''((1-t)a+t\frac{a+b}{2})|^q dt \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\int_0^1 t|2\lambda-t|^p dt \right)^{\frac{1}{p}} \left(\int_0^1 t|f''(t\frac{a+b}{2}+(1-t)b)|^q dt \right)^{\frac{1}{q}} \right] \\ & = \frac{(b-a)^2}{16} (H_p(\lambda))^{\frac{1}{p}} \left[\left(\int_0^1 t|f''((1-t)a+t\frac{a+b}{2})|^q dt \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\int_0^1 t|f''(t\frac{a+b}{2}+(1-t)b)|^q dt \right)^{\frac{1}{q}} \right] \end{aligned}$$

for $p, q > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$ and $H_p(\lambda) = \int_0^1 t|2\lambda-t|^p dt$. Therefore, by the quasi-convexity of $|f''|^q$, we obtain

$$\begin{aligned} & \left| \lambda \frac{f(a)+f(b)}{2} + (1-\lambda)f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(x)dx \right| \\ & \leq \frac{(b-a)^2}{16} (H_p(\lambda))^{\frac{1}{p}} \left(\frac{1}{2} \right)^{\frac{1}{q}} \left[\left(\max \left\{ |f''(a)|^q, |f''\left(\frac{a+b}{2}\right)|^q \right\} \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\max \left\{ |f''\left(\frac{a+b}{2}\right)|^q, |f''(b)|^q \right\} \right)^{\frac{1}{q}} \right] \\ & = \frac{(b-a)^2}{16} (H_p(\lambda))^{\frac{1}{p}} \left(\frac{1}{2} \right)^{\frac{p-1}{p}} \left[\left(\max \left\{ |f''(a)|^{\frac{p}{p-1}}, |f''\left(\frac{a+b}{2}\right)|^{\frac{p}{p-1}} \right\} \right)^{\frac{p-1}{p}} \right. \\ & \quad \left. + \left(\max \left\{ |f''\left(\frac{a+b}{2}\right)|^{\frac{p}{p-1}}, |f''(b)|^{\frac{p}{p-1}} \right\} \right)^{\frac{p-1}{p}} \right], \end{aligned}$$

where $H_p(\lambda)$ as given in (17). Thus, we complete the proof of Theorem 2.2. \square

References

- Alomari M., Darus M. & Dragomir S. S. 2010. New Inequalities of Hermite-Hadamard type for functions whose second derivatives absolute values are quasi-convex. *Tamkang Journal of Mathematics* **41**(4): 357-359.
- Ciurdariu L. 2012. A note concerning several Hermite-Hadamard type inequalities for different types of convex functions. *International Journal of Mathematical Analysis* **6**(33): 1623-1637.
- Dragomir S. S & Argawal R. P. 1998. Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula. *Applied Mathematics Letter* **11**(5): 91-95.
- González L., Merentes N. & Valera-López. 2015. Some estimates on the Hermite-Hadamard inequality through convex and quasi-convex stochastic processes. *Mathematica Aeterna* **5**(5): 745-767.
- Ion D. A. 2007. Some estimates on the Hermite-Hadamard inequality through quasi-convex functions. *Annals of University of Craiova, Mathematics and Computation Science Ser.* **34**: 82-87.

- Iqbal M., Bhatti M. I. & Dragomir S. S. 2012. Hadamard-type inequalities through h -convexity. *Proceedings of the Pakistan Academy of Sciences* **49**(4): 301-307.
- Iscan I. 2013. Some generalized Hermite-Hadamard type inequalities for quasi-geometrically convex functions. *American Journal of Mathematical Analysis* **1**(3): 48-52.
- Khan M. A., Khurshid Y., Du T-S. & Chu Y-M. 2018. Generalization of Hermite-Hadamard type inequalities via conformable fractional integrals. *Journal of Function Spaces* **2018**: 1-12.
- Niculescu C. P. & Persson L-E. 2003/2004. Old and new on the Hermite-Hadamard inequality. *Real Analysis Exchange* **29**(2): 663-685.
- Pearce C. E. M & Pečarić J. 2000. Inequalities for differentiable mappings with applications to special means and quadrature formulae. *Applied Mathematics Letter*. **13**(2): 51-55.
- Sarikaya M. Z. & Aktan N. 2011. On the generalization of some integral inequalities and their applications. *Mathematical and Computer Modelling* **54**: 2175-2182.
- Xi B-Y. & Qi A. F. 2013. Some inequalities of Hermite-Hadamard type for h -convex functions. *Advances in Inequalities and Applications* **2**(1): 1-15.

*Centre for Modelling and Data Science
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
43600 UKM Bangi
Selangor DE, MALAYSIA
E-mail: alaibra@ukm.edu.my**

*Corresponding author