

THE UNIVERSITY OF WARWICK

Original citation:

Gould, Phillip Spencer, Dyer, Nigel, Croft, Wayne D., Ott, Sascha and Easton, Andrew. (2014) Cellular mRNAs access second ORFs using a novel amino acid sequence-dependent coupled translation termination-reinitiation mechanism. RNA . ISSN 1355-8382

Permanent WRAP url:

<http://wrap.warwick.ac.uk/58872>

Copyright and reuse:


The Warwick Research Archive Portal (WRAP) makes this work of researchers of the University of Warwick available open access under the following conditions.

This article is made available under the Creative Commons Attribution 3.0 (CC BY 3.0) license and may be reused according to the conditions of the license. For more details see: <http://creativecommons.org/licenses/by/3.0/>

A note on versions:

The version presented in WRAP is the published version, or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: publications@warwick.ac.uk



<http://wrap.warwick.ac.uk>



RNA

A PUBLICATION OF THE RNA SOCIETY

Cellular mRNAs access second ORFs using a novel amino acid sequence-dependent coupled translation termination –reinitiation mechanism

Phillip S. Gould, Nigel P. Dyer, Wayne Croft, et al.

RNA published online January 10, 2014

Supplemental Material <http://rnajournal.cshlp.org/content/suppl/2014/01/06/rna.041574.113.DC1.html>

P<P Published online January 10, 2014 in advance of the print journal.

Open Access Freely available online through the *RNA* Open Access option.

Creative Commons License This article, published in *RNA*, is available under a Creative Commons License (Attribution 3.0 Unported), as described at <http://creativecommons.org/licenses/by/3.0/>.

Email Alerting Service Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

**Exiqon Grant
Program 2014**

Accelerate your RNA discoveries
with a grant from Exiqon

EXIQON

To subscribe to *RNA* go to:
<http://rnajournal.cshlp.org/subscriptions>

Cellular mRNAs access second ORFs using a novel amino acid sequence-dependent coupled translation termination–reinitiation mechanism

PHILLIP S. GOULD,¹ NIGEL P. DYER,² WAYNE CROFT,² SASCHA OTT,² and ANDREW J. EASTON^{1,3}

¹School of Life Sciences, ²Warwick Systems Biology Centre, University of Warwick, Coventry CV4 7AL, United Kingdom

ABSTRACT

Polycistronic transcripts are considered rare in the human genome. Initiation of translation of internal ORFs of eukaryotic genes has been shown to use either leaky scanning or highly structured IRES regions to access initiation codons. Studies on mammalian viruses identified a mechanism of coupled translation termination–reinitiation that allows translation of an additional ORF. Here, the ribosome terminating translation of ORF-1 translocates upstream to reinitiate translation of ORF-2. We have devised an algorithm to identify mRNAs in the human transcriptome in which the major ORF-1 overlaps a second ORF capable of encoding a product of at least 50 aa in length. This identified 4368 transcripts representing 2214 genes. We investigated 24 transcripts, 22 of which were shown to express a protein from ORF-2 highlighting that 3' UTRs contain protein-coding potential more frequently than previously suspected. Five transcripts accessed ORF-2 using a process of coupled translation termination–reinitiation. Analysis of one transcript, encoding the CASQ2 protein, showed that the mechanism by which the coupling process of the cellular mRNAs was achieved was novel. This process was not directed by the mRNA sequence but required an aspartate-rich repeat region at the carboxyl terminus of the terminating ORF-1 protein. Introduction of wobble mutations for the aspartate codon had no effect, whereas replacing aspartate for glutamate repeats eliminated translational coupling. This is the first description of a coordinated expression of two proteins from cellular mRNAs using a coupled translation termination–reinitiation process and is the first example of such a process being determined at the amino acid level.

Keywords: coupled translation; translation initiation; second ORF

INTRODUCTION

The eukaryotic translation machinery exploits a number of processes to control gene expression in a wide range of fundamental cellular processes (Aitken and Lorsch 2012). The majority of eukaryotic mRNAs are monocistronic, expressing a single polypeptide from the 5' proximal open reading frame (ORF). If the sequence surrounding the first AUG is not favorable the ribosome may use leaky scanning to initiate translation at the next AUG to generate additional proteins (Kozak 1997). Ribosomes can be detected translating alternative reading frames (Wilson and Masel 2011; Michel et al. 2012). Additional translational initiation mechanisms include the use of internal ribosome entry sites (IRES), ribosomal shunting, and coupled translation (Ahmadian et al. 2000; Rogers et al. 2004; Spriggs et al. 2008). These mecha-

nisms have the potential to expand the coding capacity of the genome.

In bacteria, where polycistronic mRNAs are common, ribosomes can scan bidirectionally around termination codons prior to reinitiation on upstream or downstream AUG codons (Adhin and van Duin 1990). In eukaryotes, termination of translation of a large 5' proximal ORF followed by reinitiation of translation on the same mRNA to access a second ORF has been considered a rare event, first demonstrated with the hepatitis B virus P mRNA and also with artificially made mRNAs (Peabody and Berg 1986; Kozak 1989). In the other cases where reinitiation has been seen the upstream ORF frequently does not encode a substantive protein with a defined function and its presence generally reduces translation of the downstream ORF (Pöyry et al. 2004; Jackson et al. 2010). Examples have also emerged where the small upstream ORF has a biological role (Tautz 2009). The M2 mRNAs of all pneumoviruses contain two open reading

³Corresponding author

E-mail a.j.easton@warwick.ac.uk

Article published online ahead of print. Article and publication date are at <http://www.rnajournal.org/cgi/doi/10.1261/rna.041574.113>. Freely available online through the RNA Open Access option.

© 2014 Gould et al. This article, published in RNA, is available under a Creative Commons License (Attribution 3.0 Unported), as described at <http://creativecommons.org/licenses/by/3.0/>.

frames, conserved in location though not in sequence. ORF-1 utilizes 60%–75% of the coding capacity of the mRNA and encodes the M2-1 protein product, the virus transcriptional activator (Collins and Wertz 1985; Ling et al. 1992; Ahmadian et al. 1999; Fearn and Collins 1999). We have demonstrated that ribosomes access and translate the second ORF in vivo in a controlled process by utilizing the three AUG codons located upstream of the ORF-1 termination codon, and expression from these initiation codons requires the prior termination of M2 ORF-1 translation (Ahmadian et al. 2000; Gould and Easton 2005). The RSV M2-2 protein produced by coupled translation is thought to be involved in control of the switch between virus RNA transcription and replication (Bermingham and Collins 1999). Extending the distance between the M2 ORF-1 termination codon and the M2 ORF-2 initiation codon from the 32-nt maximum observed in vivo to the 72-nt ablated translation of ORF-2. These data also demonstrated that translation of the second ORF is not due to the presence of an IRES sequence. The region of overlap of the M2 ORF-1 and ORF-2 alone is not sufficient to achieve coupled expression but also requires additional sequences located within the M2-1 ORF (Gould and Easton 2005). Importantly, in all of these studies the data were generated in cells in which no other virus genes were being expressed, indicating that the coupled translation process is, in principle, an option available to all cells. Coupled translation mechanisms have also been described in other pneumoviruses and in a number of caliciviruses and influenza B virus (Luttermann and Meyers 2007, 2009; Powell et al. 2008, 2011). The mechanism directing the coupling process differs in caliciviruses and influenza B. Here both require a short section of the mRNA, including a motif that binds to 18S rRNA, just upstream of the overlap (Luttermann and Meyers 2007, 2009; Powell et al. 2008, 2011).

Since the coupled translation process functions in the absence of any viral proteins we considered the possibility that coupled translation may occur with cellular mRNAs. We have screened the human genome for mRNAs containing overlapping ORFs where the second ORF was at least 150 nt in length and contained at least one AUG codon upstream of the ORF-1 stop codon, as seen with the pneumovirus M2 mRNAs. The algorithm would also identify candidate mRNAs that could utilize the mechanism seen in caliciviruses. This identified 4368 transcripts representing 2214 genes. We have demonstrated that the majority of the transcripts analyzed (22 of 24 tested) express proteins from ORF-2 and that five of these genes achieve this using a coupled translation process previously described only in viral transcripts.

The mechanism identified here for the five human transcripts was different from the two previously characterized viral mechanisms. Here, the amino acid sequence at the carboxyl terminus of the protein encoded by ORF1 modulates the coupling processes.

RESULTS

Identification of additional coding capacity within cellular transcripts

We selected 24 human transcripts from a candidate list for analysis (full details supplied in Supplemental Table 1). The selection was based on the gene demonstrating a high probability of translational coupling using the scoring algorithm. The transcript encoding MOCS2 has previously been shown to access ORF-2 using leaky scanning to synthesize a component of the functional molybdopterin synthase enzyme and this was included as a control (Sloan et al. 1999; Stallmeyer et al. 1999). All other second ORFs were previously uncharacterized and no protein products from these ORFs have been described. Expression from ORF-2 was investigated by insertion of a CAT reporter gene lacking its endogenous AUG initiation codon and detection using an ELISA as described previously (Fig. 1A; Ahmadian et al. 2000; Gould and Easton 2005, 2007). This showed that 22 of the initial 24 transcripts studied expressed the CAT protein product while two transcripts, *MYADM* and *ARSD*, did not (Supplemental Table 1). Thus 92% of transcripts screened were able to express a protein from the additional ORF.

Translational regulation of ORF-2

To screen for the utilization of coupled translation to access the ORF-2 the first nucleotide of the stop codon of ORF-1 in each construct was mutated. The next in-frame stop codon was 36 nt further downstream within the CAT ORF, resulting in a larger ORF-1 product (Fig. 1A). These were called STOP mutants. In previously characterized mRNAs where coupled translation termination–reinitiation occurs the expression level from ORF-2 is severely reduced when the ORF-1 stop codon is mutated as the ribosome must translocate upstream following termination of translation of ORF-1 and this is a distance-dependent phenomenon (Ahmadian et al. 2000; Luttermann and Meyers 2007). The effect of the STOP mutation also demonstrates that neither ribosomal scanning nor an IRES are responsible for the translation of ORF-2 as neither process would be affected by a mutation downstream from the translation initiation codon of ORF-2. Similarly, the process cannot be due to translation of degraded mRNA as this would not be affected by the STOP mutations. The average level of CAT protein expression of the appropriate wild-type control wells was set at 100% and the average level of expression for the relevant STOP mutants was compared with this. Analysis of the data from the STOP mutants identified five mRNAs in which the mutant showed a significant reduction in expression from ORF-2 to 10% to 41% of the original high levels demonstrating that coupled translation occurs in these cases. This is consistent with previous observations with the RSV M2 mRNA in which the level of reduction depends on the distance between the ORF-1 stop codon and the 5'

CASQ2

ORF 1---I N T E **D D D E D D D D D N S D E E D N D D S D D D D D E**
 UGGAAAGAUAAACACUGAAGUAAGAUGAUGAUGAUGAUAUUCUGAUGAAGAGGAUAAUGAUGACAGUGAUGACGAUGAUGAUGAA**UAG...**
M M M K M M M M M I I L M K R I M M T V M T M M M N---ORF 2

C22 ORF32

ORF 1---L Y V G T L I S K N F A A L L E E H **D I F V P E D D D D D D**
 CCCCUIUCUUAUGUCGGGACACUCAUAGCAAGAACUUUGCUCUCUACUUGAGGAACAUGACAUUUUUGUCCAGAGGAUGAUGAUGAUGAUGAC**UAA...**
M S G H S L A R T L L L Y L R N M T F L F Q R M M M M M T---ORF 2

CCDC36

ORF 1---L G C S E T P L C K E A G K N L L Y **D L G F D S S D D D G F**
 CCUCAUUCUGGAUGUUCAGAGACCCUCUAUGCAAGGAGGAGGAAAGAAUUUGCUCUUAUGACCUGGUUUUGAUGACAGUGAUGAUGAUGGCUUC**UGA...**
M T W V L I A V M M M A S---ORF 2

WDR45

ORF 1---T F H K Y V F T P **D G N C N R E A F D V Y L D I C D D D D F**
 AGAUGGGACCUUCCAAAUJAUUGUCUUCACUCCUGAUGGAAACUGCAACAGAGAGGCUUUCGACGUGUACCUUGACAUCUGUGAUGAUGAUGACUU**UAA...**
M G P S T N M S S L L M E T A T E R L S T C T L T S V M M M T F---ORF 2

CASQ1

ORF 1---V N V T **D A D S V W M E M D D E E D L P S A E E L E D W L E D**
 AGGAGUCUGCAUUGUUCGGAUGCGGAUAGCGUAUGGAAUUGGACGAUGAGGAGGACCUUGCUGUGAGGACUGGAGGACUGGCGGAGGAGG
M L L M R I A Y G W K W T M R R R T C L L L R S W R T G W R M
 V L E G E I N T E **D D D D D D D D**
 GUCCUGGAGGGCGAGAUCAACACAGAGGACGAUGACGAUGAUGAUGAUGAC**UAG...**
S W R A R S T Q R T M T M M M M T---ORF 2

FIGURE 2. The sequences and coding capacity of the regions of overlap between ORF-1 and ORF-2 of the five genes shown to access ORF-2 using a coupled translation termination–reinitiation process. The multiple aspartate residues encoded by ORF-1 are shown in bold and italics and the potential AUG initiation codons for ORF-2 are shown in bold and underlined.

monoclonal antibodies. Figure 3C shows that the eGFP-CASQ2 fusion protein of 73.5 kDa from eGFP-FLCASQ2 wt was detected (ORF-1GFP). Mutation of the stop codon increased the size of eGFP by 1.7 kDa as expected. The CAT protein expression from ORF-2 in the wild-type construct was dramatically reduced in the eGFP-FLCASQ2/STOP mutant, indicating that ORF-2 was accessed by coupled translation termination–reinitiation (Fig. 3C, ORF-2 CAT). To determine whether the sequences within the CASQ2 gene are required for the coupling process a further construct was generated in which the region of the CASQ2 ORF-1 up to the position of the first ORF-2 AUG codon was entirely replaced with the gene encoding eGFP and the ORF-2 sequence was replaced with the CAT reporter gene. This left only the 81-nt CASQ2 ORF-1/ORF-2 overlap region flanked by the two reporter genes (eGFP-81CASQ2 wt) (Fig. 3A). An eGFP-81CASQ2/STOP mutant was also generated. The eGFP and CAT proteins were detected by Western blot and the eGFP expressed from the STOP mutant was increased in size demonstrating that the mutation had extended ORF-1 (Fig. 3C). The eGFP-81CASQ2 fusion protein was more readily detected in comparison to the eGFP-FLCASQ2 fusions. This is likely the result of the folded full-length chimeric protein masking the epitope sites. CAT protein production was not affected indicating that

the mRNA was not being degraded (Fig. 3C). Importantly, ORF-2 CAT expression was reduced significantly in the STOP codon mutant (Fig. 3C, ORF-2 CAT). The level of reduction in CAT expression seen when comparing transfection with eGFP-81CASQ2 and eGFP-81CASQ2/STOP was consistent with the data showing a 90% reduction for the construct containing the authentic CASQ2 ORF-1 (Fig. 1B). This confirms that the overlap region of the CASQ2 ORF-1 and ORF-2 alone is sufficient to direct coupled translation termination–reinitiation. The sizes of the various GFP proteins also confirmed that the expression of the CAT protein did not arise as a result of a translational frame-shifting event in any of the constructs. It was possible to utilize the construct eGFP-81 CASQ2 to quantify the coupling efficiency of the CASQ2 overlap as purified eGFP and CAT protein were available. Over three experimental repeats we found that CAT protein levels were 82-fold lower than eGFP (Supplemental Fig. 1) which is consistent with that seen for pneumoviral M2 mRNAs although reinitiation on calicivirus subgenomic mRNAs is an order of magnitude more efficient at 10%–20%.

To further investigate the mechanism of coupled translation used by the overlap region of the CASQ2 gene, specific mutations were introduced into the region of overlap between ORF-1 and ORF-2 in the construct expressing GFP

appears as a discrete product despite variation in the number of potential AUG initiation codons for ORF-2 that might be expected to generate a heterogeneous set of proteins. This suggests that the reinitiation event may preferentially use one initiation codon. To investigate this we carried out further mutational analysis on the CASQ2 GAU/GAC mutant that contains three AUG initiation codons. The initiation codons were each mutated singly or in all possible combinations, as summarized in Figure 3D, and matched STOP mutants in which the ORF-1 stop codon was also mutated were also generated. Western blot analysis showed that termination–reinitiation of ORF-2 occurred readily when either only AUG 2 (in Mut 5) or only AUG 3 (in Mut 4) was present (Fig. 3D). The CAT protein synthesized from AUG2 migrated marginally slower than that from AUG3 as expected from the predicted sizes of the proteins. When only AUG 1 was present (in Mut 6) the level of expression from ORF-2 was the same as seen in the absence of any AUG codons (Mut 7) (Fig. 3D). The intensity of the CAT protein product was consistently greatest with Mut 4, than Mut 5 and this, together with the absence of expression from Mut 6 strongly suggests that there is a preference for the most proximal AUG initiation codon to the ORF-1 translation termination site.

To unambiguously show that it was the aspartate-rich region that promotes coupling two final pairs of constructs were generated. As with other constructs the first and last aspartic acids that define the start and finish of the overlap region and provide start codons for ORF2 along the central AUG were maintained. The construct CASQ2 D/E replaced the aspartates with glutamates, thus producing a negatively charged homopolymer carboxyl terminus (Fig. 4A). An additional construct (CASQ2 D/E plus) maintained nine aspartate residues at the amino-terminal of the overlap region followed by the glutamic acid motif used in CASQ2 D/E. STOP mutants of both constructs were also generated to screen for coupled translation. The mutations had no effect on ORF1 (ORF-1 GFP) expression (Fig. 4B). Neither of the D/E construct pairs were able to express the second ORF protein (Fig. 4B). This suggests that the aspartic acid motif must be located at or near the carboxyl terminus of the ORF-1 protein within the overlap.




FIGURE 4. (A) The sequences of the calsequestrin 2 (CASQ2) 81-nt sequence from the region of overlap between ORF-1 and ORF-2 subjected to mutation. The multiple aspartate residues of ORF-1 are shown in bold and italics and the potential initiation methionine residues of ORF-2 are shown in bold and underlined. The wild-type unaltered sequence is shown (CASQ2 wt) together with the CASQ2 D/E mutant overlap in which the multiple aspartate residues were altered to glutamate residues and the CASQ2 D/E plus mutant in which nine aspartate residues at the amino terminal of the overlap region were maintained followed by the glutamic acid motif used in the CASQ2 D/E construct. (B) Detection of GFP and CAT protein in lysates of cells transfected with expression plasmids by Western blot. GFP-specific and CAT-specific monoclonal antibodies were used to detect protein expression. Cells were transfected with the plasmid construct indicated. Lanes on the *left* of each pair are the constructs identified and lanes on the *right* of each pair are the appropriate STOP mutant. Controls were GFP and CAT protein standards and lysate from cells transfected with empty expression vector, as indicated.

DISCUSSION

The data presented here demonstrate for the first time that cellular genes express proteins from second ORFs in mRNAs using a coupled translation termination–reinitiation process. The process of coupled translation termination–reinitiation requires that a ribosome that completes translation of a 5' proximal ORF translocates to an upstream AUG initiation codon to reinitiate translation of a second ORF. To date this process has been identified only in a limited number of virus-encoded mRNAs that use one of two possible mechanisms to achieve expression of the second ORF: In the RSV M2 mRNA the mechanism requires the presence of a highly structured region ~150 nt upstream of the second ORF and in the caliciviruses the mechanism requires the presence of a short sequence homologous to the

18S rRNA ~70 nt upstream of the second ORF. The coupled translation achieves two results, firstly it increases the coding capacity of the genome, and secondly the proportions of the two proteins produced from the mRNA and synthesized in stoichiometrically regulated amounts determined by the efficiency of the coupling process. Viruses use cellular translation machinery during infection and this raises the possibility that cellular genes may also use coupled translation to direct the synthesis of additional proteins.

The analysis described here investigated 24 mRNA transcripts representing 4368 candidate transcripts containing key features found in the viral mRNAs using coupled translation termination–reinitiation (Ahmadian et al. 2000; Gould and Easton 2005, 2007; Luttermann and Meyers 2007, 2009). Expression from second ORFs present in 22 of the 24 (92%) transcripts demonstrates that many more cellular transcripts access second ORFs to produce novel proteins than has previously been suggested. Expression levels from ORF-2 varied between the transcripts with a range of from 200 pg to 40 ng per 10^6 cells (Supplemental Table 1). Several mechanisms have been described by which ribosomes can access second ORFs such as leaky scanning, ribosomal shunting, or by use of an upstream IRES and we identified a proportion (9/24) of the transcripts analyzed where the data were compatible with one or more of these processes (Fig. 1B). However, a further subset showed an unexpected profile in which there was a marked increase in ORF-2 expression when the ORF-1 stop codon was mutated (Fig. 1B). The reasons for this are not yet clear but it may be due to stalled termination of translation of ORF-1 inhibiting reinitiation. This may be similar to the situation described for the cytomegalovirus UL4 gene in which a delay in cleavage of the final aminoacyl tRNA peptidyl bond in a protein encoded by a short ORF upstream of the main ORF results in stalling of termination and subsequent reduction in translation of the primary product (Degnin et al. 1993; Janzen et al. 2002). In the situation of an overlapping ORF-1 and ORF-2 this may also lead to physical occlusion of the ORF-2 initiation codon to ribosomes that would generate the result seen.

In the pneumovirus M2 mRNA the two proteins produced by the single mRNA through the coupling process are functionally linked with one being a transcriptional activator and the other an inhibitor with the linked expression providing a level of previously unknown control of coordinated expression (Birmingham and Collins 1999; Fearn and Collins 1999). Further investigation of the cellular ORF-2 protein products may provide interesting insights into similar control processes. It is also possible that the second ORFs in these transcripts are involved in the process of de novo gene birth (Carvunis et al. 2012).

Five of the transcripts accessed ORF-2 by a process of coupled translation termination reinitiation (Fig. 1B). None of the genes shown to access ORF-2 by coupled translation had any sequence identity with the various virus mRNAs shown to use coupled translation termination–reinitiation (Kupfermann

et al. 1996; Ahmadian et al. 2000; Gould and Easton 2005, 2007; Luttermann and Meyers 2007, 2009). Most strikingly, all five transcripts contained multiple GAUGAU repeats in the region of overlap between ORF-1 and ORF-2 with the AUGs forming the initiation codons of ORF-2 and the carboxyl terminus of the ORF-1 proteins containing multiple aspartic acid residues (Fig. 2). This, together with the observation that the 81-nt overlap region of the CASQ2 mRNA alone is capable of directing the coupling process indicates that the cellular mRNAs use a novel mechanism to direct the coupling process (Fig. 3). In the CASQ2 mRNA the data in Figure 3D strongly suggest that the initiation of ORF-2 preferentially occurs at a specific, or limited number, of the available AUG initiation codons and while the 5' proximal ORF-2 AUG codon can be used there is a preference for the codon(s) nearest to the ORF-1 stop codon. The process is therefore a length-dependent one, as is seen with the RSV M2 mRNA.

The presence of homopolymer runs of aspartate of the CASQ2 transcript is essential for coupled translation to occur for this mRNA, further confirming that the process used is novel. The data presented in Figure 4 suggest that the aspartate motif must be located at or near the carboxyl terminus of the ORF-1 protein. The aspartate motif is located in this region in all of the mRNAs shown to direct coupled translation (Fig. 2). The presence of such extensive homopolymer runs of amino acids in proteins using the same codon is extremely unusual and is likely to have consequences for the translation of the mRNA. One possible consequence is that translation may be slowed if charged tRNAs cannot be provided rapidly. Also, the interaction between the nascent polypeptide and the ribosomal exit tunnel can directly affect translation, including causing stalling (Kramer et al. 2009). Stalling the ribosomes translating ORF-1 may be a necessary requirement to ensure that the terminating ribosomes are able to move in a 5' direction before reinitiating translation at the start codon for ORF-2. The possibility that the presence of sequences rich in other single amino acids are sufficient to direct coupled translation was excluded, as replacement of the aspartate residues with multiple serine or glutamate residues eliminated the CASQ2 translational coupling (Figs. 3B,C, 4). Similarly, in the *NT5C2* and *TMEM97* genes (Fig. 1B) the overlapping regions between the ORF-1 and ORF-2 were rich in glutamate and lysine, respectively (Fig. 5), and while both expressed CAT protein from ORF-2, neither showed evidence of coupled translation. Taken together these data indicate that it is the presence of multiple aspartic acid residues in the carboxyl terminus of the ORF-1 protein and not the nucleotide sequence of the overlap region that is critical for coupled termination–reinitiation in the CASQ2 gene.

These data indicate that our understanding of the coding capacity of the human genome is not yet complete and that if the high proportion of mRNAs identified in this study utilizing second ORFs to produce protein products is representative of the several thousand genes identified as containing overlapping cellular ORFs, the scale of this is likely to be

NT5C2

ORF 1---C H D **E** D D D **E E E E E E E E**
 UACACACUGCCAUGACGAAGAUGAUGAUGAAGAGGAGGAGGAGGAGGAAGAAUAAGGAGGAAAACCAA...
M T K **M M M** K R R R R R R K N K E E N Q ---ORF 2

TMEM97

ORF 1---P Y Y **K** Y E E **K R K K K**
 GCGGAGCCCUACUACAAGUAAGAAGAGAAAAGAAAAAUAUGAAGGAACAACCCUGGCCAGGG...
M K R K E K K N E G N N H W P R ---ORF 2

FIGURE 5. Sequences in the overlap region between ORFs 1 and 2 of the *NT5C2* and *TMEM97* genes. The glutamate and lysine-rich regions of the terminal regions of the ORF-1 proteins are highlighted in bold and italics. The potential initiation methionine residues of ORF-2 are shown in bold and underlined.

considerably higher than previously suspected. Many of these second ORF proteins will be produced by known processes such as leaky scanning, internal initiation with or without the use of IRES sequences, or ribosomal shunting. However, the data here suggest that coupled translation termination–reinitiation is also a significant translational control mechanism available to eukaryotic cells that will direct the synthesis of two proteins simultaneously in stoichiometrically regulated amounts. This suggests that the protein products are likely to be involved in related functions as seen with the RSV M2 proteins. The data also demonstrate that cells use at least two distinct mechanisms in which the overlap region between two ORFs with or without the need for additional upstream sequences can be utilized.

MATERIALS AND METHODS

An algorithm was generated to search the complete list of alternative transcripts from the Ensembl release 55 of the human genome for transcripts where there were two ORFs in separate reading frames that were at least 150 nt in length that overlapped by between 4 and 120 nt. By definition the second ORF contained at least one AUG codon. If multiple start codons existed within the 120-nt overlap, then the start codon giving the smallest overlap was used to define the start of the second ORF. A BLAST search (release version 2.2.26) was performed of the protein sequences for each pair of ORFs against release 2012_3 of the UniProt/Swiss-Prot reference set of nonredundant protein sequences from multiple organisms. Transcripts where both ORFs matched a single known protein with E-values of $<10^{-5}$, and transcripts where the first ORF did not match a known sequence with less than an E-value of 10^{-5} were rejected on the basis that there is likely to have been a misannotation or a misidentification of the first ORF. Transcripts were assigned a probability score based on the joint probability of a number of features that are believed to be associated with coupled translation based on the features of the RSV M2 mRNA. The less likely the feature, the greater the assumed likelihood of coupled translation. However, the algorithm would also identify candidate transcripts that utilize an alternative coupled translation termination–reinitiation mechanism such as that used by the calciviruses. It was believed that the number of start codons, and the number of such codons followed by an A or C both influence the likelihood of coupled translation so that two of the factors contributing to the score were the probability of seeing at least the ob-

served number of start codons in the 120 nt preceding the end of the second ORF and the probability of there being at least the observed number of trailing A and C residues associated with these start codons. It was assumed that longer second ORFs are an indication that the ORF corresponds to a biologically active protein, so a further factor is the probability of an ORF of that length or longer. Finally, it was assumed that the shorter overlaps are more likely to result in coupled translation so the final factor is the probability of an overlap length that is less than the one observed. These scores were used to order the candidate transcripts. The transcripts examined include

some that were identified by early versions of the program but not by the current version as a result of updates to the reference sequences and the BLAST program. The current program identifies all of the transcripts where translational coupling has been verified experimentally as described below. The algorithm and software are available from the investigators.

The transcripts used in the study were those for calsequestrin 2 (cardiac muscle) (*CASQ2*: GenBank accession number NM_001232.3), chromosome 22 ORF 32 (*C22ORF32*: NM_033318.4), coiled-coil domain containing 36 (*CCDC36*: NM_001135197.1), WD repeat domain 45 (*WDR45*: NM_007075), calsequestrin 1 (*CASQ1*: NM_001231), amino-terminal enhancer of split (*AES*: NM_198969), t-complex 1 (*TCPI*: NM_030752), SMT3 suppressor of mif two 3 homolog 2 (*SUMO2*: NM_006937), molybdenum cofactor synthesis 2 (*MOCS2*: NM_176806), heterogeneous nuclear ribonucleoprotein C (C1/C2) (*HNRNPC*: NM_031314), chronic lymphocytic leukaemia up-regulated one opposite strand (*CLU1OS*: NM_001025232), basic transcription factor 3 (*BTF32*: NM_001207), v-Ha-ras Harvey rat sarcoma viral oncogene homolog (*HRAS*: NM_176795), staphylococcal nuclease, and tudor domain containing 1 (*SND1*: NM_014390), Transmembrane protein 97 (*TMEM97*: NM_014573.2), chloride channel, nucleotide-sensitive, 1A (*CLNS1A*: NM_001293), RNA polymerase II polypeptide M (*POLR2M*: NR_027390.1), basic transcription factor 3 (*BTF31*: NM_001037637), debranching enzyme homolog 1 (*DBRI*: NM_016216), lymphocyte-specific protein 1 (*LSP1*: NM_002339), family with sequence similarity 127, member A (*FAM127A*: NM_001078171), 5'-nucleotidase, cytosolic II (*NT5C2*: NM_012229), myeloid-associated differentiation marker (*MYADM*: NM_001020818), arylsulfatase D (*ARSD*: NM_001669.2).

For cloning, PCR reagents were obtained from Promega and reactions were carried out using Pfu. Primers were obtained from IDT and are listed in Supplemental Tables 2 and 3. The cDNA clones of the genes of interest were obtained from commercial suppliers (Origene or Source Bioscience IMAGE clones) or synthesized from cDNA generated from HEP2 cell lines in our laboratory using standard protocols. The first ORF from each gene was amplified by PCR and the CAT reporter gene was inserted to replace the coding region of the second ORF in each gene immediately downstream from the overlap between ORF-1 and ORF-2. The entire construct was inserted into pBlueScribe as previously described (Ahmadian et al. 2000; Gould and Easton 2005). This general structure of the constructs is illustrated in Figure 1A including the stop mutant. For each construct an associated STOP mutant was generated by using an alternative reverse primer to that used to clone the wild type designed to mutate

the ORF-1 translation termination codon, altering the sequence such that termination of ORF-1 at that point was eliminated and translation then terminated 36 nt further downstream within the CAT gene ORF. In total, 48 plasmid constructs were prepared and the sequences of all plasmids were verified prior to use.

The eGFP gene was obtained from plasmid pEGFP (Clontech) and was cloned upstream and in-frame with either the full-length CASQ2 ORF or the CASQ2 81-nt overlap. The CAT reporter was cloned downstream in frame with the ORF2 start codons generating plasmids eGFP-FLCASQ2 wt and eGFP81CASQ2, respectively (Fig. 3A). Mutations of the 81-nt CASQ2 overlap region were generated using synthetic primers which when annealed in overlapping PCR created the novel sequence as shown in Figure 3B replacing the wild-type 81-nt CASQ2 overlap sequence. Reciprocal STOP mutants were generated for all constructs.

Transfections of HEp2 cells were carried out in triplicate and repeated a minimum of three times and CAT protein detection by ELISA was carried out as described previously (Gould and Easton 2007). The antibodies used for direct detection of proteins in Western blots were for CAT #ab50151 and for GFP #ab290 (AbCAM). All lanes received the same amount of protein lysate. All Western blots were repeated in three independent experiments and a representative example is shown.

SUPPLEMENTAL MATERIAL

Supplemental material is available for this article.

ACKNOWLEDGMENTS

This work was funded by the Biotechnology and Biological Sciences research Council grant number BB/I022880/1. S.O. was funded by an RCUK Academic Fellowship during part of his work. W.C. was supported by funding from the Warwick University BBSRC/EPSC Doctoral Training Centre. The authors would like to thank the reviewers for their constructive criticism that enhanced the paper.

Received July 23, 2013; accepted December 3, 2013.

REFERENCES

- Adhin MR, van Duin J. 1990. Scanning model for translational reinitiation in eubacteria. *J Mol Biol* **213**: 811–818.
- Ahmadian G, Chambers P, Easton AJ. 1999. Detection and characterization of proteins encoded by the second ORF of the M2 gene of pneumoviruses. *J Gen Virol* **80** (Pt 8): 2011–2016.
- Ahmadian G, Randhawa JS, Easton AJ. 2000. Expression of the ORF-2 protein of the human respiratory syncytial virus M2 gene is initiated by a ribosomal termination-dependent reinitiation mechanism. *EMBO J* **19**: 2681–2689.
- Aitken CE, Lorsch JR. 2012. A mechanistic overview of translation initiation in eukaryotes. *Nat Struct Mol Biol* **19**: 568–576.
- Birmingham A, Collins PL. 1999. The M2-2 protein of human respiratory syncytial virus is a regulatory factor involved in the balance between RNA replication and transcription. *Proc Natl Acad Sci* **96**: 11259–11264.
- Carvunis AR, Rolland T, Wapinski I, Calderwood MA, Yildirim MA, Simonis N, Charloteaux B, Hidalgo CA, Barbette J, Santhanam B, et al. 2012. Proto-genes and *de novo* gene birth. *Nature* **487**: 370–374.
- Collins PL, Wertz GW. 1985. The 1A protein gene of human respiratory syncytial virus: Nucleotide sequence of the mRNA and a related polycistronic transcript. *Virology* **141**: 283–291.
- Degnin CR, Schleiss MR, Cao J, Geballe AP. 1993. Translational inhibition mediated by a short upstream open reading frame in the human cytomegalovirus gpUL4 (gp48) transcript. *J Virol* **67**: 5514–5521.
- Fearn R, Collins PL. 1999. Role of the M2-1 transcription antitermination protein of respiratory syncytial virus in sequential transcription. *J Virol* **73**: 5852–5864.
- Gould PS, Easton AJ. 2005. Coupled translation of the respiratory syncytial virus M2 open reading frames requires upstream sequences. *J Biol Chem* **280**: 21972–21980.
- Gould PS, Easton AJ. 2007. Coupled translation of the second open reading frame of M2 mRNA is sequence dependent and differs significantly within the subfamily *Pneumovirinae*. *J Virol* **81**: 8488–8496.
- Jackson RJ, Hellen CU, Pestova TV. 2010. The mechanism of eukaryotic translation initiation and principles of its regulation. *Nat Rev Mol Cell Biol* **11**: 113–127.
- Janzen DM, Frolova L, Geballe AP. 2002. Inhibition of translation termination mediated by an interaction of eukaryotic release factor 1 with a nascent peptidyl-tRNA. *Mol Cell Biol* **22**: 8562–8570.
- Kozak M. 1989. The scanning model for translation: An update. *J Cell Biol* **108**: 229–241.
- Kozak M. 1997. Recognition of AUG and alternative initiator codons is augmented by G in position +4 but is not generally affected by the nucleotides in positions +5 and +6. *EMBO J* **16**: 2482–2492.
- Kramer G, Boehringer D, Ban N, Bukau B. 2009. The ribosome as a platform for co-translational processing, folding and targeting of newly synthesized proteins. *Nat Struct Mol Biol* **16**: 589–597.
- Kupfermann H, Thiel HJ, Dubovi EJ, Meyers G. 1996. Bovine viral diarrhoea virus: Characterization of a cytopathogenic defective interfering particle with two internal deletions. *J Virol* **70**: 8175–8181.
- Ling R, Easton AJ, Pringle CR. 1992. Sequence analysis of the 22K, SH and G genes of turkey rhinotracheitis virus and their intergenic regions reveals a gene order different from that of other pneumoviruses. *J Gen Virol* **73** (Pt 7): 1709–1715.
- Luttermann C, Meyers G. 2007. A bipartite sequence motif induces translation reinitiation in feline calicivirus RNA. *J Biol Chem* **282**: 7056–7065.
- Luttermann C, Meyers G. 2009. The importance of inter- and intramolecular base pairing for translation reinitiation on a eukaryotic bicistronic mRNA. *Genes Dev* **23**: 331–344.
- Michel AM, Choudhury KR, Firth AE, Ingolia NT, Atkins JF, Baranov PV. 2012. Observation of dually decoded regions of the human genome using ribosome profiling data. *Genome Res* **22**: 2219–2229.
- Peabody DS, Berg P. 1986. Termination-reinitiation occurs in the translation of mammalian cell mRNAs. *Mol Cell Biol* **6**: 2695–2703.
- Powell ML, Naphine S, Jackson RJ, Brierley I, Brown TD. 2008. Characterization of the termination-reinitiation strategy employed in the expression of influenza B virus BM2 protein. *RNA* **14**: 2394–2406.
- Powell ML, Leigh KE, Pöyry TA, Jackson RJ, Brown TD, Brierley I. 2011. Further characterisation of the translational termination-reinitiation signal of the influenza B virus segment 7 RNA. *PLoS One* **6**: e16822.
- Pöyry TA, Kaminski A, Jackson RJ. 2004. What determines whether mammalian ribosomes resume scanning after translation of a short upstream open reading frame? *Genes Dev* **18**: 62–75.
- Rogers GW, Edelman GM, Mauro VP. 2004. Differential utilization of upstream AUGs in the β -secretase mRNA suggests that a shunting mechanism regulates translation. *Proc Natl Acad Sci* **101**: 2794–2799.
- Sloan J, Kinghorn JR, Unkles SE. 1999. The two subunits of human molybdopterin synthase: Evidence for a bicistronic messenger RNA with overlapping reading frames. *Nucleic Acids Res* **27**: 854–858.
- Spriggs KA, Stoneley M, Bushell M, Willis AE. 2008. Re-programming of translation following cell stress allows IRES-mediated translation to predominate. *Biol Cell* **100**: 27–38.
- Stallmeyer B, Drugeon G, Reiss J, Haenni AL, Mendel RR. 1999. Human molybdopterin synthase gene: Identification of a bicistronic transcript with overlapping reading frames. *Am J Hum Genet* **64**: 698–705.
- Tautz D. 2009. Polycistronic peptide coding genes in eukaryotes—how widespread are they? *Brief Funct Genomic Proteomic* **8**: 68–74.
- Wilson BA, Masel J. 2011. Putatively noncoding transcripts show extensive association with ribosomes. *Genome Biol Evol* **3**: 1245–1252.