

New distribution data of the least weasel *Mustela nivalis* in Castilla y León, Spain

Nuevos datos sobre la distribución de la comadreja *Mustela nivalis* en Castilla y León, España

Julio C. Domínguez^{1*}, Francisco Díaz-Ruiz^{1,2}, Javier Viñuela¹, Noelia de Diego³, Sonia Illanas³, Pedro P. Olea³, Ana E. Santamaría¹, Juan J. Oñate³, Jesús Herranz³, Pablo Acebes³, Pablo Ferreras¹, Juan E. Malo³, Xurxo Piñeiro¹, Alfonso Paz^{4,5}, Carlos Cuéllar⁴, Patricia Mateo-Tomás^{6,7}, Isabel Barja⁸, Daniel Jareño¹, Ana Piñeiro⁹ & Jesús T. García¹

1. IREC, Instituto de Investigación en Recursos Cinegéticos (CSIC-UCLM-JCCM), Ronda de Toledo s/n, 13071 Ciudad Real, Spain.
2. Universidad de Málaga, Departamento de Biología Animal, 29071 Málaga, Spain.
3. Terrestrial Ecology Group (TEG), Departamento de Ecología, Facultad de Ciencias, Universidad Autónoma de Madrid, C/Darwin 2, 28049 Madrid, Spain.
4. GREFA. Grupo de Rehabilitación de la Fauna Autóctona y su Hábitat. Monte del Pilar s/n Majadahonda, 28220 Madrid, Spain.
5. Department of Zoology, Stockholm University, Zoologiska Institutionen, 106 91 Stockholm, Sweden.
6. PMT, Centro de Ecología Funcional, Departamento de Ciências da Vida, Universidade de Coimbra, calçada Martim de Freitas, 3000-456 Coimbra, Portugal.
7. Research Unit of Biodiversity (UO/CSIC/PA), Campus de Mieres, Universidad de Oviedo, C/ Gonzalo Gutiérrez Quirós s/n, 33600 Mieres, Spain.
8. IB, Departamento de Biología, Unidad de Zoología, Facultad de Ciencias, Universidad Autónoma de Madrid, C/ Darwin 2, 28049 Madrid, Spain.
9. Escuela de Medicina Veterinaria, Universidad Andres Bello, Avenida República 440, Santiago centro, 7591538 Santiago de Chile, Chile.

* Corresponding author: jcesardv@gmail.com

The least weasel (*Mustela nivalis* Linnaeus, 1766) is the world's smallest carnivore (King & Powell 2007) and probably one of the most difficult to detect (García & Mateos 2009). It is widely distributed throughout the northern hemisphere (McDonald *et al.* 2016), and seems to be a common and relatively abundant species, at least in some areas of the European continent (Larivière & Jennins 2009, Tikhonov *et al.* 2010). However, many basic aspects of its biology, distribution, population dynamics and ecology remain largely unknown in most areas of its range, a lack of information that is especially striking in Spain (Palazón 2017).

Most of the scarce background knowledge available in Spain comes from general studies about carnivores (Millán *et al.* 2001, Barea-Azcón *et al.* 2007) or cases in which weasels are predated by

other species (Gil & Pleguezuelos 2001, Álvarez 2003, Palazón *et al.* 2016). In addition, these studies are made at small spatial scale (Alcover & Jaume 1983, Camps & Llimona 2000, Gil-Sánchez *et al.* 2001, Torre *et al.* 2003, García & Mateos 2009), since there has never been a national (standardized) census of the species. The few available data are compiled in specific chapters of two general compendia about Spanish Mammals (Gisbert & Santos-Reis 2007, Palazón 2017), showing a nearly ubiquitous presence in peninsular Spain, Mallorca and Menorca islands and Ceuta and Melilla, albeit unevenly distributed over the territory (Fig. 1).

In Castilla y León (the largest autonomous region in Spain, ~93,000 km², 18.5 % of the country's total area) it has a widespread distribution and it has been reported in almost any type of




Figure 1. Map of Castilla y León depicting the previous known distribution of least weasel *Mustela nivalis* (light grey), with black and white dots representing UTM cells with and without captures during our study, respectively. Duero river (bold line) and province names are shown for display purposes. Inset shows the Spanish distribution of least weasel *Mustela nivalis* (10x10 km UTM-grids; Gisbert & Santos-Reis 2007).

environment (riverside, forest, agricultural land, *dehesas*, shrubland, pastures and meadows, human settlements, etc; Peris *et al.* 1999, Gisbert & Santos-Reis 2007, Palazón 2017), so most of the region can be regarded as potential habitat for the species. However, the existence of some noticeable gaps in an otherwise continuous distribution across the region is difficult to explain. This probably reflects an inadequate sampling of UTM cells and/or geographical differences in occurrence records reported over the map (Gisbert & Santos-Reis 2007), as shown by its continuous distribution in some provinces (e.g. Salamanca, Fig. 1, Peris *et al.* 1999) and its generalized absence in others (e.g. Segovia and León provinces).

The elusive and secretive habits of the least weasel and its small body size (92-263 g body mass in males and 65-90.5 g in females; Palazón 2017) lead to reduced detectability. Thus, the lack of species-specific methods in detecting (Torre *et al.* 2003, Barea-Azcón *et al.* 2007), capturing (Gorini *et al.* 2013) or quantifying the species populations

(García & Mateos 2009) may contribute to explain the observed distribution gaps, which is a common issue in the occurrence records of several rare and elusive species at large scale (O'connell *et al.* 2006, Croft *et al.* 2017).

The least weasel is listed as Least Concern in the IUCN, both globally (McDonald *et al.* 2016), and in Spain (Blanco 2007), but faces mayor threats along its continental distribution (McDonald *et al.* 2016). Incidental poisoning with rodenticides could also affect the species in agricultural areas, given its rodent-based diet, particularly microtines (Sheffield & King 1994, Gisbert & Santos-Reis 2007). Outbreaks of vole populations have been recurrent in recent years in Castilla y León, and hot debates have taken place around the use of poison to control them due to side effects on other species (Olea *et al.* 2009). Therefore, a better knowledge of the least weasel distribution would be essential for the design of successful conservation and management strategies. Further, a better estimation of the species' real distribution will allow more reliable

predictions under future scenarios of climate/land use change (Araújo *et al.* 2011), now considered two major threats to open agricultural habitats, which are the preferred by least weasels (McDonald *et al.* 2016). Other potentially important threats for weasels come from habitat loss and fragmentation (Swihart *et al.* 2003), road-kill (Cervinka *et al.* 2015, Simpson *et al.* 2016), infectious diseases and pathologies (Simpson *et al.* 2016), or the rapid expansion of new invasive competitors such as the American mink (*Neovison vison*) (Pódra & Gómez 2018).

We present here new data of the least weasel distribution in Castilla y León and surrounding areas (2 UTM-grids in La Rioja and 1 UTM-grid in Aragón). Castilla y León mostly comprises the northern plateau of the Iberian Peninsula, an area dominated by agricultural landscapes (3.7 million ha; 37.81% of the total area of Castilla y León; MAGRAMA, 2012) over the eastern basin of the Duero River. The known distribution of the species in Castilla y León includes 537 10x10 km UTM-grids, corresponding to the 50.5% of the total UTM cells in the region (Gisbert & Santos-Reis 2007). Data on weasel distribution gathered here are the results of trapping data (Sherman traps) collected between 2007 and 2017 by our research group. Weasels were incidentally trapped in surveys conducted in agro-ecosystems to understand different aspects of the population dynamics and dispersal of common vole *Microtus arvalis* (Pallas, 1778) and its predators, funded by different projects. Therefore, presence of the species in each UTM cell was confirmed by different trapping protocols, variable number of trapping sessions per site, variable trapping effort and different spatiotemporal coverage, factors that can influence the number of UTM with recorded presence and absence. However, some presences obtained during 2016 and 2017 (N = 21; 7 UTM cells), were recorded by the use of specific sampling methods (live traps, track traps, hair traps and video-trapping) aimed to develop a methodology to monitor weasel populations in agricultural landscapes (Díaz-Ruiz *et al.* 2017a, 2017b). Individuals were always released at their respective capture site and all handling procedures were performed in accordance with the Spanish and European regulations for animal protection and experimentation and were approved by the Ethics Committee of the University of Castilla-La Mancha (reference number CEEA: PR20170201).

Overall, 168 weasels were captured in 252,385 trap-nights during the period 2007-2017 in all nine provinces of Castilla y León, and one weasel was captured in a nearby locality (Haro, La Rioja, Fig. 1). Our trapping work was carried out in 64 UTM-grids (5.9% of the UTM grid in Castilla y León), and the species was captured in 38 of them (59.4%). Weasels were captured in 16 new UTM-grids where it had not been previously reported (Gisbert & Santos-Reis 2007), now totalling 553 UTM-grids with confirmed presence of the species in the region. Finally, the species was not detected in 19 out of the 26 UTM-grids where it had previously been reported.

Taken together, these results suggest that the species is actually present in a larger number of UTM cells than those reported in the Spanish Mammal Atlas (Palomo *et al.* 2007) and that many empty UTM cells likely correspond with false absences, in line with the low detectability of the species. However, we cannot rule out the possibility of real absences, given that quick colonisations or local extinctions are well known to occur in this species (Meijide *et al.* 1996, Palazón 2017), perhaps associated with prey fluctuations (coupled predator-rodent cycles; Krebs 2013). We captured weasels in several types of habitat within the agricultural landscape, including rural tracks, riversides and different crops (both irrigated and non-irrigated) and edge types, according to a wide range of habitats in well-surveyed provinces where the species has been reported (Peris *et al.* 1999). All of this suggests that the species could be present in most of the UTM cells of central Duero valley, where vole plagues are recurrent (Luque-Larena *et al.* 2013), but also warns about the potential recent impact of management and treatments against these plagues on this specialist predator.

The development and application of a standardized method to sample weasel populations is recommended, since everything suggest an apparent absence of the species in large areas of Castilla y León (more than the 50% of the territory) due to the lack of specific studies on this small carnivore. Therefore, a national census would be desirable for a better understanding of its distribution as well as to ascertain to which degree conservation and management measures should be implemented. It should be noticed that the species looks particularly rare in the southern half of Spain (Fig. 1), where additional sampling should be considered a priority in the current scenario of global warming.

Acknowledgments

We appreciate the human and logistic cooperation of GREFA, and support of city villages of Villalar de los Comuneros, Boada de Campos, San Martín de Valderaduey and Fuentes de Nava. We would also like to thank the dozens of students and friends that helped us during surveys and trapping work along these years. David Camps and an anonymous reviewer provided valuable comments on the manuscript. Julio Domínguez was supported by a predoctoral grant: “Programa Talento Formación” funded by Fondo Social Europeo (FSE) and Castilla La Mancha regional government (JCCM). Francisco Díaz-Ruiz was supported by a Juan de la Cierva postdoctoral contract (Ministerio de Ciencia e Innovación, Funder Id: 10.13039/501100004837, FJCI-2015- 24949) from the Spanish Ministry of Economy, Industry and Competitiveness. Financial support was provided by MINECO (CGL2011-30274, CGL2015-71255-P), Fundación Biodiversidad and Fundación BBVA (TOPIGEPLA).

References

- Alcover J.A. & Jaume D. 1983. Sobre el mostel *Mustela nivalis* Linnaeus 1758 de les Balears (Carnivora, Mustelidae). *Bolletí de la Societat d'Historia Natural de les Balears*, 27: 145-164.
- Álvarez L. 2003. Un nuevo caso de predación de lechuga común (*Tyto alba*) sobre un carnívoro. *Galemys, Spanish Journal of Mammology*, 15 (1): 65-66.
- Araújo M. B., Guilhaumon F., Neto D. R., Pozo I. & Calmaestra R. 2011. *Impactos, Vulnerabilidad y Adaptación al Cambio Climático de la Biodiversidad española. 2 Fauna de vertebrados*. MMARM, Madrid. 644 pp.
- Barea-Azcón J.M., Virgós E., Ballesteros-Duperon E., Moleón M. & Chiroso M. 2007. Surveying carnivores at large spatial scales: a comparison of four broad-applied methods. *Biodiversity and Conservation*, 16: 1213-1230. DOI: [10.1007/s10531-006-9114-x](https://doi.org/10.1007/s10531-006-9114-x)
- Blanco J.C. 2007. Estado de conservación de los mamíferos de España. Pp. 66-70. En: Palomo L.J., Gisbert J., Blanco J.C. (Eds.) *Atlas y libro rojo de los mamíferos terrestres de España*. Dirección General para la Biodiversidad SECEM y SECEMU, Madrid.
- Camps D. & Llimona F. 2000. Primeres dades sobre l'estudi de la mostela (*Mustela nivalis*) al Parc de Collserola. Pp. 111-114. En: Llimona F., Espelta J.M., Guix J.C., Mateos E., Rodríguez-Teijeiro J.D. (Eds.). *I Jornades sobre la Recerca en el sistemes naturals de Collserola: aplicacions a la gestió del Parc*. Consorci Parc de Collserola, Barcelona.
- Červinka J., Riegert J., Grill S. & Šálek M. 2015. Large-scale evaluation of carnivore road mortality: the effect of landscape and local scale characteristics. *Mammal Research*, 60 (3): 233-243. DOI: [10.1007/s13364-015-0226-0](https://doi.org/10.1007/s13364-015-0226-0)
- Croft S, Chauvenet A.L.M. & Smith G.C. 2017. A systematic approach to estimate the distribution and total abundance of British mammals. *PLoS ONE*, 12 (6): e0176339. DOI: [10.1371/journal.pone.0176339](https://doi.org/10.1371/journal.pone.0176339)
- Díaz-Ruiz F., Jareño D., de Diego N., Ferreras P, García J.T., Olea P.P., Viñuela J., Oñate J.J., Herranz J., Acebes P., Santamaría A.E. & Domínguez J.C. 2017a. Comparación de métodos para el seguimiento de las poblaciones de comadreja *Mustela nivalis* en ambientes agrícolas. Resúmenes XIII Congreso SECEM, Guadalajara, pp. 40.
- Díaz-Ruiz F., de Diego N., Santamaría A. E., Domínguez J.C., Galgo A., García J.T., Olea P.P. & Viñuela J. 2017b. Direct evidence of scavenging behaviour in the garden dormouse (*Eliomys quercinus*). *Mammalia*, 20170087. DOI: [10.1515/mammalia-2017-0087](https://doi.org/10.1515/mammalia-2017-0087)
- García P. & Mateos I. 2009. Evaluation of three indirect methods for surveying the distribution of the Least Weasel *Mustela nivalis* in a Mediterranean area. *Small Carnivore Conservation*, 40: 22-26.
- Gil J.M. & Pleguezuelos J.M. 2001. Prey and prey-size selection by the short-toed eagle (*Circus gallicus*) during the breeding season in Granada (south-eastern Spain). *Journal of Zoology*, 255: 131-137. DOI: [10.1017/s0952836901001182](https://doi.org/10.1017/s0952836901001182)
- Gil-Sánchez J.M., Moleón M., Molino F.M. & Valenzuela G. 2001. Distribución de los mamíferos carnívoros en la provincia de Granada. *Galemys, Spanish Journal of Mammology*, 13: 37-46.
- Gisbert J. & Santos-Reis M. 2007. *Mustela nivalis* Linnaeus, 1766. Pp. 283-286. En: Palomo L.J., Gisbert J., Blanco J.C. (Eds.) *Atlas y libro rojo de los mamíferos terrestres de España*. Dirección General para la Biodiversidad SECEM y SECEMU, Madrid.
- Gorini L., Wedul S.J., Arnemo J.M., Linnell J.D.C., Boitiani L. & Nielsen E.B. 2013. Field anesthesia of least weasels (*Mustela nivalis nivalis*) with isoflurane. *Wildlife Biology in Practice*, 9 (1): 7-13.
- King C.M. & Powell R.A. 2007. *The Natural History of Weasels and Stoats: Ecology, Behavior, and Management*. 2nd edition. Oxford University Press, New York.
- Krebs C.J. 2013. Predation as The Explanation for Fluctuations. Pp. 143-161 In: *Population fluctuations in rodents*. University of Chicago Press.
- Larivière S. & Jennings A.P. 2009. Family Mustelidae. Pp. 564- 655 In: D.E. Wilson & R.A. Mittermeier (eds). *Handbook of the Mammals of the World. Vol.1. Carnivores*. Lynx Edicions, Barcelona.
- Luque-Larena J.J., Mougeot F., Viñuela J., Jareño D., Arroyo L., Lambin X & Arroyo, B. 2013. Recent large-scale range expansion and outbreaks of the common vole (*Microtus arvalis*) in NW Spain. *Basic and Applied Ecology*, 14 (5): 432-441. DOI: [10.1016/j.baae.2013.04.006](https://doi.org/10.1016/j.baae.2013.04.006)

- McDonald R.A., Abramov A.V., Stubbe M., Herrero J., Maran T., Tikhonov A., Cavallini P., Kranz A., Giannatos G., Krytufek B. & Reid F. 2016. *Mustela nivalis*. In: IUCN 2016. The IUCN Red List of Threatened Species. Version 2017.3 <www.iucnredlist.org>. Downloaded on 19 March 2018.
- MAGRAMA 2012. *Encuesta sobre Superficies y Rendimientos de Cultivos 2012*. Ministerio de Agricultura Alimentación y Medio Ambiente. Secretaría General Técnica. Centro de Publicaciones. Madrid, 166 pp.
- Meijide M., Meijide F., Clavel F. & García J.M. 1996. Atlas preliminar de los mamíferos de Soria (España). *Doñana, Acta Vertebrata*, 23 (2): 253-282.
- Millán J., Gortázar C., Marco J. & Escudero M. 2001. Carnivores detected by night survey in Aragón. *Galemys, Spanish Journal of Mammology*, 13: 25-36.
- O'Connell Jr A. F., Talancy N. W., Bailey L. L., Sauer J. R., Cook R. & Gilbert A. T. 2006. Estimating site occupancy and detection probability parameters for meso-and large mammals in a coastal ecosystem. *Journal of Wildlife Management*, 70 (6), 1625-1633. DOI: [10.2193/0022-541X\(2006\)70\[1625:ESOADP\]2.0.CO;2](https://doi.org/10.2193/0022-541X(2006)70[1625:ESOADP]2.0.CO;2)
- Olea P.P., Sánchez-Barbudo I.S., Viñuela J., Barja I., Mateo-Tomás P., Piñeiro A., Mateo R. & Purroy F.J. 2009. Lack of scientific evidence and precautionary principle in massive release of rodenticides threatens biodiversity: old lessons need new reflections. *Environmental Conservation*, 36: 1-4. DOI: [10.1017/s0376892909005323](https://doi.org/10.1017/s0376892909005323)
- Palazón S. 2017. Comadreja – *Mustela nivalis*. En: Enciclopedia Virtual de los Vertebrados Españoles. Salvador A. & Barja I. (eds.). Museo Nacional de Ciencias Naturales, Madrid. <http://www.vertebradosibericos.org/>
- Palazón S., Camps D., Carbonell F. & Grajera J. 2016. Predation of the Weasel (*Mustela nivalis*) by diurnal raptors. *Galemys, Spanish Journal of Mammology*, 28: 66-67. DOI: [10.7325/Galemys.2016.N5](https://doi.org/10.7325/Galemys.2016.N5)
- Palomo L.J., Gisbert J. & Blanco J.C. 2007. *Atlas y Libro Rojo de los Mamíferos Terrestres de España*. Dirección General para la Biodiversidad-SECEM-SECEMU, Madrid, 588 pp.
- Peris S.J., Reyes E., Hernández L. 1999. *Atlas de mamíferos silvestres de la provincia de Salamanca*. Excma. Diputación Provincial de Salamanca, Salamanca. 159 pp.
- Pödra M. & Gómez A. 2018. Rapid expansion of the American mink poses a serious threat to the European mink in Spain. *Mammalia*, 20170013. DOI: [10.1515/mammalia-2017-0013](https://doi.org/10.1515/mammalia-2017-0013)
- Sheffield S.R. & King C.M. 1994. *Mustela nivalis*. *Mammalian Species*, 454: 1-10. DOI: [10.2307/3504183](https://doi.org/10.2307/3504183)
- Simpson V. R., Tomlinson A. J., Stevenso K., McLuckie J. A., Benavides J. & Dagleish M. P. 2016. A post-mortem study of respiratory disease in small mustelids in south-west England. *BMC Veterinary Research*, 12, 72. DOI: [10.1186/s1291701606939](https://doi.org/10.1186/s1291701606939)
- Swihart R. K., Gehring T. M., Kolozsvary M. B. & Nupp T. E. 2003. Responses of 'resistant' vertebrates to habitat loss and fragmentation: the importance of niche breadth and range boundaries. *Diversity and Distributions*, 9 (1): 1-18.
- Tikhonov A., Cavallini P., Maran T., Kranz A., Herrero J., Giannatos G., Stubbe M., Conroy J., Krytufek B., Abramov A., Wozencraft C., Reid F. & McDonald R. 2010. *Mustela nivalis*. The IUCN Red List of Threatened Species 2007. Version 2017.3 <www.iucnredlist.org>. Downloaded on 19 March 2018.
- Torre I., Arrizabalaga A. & Flaquer C. 2003. Estudio de la distribución y abundancia de carnívoros en el Parque Natural del Montnegre-Corredor mediante trampeo fotográfico. *Galemys, Spanish Journal of Mammology*, 15: 31-44.

Submitted 9 April 2018.

Accepted 29 June 2018

Associate editor was José María López Martín