



Journal of Peptide Science

## The mapping of linear B-cell epitope regions in desmoglein 1 and 3 proteins: Recognition of immobilized peptides by pemphigus patients' serum autoantibodies

Journal:	<i>Journal of Peptide Science</i>
Manuscript ID:	Draft
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Szabados, Hajnalka; Hungarian Academy of Sciences, Eötvös Loránd University, Research Group of Peptide Chemistry Bősze, Szilvia; Hungarian Academy of Sciences, Eötvös Loránd University, Research Group of Peptide Chemistry Silló, Pálma; Semmelweis University, Department of Dermatology, Venereology and Dermato-Oncology Hudecz, Ferenc; Hungarian Academy of Sciences, Eötvös Loránd University, Research Group of Peptide Chemistry; Eötvös Loránd University, Institute of Chemistry Kárpáti, Sarolta; Semmelweis University, Department of Dermatology, Venereology and Dermato-Oncology Uray, Katalin; Hungarian Academy of Sciences, Eötvös Loránd University, Research Group of Peptide Chemistry
Keywords:	autoantibody epitopes, pin-bound peptides, modified ELISA, desmoglein 1 and 3, pemphigus

SCHOLARONE™  
Manuscripts

1  
2  
3 The mapping of linear B-cell epitope regions in desmoglein 1 and 3 proteins: Recognition of  
4 immobilized peptides by pemphigus patients' serum autoantibodies

5  
6 Hajnalka Szabados; Szilvia Bősze; Pálma Silló; Sarolta Kárpáti; Ferenc Hudecz; Katalin Uray

7  
8 Research Group of Peptide Chemistry, Hungarian Academy of Sciences, Eötvös L.

9 University, Budapest 112, POBox 32, H-1518, HUNGARY

10 Department of Dermatology, Venereology and Dermato-Oncology, Semmelweis University,  
11 Budapest, HUNGARY

12 Institute of Chemistry, Eötvös L. University, Budapest, HUNGARY  
13  
14  
15

16  
17 Abstract

18  
19 Desmosomal transmembrane glycoproteins desmoglein 1 and desmoglein 3 are targets of life-  
20 threatening autoimmune blistering disorders such as *Pemphigus vulgaris* (PV) and *Pemphigus*  
21 *foliaceus* (PF). In these diseases pemphigus autoantibodies are produced against Dsg1 and  
22 Dsg3 proteins. The autoantibodies bind to these transmembrane elements leading to a loss of  
23 desmosomal cell-cell adhesion, and clinically, to the presence of blisters and erosions.  
24 Identification, characterization and detailed analysis of the binding sites of autoantibodies  
25 have an outstanding importance in understanding the immunopathology of the disease and  
26 also in the design of novel diagnostics.  
27  
28

29  
30 Here we describe the localization of the B-cell epitope regions of Dsg1 and Dsg3 proteins  
31 extracellular parts recognized by IgG-type serum autoantibodies of patients with PV and PF.  
32 In our study overlapping pentadecapeptides were synthesized on hydroxypropylmethacrylate  
33 pins based on the results of *in silico* predictions. To detect the interaction between the serum  
34 autoantibodies and the immobilized synthetic peptides, modified ELISA (Enzyme Linked  
35 Immunosorbent Assay) was performed with pin-attached peptides testing the serum samples  
36 of ten patients and four healthy donors. We identified five possible epitope regions (aa86-110,  
37 aa196-220, aa226-250, aa326-340, and aa486-520) within the Dsg1 protein sequence and four  
38 possible epitope regions (aa64-78, aa330-344, aa375-399, aa446-460) within the Dsg3 protein  
39 sequence using these methods. Our data showed that serum autoantibodies of patients,  
40 previously identified as Dsg1 and Dsg3 positive, are able to recognize continuous linear  
41 epitope regions of both Dsg1 and Dsg3 proteins using pin-bound overlapping peptides in  
42 modified ELISAs.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## 1. Introduction

Desmoglein (Dsg), a desmosomal  $\text{Ca}^{2+}$ -dependent transmembrane glycoprotein, belongs to the cadherin superfamily of cell-cell adhesion molecules with four isotypes: Dsg1, Dsg2, Dsg3 and Dsg4 [1, 2]. These adhesion units specialize in conferring epidermal keratinocyte cohesion and are linked to intercellular molecules of the desmosomal plaque, which in turn interact with components of the cytoskeleton [3]. These glycoproteins are involved in the development of autoimmune diseases. Dsg1 (1049 aa) and Dsg3 (999 aa) are targets of *Pemphigus vulgaris* (PV), while Dsg1 is target of *Pemphigus foliaceus* (PF) also. These, and especially PV, are life-threatening, autoimmune blistering disorders that affect the skin and mucous membranes [4, 5].

There is no common theory for the pathomechanism of these diseases yet, however, it is known that pemphigus autoantibodies are expressed against Dsg1 and Dsg3 proteins and bind to them leading to a loss of desmosomal cell-cell adhesion, and clinically, to the presence of blisters and erosions [6-10].

Although there are several diagnostic tools for autoimmune blistering diseases using monkey esophagus and recombinant proteins, identification, characterization and detailed analysis of the binding sites of autoantibodies has an outstanding importance in understanding the immunopathology and development as well as in the design of novel diagnostics [11, 12].

Antibodies recognize parts of the protein antigen (antigenic determinants or epitopes) which are created by the primary sequence of residues in the protein (called linear epitopes/determinants) and/or by the secondary, tertiary or quaternary structure of the molecule (called conformational epitopes/determinants) [13, 14]. Antigenic determinants are usually limited to those portions of the antigen that are accessible to antibodies. Generally these are small, 4-8 amino acid long residues frequently localized in or near to  $\beta$ -turns or loop regions of a protein [15].

Antigenic determinants can be localized in theoretical ways using *in silico* secondary structure prediction or 3D homology modeling of proteins [16], and experimental ways using domain-swapped and point mutated recombinant proteins [17], synthetic protein fragments [18], X-ray crystallography and NMR spectroscopy [19]. The use of small synthetic peptides of target proteins can be an alternative way to identify epitope regions because of the sufficient similarity to the native antigen which allows the binding of antibodies. This is a rapid, practical, and cost-effective method for linear epitope region identification [20, 21]. B-cell epitope mapping using a series of pin-attached overlapping synthetic peptides can be a very efficient way to identify linear antigenic determinants recognized by serum antibodies or a protein specific antibody in an immunoassay [22]. This strategy has successfully been used in several studies [23-27] to identify antigenic epitope regions of viral, bacterial and other target proteins.

Attempts to localize potentially antigenic regions within the desmoglein proteins have been made previously with inconsistent results. Lucchese et al. have found a 12 amino acid long peptide (aa49-60) on the extreme *N*-terminal region of the Dsg3 protein which was recognized

1  
2  
3 by the sera of PV and PF patients, using a small number of synthetic peptides (Dsg1 aa49-60,  
4 Dsg3 aa36-44, Dsg3 aa49-60, Dsg3 aa190-204, Dsg3 aa373-380, Dsg3 aa518-525) based on  
5 the search of the least redundant peptide sequences [16]. In the domain swapping experiments  
6 of Kawasaki et al. 87-194 residue long protein fragments were swapped between Dsg1 and  
7 Dsg3 proteins. These studies showed the importance of the 1-88 [28] or 1-162 [17] region of  
8 desmogleins. The same group performed point-mutations as well on human Dsg3 protein, and  
9 found two short sequences (aa74-78 and aa102-105, recognized by Kawasaki's mouse  
10 monoclonal antibody) of which the amino acid residues were responsible for antibody  
11 recognition [17]. Hacker-Foegen et al. identified a conformational epitope in the first 87 N-  
12 terminal residues of the Dsg1 protein using domain-swapped proteins and smaller  
13 recombinant protein fragments [29]. Bhol et al. identified two antigenic protein fragments  
14 BOS1 (aa50-79) and BOS6 (aa200-229) from Dgs3 protein sequence [18] with the usage of  
15 12 synthetic 30mer peptides nearly covering the Dsg3 aa50-468 region. Dworschak et al.  
16 analyzed 254 14mer peptides from the aa50-566 region of the Dsg3 protein and selected 5  
17 larger peptides about 17-33 amino acid length and 2 original 14mer for further synthesis and  
18 studies. These peptides were the following: aa50-78, aa243-255, aa273-293, aa304-330,  
19 aa333-365, aa371-383, aa431-452. The N-termini of the peptides were biotinylated and  
20 coupled to streptavidine containing ELISA plates. They identified the aa333-365 region as  
21 major antigenic site and did not find any difference between normal human sera and PV  
22 patient sera using the other peptides. [30].  
23  
24  
25  
26  
27  
28

29 The aim of the present study is to systematically localize linear B-cell epitope regions of the  
30 extracellular parts of Dsg1 (aa50-550) and Dsg3 (aa50-625) proteins recognized by IgG-type  
31 serum autoantibodies of patients with *Pemphigus vulgaris* and *Pemphigus foliaceus* using  
32 theoretical and experimental methods as well. Therefore we have performed Chou-Fasman  
33 secondary structure prediction studies [31] and Eisenberg hydrophobicity predictions [32] on  
34 the extracellular part of Dsg1 and Dsg3 proteins. These methods have successfully been used  
35 in several studies to identify hydrophilic  $\beta$ -turn segments of proteins [23, 24, 33]. For binding  
36 studies, based on the results of predictions, overlapping pentadecapeptides were synthesized  
37 on hydroxypropylmethacrylate pins based on the results of predictions. Finally, a modified  
38 ELISA procedure has been optimized and ELISA experiments with eleven sera of ten patients  
39 and with four healthy donors' sera were performed on the pin-bound peptides.  
40  
41  
42  
43

44 We have defined five possible epitope regions (aa86-110, aa196-220, aa226-250, aa326-340,  
45 and aa486-520) within the Dsg1 protein sequence and four possible epitope regions (aa64-78,  
46 aa330-344, aa375-399, aa446-460) within the Dsg3 protein sequence using these methods.  
47 The defined epitope regions were subjected to homology search.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## 2. Materials and methods

### 2.1. Chemicals and Reagents

#### Multipin peptide synthesis using Fmoc/*t*Bu strategy

Multipin systems were obtained from Mimotopes (Clayton, Victoria, Australia). Fmoc protected amino acid residues (Fmoc-AA) were purchased from Iris Biotech (Marktredwitz, Germany). 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), *N,N'*-diisopropylcarbodiimide (DIC), thioanisole, 1,2-ethanedithiol (EDT), phenol, piperidine, 1-hydroxybenzotriazole (HOBt), *N,N*-diisopropylethylamine (DIEA) and trifluoroacetic acid (TFA) were purchased from Sigma-Aldrich (Budapest, Hungary). *N,N*-dimethylformamide (DMF), methanol and acetic anhydride (Ac<sub>2</sub>O) were obtained from Molar (Budapest, Hungary). *N*-methyl-2-pyrrolidone (NMP) was purchased from Merck (Budapest, Hungary).

#### ELISA

Phosphate buffered saline (0.1M PBS, pH=7.4) was prepared by dissolving 150 mmole NaCl, 8 mmole Na<sub>2</sub>HPO<sub>4</sub>×2H<sub>2</sub>O, 2.7 mmole KCl and 15 mmole KH<sub>2</sub>PO<sub>4</sub> (products of Sigma-Aldrich, Budapest, Hungary) in 1000 mL of deionized water. Gelatin, Tween-20, *o*-phenylenediamine/H<sub>2</sub>O<sub>2</sub> (SigmaFast), 2-mercaptoethanol and sodium dodecyl sulfate (SDS) were also purchased from Sigma-Aldrich (Budapest, Hungary). Rabbit anti-human  $\gamma$ -immunoglobulin conjugated to horseradish peroxidase (IgG/HRP) secondary antibody was obtained from Dako (Glostrup, Denmark). 96-well microtest plates were the products of Sarstedt (Nümbrecht, Germany).

### 2.2. B-cell epitope prediction

The human Dsg1 and Dsg3 sequences used for prediction were obtained from the protein sequence database [34, 35] (<http://www.ncbi.nlm.nih.gov/protein> [last accessed June 2012]).

As antibody epitopes are located mainly on the hydrophilic surface of proteins, and many of them at or near  $\beta$ -turn structures, we have predicted  $\beta$ -turns within the sequences of proteins Dsg1 and Dsg3 using the Chou-Fasman secondary structure prediction method [31] and performed Eisenberg hydrophobicity prediction [32]. In both cases, windows of seven amino acids were used in MS Excel applying their original matrices. Segments with high probability of  $\beta$ -turn secondary structure ( $P_{\beta\text{-turn}} > 1$ ) and low probability of hydrophobicity ( $P_{\text{hydrophobicity}} < 0$ ) were selected. We have supported our results using the PredictProtein website [36] (<http://www.predictprotein.org/> [last accessed June 2012]).

Geno3D (<http://geno3d-pbil.ibcp.fr> [last accessed June 2012]), an automatic web server for protein molecular modelling was used to obtain 3D model of human Dsg1 and Dsg3 proteins, based on a cell adhesion (pdb3q2wA-0) and a structural protein (pdb1q55A-0) template [37].

### 2.3. Multipin peptide synthesis on a Mimotopes block

Forty-five Dsg1 and forty-seven Dsg3 acetyl-pentadecapeptides overlapping in five amino acid residues were synthesized in duplicates on Mimotopes's hydroxypropylmethacrylate non-cleavable pins (Figure 1, the gear contains the functional sites, nominal loading: 66 nmol) with Fmoc/tBu chemistry according to Geysen's method [22]. We used tBu (Thr, Ser, Tyr), OtBu (Asp, Glu), Trt (His, Gln, Asn), Pbf (Arg), AcM (Cys) and Boc (Lys, Trp) as side chain protecting groups. The Fmoc  $\alpha$ -amino protecting group was removed with 2% piperidine, 2% DBU/DMF (v/v) in 20 minutes. Then the pins were washed four times with DMF and twice with methanol. Coupling was performed with 100 equivalent Fmoc-AA/DIC/HOBt 1:1:1 (n/n/n) in NMP for 2x60 min and monitored with bromophenol blue (0.1mM/DMF) added to the coupling mixture [38]. At the end, the *N*-termini of the peptides were acetylated with Ac<sub>2</sub>O/DIEA/DMF 5:1:50 (v/v/v) for 90 min, then after washing the pins the side chain protecting groups (except AcM) were removed with TFA/thioanisole/phenol/water/EDT 82.5:5:5:5:2.5 (v/v/v/v/v) for 150 min, but the unprotected peptides remained covalently attached to the pins.

#### 2.4. Amino Acid Analysis

During and after the completions of the synthesis three-three pins were chosen as test samples. Amino acid analysis was performed on Sykam Amino Acid S433H analyser (Eresing, Germany) equipped with an ion exchange separation column and postcolumn derivatization. Prior to analysis the gear was removed from the polyethylene pin (or rod, Figure 1) and chopped; the gear-bound peptides were hydrolyzed with 6 M HCl in sealed and evacuated tubes at 110°C for 24 h. For post-column derivatization the ninhydrin-method was used. Amino acid analysis data are summarized in Table 1.

#### 2.5. Human serum samples

Patients have been characterized as Dsg1 or Dsg3 positive according to the results of Mesacup Dsg-1 (MB-RG7680EC-D) and Dsg-3 (MB-RG7685EC-D) ELISAs (MBL CO., LTD, Naka-ku, Nagoya, Japan) [4, 39, 40, 41]. Disease types have been identified with direct and indirect immunofluorescence techniques (Department of Dermato-Venereology and Skin Oncology, Semmelweis University). Characteristics of the serum samples are summarized in Table 2. Eleven sera of ten patients of ages 23-89 (6♂, 4♀) were collected. Serum samples of patients with mucosal PV (3 Dsg3 positive cases), sera of patients with PF (5 Dsg1 positive cases), samples of patients with mucocutan PV (3 samples of both Dsg1 and Dsg3 positivity), and four serum samples of healthy individuals were tested. Blood samples were centrifuged on 4000 rpm for 15 minutes than the sera were collected and stored at -20°C before ELISA.

## 2.6. Immunoserological epitope mapping using a modified ELISA with pin-bound peptides

To detect the interaction between the serum autoantibodies and the synthetic peptides, modified ELISA (Enzyme Linked Immunosorbent Assay) was performed with pin-attached peptides. First, the pins were incubated in 0.01M PBS (pH=7.4) containing 0.5% gelatin for 60 min at room temperature for saturation of non-specific binding sites and then washed twice with 0.01M PBS. The serum samples were added to the ELISA plates in a dilution of 1:500 (diluted in 0.01M PBS containing 0.5% gelatin and 0.05% Tween-20). After overnight incubation at room temperature, the pins were washed again for 5×5 min with 0.01M PBS. Finally, the pins were incubated with rabbit anti-human IgG/HRP secondary antibody in a dilution of 1:2000 (diluted in 0.01M PBS containing 0.5% gelatin and 0.05% Tween-20) for 60 min at room temperature. After washing with 0.01M PBS (5×5 min) the color reaction was developed with *o*-phenylenediamine/H<sub>2</sub>O<sub>2</sub> dissolved in deionized water and measured on iEMS MF ELISA reader (Labsystem, Helsinki, Finland) at 492 and 620 nm 10 minutes later. After the ELISA, the pins were regenerated by incubation in 0.1 M phosphate buffer, with 1% (w/v) sodium dodecyl sulphate (SDS) pH 7.2 and 0.1% 2-mercaptoethanol. This was performed at 60 °C in a sonicator bath (ULTRASONIK 3QT, 300ULTRASONIKA, 11,5 kW, Pro-Analitika, Budapest, Hungary), for 5 minutes, twice. The pins were rinsed for 30 s in MilliQ purified water pre-heated to 60 °C, followed by another rinse for at least 30 min in purified water at an initial temperature of 60 °C, on a shaker table. To prepare the pins for subsequent tests, they were immersed in methanol for 15 s and then air dried.

## 2.7. Homology search

We have compared the sequence of epitope regions of human Dsg1 (aa86-110, aa196-220, aa226-250, aa326-340, aa486-520) and Dsg3 (aa64-78, aa330-344, aa375-399, aa446-460) with that of proteins present in the non-redundant protein database by BLASTP 2.2.25+ program and used the following parameters according to the reference: Expect = 10; Matrix: PAM30; Gap cost Existence: 9 Extension: 1; Word size = 2; filter: off; composition based statistics: off [42] (<http://www.ncbi.nlm.nih.gov/blast/Blast.cgi> [last accessed June 2012]).

For further analysis of the determined regions we used the Peptide Match Program (<http://www.pir.georgetown.edu/pirwww> [last accessed June 2012]) to search for identical peptide sequences (100% Blast) in the human proteome.

The Immune Epitope Database and Analysis Resource (<http://www.immuneepitope.org> [last accessed June 2012]) have been also used with 70% minimum Blast to search for homologous motifs in known epitopes of other proteins.

### 3. Results and Discussion

#### 3.1. B-cell epitope prediction

As antibody epitope regions are usually found at or near to hydrophilic  $\beta$ -turn segments of proteins, 5 and 8 regions with high probability of  $\beta$ -turn secondary structure ( $P_{\beta\text{-turn}} > 1$ ) and low probability of hydrophobicity ( $P_{\text{hydrophobicity}} < 0$ ) were selected within the extracellular part of Dsg1 (aa50-550) and Dsg3 (aa50-625) proteins, respectively. The selection is based on the results of Chou-Fasman secondary structure prediction, Eisenberg hydrophobicity prediction and the results of the prediction provided by the PredictProtein website. These regions cover the following sequences: aa56-110, aa126-305, aa306-370, aa381-405, aa406-550 from Dsg1, and aa54-98, aa100-141, aa150-194, aa200-364, aa375-399, aa416-460, aa471-565, aa571-625 from Dsg3 protein (Figure 2.).

#### 3.2. Non-cleavable Multipin peptide synthesis

For immunoserological examination of the predicted regions, multipin systems with two sets of overlapping pentadecapeptides covering regions I-V (Dsg1) and I-VIII (Dsg3) were synthesized with solid phase Fmoc/tBu chemistry. Forty-five Dsg1 and forty-seven Dsg3 pentadecapeptides overlapping in five amino acid residues were prepared in duplicates on hydroxypropylmethacrylate pins. Peptides used for B-cell epitope mapping were acetylated at the *N*-terminus to avoid non-specific ionic interactions. The side chain protecting groups of the amino acids were removed at the end of the synthesis, except AcM from Cys (mimicking disulfide bridges in the protein). During and after the completions of the synthesis three-three pins were chosen as test samples for amino acid analysis. Amino acid analysis of peptides synthesized on pins yielded the ratios summarized in Table 1 (the number in parentheses is the theoretical value) and confirmed that the synthesis of pin-attached peptides was successful. It should be noted that these analyses were carried out on the side-chain protected peptides and this may affect the measured values. The amino acid analysis also confirmed that the amount of peptide on each tested pin was similar, approximately 40 nmol/pin (detailed data not shown), which is a sufficient amount of pin-attached peptide for the majority of antigen – antibody interactions.

#### 3.3. Immunoserological epitope mapping using pin-attached synthetic peptides

The pin-attached overlapping peptides were tested with ELISA optimized by our laboratory. The peptides were synthesized on non-cleavable functionalized gears attached to polyethylene pins. These pins are arranged in a modular unit which complements the 96-well format of a microtiter plate. The pins with the gears are of such configuration that they fit into microtiter wells. This feature greatly simplifies the handling of the pins. All pins with peptides are contacted with serum samples and autoantibodies from the serum were bound to certain peptides. After the pins were taken out of the serum sample and washed, the autoantibodies remained bound to the peptides and were tested with enzyme-labeled, human IgG specific



antibody conjugates for the presence of the peptide bound autoantibody. These IgG type Dsg protein specific autoantibodies were detected from the sera of ten patients (eleven samples; mucosal PV, mucocutan PV, PF) and four healthy donors. The serum antibodies of the healthy individuals did not show any specific recognition in the tests. Therefore, the following recognition criteria were chosen as positive recognition: patients' measured OD  $\geq$  mean OD + 3 $\times$  SD value of healthy donors.

The results of Dsg1 and Dsg3 protein epitope mapping are summarized in Table 3. The forty-five synthetic pin-bound pentadecapeptides covered 83% of the extracellular region of Dsg1 protein. From forty-five pentadecapeptides thirteen were recognized (29%) by the majority of the autoantibodies of both Dsg1 positive and Dsg3 positive sera, twelve were recognized (27%) only by the majority of the autoantibodies of Dsg3 positive sera, while twenty did not have any recognition (44%).

The forty-seven pentadecapeptides studied from the Dsg3 protein sequence also covered 83% of the extracellular region of the protein. Among the forty-seven peptides eight were recognized (17%) by the majority of the serum autoantibodies of the patients, six were recognized (13%) only by the majority of the autoantibodies of Dsg3 positive sera and one were recognized (2%) only by the majority of the autoantibodies of Dsg1 positive sera. The remaining thirty-two peptides did not have any recognition (68%).

According to the immunoserological analysis of the synthetic overlapping peptides we have determined five possible epitope regions (aa86-110, aa196-220, aa226-250, aa326-340, aa486-520) within the Dsg1 protein sequence and four of those (aa64-78, aa330-344, aa375-399, aa446-460) within the Dsg3 protein sequence (Figure 3) with high recognition. Recognition of these epitope regions by autoantibodies from serum samples are summarized in Table 4. To exemplify these data, detailed analysis of the two peptides within the Dsg1 aa196-220 region (Figure 4a,b) and of peptide Dsg3 aa64-78 (Figure 4c) is presented in Figure 4. Our results with the highlighted samples (#11185, #11524, #11873, #11096, #11696, #11608, #11689) show good matches with the clinical characteristics. Serum samples #11096 and #11696 are from the same patient collected at two different time points. In case of the latter sample (#11696) lower values were measured with the Mesacup Dsg-3 (MB-RG7685EC-D) ELISA, corresponding to the lower recognition of individual peptides. Serum autoantibodies of the Dsg1 positive samples #11542, #11190 recognized epitope regions from the Dsg3 protein as well, while autoantibodies from Dsg3 positive samples #10915, #11205 showed recognition with not only Dsg3, but also Dsg1 peptides.

### 3.4. Homology studies of the identified epitope regions

The results of the homology search of the identified epitope regions are presented in Table 5. As we expected, immunologically relevant hits showed homology mostly with proven or predicted Dsg1, Dsg3, Dsg2, Dsg4 proteins, although some hypothetical and bacterial protein hits were found as well.

1  
2  
3 Further analysis of the identified regions we used the Peptide Match Program to search for  
4 identical peptide sequences (100% Blast) in the human proteome. No match was found for  
5 any of the determined regions in the database.  
6

7  
8 The Immune Epitope Database and Analysis Resource have been also used with 70%  
9 minimum Blast to search for homologous motifs in known epitopes of other proteins. From  
10 the possible antigenic regions of Dsg1 protein sequence aa196-220 showed homology with a  
11 human transaldolase 1 epitope (aa231-245). We have to consider this in the further  
12 examination of that sequence. Other identified regions did not show any homology with  
13 epitopes from the human proteome.  
14  
15

#### 16 17 18 4. Conclusion 19

20 Mapping of B-cell epitopes (antigenic determinants, antibody combining sites) of a protein is  
21 the essential part of the design of synthetic diagnostics. These epitopes can be mapped using  
22 numerous methods: they can be predicted from the amino acid sequence of a protein. They  
23 can be assessed using synthetic peptides. Short synthetic peptides are suitable tools for  
24 locating the epitope regions of larger proteins recognized by autoantibodies. In this paper we  
25 presented that the multipin synthesis of overlapping peptides combined with modified ELISA  
26 is a suitable method to find continuous linear epitope regions of immunodominant proteins in  
27 PV. Also, analytical information was given in this study as to the amounts of amino acids  
28 present on the selected individual pins.  
29

30  
31 In our study with overlapping pentadecapeptides tested in modified ELISAs we identified  
32 epitope regions in the extracellular region of the native Dsg1 and Dsg3 proteins. In contrast to  
33 Hacker-Foegen's study [29] we identified five linear epitope regions (aa86-110, aa196-220,  
34 aa226-250, aa326-340, and aa486-520) within the Dsg1 protein sequence and demonstrated  
35 that not only conformational, but linear epitope regions are spread in the entire extracellular  
36 region of the protein as well. From the Dsg3 protein sequence four linear epitope regions  
37 (aa64-78, aa330-344, aa375-399, aa446-460) were determined containing two new linear  
38 epitope regions (aa375-399 and aa446-460). In contrast to Dworschak's study [30] our results  
39 demonstrated that we found a smaller antigenic site (aa330-344) than aa333-365 and in our  
40 experiments aa64-78 and aa375-399 were recognized by the antibodies of the patients as well.  
41 Our experiments showed that serum autoantibodies of patients, previously identified as Dsg1  
42 and Dsg3 positive, are able to recognize sequences of both Dsg1 and Dsg3 proteins. This  
43 result may be in accordance with the fact that more proteins will be affected in the progress of  
44 these autoimmune blistering skin disorders, which can be explained with the high homology  
45 of these proteins. The identified regions showed homology mostly with desmogleins,  
46 although some hypothetical and bacterial protein hits were found as well. One possible  
47 epitope region within the Dsg1 protein sequence (aa196-220) showed homology with another  
48 epitope peptide from the human proteome. Further analysis of the possible epitope regions is  
49 planned in order to locate the exact antibody epitopes of Dsg1 and Dsg3 proteins.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

- [1] Koch PJ, Franke WW. Desmosomal cadherins: another growing multigene family of adhesion molecules. *Curr. Opin. Cell Biol.* 1994. **6**: 682-687
- [2] Whittock NV, Bower C. Genetic Evidence for a Novel Human Desmosomal Cadherin, Desmoglein 4. *J. Invest. Dermatol.* 2003. **120**: 523-530
- [3] Green KJ, Simpson CL. Desmosomes: new perspectives on a classic. *J. Invest. Dermatol.* 2007. **127**: 2499-2515
- [4] Stanley JR. Cell adhesion molecules as targets of autoantibodies in pemphigus and pemphigoid, bullous diseases due to defective epidermal cell adhesion. *Adv. Immunol.* 1993. **53**: 291-325
- [5] Amagai M. Adhesion molecules. I. Keratinocyte-keratinocyte interactions, cadherins and pemphigus. *J. Invest. Dermatol.* 1995. **104**: 146-152
- [6] Schiltz JR, Michel B. Production of epidermal acantholysis in normal human skin in vitro by the IgG fraction from pemphigus serum. *J. Invest. Dermatol.* 1976. **67**: 254-260
- [7] Anhalt GJ, Labib RS, Voorhees JJ, Beals TF, Diaz LA. Induction of pemphigus in neonatal mice by passive transfer of IgG from patients with the disease. *N. Engl. J. Med.* 1982. **306**: 1189-1196
- [8] Hashimoto K, Shafran KM, Webber PS, Lazarus GS, Singer KH. Anti-cell surface pemphigus autoantibody stimulates plasminogen activator activity of human epidermal cells. *J. Exp. Med.* 1983. **157**: 259-272
- [9] Amagai M, Karpati S, Prussick R, Klaus-Kovtun V, Stanley JR. Autoantibodies against the amino-terminal cadherin-like binding domain of pemphigus vulgaris antigen are pathogenic. *J. Clin. Invest.* 1992. **90**: 919-926
- [10] Mahoney MG, Wang Z, Rothenberger KL, Koch PJ, Amagai M, Stanley JR. Explanation for the clinical and microscopic localization of lesions in pemphigus foliaceus and vulgaris. *J. Clin. Invest.* 1999. **103**: 461-468
- [11] Laver WG, Air MG, Webster RG, Smith-Gill SJ. Epitopes on protein antigens: misconceptions and realities. *Cell*, 1990. **61**: 553-556
- [12] van Regenmortel MH. The recognition of proteins and peptides by antibodies. *J. Immunoassay*, 2000. **21**: 85-108
- [13] Atassi MZ. Antigenic structures of proteins. Their determination has revealed important aspects of immune recognition and generated strategies for synthetic mimicking of protein binding sites. *Eur. J. Biochem.* 1984. **145**: 1-20
- [14] Benjamin DC, Berzofsk JA, East IJ, Gurd FRN, Hannum C, Leach SJ, Margoliash E, Michael JG, Miller A, Prager EM, Reichlin M, Sercarz EE, Smith SJ, Todd PE, Wilson AC.

1  
2  
3 Introduction the antigenic structure of proteins: a reappraisal *Ann. Rev. Immunol.* 1984. **2**: 67-  
4 101

5  
6 [15] Hudecz F. Manipulation of Epitope Function by Modification of Peptide Structure: A  
7 Minireview. *Biologicals*, 2001. **29**: 197–207

8  
9 [16] Lucchese A, Mittelman A, Tessitore L, Serpico R, Sinha AA, Kanduc D. Proteomic  
10 definition of a desmoglein linear determinant common to Pemphigus vulgaris and Pemphigus  
11 foliaceus. *J. Trans. Med.* 2006. **4**: 37-45

12  
13 [17] Kawasaki H, Tsunoda K, Hata T, Ishii K, Yamada T, Amagai M. Synergistic pathogenic  
14 effects of combined mouse monoclonal anti-desmoglein 3 IgG antibodies on Pemphigus  
15 vulgaris blister formation. *J. Invest. Dermatol.* 2006. **126**: 2621-2630

16  
17 [18] Bhol K, Natarajan K, Nagarwalla N, Mohimen A, Aokit V, Ahmed AR. Correlation of  
18 peptide specificity and IgG subclass with pathogenic and nonpathogenic autoantibodies in  
19 pemphigus vulgaris: A model for autoimmunity. *PNAS. USA.* 1995. **92**: 5239-5243

20  
21 [19] Pomés A. Relevant B cell epitopes in allergic disease. *Int. Arch. Allergy. Immunol.* 2010.  
22 **152**: 1–11

23  
24 [20] Furka Á, Sebestyén F, Asgedom M, Dibó G. General method for rapid synthesis of  
25 multi-component peptide mixtures. *Int. J Pept. Protein Res.* 1991. **37**: 487-493

26  
27 [21] Houghten RA, Pinilla C, Blondelle SE, Appel JR, Dooley CT, Cuervo JH. Generation  
28 and use of synthetic peptide combinatorial libraries for basic research and drug discovery.  
29 *Nature*, 1991. **354**: 84-86

30  
31 [22] Geysen HM, Meloen RH, Barteling SJ. Use of peptide synthesis to probe viral antigens  
32 for epitopes to a resolution of a single amino acid. *PNAS. USA.* 1984. **81**: 3998-4002

33  
34 [23] Nyárady Z, Czömpöly T, Bősze Sz, Nagy G, Petrohai Á, Pál J, Hudecz F, Berki T,  
35 Németh P. Validation of in silico prediction by in vitro immunoserological results of fine  
36 epitope mapping on citrate synthase specific autoantibodies. *Mol. Immunol.* 2006. **43**: 830-  
37 838

38  
39 [24] Uray K, Hudecz F, Füst G, Prohászka Z. Comparative analysis of linear antibody  
40 epitopes on human and mycobacterial 60-kDa heat shock proteins using samples of healthy  
41 blood donors. *Int. Immunol.* 2003. **10**: 1229-1236

42  
43 [25] Kolesanova EF, Kiselar JG, Jung C, Kozin SA, Hui Bon Hoa G, Archakov AI. Antigenic  
44 mapping of bacterial and animal cytochromes P-450. *Biochimie*, 1996. **78**: 752-762

45  
46 [26] AnandaRao R, Swaminathan S, Khanna N. The identification of immunodominant linear  
47 epitopes of dengue type 2 virus capsid and NS4a proteins using pin-bound peptides. *Virus*  
48 *Res.* 2005. **112**: 60-68

- 1  
2  
3 [27] Nagahira K, Fukuda Y, Terakawa M, Hashino J, Nasu T, Nakazato H, Nakanishi T.  
4 Epitope mapping of monoclonal antibodies to tumor necrosis factor- $\alpha$  by synthetic peptide  
5 approach. *Immunol. Letters*. 1995. **46**: 135-141  
6
- 7 [28] Yeh S-W, Cavacini LA, Bhol KC, Lin M-S, Kumar M, Duval M, Posner MR, Ahmed  
8 AR. Pathogenic human monoclonal antibody against desmoglein 3. *Clinical Immunol*. 2006.  
9 **120**: 68-75  
10
- 11 [29] Hacker-Foegen MK, Janson M, Amagai M, Fairley JA, Lin M-S. Pathogenicity and  
12 epitope characteristics of anti-desmoglein-1 from Pemphigus foliaceus patients expressing  
13 only IgG1 autoantibodies. *J. Invest. Dermatol*. 2003. **121**: 1373-1378  
14
- 15 [30] Dworschak J, Recke A, Freitag M, Ludwig RJ, Langenhan J, Kreuzer OJ, Zillikens D,  
16 Schmidt E. Mapping of B cell epitopes on desmoglein 3 in pemphigus vulgaris patients by the  
17 use of overlapping peptides. *J. Dermatol. Sci*. 2012. **65**: 102-109.  
18
- 19 [31] Chou PY, Fasman GD. Prediction of protein conformation. *Biochem*. 1974. **13**: 222-45  
20
- 21 [32] Eisenberg D, Wesson M, and Wilcox W. In Fasman, G.D., ed., *Prediction of protein*  
22 *structure and the principles of protein conformation*. Plenum Press, New York, **1989**. pp.:  
23 635-646  
24
- 25 [33] Almeida RA, Luther DA, Patel D, Oliver SP. Predicted antigenic regions of  
26 Streptococcus uberis adhesion molecule (SUAM) are involved in adherence to and  
27 internalization into mammary epithelial cells. *Veterinary Microbiology*, 2011. **148**: 323–328  
28
- 29 [34] Wheeler GN, Parker AE, Thomas CL, Ataliotis P, Poynter D, Arnemann J, Rutman AJ,  
30 Pidsley SC, Watt FM, Rees DA, Buxton RS, Magee AI. Desmosomal glycoprotein DGI, a  
31 component of intercellular desmosome junctions, is related to the cadherin family of cell  
32 adhesion molecules. *PNAS*. USA. 1991. **88**: 4796-4800  
33
- 34 [35] Amagai M, Klaus-Kovtun V, Stanley JR. Autoantibodies against a novel epithelial  
35 cadherin in pemphigus vulgaris, a disease of cell adhesion. *Cell*, 1991. **67**: 869-877  
36
- 37 [36] Rost B, Yachdav G, Liu J. The PredictProtein Server. *Nucleic Acids Research*, 2004.  
38 **32**(Web Server issue): W321-W326  
39
- 40 [37] Combet C, Jambon M, Deléage G, Geourjon C. Geno3D: Automatic comparative  
41 molecular modelling of protein. *Bioinformatics*, 2002. **18**: 213-214  
42
- 43 [38] Krchnák V, Vágner J, Safár P, Lebl M. Noninvasive continuous monitoring of solid-  
44 phase peptide synthesis by acid-base indicator. *Collection Czechoslovak Chem. Commun*.  
45 1988. **53**: 2542-2548  
46
- 47 [39] Amagai M. Pemphigus: autoimmunity to epidermal cell adhesion molecules. *Adv.*  
48 *Dermatol*. 1996. **11**: 319-352  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 [40] Amagai M, Hashimoto T, Green KJ, Shimizu N, Nishikawa T. Antigen-specific  
4 immunoadsorption of pathogenic autoantibodies in pemphigus foliaceus. *J. Invest. Dermatol.*  
5 1995. **104**: 895-901  
6

7 [41] Ishii K, Amagai M, Hall RP, Hashimoto T, Takayanagi A, Gamou S, Shimizu N,  
8 Nishikawa T. Characterization of autoantibodies in pemphigus using antigen-specific ELISAs  
9 with baculovirus expressed recombinant desmogleins. *J. Immunol.* 1997. **159**: 2010-2017  
10

11 [42] Altschul SF, Madden LT, Schäffer AA, Zhang Z, Miller W, Lipman DJ. Gapped BLAST  
12 and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acid Res.*  
13 1997. **25**: 3389-3402  
14

15 [43] Geysen HM. Method for determining mimotopes. *Patent US 4833092*. 1989.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 1. Amino acid composition of some peptides synthesized on pins. The number in parentheses is the theoretical value. Gly values are not determined as the spacer moiety of the gear include glycine and  $\beta$ -alanine [43].

Code	Sequence	Amino acid analysis (found [calculated])
Dsg1 391-405	TYVVTGNMGSN DKVG	D 3.07 [3]; T 1.87 [2]; S 1.12 [1]; V 2.76 [3]; Y 0.88 [1]; M 0.75 [1]; K 1.09 [1]
Dsg1 396-405	GNMGSN DKVG	D 3.07 [3]; S 1.12 [1]; V 0.97 [1]; M 0.75 [1]; K 1.09 [1]
Dsg1 396-405	GNMGSN DKVG	D 2.96 [3]; S 1.11 [1]; V 0.95 [1]; M 0.86 [1]; K 1.11 [1]
Dsg1 462-470	GTILSIDDN	D 2.85 [3]; T 0.92 [1]; S 1.21 [1]; I 1.74 [2]; L 1.27 [1]
Dsg1 462-470	GTILSIDDN	D 2.94 [3]; T 0.93 [1]; S 1.05 [1]; I 1.80 [2]; L 1.28 [1]
Dsg3 280-294	ILSSELLRFQVTDL D	D 2.25 [2]; T 0.97 [1]; S 2.05 [2]; E 2.15 [2]; V 1.12 [1]; I 1.05 [1]; L 4.28 [4]; F 0.96 [1]; R 1.07 [1]
Dsg3 280-294	ILSSELLRFQVTDL D	D 2.14 [2]; T 0.95 [1]; S 2.05 [2]; E 2.22 [2]; V 1.12 [1]; I 1.05 [1]; L 3.98 [4]; F 0.96 [1]; R 1.07 [1]
Dsg3 330-344	KVVKALDYEQ LQSVK	D 1.14 [1]; S 1.05 [1]; E 3.22 [3]; A 0.95 [1]; V 2.86 [3]; L 1.81 [2]; Y 0.97 [1]; K 3.17 [3]

Table 2. Characteristics of the serum samples.

Disease type	Positivity	Serum number	Dsg1 titer (U/ml) <sup>1</sup>	Dsg3 titer (U/ml) <sup>1</sup>
PV mucosal	Dsg3	10915	-	258
		11689	-	155
		11205	-	151
PV mucocutan	Dsg1,3	11608	195	89
		11096*	39	63
		11696*	102	27
PF	Dsg1	11185	99,6	-
		11542	328	-
		11190	273	-
		11524	149	-
		11873	102	-

<sup>1</sup>results of Mesacup Dsg-1 (MB-RG7680EC-D) and Dsg-3 (MB-RG7685EC-D) ELISAs

\*serum samples collected from the same patient at two different time points



Table 3. Peptide recognition pattern by Dsg1 and Dsg3 positive serum samples.

Region	Dsg1 peptides	Recognition of Dsg1 positive samples (n=8) <sup>a</sup>	Recognition of Dsg3 positive samples (n=6) <sup>a</sup>	Region	Dsg3 peptides	Recognition of Dsg1 positive samples (n=8) <sup>a</sup>	Recognition of Dsg3 positive samples (n=6) <sup>a</sup>
I.	<sup>56</sup> AACREGEDNSKRNP <sup>70</sup>	-	+	I.	<sup>54</sup> FAKPCREGEDNSKRN <sup>68</sup>	+	+
	<sup>66</sup> KRNPIAKIHSDCAAN <sup>80</sup>	-	-		<sup>64</sup> NSKRNP <sup>78</sup> IAKITS <sup>88</sup> SDYQ <sup>78</sup>	-	++
	<sup>76</sup> DCAANQQV <sup>90</sup> TYRISGV <sup>90</sup>	-	+		<sup>74</sup> TSDYQATQKITYRIS <sup>88</sup>	-	-
	<sup>86</sup> RISGVGIDQPPYGI <sup>100</sup>	+	++		<sup>84</sup> TYRISGVGIDQPPFG <sup>98</sup>	++	-
	<sup>96</sup> PYGIFVINQKTGEIN <sup>110</sup>	++	++	II.	<sup>100</sup> FVVDKNTGDINITAI <sup>114</sup>	-	-
II.	<sup>126</sup> YCRALNSMGQDLERP <sup>140</sup>	+	+		<sup>114</sup> IVDREETPSFLITCR <sup>128</sup>	-	-
	<sup>136</sup> DLERPLELRVRVLDI <sup>150</sup>	-	+		<sup>127</sup> CRALNAQGLDVEKPL <sup>141</sup>	-	+
	<sup>146</sup> RVLDINDNPPVFSMA <sup>160</sup>	-	-	III.	<sup>150</sup> INDNPPVFSQQIFMG <sup>164</sup>	-	-
	<sup>156</sup> VFSMATFAGQIEENS <sup>170</sup>	-	-		<sup>160</sup> QIFMGEIEENSASNS <sup>174</sup>	-	-
	<sup>166</sup> IEENSANTLMILN <sup>180</sup>	-	-		<sup>170</sup> SASNSLVMILNATDA <sup>184</sup>	-	-
	<sup>176</sup> VMILNATDADEPNL <sup>190</sup>	-	-		<sup>180</sup> NATDADEPNHLNSK <sup>194</sup>	-	-
	<sup>186</sup> EPNNLSKIAFKIIR <sup>200</sup>	-	+++	IV.	<sup>200</sup> SQEPAGTPMFLLSRN <sup>214</sup>	-	-
	<sup>196</sup> FKIIRQEPSDSPMFI <sup>210</sup>	+	++		<sup>210</sup> LLSRNTGEVRTL <sup>224</sup> TNS <sup>224</sup>	-	+
	<sup>206</sup> SPMFIINRNTGEIRT <sup>220</sup>	+++	+		<sup>220</sup> TLTNSLDREQASSYR <sup>234</sup>	-	-
	<sup>216</sup> GEIRTMNNFLDREQY <sup>230</sup>	-	+		<sup>230</sup> ASSYRLVVS <sup>244</sup> GADKDG <sup>244</sup>	-	-
	<sup>226</sup> DREQYGYALAVRGS <sup>240</sup>	++	++		<sup>240</sup> ADKDGELSTQCECN <sup>254</sup>	-	-
	<sup>236</sup> AVRGS <sup>250</sup> DRDGGADGMS <sup>250</sup>	++	++		<sup>250</sup> QCECN <sup>264</sup> IKVDVNDNF <sup>264</sup>	-	-
	<sup>246</sup> ADGMSAECECN <sup>260</sup>	-	-		<sup>260</sup> VNDNF <sup>274</sup> PMFRDSQYSA <sup>274</sup>	-	-
	<sup>256</sup> NIKILDVNDNIPYME <sup>270</sup>	-	+		<sup>270</sup> SQYSARIEENILSSE <sup>284</sup>	-	-
	<sup>266</sup> IPYMEQSSY <sup>280</sup> TIIEQ <sup>280</sup>	+	++		<sup>280</sup> ILSSELLRFQVTDLD <sup>294</sup>	-	-
	<sup>276</sup> IEIQENTLNSNLL <sup>290</sup>	-	-		<sup>290</sup> VTDLDEEYTDNLAV <sup>304</sup>	-	-
	<sup>286</sup> NLL <sup>300</sup> LEIRVIDLDEEFS <sup>300</sup>	-	-		<sup>300</sup> NWLAVYFFTS <sup>314</sup> SGNEGN <sup>314</sup>	-	-
	<sup>291</sup> RVIDLDEEFSANWMA <sup>305</sup>	-	-		<sup>310</sup> GNEGNWFEIQ <sup>324</sup> TDPR <sup>324</sup>	-	-
III.	<sup>306</sup> VIFISGNEGNWFEI <sup>320</sup>	-	-		<sup>320</sup> TDPR <sup>334</sup> TNEGILKVVKA <sup>334</sup>	-	-
	<sup>316</sup> NWFEIEMNERTNVGI <sup>330</sup>	-	++		<sup>330</sup> KVVKALDYEQLQSVK <sup>344</sup>	+	++
	<sup>326</sup> TNVGILKVVKPLDYE <sup>340</sup>	++	+++		<sup>340</sup> LQSVKLSIAVKNKAE <sup>354</sup>	-	-
	<sup>336</sup> PLDYEAMQSLQLSIG <sup>350</sup>	-	-		<sup>350</sup> KNKAEFHQS <sup>364</sup> VISRYR <sup>364</sup>	+	-
	<sup>346</sup> QLSIGVRNKA <sup>360</sup> EFHHS <sup>360</sup>	-	+	V.	<sup>375</sup> INVREGIAFRPASKT <sup>389</sup>	+	++
	<sup>356</sup> EFHHSIMSQYK <sup>370</sup> KLKAS <sup>370</sup>	-	++		<sup>385</sup> PASKTFTVQKGISSK <sup>399</sup>	+++	++
IV.	<sup>381</sup> EGPVFRPGSKTYVVT <sup>395</sup>	-	+	VI.	<sup>416</sup> TNKAASN <sup>430</sup> VKYMGRN <sup>430</sup>	-	-
	<sup>391</sup> TYVVTGNMG <sup>405</sup> SNDKVG <sup>405</sup>	+	++		<sup>426</sup> VMGRNDGGYLM <sup>440</sup> MIDSK <sup>440</sup>	+	+
V.	<sup>406</sup> DFVATDLDTGRPST <sup>420</sup>	-	-		<sup>436</sup> MIDSKTAEIKFVKNM <sup>450</sup>	-	-
	<sup>416</sup> RPSTTVRYVMGNNPA <sup>430</sup>	-	-		<sup>446</sup> FVKNMNRDSTFIVNK <sup>460</sup>	-	+
	<sup>426</sup> GNNPADLLAVDSRTG <sup>440</sup>	-	-	VII.	<sup>471</sup> EYTGKTSTGT <sup>485</sup> VYVRV <sup>485</sup>	-	+
	<sup>436</sup> DSRTGKLT <sup>450</sup> LKNKVT <sup>450</sup>	-	-		<sup>481</sup> VYVRVPDFNDNCPTA <sup>495</sup>	-	-
	<sup>446</sup> NKVTKEQYNMLGKY <sup>460</sup>	-	+++		<sup>491</sup> NCPTAVLEKDAVCSS <sup>505</sup>	-	-
	<sup>456</sup> LGGKYQGTILSIDDN <sup>470</sup>	-	-		<sup>501</sup> AVCSSSPSVVVSART <sup>515</sup>	-	+
	<sup>466</sup> SIDDNLQRTCTGTIN <sup>480</sup>	-	-		<sup>511</sup> VSARTLNNRYTGPYT <sup>525</sup>	-	-
	<sup>476</sup> TGTINIQSFGNDD <sup>490</sup>	-	++		<sup>521</sup> TGPYTFALEDQPVKL <sup>535</sup>	+	+
	<sup>486</sup> FGNDDRTNTEPNTKI <sup>500</sup>	++	++		<sup>531</sup> QPVKLPAVWSITTLN <sup>545</sup>	-	-
	<sup>496</sup> PNTKITNTGRQEST <sup>510</sup>	+	++		<sup>541</sup> ITTLNATSALLRAQE <sup>555</sup>	+	+
	<sup>506</sup> RQUESTSSTNYDTST <sup>520</sup>	+	++		<sup>551</sup> LRAQE <sup>565</sup> QIPGVYHIS <sup>565</sup>	-	-
	<sup>516</sup> DTSTTSDSSQYSS <sup>530</sup>	-	-	VIII.	<sup>571</sup> SQNNRC <sup>585</sup> EMPRSLTLE <sup>585</sup>	-	-
	<sup>526</sup> QVYSSEPGNGAKDLL <sup>540</sup>	-	-		<sup>581</sup> SLTLEVCQCDNRGIC <sup>595</sup>	-	-
	<sup>536</sup> AKDLLSDNVHFGPAG <sup>550</sup>	-	-		<sup>591</sup> NRGICGTSYPTTSPG <sup>605</sup>	-	-
					<sup>601</sup> TTPGTRYGRPHSGR <sup>615</sup>	-	-
					<sup>611</sup> PHSGRLGPAAGLLL <sup>625</sup>	-	-

<sup>a</sup> Criteria: patients' measured OD  $\geq$  mean OD + 3 $\times$  sd value of controls. +++: positive recognition by n out of n patients, ++: positive recognition by n-1 out of n patients, +: positive recognition by n-2 out of n patients.

Table 4. Recognition of the identified epitope regions by the serum samples.

Protein	Sequence <sup>b</sup>	Dsg1+ <sup>a</sup>					Dsg1,3+ <sup>a</sup>			Dsg3+ <sup>a</sup>		
		11542	11185	11190	11524	11873	11096	11696	11608	10915	11689	11205
Dsg1	86-110	-	+	+	+	+	+	+	+	+	-	+
	196-220	+	+	+	+	+	+	+	+	+	-	+
	226-250	+	+	+	+	+	+	+	+	+	-	+
	326-340	-	+	+	+	+	+	+	+	+	+	+
	486-520	-	+	+	+	+	+	+	+	+	-	+
		11542	11185	11190	11524	11873	11096	11696	11608	10915	11689	11205
Dsg3	64-78	+	-	+	-	+	+	-	+	+	+	+
	330-344	+	+	+	-	-	+	+	+	-	+	+
	375-399	+	-	+	-	+	+	+	+	-	+	+
	446-460	-	+	+	-	-	+	+	+	+	-	+

<sup>a</sup>according to the results of Mesacup Dsg-1 (MB-RG7680EC-D) and Dsg-3 (MB-RG7685EC-D) ELISAs

+: recognition; -: no recognition (criteria: patients' measured OD  $\geq$  mean OD + 3 $\times$  sd value of controls)

<sup>b</sup>sequence of the determined epitope region

All sera were tested for both Dsg1 and Dsg3 derived pin-attached peptides. Highlighted areas show good matches with the clinical characteristics.

Table 5. Homology search results. All subsequences recognized by serum autoantibodies possess high score hits to the Dsg1 (in case of Dsg1 subsequences) and Dsg3 (in case of Dsg3 subsequences) proteins first, and some other adhesion proteins, hypothetical and bacterial proteins were found with lower score as well.

<b>Dsg1 subsequences</b>	<b>Description</b>
86-110	Dsg1, other desmogleins, cadherin-8, some hypothetical proteins
196-220	Dsg1, other desmogleins, cadherins, some bacterial and virus proteins
226-250	Dsg1, Dsg4, some hypothetical and bacterial proteins
326-340	Dsg1, other desmogleins, desmocollin, cadherins
486-520	Dsg1, some hypothetical and bacterial proteins
<b>Dsg3 subsequences</b>	<b>Description</b>
64-78	Dsg3, other desmogleins
330-344	Dsg3, other desmogleins
375-399	Dsg3, other desmogleins, some hypothetical proteins
446-460	Dsg3, some hypothetical and bacterial proteins

**Figure legends**

Figure 1. The outlook of the Mimotopes non-cleavable pin.

Figure 2. Prediction profiles of Dsg1 (a) and Dsg3 (b) proteins generated by Chou-Fasman secondary structure prediction and Eisenberg hydrophobicity prediction. Dark gray: Chou-Fasman  $\beta$ -turn prediction; light gray: Eisenberg hydrophobicity prediction; black: selected regions for multipin peptide synthesis.

Figure 3. Localisation of experimentally determined epitope regions on 3D homology model of human Dsg1 (a) and Dsg3 (b) proteins. Red ribbons: 8/8 (Dsg1+ patients) positive recognition; blue ribbons: 7/8 (Dsg1+ patients) and 5/6 (Dsg3+ patients) positive recognition.

Figure 4. Binding of serum autoantibodies to pin-bound pentadecapeptides. Detailed representation of modified ELISA results in case of peptides within Dsg1 aa 196-220 (a,b) and peptide Dsg3 aa64-78 (c). D1-D4 indicate the binding of healthy control sera.




Figure 1. The outlook of the Mimotopes non-cleavable pin.  
24x33mm (300 x 300 DPI)




Figure 2. Prediction profiles of Dsg1 (a) and Dsg3 (b) proteins generated by Chou-Fasman secondary structure prediction and Eisenberg hydrophobicity prediction. Dark gray: Chou-Fasman  $\beta$ -turn prediction; light gray: Eisenberg hydrophobicity prediction; black: selected regions for multipin peptide synthesis. 25x20mm (300 x 300 DPI)




Figure 3. Localisation of experimentally determined epitope regions on 3D homology model of human Dsg1 (a) and Dsg3 (b) proteins. Red ribbons: 8/8 (Dsg1+ patients) positive recognition; blue ribbons: 7/8 (Dsg1+ patients) and 5/6 (Dsg3+ patients) positive recognition.  
32x25mm (300 x 300 DPI)




Figure 4. Binding of serum autoantibodies to pin-bound pentadecapeptides. Detailed representation of modified ELISA results in case of peptides within Dsg1 aa 196-220 (a,b) and peptide Dsg3 aa64-78 (c). D1-D4 indicate the binding of healthy control sera.  
78x152mm (300 x 300 DPI)