

PROOF COVER SHEET

Author(s): Frank Odei-Addo, Carminita Frost, Nanette Smith, Tomohisa Ogawa, Koji Muramoto, Maria Luiza Vilela Oliva, László Gráf, and Ryno Naude

Article title: Biochemical characterization of *Acacia schweinfurthii* serine proteinase inhibitor

Article no: GENZ_A_836642

Enclosures: 1) Query sheet
2) Article proofs

Dear Author,

Please check these proofs carefully. It is the responsibility of the corresponding author to check against the original manuscript and approve or amend these proofs. A second proof is not normally provided. Informa Healthcare cannot be held responsible for uncorrected errors, even if introduced during the composition process. The journal reserves the right to charge for excessive author alterations, or for changes requested after the proofing stage has concluded.

The following queries have arisen during the editing of your manuscript and are marked in the margins of the proofs. Unless advised otherwise, submit all corrections using the CATS online correction form. Once you have added all your corrections, please ensure you press the “Submit All Corrections” button.

Please review the table of contributors below and confirm that the first and last names are structured correctly and that the authors are listed in the correct order of contribution.

Contrib. No.	Prefix	Given name(s)	Surname	Suffix
1		Frank	Odei-Addo	
2		Carminita	Frost	
3		Nanette	Smith	
4		Tomohisa	Ogawa	
5		Koji	Muramoto	
6		Maria Luiza Vilela	Oliva	
7		László	Gráf	
8		Ryno	Naude	

AUTHOR QUERIES

- Q1: Please check whether the running head suggested is correct.
- Q2: Please spell out ‘RP-HPLC, SDS-PAGE, TSK ODS, BCA, MW, BLAST, and NCBI’ in full at the first instance.
- Q3: Please provide the town and state abbreviation (for the USA) or town and country of origin (for other countries) identifying the headquarters location for “Sigma Aldrich (USA), Optima Scientific (South Africa).”
- Q4: Please provide the town and state abbreviation (for the USA) or town and country of origin (for other countries) identifying the headquarters location for “Applied Biosystems”.

Q5: Please check the following sentence for clarity: “A similar finding was also observed. . .”.

Q6: Please provide three author names before “et al” in References “1, 5, 6, 9, 10, 13, 14, 16, 17, 19, 20, and 24”.

8
9 ORIGINAL ARTICLE

11 **Biochemical characterization of *Acacia schweinfurthii* serine**
12 **proteinase inhibitor**

15 Frank Odei-Addo¹, Carminita Frost¹, Nanette Smith², Tomohisa Ogawa³, Koji Muramoto³, Maria Luiza Vilela Oliva⁴,
16 László Gráf⁵, and Ryno Naude¹

18
19 ¹Department of Biochemistry & Microbiology, ²Department of Biomedical Sciences, Nelson Mandel Metropolitan University, Port Elizabeth,
20 South Africa, ³Department of Biomolecular Sciences, Graduate School of Life Science, Tohoku University, Sendai, Japan, ⁴Department of
21 Biochemistry, Universidade Federal de São Paulo, São Paulo – SP, Brazil, and ⁵Department of Biochemistry and Molecular Biophysical Research
22 Group of the Hungarian Academy of Sciences, Eötvös Loránd University, Pázmány sétány 1/c, Budapest, Hungary

24
25 **Abstract**

26 One of the many control mechanisms of serine proteinases is their specific inhibition by protein
27 proteinase inhibitors. An extract of *Acacia schweinfurthii* was screened for potential serine
28 proteinase inhibition. It was successfully purified to homogeneity by precipitating with 80%
29 (v/v) acetone and sequential chromatographic steps, including ion-exchange, affinity purification
30 and RP-HPLC. Reducing SDS-PAGE conditions revealed an inhibitor (ASTI) consisting of two
31 polypeptide chains A and B of approximate molecular weights of 16 and 10 kDa, respectively,
32 and under non-reducing conditions, 26 kDa was observed. The inhibitor was shown to inhibit
33 bovine trypsin (K_i of 3.45 nM) at an approximate molar ratio of inhibitor: trypsin (1:1). The A- and
34 B-chains revealed complete sequences of 140 and 40 amino acid residues, respectively.
35 Sequence similarity (70%) was reported between ASTI A-chain and ACT1 A-chain (*Acacia*
36 *leucaena leucocephala* trypsin inhibitor).

37
38
39 **Introduction**

40 Proteinases are the enzymes that catalyse the hydrolytic cleavage
41 of specific peptide bonds in their target proteins. These enzymes
42 are widely distributed in nearly all plants, animals and micro-
43 organisms and are very important for the maintenance and
44 survival of their host organism and play key roles in many
45 biological processes. The proteolytic events catalysed by these
46 enzymes serve as mediators of signal initiation, transmission and
47 termination in many of the cellular events, such as inflammation,
48 apoptosis, blood clotting and hormone processing pathways¹.

49 Despite the fact that these enzymes are indispensable to the
50 cells and organisms that host them, they may be potentially
51 damaging when overexpressed or present in high concentrations.
52 For this reason, the activities of these enzymes need to be strictly
53 regulated and controlled. The synthesis of these enzymes as
54 inactive pre-proteins, and their substrate specificity exert a control
55 on their activities, but it does not fulfil the desired level of
56 regulation, and the fact remains, that cells and organisms require
57 additional means of control².

58 One important control mechanism involves interaction of the
59 active enzymes with proteins that inhibit their activities. These
60 inhibitors form less active or fully inactive complexes with their

91 **Keywords**

92 *Acacia schweinfurthii*, amino acid sequence,
93 proteinase inhibitors, trypsin inhibitor

94
95 **History**

96 Received 27 May 2013
97 Accepted 10 August 2013
98 Published online ■ ■ ■

100
101
102
103
104
105 cognate enzymes. Interest in understanding the physiological
106 significance of proteinase inhibitors (PIs) has increased due
107 to their regulatory action in different processes related to the
108 proteinase-PI balance. The PIs are important tools to achieve a
109 better understanding of fundamental principles of protein inter-
110 action and can be used to design new substances for the control
111 of diseases and pathologic processes. PIs have drawn the attention
112 of many researchers due to their potential medical value, for
113 example, human immunodeficiency virus (HIV) PIs and severe
114 acute respiratory syndrome (SARS) coronavirus PIs may be used
115 to combat HIV and SARS virus, respectively³.

116 Most PIs interact with their target proteinases by contact
117 with the active (catalytic) site of the proteinase, resulting in the
118 formation of a stable PI complex that is incapable of enzymatic
119 activity⁴. With the exception of macroglobulin (720 kDa), which
120 inhibits proteinases of all classes, individual protein inhibitors
121 frequently inhibit proteinases belonging to a single mechanistic
122 class, although some inhibitors inhibit two different classes of
123 enzymes^{5–7}. Of these inhibitors, the most studied classes are
124 the inhibitors of serine (Ser) proteinases. These inhibitors are also
125 very unique in that they retain their inhibitory activity after
126 replacement of their reactive site residue (P1) by another residue⁸.

127 PIs of plants are proteins that are natural, defence-related
128 proteins, often present in seeds and induced in certain tissues by
129 herbivory or wounding. Thus, their main function is thought to be
130 in plant defence, the regulation of endogenous proteinases,
131 as storage proteins, as well as the prevention of unwanted
132 proteolysis^{8,9}. Plant PIs are shown to also possess anti-insect and

63 Address for correspondence: Ryno Naude, Department of Biochemistry &
64 Microbiology, Nelson Mandela Metropolitan University, PO Box 77 000,
65 Port Elizabeth 6031, South Africa. Tel: +27 41 504 2441. Fax: +2741
66 504 2814. E-mail: ryno.naude@nmmu.ac.za

133 anti-fungal activities, and have been reported to inhibit a variety
134 of Ser proteinases, including enzymes of the blood coagulation
135 cascade. They may also be involved in the regulation of
136 programmed cell death in plants¹⁰. Among the seed legumes,
137 the Ser proteinase Bowman–Birk inhibitor and Kunitz trypsin
138 inhibitor (KTI) have been studied extensively. These families
139 differ from each other in size, cysteine content and the number
140 of reactive sites. KTIs are proteins (18 000–24 000 Da), with one
141 or two polypeptide chains and low Cys content, usually with
142 four Cys residues connected by two disulphide bridges and a
143 single reactive site³.

144 Since the isolation of soybean trypsin inhibitor (SBTI) by
145 Kunitz¹¹, it has been found that PIs are widely distributed
146 in seeds of the *Leguminosae*. Thus, *Acacia schweinfurthii* (AS)
147 var *schweinfurthii* belonging to the family, *Leguminosae*, and
148 sub family *Mimosoideae*, indigenous to South Africa, was
149 chosen for this study. Seeds from this plant served as a source
150 of potential inhibitor of Ser proteinases. We report in this
151 paper the purification and characterization of the inhibitor from
152 AS seeds.

154 Materials and methods

155 Materials

156 AS seeds were purchased from Kirstenbosch botanical garden
157 (Cape Town, South Africa). Trypsin, chymotrypsin and trypsin-
158 agarose resin were purchased from Sigma Aldrich (USA), as well
159 as the substrates Z-Gly-Pro-Arg-pNA and N-Suc-Ala-Ala-
160 Pro-Phe-pNA. Toyopearl Super-Q 650 S and TSK ODS 120 T
161 column were from Tosoh (Japan). Molecular weight markers were
162 from Optima Scientific (South Africa). All other reagents were
163 of the highest analytical grade available.

166 Preparation of crude extract

167 Seeds were milled into a fine powder with a grinder. Twenty
168 grams of the milled seeds were extracted in 200 mL 0.15 M NaCl
169 as described by Joubert et al.¹². The proteins were extracted by
170 homogenizing for approximately 10 s. The extract was stirred
171 slowly overnight at 4 °C, followed by centrifugation at 5000 × g
172 for 15 min at 4 °C and the supernatant was kept for further
173 analysis. Proteins were re-extracted from the pellet as discussed
174 above and the supernatants were combined (crude extract).
175 Proteins in the supernatant were precipitated with 80% (v/v)
176 acetone at 4 °C. Protein precipitates were recovered by centrifuga-
177 tion at 5000 × g for 10 min at 4 °C. The pellets were
178 re-suspended in water and lyophilized for further studies.


180 Purification by chromatography

181 Protein from the acetone precipitated material (20 mg) was loaded
182 at 5 mL/h onto an ion exchange chromatography (IEC), Toyopearl
183 Super Q column (29 × 1.6 cm), equilibrated with 0.05 M Tris-HCl
184 buffer, pH 7.5. After extensive washing at 40 mL/h, bound
185 material was eluted with a linear NaCl gradient (0–0.6 M) (total
186 volume, 200 mL), followed by 1 M NaCl to remove any bound
187 material left on the column. Fractions (3 mL) were collected
188 and the absorbances at 280 nm were monitored. The peaks that
189 contained inhibitory activity were pooled and a Sephadex G-10
190 column (64 × 1.6 cm) was used to desalt the fractions, equili-
191 brated with 0.5 M ammonium acetate buffer, pH 8. The sample
192 was loaded at 10 mL/h and chromatography was performed at
193 40 mL/h. Fractions (3 mL) were collected and their absorbance
194 monitored at 280 nm. A trypsin-agarose resin (1.5 mL column)
195 was equilibrated with 10 mM acetic acid, pH 3.2. Protein was
196 applied at 0.5 mL/min and the column was washed with distilled
197 H₂O, and eluted with 20 mM HCl, with the absorbance at 280 nm

being monitored. Fractions (1 mL) were collected. AS inhibitor
was further purified by RP-HPLC¹³ using a Grace Vydac C-18 RP
column. The column was equilibrated with 0.1% trifluoroacetic
acid (TFA)/water and eluted using an acetonitrile gradient
(0–20%, 5 min; 20–60%, 35 min; 60–100%, 15 min) in 0.1%
TFA at a flow rate of 1.0 mL/min. Fractions were collected and
freeze dried.

207 Protein quantification and SDS-PAGE analysis

208 The protein content of the various fractions was determined
209 by using the BCA protein determination assay according to the
210 method of Smith et al.¹⁴. The purity and molecular weight of
211 ASTI were determined using SDS-PAGE (15% resolving gel)
212



252
253
254
255
256
257
258
259
260
261
262
263
264
 Figure 1. (a) Toyopearl Super-Q chromatography of acetone extract of AS (15 mg). The column (29 × 1.6 cm) was equilibrated with 0.05 M Tris-HCl buffer, pH 8.0 and a linear NaCl gradient (0–0.6 M) in 0.05 M Tris-HCl buffer was used for elution. Flow rate was 0.6 mL/min. ASa: tubes 5–12, ASb: tubes 110–135 and ASc: tubes 143–155. Approximately over 95% inhibition of trypsin was observed for fraction ASb and less than 10% was observed for ASa and ASc. (b) Affinity chromatography of fraction ASb using a trypsin-agarose column (1 mL). The column was equilibrated with 10 mM acetic acid and 20 mM HCl was used for elution. Flow rate was 0.5 mL/min. ASb1: tubes 2–5, ASb2: tubes 8–11. (c) RP-HPLC of fraction ASb2 (500 µg) on a Grace Vydac C-18 RP column. The column was equilibrated with 0.1% (v/v) TFA/water. Elution was performed with an acetonitrile gradient at a flow rate of 1 mL/min.

as described by Laemmli¹⁵, under reducing and non-reducing conditions. Peq GOLD Protein-Marker II was used as MW markers (10 000–200 000 Da) and bands were visualized by staining with Coomassie brilliant blue R-250.

Amino acid sequence and phylogenetic analysis

Purified, intact ASTI was subjected to reduction, alkylation and RP-HPLC¹⁶, allowing the two subunits to be separated. S-carboxamidomethylated (CAM)-ASTI (A-chain) (0.5 mg) was digested separately with endoproteinase Arg-C (5 µg, S/E = 100:1) and *Achromobacter* protease I (10 µg, S/E = 50:1) in 0.5 mL 0.1 M Tris-HCl buffer (pH 7.6) containing 10 mM CaCl₂ at 37 °C for 16 h. CAM-ASTI (A-chain) (0.5 mg) was also digested with *Staphylococcus aureus* V8 protease (10 µg, S/E = 50:1) in 0.5 mL 20 mM Tris-HCl buffer (pH 8.0) at 37 °C for 16 h¹⁷. Each digest was separated by RP-HPLC on a TSK gel ODS 120 T column (5 µm, 4.6 × 250 mm) using a gradient of acetonitrile in 0.1% TFA. The amino acid sequences of intact subunits and peptides were determined by a gas-phase protein sequencer (PPSQ-10; Shimadzu, Japan)¹⁷. Sequence similarities between ASTI and other trypsin inhibitors were performed using BLAST tools available on the NCBI website (<http://blast.ncbi.nlm.nih.gov/>). Multiple sequence alignment was obtained using ClustalW2 (<http://www.ebi.ac.uk/Tools/msa/clustalw2/>), and the phylogenetic tree was viewed using TreeView.

Table 1. Purification table of ASTI from 5 g starting material.

Step (fraction)	Total protein (mg)	Total activity (U)*	Specific activity (U/mg)	Purification (fold)	Yield (%)
Acetone (AS)	90	12.42	0.14	1	100
IEC (ASb)	15.9	4.20	0.26	1.91	33.8
Affinity (ASb2)	2.1	1.22	0.58	4.22	9.8
RP-HPLC (ASb2b)	1	1.00	1.00	7.22	8.0

*Inhibitory activity: one unit was defined as the amount of protein needed to inhibit one unit of trypsin activity. One unit of trypsin activity was defined as the enzyme activity that hydrolyses 25 mmol of Z-Gly-Pro-Arg-pNA/min under specified conditions.




Figure 2. Reducing SDS-PAGE patterns of ASTI purified by affinity chromatography and RP-HPLC. Lane 1, MW markers; lane 2, affinity purified fraction (ASb2) and lane 3, RP-HPLC purified fraction (ASb2b). 50–10 refer to kilo Dalton values of molecular weight markers.

Molecular mass measurements of proteolytic products

The molecular masses of proteolytic products resulting from fragmentation of ASTI were measured by matrix-assisted laser desorption/ionization time of flight (MALDI-TOF) mass spectrometry (Voyager-DETM STR, Applied Biosystems)¹⁷.

Trypsin and chymotrypsin inhibition assays

The inhibitory effects of ASTI on bovine trypsin and chymotrypsin were analysed using Z-Gly-Pro-Arg-pNA and N-Suc-Ala-Ala-Pro-Phe-pNA as substrates, respectively. The effect of two substrate concentrations with bovine trypsin, in the presence of varying concentrations of ASTI was studied. The assays were performed according to the method of Smith et al.¹⁸ using 50 mM Tris-HCl, pH 8.2, containing 10 mM CaCl₂, and the absorbance was recorded at 412 nm. The modes of inhibition and the




Figure 3. (a) Separation of peptides generated by digestion of ASTI A-chain with *S. aureus* V8 protease. Peptides were separated by RP-HPLC on a TSK gel ODS 120 T column using a gradient of acetonitrile in 0.1% TFA. Flow rate was 1.0 mL/min. (b) Separation of peptides generated by digestion of ASTI A-chain with endoproteinase Arg-C. Peptides were separated by RP-HPLC on a TSK gel ODS 120 T column using a gradient of acetonitrile in 0.1% TFA. Flow rate was 1.0 mL/min. (c) Separation of peptides generated by digestion of ASTI A-chain with *Achromobacter* protease I. Peptides were separated by RP-HPLC on a TSK gel ODS 120 T column using a gradient of acetonitrile in 0.1% TFA. Flow rate was 1.0 mL/min.

397 inhibition constants were calculated using double Dixon and
 398 enzyme-inhibitor molar ratio plots. For both assays, ASTI was
 399 incubated with the active enzyme for 3 min before the substrate
 400 was added.

401 Results and discussion

402 Purification of AS inhibitor


403 The past few decades have seen a growing interest in the
 404 identification, purification and characterization of novel PIs
 405 from various sources, because of their potent activity in
 406 preventing carcinogenesis in a wide range of *in vivo* and
 407 *in vitro* systems, and their use in developing pest resistance
 408 in otherwise susceptible plants^{5,19}. ASTI was purified to a
 409 final homogeneous product via a few chromatographic steps.
 410 The acetone fraction of the inhibitor was subjected to IEC
 411 (Figure 1a), affinity chromatography (Figure 1b) and RP-HPLC
 412 (Figure 1c), yielding the active inhibitor, fraction ASb2b
 413 (ASTI). The four-step purification procedure of ASTI is
 414 shown in Table 1. A 7.2-fold purification of the inhibitor was
 415 achieved with a specific activity of 1.0 U/mg protein. From
 416 the purification table, it can be concluded that the inhibitor
 417 constituted 8% of the total trypsin inhibitory activity of the
 418 seeds of AS.

402 Molecular mass estimation and homogeneity of inhibitor

403 Electrophoresis was performed under reducing conditions and
 404 ASTI was identified as two molecular weight bands of approxi-
 405 mately 10 and 16 kDa (Figure 2). Under non-reducing conditions,
 406 one band was visible, indicating an intact protein with a molecular
 407 weight of 26 kDa (figure not shown). It can be concluded from
 408 the results that the inhibitor consisted of two subunits, possibly
 409 linked by disulfide bonds. An inhibitor from *Acacia confusa*²⁰
 410 displayed two subunits under reducing SDS-PAGE conditions
 411 (15 and 6 kDa) and a single band (21 kDa) under non-reducing
 412 conditions. Similar results (i.e. a two subunit inhibitor) have been
 413 published for other *Acacia* species. Kortt and Jermyn²¹ purified
 414 an inhibitor from *Acacia elata* seeds and obtained two subunits,
 415 15 kDa for the A-chain and 5 kDa for the B-chain, with a single
 416 component under non-reducing SDS-PAGE conditions (20 kDa).

402 Primary structure

403 The enzymatic digests of CAM-ASTI (A-chain) with *S. aureus* V8
 404 protease, endoproteinase Arg-C and *Acromobacter* protease
 405 I were separated by RP-HPLC, yielding E1-E7, R1-R8 and
 406 K1-K7 (Figure 3a-c, respectively). Amino acid sequencing and
 407 MALDI-TOF mass spectrometry of the peptide fragments derived
 408 from the three enzymatic digestions allowed the determination



461 Figure 4. Amino acid sequence of the A- and B-chain of ASTI, deduced from protease digestions, RP-HPLC and sequence analysis. (E), *S. aureus* V8
 462 protease digest of A-chain; (K), *Achromobacter* protease I digest of A-chain and (R), endoproteinase Arg-C digest of A-chain.

of the complete amino acid sequence of ASTI, shown in Figure 4. The sequences of the A- and B-chain revealed 140 and 40 amino acids, respectively. There are five positions showing two amino acid residues, namely positions 52, 62, 67, 70 and 122 (Figure 4). The two residues at the first four positions are accounted for via the following sequences: E6: TQNGMPAKIWTPPRIGFLS; E7: QQNGMPAKIWSPPRIAFLS; R8: IGFLTPAFYLNIEFQPR; K7: IWSPPRIGFLSPAFYLNIEFQPRDPPSCL; and for position 122: K3: IMPYRDDYK; R2: IKPYR; E5: DRVFGPFKIMPYRD DYKIVYCE. The presence of more than one residue at certain positions in the ASTI A-chain, referred to as a multiplicity of inhibitor molecular forms, may be ascribed to isoforms. Several studies on Ser PIs have confirmed this finding²⁰. The sequences of K5 and R5 showed that endoproteinase Arg-C may also cleave Lys residues. Arg 65 is the amino acid at the P1 position of the reactive site of ASTI (Arg 65-Ile 66), being a characteristic of a trypsin inhibitor²². ClustalW was used to obtain a sequence comparison with other similar inhibitors from plant seeds (Figure 5). Sequence identity of approximately 70% was calculated between the A-chains of ASTI and ACTI²². Comparing the B-chain sequences of ASTI and *Leucaena leucocephala* trypsin inhibitor (LLTI), a 76% similarity was obtained. Taking into consideration the molecular weight of intact ASTI from SDS-PAGE analysis (26 kDa), the low Cys content (four Cys residues) and the molecule being a two chain structure, ASTI can be considered to be a Kunitz-type inhibitor²³. It has similar properties to a complete Kunitz-type trypsin inhibitor alpha chain, which contains only three Cys residues and thus also making it a Kunitz-type inhibitor, as well as 140 amino acid

residues of the A-chain of ASTI, a characteristic of Kunitz inhibitors. ACTI²² displayed a high sequence identity to the N-terminal sequence of ASTI (84% comparing residues 1–50), and it is also a Kunitz-type inhibitor. Kortt and Jermyn²¹ isolated two trypsin inhibitors from *A. elata* seeds, being homogeneous as judged by gel electrophoresis at pH 4.3 and 8.8, and both inhibitors co-eluted from a Sephadex G-100 column and had the same molecular weight of 25 kDa. It can thus be concluded that properties of ASTI are similar to those of inhibitors from comparable species. The phylogenetic comparison of ASTI with other trypsin inhibitors (Figure 6) showed similarities to other closely related inhibitors, such as ACTI and LLTI. These results were also confirmed when related to the multiple sequence alignments (Figure 5).

Inhibition of serine proteinases

For ASTI a K_i of 3.45×10^{-9} M was obtained with bovine trypsin (Figure 7). The K_i value obtained for ASTI can be compared with that of a trypsin inhibitor purified from *Entada scandens* seeds (K_i of 4.9×10^{-9} M)³, indicating a very strong and potent inhibitor. The K_i value of SBTI interacting with bovine trypsin is in the range of 10^{-9} M²⁴, similar to a K_i of 3.5×10^{-9} M obtained (result not shown). The results suggest a high affinity of ASTI, as well as SBTI, for trypsin. The molar ratio of ASTI: enzyme showed a value of 0.8:1 for near complete inhibition of trypsin (result not shown). The K_i value reported for ACTI was 0.294×10^{-9} M²⁰. The latter result indicates that ACTI has a stronger binding affinity than ASTI.



Figure 5. (a) Alignment between the sequence of the A-chain of ASTI with other comparative plant seed inhibitor sequences obtained from ClustalW search. Comparative sequences are obtained with Kunitz-type trypsin inhibitor A chain, ACTI-A (*A. confusa*), gi|299509|; Kunitz-type trypsin inhibitor A-chain, LLTI-A (*L. leucocephala*), gi|18202442|; trypsin inhibitor A-chain BVTI-A (*Bauhinia variegata*), gi|15082208| and KTI A-chain, KTI-A gi|162138868|. (*), Identical sequences; (:), similarities of three or four amino acids. (b) Alignment between the sequence of the B-chain of ASTI with other comparative plant seed inhibitor sequences obtained from ClustalW search. Comparative sequences are obtained with Kunitz-type trypsin inhibitor B-chain, ACTI-B (*A. confusa*), gi|299508|; SBTI and Kunitz-type trypsin inhibitor B-chain, LLTI-B (*L. leucocephala*), gi|18202443|. (*), Identical sequences; (:), similarities of three or four amino acids.




Figure 6. A phylogenetic tree analysis of ASTI with other inhibitors. Protein sequences were obtained from <http://blast.ncbi.nlm.nih.gov/> by a BLAST search of ASTI. Accession numbers for the sequences used are as follows: trypsin iso-inhibitor (*Adenanthera pavonina*), APTI, gi|225058; LLTI, gi|18202442; *Prosopis juliflora* trypsin inhibitor, PJTI, gi|417176; ACTI, gi|299509; *Tamarindus indica* trypsin inhibitor, TITI, gi|308756025; BVTI, gi|32363181; *Bauhinia unguilata* Factor X inhibitor, BUXI, gi|32363179; *Psophocarpus tetragonolobus* trypsin inhibitor, PTI, gi|86450987, chymotrypsin inhibitor (*Erythrina variegata*), gi|265716. The scale bar shows a branch length of 0.1 (i.e. a 10% difference in amino acids).




Figure 7. Double-Dixon plot to determine the inhibition constant (K_i) of ASTI interacting with bovine trypsin. Substrate concentrations of 25 and 5 mM were used. Bovine trypsin concentration was 1 nM. 25 mM ($R^2 = 0.913$) and 5 mM ($R^2 = 0.920$) (\circ).

Chymotrypsin inhibition was determined for ASTI and at an enzyme: inhibitor molar ratio of 1:1 weak/insignificant inhibition was observed (result not shown). A similar finding was also observed with inhibitors purified from seeds from *Mimosoideae* families, where inhibitors of trypsin were found to also weakly inhibit chymotrypsin²¹ and with inhibitors purified from *A. elata* seeds²¹, *A. confusa* seeds²⁰ and *E. scandens* seeds³. However, recently Lam and Ng²⁵ purified a 70 kDa chymotrypsin inhibitor from *A. confusa* seeds that had no trypsin inhibitory activity and was highly potent in inhibiting HIV-1 reverse transcriptase.

Conclusion

The results revealed that we purified a new PI from seeds of the *Leguminosae*, and in terms of the physicochemical and kinetic properties, ASTI can be considered a Kunitz-type trypsin inhibitor. In ongoing experiments the effect of ASTI on the blood coagulation system is studied.

Declaration of interest

This work was supported by the National Research Foundation (South Africa), the Hungarian Science Foundation (Hungary), the

FAPESP (09/53766-9) (Brazil) and the Nelson Mandela Metropolitan University.

References

- Ivanov D, et al. A serpin from the gut bacterium *Bifidobacterium longum* inhibits eukaryotic elastase-like serine proteases. *J Biol Chem* 2006;281:17246–52.
- Habib H, Fazili KM. Plant protease inhibitors: a defense strategy in plants. *Biotech Mole Biol Rev* 2007;2:68–85.
- Lingaraju MH, Gowda LR. A Kunitz trypsin inhibitor of *Entada scandens* seeds: another member with single disulfide bridge. *Biochim Biophys Acta* 2008;1784:850–5.
- Birk Y. Plant protease inhibitors – significance in nutrition, plant protection, cancer prevention and genetic engineering. Berlin: Springer Publishers; 2003.
- Oliva MLV, et al. A novel subclassification for Kunitz proteinase inhibitors from leguminous seeds. *Biochim* 2010;92:1667–73.
- De Oliveira C, et al. Effect of plant neutrophil elastase inhibitor on leucocyte migration, adhesion and cytokine release in inflammatory conditions. *Br J Pharmacol* 2010;161:899–910.
- Oliva ML, Sampaio UM. *Bauhinia* Kunitz-type proteinase inhibitors: structural characteristics and biological properties. *Biol Chem* 2008;389:1007–13.
- Laskowsky MJ, Kato I. Protein inhibitors of proteinases. *Ann Rev Biochem* 1980;49:593–626.
- Konarev AV, et al. The distribution of serine protease inhibitors in seeds of the Astridae. *Phytochem* 2004;65:3003–20.
- Ee KY, et al. Characterisation of trypsin and α -chymotrypsin inhibitors in Australian wattle seed (*Acacia victoriae* Benth). *Food Chem* 2008;107:337–43.
- Kunitz M. Crystalline soybean trypsin inhibitor. *J Gen Physiol* 1947;30:291–310.
- Joubert FJ, Merrifield EH, Dowdle EBD. The reactive sites of proteinase inhibitors from *Erythrina* seeds. *Int J Biochem* 1987;19:601–6.
- Paiva PMG, et al. Purification and primary structure determination of two Bowman-Birk type trypsin iso-inhibitors from *Cratylia mollis* seeds. *Phytochem* 2006;67:545–52.
- Smith PK, et al. Measurement of protein using bicinchoninic acid. *Anal Biochem* 1985;150:76–85.
- Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970;227:680–5.
- Kabuto S, et al. The amino acid sequence of pancreatic alpha-amylase from the ostrich, *Struthio camelus*. *Comp Biochem Physiol Part B* 2000;127:481–90.
- Terada T, et al. Structural characterization of a rhamnose-binding glycoprotein (lectin) from Spanish mackerel (*Scomberomorus niphonius*) eggs. *Biochim Biophys Acta* 2007;1770:617–29.
- Smith N, Naudé RJ, Oelofsen W. Kinetic and thermodynamic characterization of ostrich (*Struthio camelus*) trypsin and chymotrypsins. *Int J Biochem* 1993;25:367–72.
- Schuler TH, et al. Insect-resistant transgenic plants. *Trends Biotechnol* 1998;16:168–75.
- Lin J-Y, et al. Trypsin inhibitor from the seeds of *Acacia confusa*. *J Biochem* 1991;110:879–83.
- Kortt AA, Jermyn MA. *Acacia* proteinase inhibitors – purification and properties of the trypsin inhibitors from *Acacia elutu* seeds. *Eur J Biochem* 1981;115:551–7.
- Wu H-C, Lin J-Y. The complete amino acid sequence of a Kunitz family trypsin inhibitor from seeds of *Acacia confusa*. *J Biochem* 1993;113:258–63.
- Richardson M. The proteinase inhibitors of plants and micro-organisms. *Phytochem* 1977;16:159–69.
- Oliva MLV, et al. *Bauhinia* serine proteinase inhibitors: effect on factor X, factor XII and plasma kallikrein. *Immunopharmacol* 1996;32:85–7.
- Lam SK, Ng TB. A dimeric high-molecular-weight chymotrypsin inhibitor with antitumor and HIV-1 reverse transcriptase inhibitory activities from seeds of *Acacia confusa*. *Phytomed* 2010;17:621–5.