

1 **Cold-induced changes affect survival after exposure to vitrification solution during**  
2 **cryopreservation in the south-west Australian Mediterranean climate species *Lomandra sonderi***  
3 **(Asparagaceae).**

4 Akshay Menon<sup>1,2</sup>, Bryn Funnekotter<sup>1,2</sup>, Anja Kaczmarczyk<sup>1,2</sup>, Eric Bunn<sup>2,3</sup>, Shane Turner<sup>2,3</sup> and Ricardo L.  
5 Mancera<sup>1\*</sup>

6  
7 <sup>1</sup> School of Biomedical Sciences, CHIRI Biosciences, Curtin University, GPO Box U1987, Perth WA  
8 6845, Australia.

9 <sup>2</sup> Botanic Gardens and Parks Authority, Fraser Avenue, West Perth WA 6005, Australia.

10 <sup>3</sup> School of Plant Biology, Faculty of Natural and Agricultural Sciences, University of Western Australia,  
11 Crawley WA 6009, Australia.

12

13 \* **Corresponding author:** R.Mancera@curtin.edu.au

14 Tel: +618 9266 1017

15 Fax: +618 9266 2342

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

**Abstract**

There is limited knowledge of the effects of exposure to low temperatures in the unique Mediterranean climate plant species of Western Australia. We have thus investigated the effect of low temperature on cryogenic tolerance in *Lomandra sonderi*, an endemic perennial species of southwest Western Australia. *Lomandra sonderi* plants were preconditioned in tissue culture at constant 23 °C (12h light/dark cycle) or alternating 20/-1 °C (16h light and 8h dark cycle). Shoot tips from both conditions were analysed for their phospholipid, sterol and soluble sugar compositions. Shoot tips were also cryoexposed *via* a droplet-vitrification protocol. Survival in both preconditioning regimes for cryoexposed and non-cryoexposed samples was the same, but plants from the 20/-1 °C regime displayed an improved tolerance to the overall cryopreservation process in both cryoexposed and non-cryoexposed samples, thereby eliminating exposure to liquid nitrogen as a primary cause of reduced post-cryogenic viability. Preconditioning of *in vitro* shoots of *L. sonderi* at 20/-1 °C induced significant increases in phosphatidylcholine (from  $7.30 \pm 3.46$  to  $22.2 \pm 7.80$  ng mg<sup>-1</sup> FW) and increases in several soluble sugars (fructose, galactose, glucose, sucrose) compared to shoots incubated at 23 °C - changes consistent with known cold acclimation responses in plant species generally - but sterol content remained largely unchanged. Analysis of electrolyte leakage in shoot tips from both preconditioning regimes generated a significantly lower LT<sub>50</sub> value in the 20/-1 °C samples ( $-5.45 \pm 0.53$  °C) over the 23 °C samples ( $-2.5 \pm 0.08$  °C). Increased tolerance to cryoexposure in *L. sonderi* appears to lie mainly with acclimation-induced changes in membrane composition and promotion of membrane stability and hence increased resistance to freeze damage.

**Keywords**

*Lomandra sonderi*, phospholipid, sugar, cold acclimation, electrolyte leakage, cryopreservation

## 69 **Introduction**

70 Continuous maintenance of valuable germplasm material of propagation-recalcitrant or rare and  
71 endangered plants requires considerable resource allocation on the part of conservation and restoration  
72 practitioners. Such material requires reliable long-term storage methods while awaiting utilisation in  
73 conservation and restoration projects. Plants exhibiting low seed quality, or poor seed germination due to  
74 complex seed dormancy are often propagated *via in vitro* cultures of plantlets at considerable expense.  
75 Consequently, the ongoing protection of each genetic line in culture is paramount. The high level of  
76 manual input involved in a continuous micropropagation enterprise is prone to accidental contamination  
77 through operator error and/or equipment failure, while other less common (but potentially catastrophic)  
78 hazards such as fire may result in losses of irreplaceable material (Offord et al. 2009). Cryopreservation is  
79 an efficient and ultimately safer alternative for the long-term storage and biosecurity of valuable culture  
80 lines of plants (coupled with off-site storage). Agricultural, horticultural and endangered plants with low-  
81 seed production, complex dormancy or desiccation-intolerance properties that cannot otherwise be stored  
82 in conventional seed banks can all benefit from cryogenic storage technology (Kaczmarczyk et al. 2011;  
83 Kaczmarczyk et al. 2012; Food and Agriculture Organisation of the United Nations [FAO] 2013).

84 Cryopreservation involves the storage of a range of plant tissues (meristems, excised zygotic  
85 embryos, cell suspensions or callus tissues) at ultra-low temperatures in liquid nitrogen (LN) either at -  
86 135 °C in the vapour phase or -196 °C in the liquid phase (Kaczmarczyk et al. 2012). This process halts  
87 all cellular metabolic processes and preserves viability over extended periods of time (Kaczmarczyk et al.  
88 2012). However, cryopreservation is still limited by a lack of knowledge of how different plant species  
89 respond to the cryoprotective agents (CPAs) or any other pre-cryopreservation treatment (Kaczmarczyk et  
90 al. 2012). Each step in the cryopreservation process can potentially impact on the survival of the  
91 cryopreserved material for better or worse, depending on the cryo-capability of each species, particularly  
92 as regards desiccation tolerance and ability to tolerate CPA toxicity (Arakawa et al. 1990; Sakai et al.  
93 1990; Nishizawa et al. 1993).

94 The cell membrane is the primary site of cryo-injury when plant tissues are exposed to sub zero  
95 temperatures (Steponkus 1984). In addition to intracellular ice formation and the mechanical damage  
96 caused by this process, expansion-induced lysis, loss of osmotic responsiveness, membrane fusion and  
97 altered osmotic behaviour are other types of low temperature injury related to the cell membrane  
98 (Steponkus 1984; Wolfe and Bryant 1999). The preconditioning of plants by exposure to low non-  
99 freezing temperatures has been shown to improve survival following cryostorage in various plant species,  
100 with studies focusing on agriculturally important species or those native to subtropical or temperate  
101 climates, regardless of their ability to cold acclimate under natural conditions (Chang et al. 2000;  
102 Leunufna and Keller 2005; Kaczmarczyk et al. 2008; Kushnarenko et al. 2009). Additionally, sub-lethal

103 exposure to low temperature has been shown to cause changes in the chemical composition of cell  
104 membranes that coincides with improved resistance to cryo-injury (Steponkus 1984; Palta et al. 1993;  
105 Uemura et al. 1995). However, the cold acclimation ability of endemic Australian plants, such as those  
106 from the hot-summer Mediterranean climate regions of southwest Western Australia, has seldom been  
107 investigated to the same extent (Funnekotter et al. 2013). Plant species in this region of Australia can  
108 indeed be difficult to cryopreserve successfully (Kaczmarczyk et al. 2013).

109 A commonly used method of determining the relative stability or permeability state of the cell  
110 membrane is by the measurement of electrolyte leakage as a consequence of low temperature cellular  
111 damage (Campos et al. 2003; Mancuso et al. 2004). Low, lethal temperatures cause a change in the semi-  
112 permeable nature of the cell membrane, which results in a large, irreversible loss of electrolytes from  
113 plant tissues (Campos et al. 2003; Mancuso et al. 2004). Plants able to acclimate *via* changes that stabilize  
114 the cell membrane, can therefore tolerate lower temperatures by making their cell membranes “less  
115 leaky”, a response that can be easily measured and quantified using a conductivity probe and meter  
116 (Maier et al. 1994; Campos et al. 2003). The cell membrane typically consists of a lipid bilayer made up  
117 primarily of phospholipids (PLs). The type of PL headgroup and fatty acid (diglyceride) chains, as well as  
118 the level of unsaturation in these chains affect the stability, fluidity and permeability of the membrane  
119 (van Meer et al. 2008). Major PL headgroup types found in plants include phosphatidylcholine (PC),  
120 phosphatidylserine (PS), phosphatidylglycerol (PG), phosphatidylinositol (PI), phosphatidylethanolamine  
121 (PE) and phosphatidic acid (PA) (Uemura and Steponkus 1994; Uemura et al. 1995). Cold acclimation  
122 has shown to increase the concentration of PE and PC with unsaturated fatty acids as well as increased  
123 total PL levels, which reduce the occurrence of deleterious membrane phase transitions and improve  
124 freeze tolerance (Uemura and Steponkus 1994; Uemura et al. 1995).

125 Sterols or steroid alcohols are also an integral component of cell membranes that restrict the  
126 motion of the PL fatty acyl chains and thus control the fluidity and mobility of the membrane (Hartmann  
127 1998; Dufourc 2008); this also reduces the likelihood of phase separation occurring in the PLs (Dufourc  
128 2008). An increase in membrane sterol concentrations can also bring PL molecules closer together, thus  
129 reducing the permeability of the membranes (Demel and De Kruffyff 1976). The concentration of different  
130 types of sterols also has an effect on cell membranes.  $\beta$ -sitosterol has a greater effect on membrane  
131 permeability and ordering than stigmasterol (Schuler et al. 1991; Hartmann 1998). Increased chill  
132 tolerance in banana meristems and mung beans has been correlated to an increase in the  $\beta$ -sitosterol to  
133 stigmasterol ratio (Guye 1989; Zhu et al. 2006). Intracellular sugars or sugar alcohols also have a  
134 stabilising effect on cell membranes (Crowe et al. 1988; Wolfe and Bryant 1999; Turner et al. 2001).  
135 Soluble sugars reduce the gel phase to fluid phase transition temperature ( $T_m$ ) during dehydration of  
136 cellular membranes (Wolfe and Bryant 1999; Koster et al. 2000; Lenné et al. 2010). These are known to

137 occur *via* osmotic and volumetric effects of the sugars on membranes (Lenné et al. 2010). In addition,  
138 high intracellular sugar concentrations promote a vitrified state of water during cooling due to the highly  
139 viscous nature of their aqueous solutions (Wolfe and Bryant 1999).

140 Using the south-west Australian endemic *Lomandra sonderi* as a model species (Menon et al.  
141 2012) the overall aim of this study was to understand how low temperature preconditioning treatment  
142 might induce cold acclimation and potentially improve post-cryogenic survival in a Mediterranean species  
143 that might be expected to have a limited capacity to respond to low temperature acclimation. In this  
144 respect, changes to the temperature at which freeze damage occurs as well as the overall membrane PL  
145 and sterol and intracellular sugar compositions were determined during this study. These attributes were  
146 selected as indicators of cold acclimation in response to low temperature preconditioning due to their role  
147 in membrane stability and freeze damage mitigation. We discuss how the changes in composition  
148 observed relate to changes in post-cryogenic survival.

149

## 150 **Methods**

### 151 *Plant material*

152 *In vitro* plants of *Lomandra sonderi* were made available from the tissue culture collection at  
153 Kings Park and Botanic Garden, Western Australia (originally sourced from BHP Billiton Worsley  
154 Alumina Pty. Ltd., Boddington, Western Australia). Shoots of *L. sonderii* were maintained on a basal  
155 tissue culture medium (BM), which consisted of half strength Murashige and Skoog (MS; Murashige and  
156 Skoog 1962) macro- and micro-elements with 500  $\mu\text{M}$  myo-inositol, 500  $\mu\text{M}$  4-morpholineethanesulfonic  
157 acid, 100  $\mu\text{M}$  NaFeEDTA, 4  $\mu\text{M}$  niacin, 3  $\mu\text{M}$  thiamine-HCl, 2.5  $\mu\text{M}$  pyridoxine-HCl, 0.06 M sucrose,  
158 0.8% (w/v) agar and pH adjusted to 6.5 prior to autoclaving (20 min at 121 °C). BM with 0.2  $\mu\text{M}$  6-  
159 benzylaminopurine (BAP) was used for shoot maintenance and multiplication, and as a pre-culture  
160 treatment just prior to cryopreservation. Heat labile plant hormones including gibberellic acid (GA<sub>3</sub>) and  
161 zeatin (Z) [6-(4-hydroxy-3-methylbut-2-enylamino) purine] were filter-sterilised and added to BM  
162 following autoclaving to prepare recovery medium (RM). Maintenance medium (BM + 0.2  $\mu\text{M}$  BAP) was  
163 dispensed into 250 ml polycarbonate containers (~40-50ml per container) with polypropylene lids (each  
164 with 1 x 9 mm diam. vent covered with 0.2 micron micropore filter) then autoclaved; while RM was  
165 autoclaved in 500 ml media bottles and dispensed into sterile 55 mm diam. Petri plates at ~ 10 ml per  
166 plate. All media plate pouring and manipulations of *in vitro* plant material (for maintenance and  
167 cryopreservation) were carried out under sterile conditions in a laminar flow cabinet.

168

### 169 *Temperature preconditioning*

170 *L. sonderi* plants were preconditioned *in vitro* on BM (+ 0.2  $\mu\text{M}$  BAP) for 3-week incubation

171 periods using two temperature regimes: 20/-1 °C alternating (20°C during 16 h light period and -1°C  
172 during 8 h dark period) and constant 23 °C with a 12 h photoperiod, prior to any cryoexposure  
173 experiments. Preconditioning treatments were carried out in temperature controlled incubators with  
174 lighting supplied by 36 W cool white fluorescent tubes (Philips Alto; photon flux density of ~30  $\mu\text{mol m}^{-2}$   
175  $\text{s}^{-1}$ ).

176

#### 177 *Cryoexposure*

178 *Lomandra sonderi* shoot tips were extracted from *in vitro* grown shoots and cryostored using the  
179 method previously developed by Menon et al. (2012). Shoot tips from the two preconditioning regimes  
180 (20/-1 °C and 23 °C) were transferred onto BM containing 0.2  $\mu\text{M}$  BAP for 48 h as a pre-culture step,  
181 prior to loading treatment. Shoot tips were placed in ~ 10 ml of a loading solution (LS) comprised of 2 M  
182 glycerol and 0.4 M sucrose in liquid BM (as described above, without agar) in a 55 mm sterile plate for  
183 20 min, then treated in PVS2 [30% (w/v) glycerol, 15% (w/v) ethylene glycol, 15% (w/v) dimethyl  
184 sulfoxide (DMSO) in liquid BM with 0.4 M sucrose; (Sakai et al. 1990)] for 10 min at 23 °C. Each shoot  
185 tip was individually transferred to a 1  $\mu\text{L}$  droplet of PVS2 on a sterilised aluminium foil strip (~3 mm  
186 wide x 20 mm long; 5 shoot tips per strip). Two foil strips were placed in an empty, pre-chilled cryovial,  
187 plunged into LN and stored for a minimum of one hour.

188

#### 189 *Rewarming and recovery*

190 Cryovials containing *L. sonderi* shoot tips were rewarmed by immersion in a 40 °C water bath for  
191 10 seconds. The foil strips with shoot tips were then quickly removed from vials and shoot tips released  
192 into washing solution (1M sucrose in liquid BM) and incubated for 20 min at ~23°C. Control shoot tips  
193 were transferred directly after PVS2 treatment to this washing solution (no immersion in LN). All shoot  
194 tips were then placed onto a recovery medium (solid BM containing 1.0  $\mu\text{M}$  Z and 0.5  $\mu\text{M}$  GA<sub>3</sub>) for two  
195 weeks in darkness at ~25 °C, and then transferred to 23 °C/12 h photoperiod (as previously described).  
196 The survival of shoot tips on RM was determined using visual indicators 6 weeks after the date of  
197 rewarming. Green shoot tips showing leaf growth and increase in size were considered to be alive. Other  
198 shoot tips that did not show any signs of greening or growth after six weeks were considered non-viable.  
199 Percentage survival rates of the shoot tips were then determined.

200

#### 201 *Electrolyte leakage and LT<sub>50</sub> calculation following staged cooling*

202 Approximately 2 g of *L. sonderi* leaf material from the two temperature preconditioning regimes  
203 were excised per sample. The material per sample was placed in a glass test tube, and 200  $\mu\text{L}$  of deionised  
204 water was added to it. A total of 8 samples per preconditioning regime were placed into a cooling bath

205 (Model 9512 Refrigerating/Heating Circulator, PolySciences, Inc.), preset at -1 °C. The control samples  
206 were placed directly on ice. After one hour at -1 °C, the temperature of the cooling bath was gradually  
207 reduced at the rate of -1.5 °C h<sup>-1</sup> to -10 °C, with samples taken out every hour and stored on ice (8 hour  
208 procedure). 500 µL of deionised water was added to each sample before storage overnight at 4 °C. After  
209 24 h, 5 mL of deionised water was added to each sample before transferring to a shaker at 100 rpm  
210 overnight. 1 mL of each sample was transferred to 10 mL of deionised water and the conductivity of the  
211 diluted solution was measured (C<sub>1</sub>) using a 900-C Conductivity Meter (TPS Australia). The samples were  
212 boiled at 100 °C for 30 min. 1 mL of each boiled sample was then transferred to 10 mL of deionised  
213 water and the conductivity of the diluted solution was measured (C<sub>2</sub>). The electrolyte leakage during  
214 cooling was calculated as a percentage of total electrolyte concentration via the following formula:

215

$$216 \quad \% \text{ loss of electrolytes} = (C_1/C_2) * 100$$

217

218 To account for the loss of electrolytes for reasons other than freeze damage, this value was then  
219 zero corrected via the following formula:

220

$$221 \quad \text{Zero correction} = \% \text{ leakage of sample} - \% \text{ leakage of control}$$

222

223 These values were then normalized to the highest % electrolyte leakage value, to prevent  
224 percentage values above 100%, using the following formula:

225

$$226 \quad \text{Normalized value} = ([\text{zero correction}]/[\text{highest \% leakage value}]) * 100$$

227

228 The normalized values of percentage electrolyte leakage were used to construct a sigmoidal curve  
229 from which the LT<sub>50</sub> was calculated (the temperature at which 50% of total ion leakage occurs) using  
230 GraphPad Prism version 5.03 for Windows, GraphPad Software, San Diego California USA.

231

### 232 *Extraction of sterols and phospholipids*

233 Sterols and phospholipids were analysed as described by Funnekotter et al. (2013). *L. sonderi*  
234 shoot tips from the two temperature preconditioning regimes were isolated (9-11 mg, ~12 shoot tips).  
235 Shoot tips were homogenized in LN and 1 µg cholestanol and 1 ng 14:0/14:0-PC and 14:0/14:0-PG were  
236 added as sterol and PL internal standards. In order to dissolve the samples, 400 µL of methanol was added  
237 followed by incubation at 75°C for 20 min. 400 µL of chloroform and 400 µL of water were added to  
238 each followed by centrifugation at 10,000 rpm for 3 min. The chloroform phase containing the lipids was

239 then extracted. Solid phase extraction using Supelclean™ LC-Si 1 ml columns (Supelco, Bellefonte,  
240 USA) was performed on the chloroform phase. Neutral lipids were extracted using chloroform,  
241 glycolipids were extracted using acetone and the PLs were extracted using methanol. The neutral lipid  
242 and PL phases were dried under nitrogen gas, and the glycolipid phase was discarded.

243

#### 244 *Derivatisation and gas chromatography mass spectrometry (GC/MS) of sterol phase*

245 A solution of 10% potassium hydroxide (KOH) dissolved in methanol (500 µl) was used to  
246 dissolve the extracted neutral lipid phase. This was followed by incubation at 75°C for one hour and then  
247 followed by a triple hexane extraction. Sterols were derivatised by incubation in 20 µL methoxy pyridine  
248 at 65°C with shaking at 750 rpm for one hour and then incubation in 30 µL bis-  
249 (trimethylsilyl)trifluoroacetamide (BSTFA) at 65°C with shaking for one hour. An Agilent 7890A GC  
250 system coupled to an Agilent 5975C inertXL MSD (Agilent Technologies; Palo Alto, CA, USA) was  
251 utilised for the identification of sterols. A Varian VF-5MS column (30 m x 250 µm x 0.25 µm; Agilent  
252 Technologies) was used for separation of the samples. A 1 µL splitless injection was used to load the  
253 derivatised sterol extracts. The pressure of helium gas was 21.28 psi with a flow rate of 2.34 mol min<sup>-1</sup>  
254 within the column. Temperature of the column was increased from 70 °C to 325 °C. Agilent Technologies  
255 MSD ChemStation G1701EA E.02.00.493 was used for identification and quantitation of sterol species.

256

#### 257 *Liquid chromatography electrospray ionisation tandem mass spectrometry (LC-ESI-MS-MS) of* 258 *phospholipids*

259 100 µL of 50% v/v acetonitrile (in water) was used to dissolve the PL samples obtained from the  
260 solid phase extraction. An Agilent ZORBAX 300 Extended-C18 capillary column attached to an Agilent  
261 6340 ion trap LC/MS was used for the separation of PLs. A model 1200 capillary pump system from  
262 Agilent Technologies (Palo Alto, CA, USA) was used to inject the samples into the column. The method  
263 from Funnekotter et al. (2013) modified from Kim et al. (2009) and Min et al. (2010) was used for the  
264 runs. The solvents components of the mobile binary phase used were 50% v/v acetonitrile in water (A)  
265 and 90:10 (v/v) isopropanol/acetonitrile (B). The solvents contained 0.1% (v/v) formic acid and 0.05%  
266 ammonia for positive ion and negative ion detection, respectively. A separation gradient was used to  
267 separate the PL species. The gradient equilibrated to 55% solvent B as the column was loaded with the  
268 sample. The gradient ramped to 70% solvent B over 40 min and was held at 70% solvent B for an  
269 additional 5 min. Re-equilibration occurred *via* a 10 min post-run. A capillary voltage of 3500 V with a  
270 nebulizer pressure of 15 psi and dry nitrogen gas flow of 7 L min<sup>-1</sup> at 325 °C was used for electrospray  
271 ionisation. The ion scanning range for the tandem MS runs was 170-860 amu and 220-950 amu for the  
272 positive and negative modes, respectively. The scanning range for the quantitative MS runs was 620-860



273 amu for both the positive and negative modes. Agilent Technologies Data Analysis for 6300 Series Ion  
 274 Trap LC/MS v3.4 (Build 175) was used for identification and quantification of PL species. Calibration  
 275 standards of 18:0/18:0-PA, 18:0/18:0-PC, 18:0/18:0-PE, 18:0/18:0-PG, 18:1/18:1-PI and 16:0/16:0-PS  
 276 (Avanti Polar Lipids, Alabaster, AL, USA) were used to determine the quantities of species identified.

277

### 278 *Double-bond Index (DBI)*

279 The DBI was calculated for the fatty acyl chains of the PLs detected, with a formula modified from  
 280 (Quartacci et al. 2001):

281

$$282 \quad \text{DBI} = \frac{[(1 \times \% \text{monoenes}) + (2 \times \% \text{dienees}) + (3 \times \% \text{trienees}) + (4 \times \% \text{tetraenees}) + (5 \times \% \text{pentaenees})]}{\Sigma \% \text{ saturated fatty acids}}$$

283

284

### 284 *Extraction of sugars*

285 *L. sonderi* shoot tips from the two temperature preconditioning regimes were isolated (9-11 mg).  
 286 Shoot tips were homogenised in LN and sugars were extracted in 500  $\mu\text{L}$  20:2:0.2 methanol/water/ribitol  
 287 solution. Ribitol was used as an internal standard with a concentration of 0.2 mg mL<sup>-1</sup>. Samples were then  
 288 incubated at 75°C for 20 min. 100  $\mu\text{L}$  of the samples were then transferred to GC vials and dried in a  
 289 Speed Vac®.

290

### 291 *Derivatisation and GC/MS of sugars*

292 Derivatisation of sugars was also conducted in methoxypyridine and BSTFA as per the sterols.  
 293 Sugars were also identified using the Agilent 7890A GC system coupled to the Agilent 5975C inert XL  
 294 MSD as per the sterols. Flow rate of helium gas in the column was modified to 1 mol min<sup>-1</sup> at 8.8085 psi.  
 295 All other parameters remained unchanged from the sterol runs. Fructose, galactose, sucrose, glucose,  
 296 sorbitol, trehalose and raffinose (Sigma-Aldrich, St. Louis, MO, USA) were used as calibration standards  
 297 for the quantification of data. Agilent Technologies MSD ChemStation G1701EA E.02.00.493 was used  
 298 for identification and quantitation of sugar species.

299

### 300 *Statistics*

301 The cryoexposure experiments were performed using six replicates (6 x 15 shoot tips) for each  
 302 treatment. The cooling bath experiments were performed with three replicates per treatment. The sterol  
 303 and sugar experiments were performed with three replicates each (three times ~12 shoot tips) per  
 304 treatment. PL experiments were performed with four replicates (four times ~12 shoot tips) per treatment.  
 305 The Shapiro-Wilk test was used to check normality of data. No transformations were applied if data was

306 statistically normal. Statistical comparisons were made using independent t-tests for two-condition  
307 comparisons using SigmaPlot for Windows Version 11.0 Build 11.0.0.77 (Systat Software, Inc. Chicago,  
308 IL, USA).

309

## 310 **Results**

### 311 *Cryoexposure*

312 As previously determined (Menon et al. 2012), there was a significant difference in post-  
313 cryoexposure survival of shoot tips derived from the two preconditioning temperatures investigated (23  
314 and 20/-1 °C). Plants that had undergone the 20/-1 °C preconditioning had shoot tip survival of ~32.2%  
315 for samples not exposed to LN (-LN) and samples exposed to LN (+LN). Samples from the 23 °C  
316 preconditioning regime showed only ~14.4% survival for both -LN and +LN samples (Menon et al.  
317 2012). However, within each regime, there was no significant difference between -LN and +LN samples  
318 (Menon et al. 2012).

319

### 320 *LT<sub>50</sub>*

321 Alternating temperature preconditioning of plants resulted in altered electrolyte leakage readings  
322 from shoot samples, indicating that a change in cell membrane characteristics had occurred. The mean  
323 LT<sub>50</sub> value of shoot cultures incubated at 23°C was  $-2.50 \pm 0.08$  °C compared to  $-5.45 \pm 0.53$  °C for  
324 shoots incubated at 20/-1 °C.

325

### 326 *Phospholipid analysis*

327 PC and PE accounted for the highest content of PLs in both preconditioning regimes (over 97%).  
328 The remaining PL classes (PA, PG, PI and PS) were less abundant, being less than 3% of the total PLs  
329 detected. There was a significant increase ( $P < 0.05$ ) in the amount of PC in shoot tips from the 20/-1 °C  
330 preconditioning regime compared to those from the 23 °C preconditioning regime. The amounts of all the  
331 other PLs detected, particularly PE, did not significantly change between temperature treatments (Fig. 1).  
332 Due to the large increase in PC content, the total PL content in shoot tips from the 20/-1 °C  
333 preconditioning ( $25.0 \text{ ng mg}^{-1} \text{ FW}$ ) was significantly higher ( $P < 0.05$ ) than in those from the 23 °C  
334 preconditioning ( $9.8 \text{ ng mg}^{-1} \text{ FW}$ ). See Table S1 in Supporting Information for content of individual PL  
335 classes measured.

336 A total of 51 different PL species were identified and quantified (see Table S2 in Supporting  
337 Information). The widest range of PL species was detected in PC and PE, with only one type of PA  
338 species detected. Table 1 reports the amounts of each individual fatty acyl chain type detected in the PLs.  
339 Across both preconditioning regimes, palmitoleic acid (16:1) was the most common fatty acid detected,

340 constituting 20.3% and 18.7% of all fatty acyl chains detected in the 23 and 20/-1 °C samples,  
341 respectively (Table 1). The other fatty acyl chains found in high amounts in the 23 °C samples were oleic  
342 acid (18:1), linoleic acid (18:2), and linolenic acid (18:3), all of which were present in significantly higher  
343 amounts in the 20/-1 °C samples (Table 1). Large, significant increases were also seen in fatty acyl chains  
344 16:0, 18:0, 18:4, 20:2, 20:3 and 20:4 in the 20/-1 °C samples. The amount of C16 chains of the PLs had  
345 the largest increase after preconditioning, rising by 333% (Table 1).

346 Table 2 shows a comparison of the degrees of saturation and unsaturation in fatty acids between  
347 preconditioning regimes. There were large, significant increases in both the saturated and the unsaturated  
348 monoenes, dienes, trienes and tetraenes in the 20/-1 °C samples (Table 2). There was also an increase in  
349 pentaenes but this was not statistically significant. The associated DBIs were computed, revealing a large  
350 (50%), significant drop in DBI after preconditioning at 20/-1 °C ( $9.93 \pm 0.81$ ) compared to the 23 °C  
351 regime ( $20.32 \pm 1.23$ ).

352

### 353 *Sterol Analysis*

354 Cholesterol, stigmaterol and  $\beta$ -sitosterol were the three sterols detected. Any other sterol types  
355 (such as campesterol) were either present in trace quantities or non-existent.  $\beta$ -sitosterol accounted for the  
356 highest content of sterols in both preconditioning regimes. There was no significant difference in the  
357 content of any of the three sterol types detected between the two temperature pre-treatments.

358 The total content of sterols measured for the 23 °C samples ( $383.9 \pm 120.6$  ng mg<sup>-1</sup> FW) was  
359 ~8.5% higher than the 20/-1 °C samples ( $353.7 \pm 80.6$  ng mg<sup>-1</sup> FW), but was not significantly different.  
360 See Table S3 in Supporting Information for content of individual sterol types measured.

361

### 362 *Soluble Sugar Analysis*

363 There were seven types of sugar identified in the shoot tips. Fructose, galactose, sucrose and  
364 glucose were the main sugars present (>30 ng mg<sup>-1</sup> FW) and their concentrations were significantly  
365 different between the two preconditioning regimes (Fig. 2). The other three sugars identified (sorbitol,  
366 trehalose and raffinose) were detected in significantly smaller amounts (<10 ng mg<sup>-1</sup> FW) (Fig. 2). See  
367 Table S4 in Supporting Information for content of individual sugar types measured.

368 The total sugar contents in the two preconditioning regimes investigated showed large variations  
369 ( $P < 0.001$ ). The total sugar content of the 20/-1 °C samples ( $310.1 \pm 65.3$  ng mg<sup>-1</sup> FW) was ~15 times  
370 higher than that of the 23 °C samples ( $20.6 \pm 8.4$  ng mg<sup>-1</sup> FW).

371

372

373

## 374 Discussion

375 Overall, the preconditioning treatment resulted in substantial increases in sugar levels and PC content  
376 of shoot tissues, accompanied by a two-fold improvement in the ability of shoot tips to survive  
377 desiccation. In a previous study following preconditioning at 20/-1 °C shoot tips of *L. sonderi* showed an  
378 optimal mean survival of  $32.2 \pm 11.5\%$  following cryopreservation (Menon et al. 2012). In comparison,  
379 only  $14.4 \pm 2.7\%$  of shoot tips derived from the 23 °C preconditioning treatment survived after the same  
380 cryogenic treatment (Menon et al. 2012). The mean survival rates of the -LN controls were very similar to  
381 the corresponding mean +LN survival rates in shoot tips with the same preconditioning regime (Menon et  
382 al. 2012), suggesting that factors other than the cryopreservation protocol used (which was identical apart  
383 from the LN step) are responsible for the differences in survival observed. As previously determined  
384 (Menon et al. 2012), the lack of difference in survival between +LN and -LN samples could be attributed  
385 to the effective elimination of ice formation through the use of CPAs. Any reduction in survival would  
386 have therefore not been directly attributable to the LN immersion step, but rather to other processes such  
387 as toxic or desiccation stresses experienced by the plant material during the preparatory steps leading to  
388 cryopreservation. To temper these deleterious effects, cold acclimation stimulates plants to develop  
389 tolerance mechanisms that enhance their capacity to survive adverse conditions such as desiccation, to  
390 which plant tissues are subject during the cryogenic steps leading up to LN immersion (Thomashow  
391 1999). These tolerance mechanisms arise primarily *via* improved stability of cell membranes and  
392 accumulation of sugars (Thomashow 1999). Sugar accumulation is one of the mechanisms of desiccation  
393 tolerance in 'resurrection plants', such as *Craterostigma plantagineum*, as well as changes to phospholipid  
394 metabolism due to differential activation of phospholipase D genes (Bartels and Salamini 2001). Similar  
395 families of genes responsible for related processes, such as sugar transporters, sugar synthases and LEA  
396 proteins are upregulated as a response to both cold and desiccation stresses (Ingram and Bartels 1996;  
397 Thomashow 1999; Chinnusamy et al. 2007). It is therefore reasonable to postulate that improving the  
398 tolerance of a plant to one kind of stress, i.e. cold stresses, *via* preconditioning, would also improve  
399 tolerance to other stresses such as desiccation.

400 The  $LT_{50}$  values for the 23 °C and 20/-1 °C preconditioning regimes differed significantly. Similar  
401 findings were reported by Mancuso et al. (2004), who investigated the freezing tolerance of the Australian  
402 genera *Grevillea* and *Callistemon* using a similar electrolyte leakage method and determined that plants  
403 that were acclimated naturally to cold winter conditions exhibited significantly lower  $LT_{50}$  values.  
404 According to the plant hardiness map of Australia, divided into seven zones (Dawson 1991), the two  
405 *Grevillea* species studied by Mancuso et al. (2004; Table 3) grow in Northwest Western Australia (near  
406 Shark Bay), i.e. zone 5, with minimum temperatures during winter from 5 °C to 0 °C, and the four  
407 *Callistemon* species grow in Eastern New South Wales, a zone 3 region, with minimum temperatures

408 during winter from 0 °C to -5 °C. *Lomandra sonderi* naturally grows in southwest Western Australia, a  
409 zone 3 region, with similar winter conditions to the *Callistemon* species. Nevertheless, a major difference  
410 between both studies is that *L. sonderi* cultures were acclimated *in vitro* under controlled laboratory  
411 conditions intended to simulate natural average winter conditions, as opposed to the acclimation of pot-  
412 grown plants tested by Mancuso et al. (2004), which were maintained in greenhouses during winter.  
413 Regardless of the differences in methodology, Table 3 shows that the LT<sub>50</sub> values of non-acclimated and  
414 acclimated *L. sonderi* plant material are not greatly different from either *Callistemon* or *Grevillea* species,  
415 as described by Mancuso et al. (2004).

416 Cold acclimation is, however, known to be more effective in species that naturally grow in low  
417 average temperature climates than those from warmer origins (Flint 1972). For instance, the electrolyte  
418 leakage analysis of xylem tissue of evergreen Peach (*Prunus persica* [L.] Batsch) sampled in May (i.e.  
419 non-acclimated or deacclimated) generated an LT<sub>50</sub> value of -3.0 compared to samples taken in January,  
420 (i.e. cold acclimated) which had an LT<sub>50</sub> value of -17.0 °C (Arora et al. 1992). The plants were grown in  
421 Kearneysville, West Virginia, USA, which has an average annual minimum temperature of -17.8 °C to -  
422 20.6 °C (USDA 2012). This indicates that although cold acclimation can occur in species from warmer  
423 Mediterranean climates, the degree of acclimation possible appears to be far lower than in species from  
424 much colder climatic regions. Our findings with *L. sonderi* tend to reflect this difference.

425 The lower LT<sub>50</sub> values for *L. sonderi* in the 20/-1 °C regime are strong indicators of reduced  
426 membrane permeability and improved membrane stability, which may partly explain the improved  
427 cryogenic tolerance of shoot tips following a suitable acclimation treatment, as found for other species  
428 (Flint et al. 1967; Thomashow 1999; Campos et al. 2003; Mancuso et al. 2004).

429 PLs are the major component of the plasma membrane (van Meer et al. 2008). The increase  
430 measured in the total PL content in the 20/-1 °C preconditioning samples was largely due to the large  
431 (two-fold) increase in the amount of PC (Fig. 1). Funnekotter et al. (2013) observed a similar trend in the  
432 PC content of the Australian species they studied where it rose by 77% in *Grevillea scapigera* and by  
433 65% in *Loxocarya cinerea*, after preconditioning at 20/10 °C for three weeks. This suggests that  
434 preconditioning using low temperature regimes seems to increase PC content to stabilise cell membranes  
435 as a mechanism of chill tolerance. It is known that strong hydrating PL classes like PC are more likely to  
436 retain a lamellar phase in the cell membrane as compared to PL classes like PE, which are less hydrating  
437 and tend to form hexagonal II phases instead (Sen and Hui 1988). The total amount of PE, which is  
438 substantially lower than that of PC, remained unchanged in the two preconditioning regimes. These  
439 findings agree with the model of strongly hydrating PL classes, such as PC, which are more likely to  
440 maintain a functional liquid crystalline (lamellar) phase during cooling. Membranes containing higher  
441 amounts of PC are therefore more likely to withstand the stresses experienced during the entire

442 cryopreservation process, which is reflected in the increase observed in both cryopreserved (+LN) and  
443 control (-LN) shoot tip survival.

444         The gel-to-liquid crystalline transition temperature, i.e. the temperature at which the gel phase of  
445 a PL bilayer melts to form a lamellar phase, is affected by the fluidity of the membrane, which in turn  
446 affects its stability. The strength of attractive van der Waals forces between lipid molecules largely  
447 controls the phase behaviour of lipid bilayers. These forces are affected by the length and saturation of  
448 fatty acid tails of PLs. Longer acyl chains result in higher transition temperatures and reduced fluidity;  
449 conversely, higher degrees of unsaturation result in lower transition temperatures and increased fluidity  
450 (Silvius 1982). Interestingly, recent studies have revealed that increases in unsaturation in model cell  
451 membranes increase resistance to the deleterious effects of CPAs (Hughes et al. 2012; Malajczuk et al.  
452 2013) We would, therefore, expect to see a rearrangement in the composition of fatty acyl chains of PLs  
453 after 20/-1 °C preconditioning, as the plant attempts to increase the fluidity (and, hence, lower the  
454 transition temperature) of its cell membranes as a stress response, either by accumulating more short-  
455 tailed fatty acids or by increasing the degree of unsaturation. Indeed, the largest increase was seen in 16C  
456 chains in the PLs detected (increased by 333%), followed by 18C chains (increased by 195%) and then  
457 20C chains (increased by 150%; no significant increases were detected in 20:0, 20:1 and 20:5 fatty acyl  
458 chains). It would appear that an accumulation of PLs containing shorter fatty acyl chains is favoured  
459 during the cold acclimation of *L. sonderi*.

460         The DBI is a measure of the overall degree of unsaturation in fatty acyl chains. Increases in DBI  
461 indicate a higher degree of unsaturation, which in turn mean increased disorder in the PL bilayer structure  
462 due to the increase in the number of kinks in the chains, which consequently leads to increased fluidity  
463 (and reduced transition temperature) (Silvius 1982; Nishida and Murata 1996). Higher DBI has been  
464 related to stress tolerance in other plants (Nishida and Murata 1996). However, in the case of *L. sonderi* a  
465 large 50% decrease in DBI was observed after preconditioning at 20/-1 °C ( $9.93 \pm 0.81$ ) compared to  
466 preconditioning at 23 °C ( $20.32 \pm 1.23$ ). This trend was similarly observed by Funnekotter et al. (2013) in  
467 *Loxocarya cinerea*, which had a significantly lower DBI after preconditioning using an alternating 20/10  
468 °C regime, compared to preconditioning using a constant 23 °C regime. Zhu et al. (2006), who utilised a  
469 sucrose-pretreatment regime to improve cryoexposure tolerance in different banana varieties, found a  
470 significant decrease of DBI in PL fatty acids in the pretreated samples and simultaneously observed  
471 significantly improved cryogenic survival as a result of this pretreatment. It becomes clear that the loss of  
472 fluidity due to the decrease in DBI is counteracted by the accumulation of short, saturated fatty acyl chain  
473 PLs, in particular the more strongly hydrating PC, as discussed above. The level of contribution of each  
474 factor to membrane fluidity is, however, unknown and merits further research.

475 Membrane sterols are known to improve membrane stability *via* ordering of PLs (Demel and De  
476 Kruyff 1976), which can also protect membranes against damage by cryosolvents such as DMSO and  
477 polyols (Hughes and Mancera 2013; Hughes et al. 2013). The sterols identified in shoot tips of *L. sonderi*  
478 include stigmasterol, cholesterol and  $\beta$ -sitosterol. There was no statistically significant change measured  
479 in total sterol content across the two preconditioning regimes. Similarly, O'Neill et al. (1981) and  
480 Funnekotter et al. (2013) detected no changes in free sterols in *Fragraria virginiana* and *Arabidopsis*  
481 *thaliana*, in their respective studies, after cold acclimation. This may not hold true for all plant species,  
482 however, as Funnekotter et al. (2013) also found that *G. scapigera* and *L. cinerea* showed significantly  
483 higher amounts of sterols after preconditioning at 20/10 °C (69% and 88% increases, respectively). This  
484 suggests that, in the case of *L. sonderi*, membrane stability and/or permeability is not necessarily  
485 regulated exclusively by or reliant on changes in sterol content, at least in response to the temperature  
486 range treatments in this study.

487 Sugars are known to have a stabilising effect on cell membranes and promote the vitrification of  
488 water at low temperature (Wolfe and Bryant 1999). Preconditioning at 20/-1 °C largely increased the total  
489 content of sugar species in shoot tips of *L. sonderi* compared to the total content of sugars after 23 °C  
490 preconditioning. This may be due to the physiological responses to cold stress when exposed to low  
491 temperature in the 20/-1 °C preconditioning regimes. Increased intracellular sugar content may be a stress  
492 response that favours the vitrification of water (and hence the reduction of ice formation). In addition, the  
493 increased solute potential within the cells as a consequence of the increased sugar concentration would  
494 greatly reduce the desiccation effects of the cryopreservation processes. This can be associated with the  
495 greater survival rates that were recorded in both the cryopreserved (+LN) and control (-LN) shoot tips  
496 exposed to 20/-1 °C preconditioning. The large change in total sugar content was caused primarily by  
497 significant increases in glucose, fructose, galactose and sucrose, with minor, insignificant changes in  
498 trehalose, raffinose and sorbitol (which were detected in relatively small amounts). These changes  
499 indicate a robust response to the low temperature preconditioning regimes. This was similarly observed  
500 by Funnekotter et al. (2013) in *G. scapigera* and *L. cinerea*, which showed increases in total sugar content  
501 after preconditioning at 20/10 °C. These observations relate to previous findings, which showed an  
502 increase in sugar content as a survival response to abiotic stresses (Sasaki et al. 1996, Zhu et al. 2006).

503 The preconditioning regimes used in this study caused several significant changes to the plant  
504 shoot tip cells that are likely to be related to the improvement in post-cryoexposure survival. Changes to  
505 membrane composition, and by extension, its stability, as demonstrated by electrolyte leakage and  
506 PL/sterol analysis, appear to be involved in the reduction of freeze, osmotic and toxic stresses (Menon et  
507 al. 2012).

508 Studies describing acclimation effects on cryostorage success have been published with temperate  
509 and sub-tropical species (Chang et al. 2000; Leunufna and Keller 2005; Kaczmarczyk et al. 2008;  
510 Kushnarenko et al. 2009), however the number of studies on Mediterranean spp. especially Australian  
511 taxa is very limited by comparison. Indeed, given that five of the 34 internationally recognised  
512 biodiversity hotspots are found in small but floristically diverse Mediterranean climate regions including  
513 southwest Western Australia, South Africa's Cape floristic region, the European Mediterranean basin, the  
514 Californian Floristic Province and the Chilean Matorral it is imperative the underlying physiology of  
515 plants from these environments be better understood as a means to aid and enhance current conservation  
516 measures (Myers et al. 2000; Mittermeier et al. 2005). It might be argued that temperate or subtropical  
517 species are 'related enough' to Mediterranean species in terms of climatic extremes (particularly  
518 temperature) to void claims of any discernable physiological differences, however in the context of south-  
519 west Australian taxa the type of hot-summer Mediterranean climate is not directly comparable to classical  
520 temperate or sub-tropical climates. In particular, summer temperatures can be extreme and drought  
521 frequent, unlike most 'temperate' climates; with short cool wet winters this is again quite different from  
522 sub-tropical climates with warm dry 'winters' and hot wet summers (McKnight and Hess 2000).  
523 Therefore it is logical to expect that the adaptive physiology of south-west Australian Mediterranean  
524 plants would also express a different range (and possibly intensity) of responses to acclimation treatments  
525 compared to temperate or sub-tropical taxa. From this perspective while the current study reflects some  
526 subtle differences in adaptive physiology for acclimated *L. sonderi* plants, it also illustrates several key  
527 similarities to other studies on cold acclimation in species from more temperate northerly regions,  
528 suggesting that basic acclimation responses are conserved but adaptations are likely to have evolved in  
529 response to unique local environmental stresses.

530 The information gained on the effects of preconditioning with respect to changes in the  
531 membrane PL and cell sugar contents provide an excellent framework for further research on membrane  
532 stability and its relation to improved stress tolerance. Overall this study provides valuable insight into the  
533 underlying processes governing effects of low temperature preconditioning on post-cryogenic survival of  
534 native Australian plant species and enhances our current cryogenic approaches for the long term storage  
535 of valuable and endangered plant germplasm, as well as other plant material currently maintained under  
536 tissue culture conditions used for post-mining restoration.

537

### 538 **Acknowledgements.**

539 The authors declare that there are no conflicts of interest. This work was funded by ARC Linkage  
540 Grant LP0884027 with further financial support from Alcoa of Australia Ltd and BHP Billiton Worsley  
541 Alumina Pty Ltd. Dr. David Willyams of Alcoa of Australia's Marrinup Nursery is acknowledged for



542 providing the plant cultures used to conduct all experiments. The support of Matthew Timmins  
543 (Metabolomics Australia) for the analytical work is appreciated.

544

545

#### 546 **References**

547 Arakawa T, Carpenter JF, Kita YA, Crowe JH (1990) The basis for toxicity of certain cryoprotectants: a  
548 hypothesis. *Cryobiology* 27:405-415. doi:10.1016/0011-2240(90)90017-X

549 Arora R, Wisniewski ME, Scorza R (1992) Cold acclimation in genetically related (sibling) deciduous  
550 and evergreen peach (*Prunus persica* [L.] Batsch) I. Seasonal changes in cold hardiness and  
551 polypeptides of bark and xylem tissues. *Plant Physiol* 99 (4):1562-1568. doi:10.1104/pp.99.4.  
552 1562

553 Bartels D, Salamini F (2001) Desiccation Tolerance in the Resurrection Plant *Craterostigma*  
554 *plantagineum*. A Contribution to the Study of Drought Tolerance at the Molecular Level. *Plant*  
555 *Physiol* 127 (4):1346-1353. doi:10.1104/pp.010765

556 Campos PS, Quartin VN, Ramalho JC, Nunes MA (2003) Electrolyte leakage and lipid degradation  
557 account for cold sensitivity in leaves of *Coffea* sp. plants. *J Plant Physiol* 160 (3):283-292.  
558 doi:10.1078/0176-1617-00833

559 Chang Y, Barker RE, Reed BM (2000) Cold acclimation improves recovery of cryopreserved grass  
560 (*Zoysia* and *Lolium* sp.). *Cryoletters* 21 (2):107-116

561 Chinnusamy V, Zhu J, Zhu J-K (2007) Cold stress regulation of gene expression in plants. *Trends Plant*  
562 *Sci* 12 (10):444-451. doi:10.1016/j.tplants.2007.07.002

563 Crowe JH, Crowe LM, Carpenter JF, Rudolph AS, Wistrom CA, Spargo BJ, Anchordoguy TJ (1988)  
564 Interactions of sugars with membranes. *Biochim Biophys Acta* 947:367-384. doi:10.1016/0304-  
565 4157(88)90015-9

566 Dawson I (1991) Plant hardiness zones for Australia. *Australian Horticulture* 90 (8):37-39

567 Demel RA, De Kruyff B (1976) The function of sterols in membranes. *Biochim Biophys Acta* 457  
568 (2):109-132. doi:10.1016/0304-4157(76)90008-3

569 Dufourc EJ (2008) Sterols and membrane dynamics. *J Chem Biol*, 1(1-4), 63-77.

570 Food and Agriculture Organisation of the United Nations (2013) Genebank Standards for Plant Genetic  
571 Resources for Food Agriculture. Rome.

572 Flint HL (1972) Cold hardiness of twigs of *Quercus rubra* L. as a function of geographic origin.  
573 *Ecology*:1163-1170

- 574 Flint HL, Boyce BR, Beattie DJ (1967) Index of Injury—A useful expression of freezing injuries to plant  
575 tissues as determined by the electrolytic method. *Can J Plant Sci* 47 (2):229-230.  
576 doi:10.4141/cjps67-043
- 577 Funnekotter B, Kaczmarczyk A, Turner S, Bunn E, Zhou W, Smith S, Flematti G, Mancera R (2013)  
578 Acclimation-induced changes in cell membrane composition and influence on cryotolerance of in  
579 vitro shoots of native plant species. *Plant Cell Tiss Org*:1-14. doi:10.1007/s11240-013-0308-0
- 580 Guye M (1989) Phospholipid, sterol composition and ethylene production in relation to choline-induced  
581 chill-tolerance in mung bean (*Vigna radiata* L. Wilcz.) during a chill-warm cycle. *J Exp Bot* 40  
582 (3):369-374. doi:10.1093/jxb/40.3.369
- 583 Hartmann M (1998) Plant sterols and the membrane environment. *Trends Plant Sci* 3 (5):170-175.  
584 doi:10.1016/s1360-1385(98)01233-3
- 585 Hughes ZE, Malajczuk CJ, Mancera RL (2013) The Effects of Cryosolvents on DOPC- $\beta$ -sitosterol  
586 Bilayers determined from molecular dynamics simulations. *J Phys Chem B* 117 (12):3362-3375.  
587 doi:10.1021/jp400975y
- 588 Hughes ZE, Mark, AE, Mancera, RL (2012) Molecular dynamics simulations of the interactions of  
589 DMSO with DPPC and DOPC phospholipid membranes. *J Phys Chem B* 116 (39):11911-11923.
- 590 Hughes ZE, Mancera RL (2013) Molecular dynamics simulations of mixed DOPC- $\beta$ -sitosterol bilayers  
591 and their interactions with DMSO. *Soft Matter* 9 (10):2920-2935. doi:10.1039/C3SM27805K
- 592 Ingram J, Bartels D (1996) The molecular basis of dehydration tolerance in plants. *Annu Rev Plant Biol*  
593 47 (1):377-403. doi:10.1146/annurev.arplant.47.1.377
- 594 Kaczmarczyk A, Funnekotter B, Menon A, Phang PY, Al-Hanbali A, Bunn E, Mancera RL (2012)  
595 Current Issues in Plant Cryopreservation. In: Katkov II (ed) *Current Frontiers in*  
596 *Cryopreservation*. InTech Open Access Publisher, pp 417-428
- 597 Kaczmarczyk A, Funnekotter B, Turner SR, Bunn E, Bryant G, Hunt TE, Mancera RL (2013)  
598 Development of cryopreservation for *Loxocarya cinerea* – an endemic Australian plant species  
599 important for post-mining restoration”, *Cryo Letters* 34 (5): 508-519.
- 600 Kaczmarczyk A, Shvachko N, Lupysheva Y, Hajirezaei M-R, Keller E (2008) Influence of alternating  
601 temperature preculture on cryopreservation results for potato shoot tips. *Plant Cell Rep* 27  
602 (9):1551-1558. doi:10.1007/s00299-008-0574-1
- 603 Kaczmarczyk A, Turner SR, Bunn E, Mancera RL, Dixon KW (2011) Cryopreservation of threatened  
604 native Australian species—what have we learned and where to from here? *In Vitro Cell Dev-Pl*  
605 47 (1):17-25
- 606 Kim H, Min H, Kong G, Moon M (2009) Quantitative analysis of phosphatidylcholines and  
607 phosphatidylethanolamines in urine of patients with breast cancer by nanoflow liquid

- 608 chromatography/tandem mass spectrometry. *Anal Bioanal Chem* 393 (6):1649-1656.  
609 doi:10.1007/s00216-009-2621-3
- 610 Koster KL, Lei YP, Anderson M, Martin S, Bryant G (2000) Effects of Vitrified and Nonvitrified Sugars  
611 on Phosphatidylcholine Fluid-to-Gel Phase Transitions. *Biophysical Journal* 78 (4):1932-1946.  
612 doi:10.1016/s0006-3495(00)76741-5
- 613 Kushnarenko SV, Romadanova NV, Reed BM (2009) Cold Acclimation Improves Regrowth of  
614 Cryopreserved Apple Shoot Tips. *Cryoletters* 30 (1):47-54
- 615 Lenné T, Garvey CJ, Koster KL, Bryant G (2010) Kinetics of the lamellar gel–fluid transition in  
616 phosphatidylcholine membranes in the presence of sugars. *Chemistry and Physics of Lipids* 163  
617 (2):236-242. doi:10.1016/j.chemphyslip.2009.12.001
- 618 Leunufna S, Keller ER (2005) Cryopreservation of yams using vitrification modified by including droplet  
619 method: effects of cold acclimation and sucrose. *Cryo Letters* 26 (2):93-102
- 620 Maier FP, Lang NS, Fry JD (1994) Evaluation of an Electrolyte Leakage Technique to Predict St.  
621 Augustinegrass Freezing Tolerance. *HortScience* 29 (4):316-318
- 622 Malajczuk CJ, Hughes ZE, Mancera RL (2013) Molecular dynamics simulations of the interactions of  
623 mono- and polyhydroxylated cryosolvents with a hydrated phospholipid bilayer. *Biochim  
624 Biophys Acta - Biomembranes* 1828 (9): 2041-2055.
- 625 Mancuso S, Nicese FP, Masi E, Azzarello E (2004) Comparing fractal analysis, electrical impedance and  
626 electrolyte leakage for the assessment of cold tolerance in *Callistemon* and *Grevillea* spp. *Journal  
627 of Horticultural Science and Biotechnology* 79 (4):627-632
- 628 McKnight TL, Hess, D (2000). *Climate Zones and Types*. In: *Physical Geography: A Landscape  
629 Appreciation*. Upper Saddle River, NJ: Prentice Hall.
- 630 Menon A, Funnekotter B, Kaczmarczyk A, Bunn E, Turner SR, Mancera RL (2012) Cryopreservation of  
631 *Lomandra sonderi* (Asparagaceae) shoot tips using droplet-vitrification. *CryoLetters* 33 (4):259-  
632 270
- 633 Min H, Kong G, Moon M (2010) Quantitative analysis of urinary phospholipids found in patients with  
634 breast cancer by nanoflow liquid chromatography–tandem mass spectrometry: II. Negative ion  
635 mode analysis of four phospholipid classes. *Anal Bioanal Chem* 396 (3):1273-1280.  
636 doi:10.1007/s00216-009-3292-9
- 637 Mittermeier RA, Gil PR, Hoffman M, Pilgrim J, Brooks T, Mittermeier CG, Lamoreux J, Da Fonseca  
638 GAB, Seligmann P (Eds) (2005) 'Hotspots revisited. *Conservation International*.' (University of  
639 Chicago Press: Chicago, IL).

- 640 Murashige T, Skoog F (1962) A Revised Medium for Rapid Growth and Bio Assays with Tobacco Tissue  
641 Cultures. *Physiologia Plantarum* 15 (3):473-497. doi:10.1111/j.1399-3054.1962.tb08052.x
- 642 Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Kent J. 2000. Biodiversity hotspots for  
643 conservation priorities. *Nature* 403: 803–808
- 644 Nishida I, Murata N (1996) Chilling sensitivity in plants and cyanobacteria: the crucial contribution of  
645 membrane lipids. *Annu Rev Plant Biol* 47 (1):541-568
- 646 Nishizawa S, Sakai A, Amano Y, Matsuzawa T (1993) Cryopreservation of asparagus (*Asparagus*  
647 *officinalis* L.) embryogenic suspension cells and subsequent plant regeneration by vitrification.  
648 *Plant Science* 91 (1):67-73. doi:10.1016/0168-9452(93)90189-7
- 649 O'Neill SD, Priestley DA, Chabot BF (1981) Temperature and Aging Effects on Leaf Membranes of a  
650 Cold Hardy Perennial, *Fragaria virginiana*. *Plant Physiol* 68 (6):1409-1415
- 651 Offord CA, Bunn E, Turner SR, Sommerville KD, Siemon J, Ashmore SE (2009) Tissue Culture. In:  
652 Offord CA, Meagher PF (eds) *Plant Germplasm Conservation in Australia*. Australian Network  
653 for Plant Conservation Inc., Sydney, pp 109-128
- 654 Palta JP, Whitaker BD, Weiss LS (1993) Plasma Membrane Lipids Associated with Genetic Variability in  
655 Freezing Tolerance and Cold Acclimation of *Solanum* Species. *Plant Physiol* 103 (3):793-803.  
656 doi:10.1104/pp.103.3.793
- 657 Quartacci MF, Cosi E, Navari-Izzo F (2001) Lipids and NADPH-dependent superoxide production in  
658 plasma membrane vesicles from roots of wheat grown under copper deficiency or excess. *Journal*  
659 *of Experimental Botany* 52 (354):77-84
- 660 Sakai A, Kobayashi S, Oiyama I (1990) Cryopreservation of nucellar cells of navel orange (*Citrus*  
661 *sinensis* Osb. var. *brasiliensis* Tanaka) by vitrification. *Plant Cell Reports* 9 (1):30-33.  
662 doi:10.1007/bf00232130
- 663 Sasaki H, Ichimura K, Oda M (1996) Changes in Sugar Content during Cold Acclimation and  
664 Deacclimation of Cabbage Seedlings. *Ann Bot* 78 (3):365-369. doi:10.1006/anbo.1996.0131
- 665 Schuler I, Milon A, Nakatani Y, Ourisson G, Albrecht AM, Benveniste P, Hartman MA (1991)  
666 Differential effects of plant sterols on water permeability and on acyl chain ordering of soybean  
667 phosphatidylcholine bilayers. *Proceedings of the National Academy of Sciences* 88 (16):6926-  
668 6930
- 669 Sen A, Hui S-W (1988) Direct measurement of headgroup hydration of polar lipids in inverted micelles.  
670 *Chemistry and Physics of Lipids* 49 (3):179-184. doi:10.1016/0009-3084(88)90005-9
- 671 Silvius J (1982) Thermotropic phase transitions of pure lipids in model membranes and their  
672 modifications by membrane proteins. *Lipid-protein interactions* 2:239-281

- 673 Steponkus PL (1984) Role of the Plasma Membrane in Freezing Injury and Cold Acclimation. *Annu Rev*  
674 *Plant Phys* 35 (1):543-584. doi:10.1146/annurev.pp.35.060184.002551
- 675 Thomashow MF (1999) PLANT COLD ACCLIMATION: Freezing Tolerance Genes and Regulatory  
676 Mechanisms. *Annu Rev Plant Phys* 50 (1):571-599. doi:doi:10.1146/annurev.arplant.50.1.571
- 677 Turner SR, Senaratna T, Touchell DH, Bunn E, Dixon KW, Tan B (2001) Stereochemical arrangement of  
678 hydroxyl groups in sugar and polyalcohol molecules as an important factor in effective  
679 cryopreservation. *Plant Science* 160 (3):489-497. doi:10.1016/s0168-9452(00)00420-9
- 680 Uemura M, Joseph RA, Steponkus PL (1995) Cold Acclimation of *Arabidopsis thaliana*. *Plant Physiol*  
681 109:15-30
- 682 Uemura M, Steponkus PL (1994) A Contrast of the Plasma Membrane Lipid Composition of Oat and Rye  
683 Leaves in Relation to Freezing Tolerance. *Plant Physiol* 104 (2):479-496
- 684 van Meer G, Voelker DR, Feigenson GW (2008) Membrane lipids: where they are and how they behave.  
685 *Nat Rev Mol Cell Biol* 9 (2):112-124
- 686 Wolfe J, Bryant G (1999) Freezing, drying and/or vitrification of membrane-solute-water systems.  
687 *Cryobiology* 39:103-129
- 688 Zhu GY, Geuns J, Dussert S, Swennen R, Panis B (2006) Change in sugar, sterol and fatty acid  
689 composition in banana meristems caused by sucrose-induced acclimation and its effects on  
690 cryopreservation. *Physiologia Plantarum* 128 (1):80-94
- 691
- 692
- 693
- 694
- 695
- 696
- 697
- 698
- 699
- 700
- 701
- 702
- 703
- 704
- 705

## 706 **Supporting Information**

707

708 **Table S1.** Phospholipids (ng mg<sup>-1</sup> FW) detected in *Lomandra sonderi* shoot tips after temperature  
709 preconditioning at 23 °C and 20/-1 °C for 3 weeks.

710 **Table S2.** Identified phospholipid species (ng mg<sup>-1</sup> fresh weight ± SD) found in *Lomandra sonderi* shoot  
711 tips after preconditioning at 23 °C or 20/-1 °C for 3 weeks.

712 **Table S3.** Sterols (ng mg<sup>-1</sup> FW) detected in *Lomandra sonderi* shoot tips after temperature  
713 preconditioning at 23 °C and 20/-1 °C for 3 weeks.

714 **Table S4.** Sugars (ng mg<sup>-1</sup> FW) detected in *Lomandra sonderi* shoot tips after temperature  
715 preconditioning at 23 °C and 20/-1 °C for 3 weeks.

716

## 717 **Figure Legends**

718 **Fig. 1** Content of phospholipid (PL) classes (ng mg<sup>-1</sup> fresh weight) analysed in *Lomandra sonderi* shoot  
719 tips exposed to three-week preconditioning at 23 °C or 20/-1 °C. Bars represent average ± SE of four  
720 replicates per treatment assessed. Bars labelled with different letters differ significantly when compared  
721 across treatments ( $P < 0.05$ ). PC - phosphatidylcholine, PE - phosphatidylethanolamine, PA -  
722 phosphatidic acid PG - phosphatidylglycerol, PI - phosphatidylinositol, PS - phosphatidylserine

723

724 **Fig. 2** Content of each sugar (ng mg<sup>-1</sup> fresh weight) analysed in *Lomandra sonderi* shoot tips exposed to  
725 three-week preconditioning at 23 °C and 20/-1 °C. Bars represent average ± SE of three replicates of  
726 treatments assessed. Bars labelled with different letters differ significantly when compared across  
727 treatments ( $P < 0.05$ )

728

729

730

731

732

733

734

735

736


737

738

739

740 **Figure 1.**

741




742

743

744 **Figure 2.**

745



746