


On: 14 November 2013, At: 05:51

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Bird Study

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tbis20>

Geographic and temporal variation in the consumption of bats by European Barn Owls

Alexandre Roulin^a & Philippe Christe^a

^a Department of Ecology and Evolution, University of Lausanne, Building Biophore, 1015 Lausanne, Switzerland

Published online: 24 Oct 2013.

To cite this article: Alexandre Roulin & Philippe Christe (2013) Geographic and temporal variation in the consumption of bats by European Barn Owls, *Bird Study*, 60:4, 561-569, DOI: [10.1080/00063657.2013.847051](https://doi.org/10.1080/00063657.2013.847051)

To link to this article: <http://dx.doi.org/10.1080/00063657.2013.847051>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

SHORT REPORT

Geographic and temporal variation in the consumption of bats by European Barn Owls

ALEXANDRE ROULIN* and PHILIPPE CHRISTE

Department of Ecology and Evolution, University of Lausanne, Building Biophore, 1015 Lausanne, Switzerland

Capsule We report a review of the occurrence of bats in the Barn Owl diet *Tyto alba* in Europe. Based on 802 studies reporting 4.02 million prey items identified in pellets, 4949 were bats (0.12%). We found that bat predation decreased during the last 150 years, is more frequent on islands than mainland, and is higher in eastern than western Europe and in southern than northern Europe. Although Barn Owls usually capture bats opportunistically, they can sometimes specialize on them.

In Europe, the Barn Owl *Tyto alba* rarely consumes bats, although individuals can sometimes specialize on them (Bauer 1956). This specialization can be a consequence of owls and bats roosting or breeding in similar habitats such as buildings or cliffs. As a consequence, bats can sometimes abandon a site if Barn Owls depredate them or simply because of their mere presence (Boireau 2009). Although few data are available on population trends over decades, bats have apparently declined in Europe (Stebbins 1988). By reviewing the literature on the Barn Owl diet in Europe, we could, therefore, test whether Barn Owls have reduced bat consumption during the last 150 years (Uhrin *et al.* 2010). This review also allowed us to investigate whether there is geographic variation in bat predation or whether some bats are more vulnerable when displaying specific flying behaviour and so whether Barn Owls capture bats opportunistically.

As described in two previous reviews on Barn Owl diet obtained from pellet analyses (Roulin & Dubey 2012, 2013), we collected all papers published in international and local journals we could find. Among the 802 studies considered in the present review, 319 (39.8%) reported at least one bat prey item (Fig. 1). The highest percentage of bats was 54.2% in a study that reported 52 out of 96 prey items (Bauer 1956). Table 1 lists, for each European country and island, the total number of bats that were found in the Barn Owl diet. In total, 4025 523 vertebrate and

invertebrate prey items were identified of which 4949 were bats (0.12%). The highest proportion of bats was found in Crete (3.59%) followed by Sardinia (1.64%).

Using the entire sample of 802 studies, the distribution of the proportion of bats in the Barn Owl diet strongly departed from normality because of the large number of studies that did not report any bats ($n = 483$, Fig. 1). We, therefore, performed non-parametric analyses. The proportion of bats did not differ between islands and mainland (Wilcoxon matched-pair signed-rank, $z = 0.17$, $P = 0.86$) and was not significantly associated with latitude (Spearman's correlation: $r_s = -0.05$, $n = 788$, $P = 0.19$) but increased with longitude ($r_s = 0.15$, $n = 788$, $P < 0.0001$) and decreased between 1860 and 2010 ($r_s = -0.13$, $n = 763$, $P = 0.0002$). When considering only studies ($n = 319$) with at least one bat prey item, the log-transformed percentage of bats was normally distributed, and hence we could perform more powerful parametric analyses. In an ANCOVA, the percentage of bats in the Barn Owl diet was significantly higher on islands than mainland ($F_{1,295} = 4.4$, $P = 0.037$), decreased with latitude ($F_{1,295} = 5.8$, $P = 0.017$, Fig. 2a), increased with longitude ($F_{1,295} = 14.5$, $P = 0.0002$; Fig. 2b) and decreased between 1860 and 2010 ($F_{1,295} = 5.52$, $P = 0.019$; Fig. 2c: note a quadratic function of year to test whether this decline has slowed in recent years – see below – did not give a better fit). In this model, we statistically controlled for log-transformed total number of prey items identified ($F_{1,295} = 167.7$, $P < 0.0001$) because lower percentages of bats were found in larger samples (Pearson's correlation:

*Correspondence author. Email: alexandre.roulin@unil.ch




Figure 1. Frequency distribution of 802 studies reporting the percentage of bats in Barn Owl diet in Europe. Number above bars indicates the absolute number of studies.

$r = -0.60$, $P < 0.0001$). Note that if we do not control for this variable, we obtain qualitatively similar results. The higher proportion of bats found in the Barn Owl

diet on islands may reflect the lower diversity and abundance of terrestrial small mammals on islands compared to the mainland (Alcover *et al.* 1998). The positive correlation between proportion of bats in owl pellets and longitude may be explained by higher bat population sizes in eastern European or/and by a closer proximity of Barn Owl roosts with bat colonies in eastern Europe, for example, if bat colonies are more often located in anthropogenic habitats than in natural caves. The higher proportion of bats reported in southern Europe may be explained by the less extreme weather conditions known to be detrimental to insectivorous bats and therefore higher density of bat populations and longer activity period during the year (Humphries *et al.* 2002 but see Sendor & Simon 2003). Clinal variation in bat depredation by Barn Owls had already been observed in Spain and Europe

Table 1. Proportion of Barn Owl diet composed of bats in different European countries and islands.

	Number of studies	Number of prey identified	Number of bats as prey	Percentage of bats in the diet
Albania	1	68	0	0
Austria	12	15 011	153	1.02
Balearic Islands	3	10 425	7	0.07
Belgium	11	159 384	64	0.04
Bosnia	2	2272	0	0
Bulgaria	5	41 787	40	0.10
Corfu	2	5888	20	0.34
Corsica	6	24 498	125	0.51
Cos	1	2277	0	0
Crete	4	4321	155	3.59
Croatia	3	8633	39	0.45
Czech Republic	27	115 516	99	0.09
Denmark	1	36 173	49	0.14
France	109	804 778	710	0.09
Germany	209	788 917	946	0.12
Greece	11	9630	2	0.02
Hungary	39	233 475	311	0.13
Ireland	19	28 442	60	0.21
Italy	93	150 954	231	0.15
Luxemburg	10	11 286	22	0.19
Malta	2	424	3	0.71
The Netherlands	13	150 131	37	0.02
Poland	28	201 870	477	0.24
Portugal	12	23 682	9	0.04
Romania	3	3283	0	0
Russia	9	2088	0	0
Sardinia	6	2619	43	1.64
Serbia	2	8576	1	0.01
Sicily	6	13 210	78	0.59
Slovakia	8	23 431	192	0.82
Slovenia	8	27 896	277	0.99
Spain	51	204 342	581	0.28
Sweden	2	3285	0	0
Switzerland	18	298 051	28	0.01
UK	61	572 714	155	0.03
Ukraine	4	34 128	35	0.10
Total	801	4 023 465	4949	0.12




Figure 2. Frequency of bats in Barn Owl diet in Europe in relation to latitude (a), longitude (b) and year (c). Predicted lines from an overall model (see text) are plotted.

(Pérez-Barbería 1991) further indicating that this finding is robust. Finally, the long-term decrease in the proportion of bats in the Barn Owl diet may reflect historical declines of bat populations during the last century (Stebbins 1988). In Poland, Lesiński (2010) showed that Tawny Owls (*Strix aluco*) consumed fewer

bats in the 1980s probably because of the intensive use of insecticides in those years. Bat populations recovered after the 1980s due to a lower use of insecticides, leading to an increase in bat consumption by Tawny Owls. We, therefore, tested whether in the Barn Owl bat consumption increased along the years after 1980. This was, however, not the case (similar model as above, year: $F_{1,110} = 0.06$, $P = 0.80$).

In many papers, the authors identified bats to the species level (supplementary online table shows the number of bats identified to the species level in different European countries and islands). Because in several papers, the authors reported only bats but not the other prey species, more bats were reported in Table 2 than in Table 1. We could thus examine whether bat depredation was related to bat body mass and measures associated with flying strategies (wing loading and aspect ratio). For instance, bats with long and narrow wings (high wing loading) fly rapidly and higher above the ground, whereas species with broad wings (low wing loading) fly slowly and are highly manoeuvrable (Norberg & Rayner 1987, Fenton 1990, Obrist *et al.* 2011). However, among the 23 species of bats identified as Barn Owl prey and for which we had data on wing loading, aspect ratio and body mass (Table 2), their log-transformed frequency in Barn Owl diet was not associated with these three variables (multiple regression analysis: P -values > 0.22). This suggests that bat flying behaviour does not play an important role on the probability to be preyed by Barn Owls. This is not surprising because Barn Owls frequently capture young bats that are not yet able to fly (Bauer 1956, Schmidt & Topal 1971, Glutz & Bauer 1980) and bats seem to be rarely captured in flight (König 1961). This suggests that most bats are captured at roosting and breeding sites in an opportunistic way, as shown in the Tawny Owl *Strix aluco* (Lesiński *et al.* 2012). In line with this statement, we found that bats foraging in open and semi-open landscapes were captured as often as bats foraging in forests (two-way ANOVA: $F_{1,27} = 0.07$, $P = 0.80$) and bats with more stable populations were more often captured ($F_{1,27} = 14.33$, $P = 0.0008$) indicating that bat abundance rather than some biological specificity is associated with their occurrence in the Barn Owl diet.

To conclude, our study suggests that Barn Owls opportunistically depredate bats. By reviewing the literature on Barn Owl diet, we could show that predation on bats decreased during the last century probably because bats declined. Predation was also

Table 2. Number of individuals and of regions (see list in Table 1) (N/#) where bats (identified to the species level) were found in the Barn Owl diet in Europe. For each identified species, we give information on hunting habitat, location of roosts and colonies, size of colonies, population status, population trend and body mass. Information on population status and trends are from IUCN 2012 (IUCN Red List of Threatened Species. Version 2012.2. <www.iucnredlist.org>). Biological data are from Dietz *et al.* (2009) and data on wing loading and aspect ratio are from Norberg & Rayner (1987).

	N/#	Wing loading (N/m ²)	Aspect ratio	Habitat	Roost colonies	Colony size	Population status	Population trend	Mean body mass (g)
Rhinolophidae									
Blasius' horseshoe bat (<i>Rhinolophus blasii</i>)	2/1			1	1, 2	30–500	1	1	12
Mediterranean horseshoe bat (<i>Rhinolophus euryale</i>)	9/4	8.1	6.2	2	1, 2	20–300	2	1	11
Greater horseshoe bat (<i>Rhinolophus ferrumequinum</i>)	133/6	12.2	6.1	2	1, 2	20–1000	1	1	21
Lesser horseshoe bat (<i>Rhinolophus hipposideros</i>)	28/8	7.1	5.7	2	1, 2	10–200	1	1	7
Mehely's horseshoe bat (<i>Rhinolophus mehelyi</i>)	1/1	11.6		1	1, 2	20–200	3	1	14
<i>Rhinolophus</i> sp.	3/2				1, 2				
Vespertilionidae									
Western barbastelle (<i>Barbastella barbastellus</i>)	47/8	9.1	6.0	2	2, 3	10–100	2	1	9
Northern bat (<i>Eptesicus nilssonii</i>)	17/4	8.1	6.6	2	2, 3	20–50	1	2	13
Serotine bat (<i>Eptesicus serotinus</i>)	920/15	12.2	6.5	1	2	10–60	1	4	25
Savi's pipistrelle (<i>Hypsugo savii</i>)	16/4			1	2	40–70	1	2	7
Bechstein's myotis (<i>Myotis bechsteini</i>)	36/5	9.0	6.0	2	3	20–50	2	1	8
Lesser mouse-eared bat (<i>Myotis blythii</i>)	153/10	10.1	6.7	1	1, 2	50–500	1	1	21
Brandt's myotis (<i>Myotis brandtii</i>)	15/4			2	3	20–60	1	2	6
Long-fingered bat (<i>Myotis capaccinii</i>)	3/2			1	1, 2	30–500	3	1	8
Pond bat (<i>Myotis dasycneme</i>)	31/6	10.4	6.8	1	2, 3	20–300	2	1	15
Daubenton's bat (<i>Myotis daubentoni</i>)	107/7	7.0	6.3	1	2, 3	20–50	1	3	8
Geoffroy's bat (<i>Myotis emarginatus</i>)	45/7	7.1	5.9	2	1, 2	20–500	1	2	7
Greater mouse-eared bat (<i>Myotis myotis</i>)	1,278/17	11.2	6.3	2	1, 2	50–1,000	1	2	23
Whiskered bat (<i>Myotis mystacinus</i>)	56/8	7.1	6.0	1	2, 3	20–60	1	4	6
<i>M. mystacinus</i> or <i>M. brandtii</i>	1/1								
Natterer's bat (<i>Myotis nattereri</i>)	512/10	6.1	6.4	1	2, 3	20–50	1	2	8
<i>Myotis</i> sp.	68/11								
Giant noctule (<i>Nyctalus lasiopterus</i>)	2/1			1	3	Up to 80	2	1	44
Leisler's bat (<i>Nyctalus leisleri</i>)	21/4	19.3	7.9	2	3	20–50	1	4	15
Noctule bat (<i>Nyctalus noctula</i>)	363/10	16.1	7.4	1, 2	2, 3	20–60	1	4	26
<i>Nyctalus</i> sp.	1/1								
Kuhl's pipistrelle (<i>Pipistrellus kuhlii</i>)	178/7	8.5	6.3	1	2	20–100	1	4	7
Nathusius' pipistrelle (<i>Pipistrellus nathusii</i>)	130/11	9.8	7.2	2	3, 2	20–200	1	4	8
Common pipistrelle (<i>Pipistrellus pipistrellus</i>)	621/17	8.1	7.5	1	2, 3	50–100	1	2	5
Pygmy pipistrelle (<i>Pipistrellus pygmaeus</i>)	50/1			1	2, 3	15–800	1	4	6
<i>P. pipistrellus</i> or <i>P. pygmaeus</i>	36/1								
<i>Pipistrellus</i> sp.	143/10								
Brown big-eared bat (<i>Plecotus auritus</i>)	323/11	7.1	5.7	2	2, 3	5–50	1	2	8
Grey long-eared bat (<i>Plecotus austriacus</i>)	242/8	7.9	6.1	1	2, 3	10–30	1	4	8
<i>Plecotus</i> sp.	31/6								
Particoloured bat (<i>Vespertilio murinus</i>)	107/7	10.2	7.0	1	2	20–60	1	2	12
Miniopteridae									
Schreiber's bat (<i>Miniopterus schreibersii</i>)	54/8	10.2	7.0	1	1	100–20,000	2	1	12
Molossidae									
European free-tailed bat (<i>Tadarida teniotis</i>)	6/3			1	2	5–50	4	4	25
Pteropodinae									
Egyptian fruit bat (<i>Rousettus aegyptiacus</i>)	79/1			1	1	50–500	1–2	4	124
Total	5868								

Habitat – hunting habitat: 1 = open and semi-open landscape; 2 = forest. Location of roosts and colonies: 1 = caves; 2 = buildings cliff, crevasses; 3 = trees. Population status: 1 = least concern, 2 = near threatened, 3 = vulnerable, 4 = unknown. Population trend: 1 = Decreasing, 2 = stable, 3 = increasing, 4 = unknown.

more intense in southern and eastern Europe than in northern and western Europe, probably due to an opportunistic response to increased abundance.

ACKNOWLEDGEMENTS

We are grateful to the Swiss Ornithological Institute for opening their library to us, and to Will Cresswell and an anonymous referee for useful comments.

SUPPLEMENTAL DATA

A supplementary online table, showing the number of bats identified to species level in the Barn Owl diet in different European countries and islands, can be accessed <http://dx.doi.org/10.1080/09593330.2013.847051>.

REFERENCES

- Alcover, J.A., Sans, A. & Palmer, M.** 1998. The extent of extinctions of mammals in islands. *J. Biogeogr.* **25**: 913–918.
- Alegre, J., Hernández, Á., Purroy, F.J. & Sánchez, A.J.** 1989. Distribucion altitudinal y patrones de afinidad trofica geografica de la lechuza comun (*Tyto alba*) en Leon. *Ardeola* **36**: 41–54.
- Alvizatos, H. & Goutner, V.** 1999. Winter diet of the barn owl (*Tyto alba*) and long-eared owl (*Asio otus*) in northeastern Greece: a comparison. *J. Rapt. Res.* **33**: 160–163.
- Aloise, G., Pelosi, M. & Ronca, M.** 1990. Small mammal communities of the 'Monte Rufeno' natural reserve (Latium, Italy): data from barn owl *Tyto alba* pellets. *Hystrix It. J. Mamm.* **2**: 23–34.
- Altum, B.** 1863. Die Nahrung unserer Eulen. *J. Ornithol.* **11**: 41–46.
- Andera, M. & Vohralík, V.** 1982. The mammals of Broumov Region (Northeastern Bohemia). *Lynx* **21**: 15–39.
- Andera, M., Hanak, V. & Kratka, D.** 1982. Small mammals of the Vihorlat mountains eastern Slovakia Czechoslovakia. *Casopis Narodního Muzea Oddíl Přírodovědný* **151**: 185–198.
- Arnold, A.** 1982. Zur Beute der Schleiereule. *Falke* **29**: 193–196.
- Aulagnier, S.** 2009. Nouvelles données sur le régime alimentaire de l'effraie des clochers, *Tyto alba*, dans les monts de Lacauene et autres localités du département du Tarn (France). *Le Pistrac, Bulletin de l'AROMP* **21**: 44–47.
- Aulagnier, S., Broyer, J., Brunet-Lecompte, P., Coquillart, H., Destre, R. & Erome, G.** 1980. Comparaison de la faune micromammalienne de la Dombes et de la plaine du Forez. *Le Bièvre* **2**: 131–142.
- Balát, F.** 1956. Beitrag zur Ernährung der Schleiereule (*Tyto alba*) in Südmähren und in der Südslowakei. *Folia Zool.* **19**: 237–258.
- Baudvin, H.** 1983. Le régime alimentaire de la chouette effraie. *Le Jean le Blanc* **22**: 1–108.
- Bauer, K.** 1956. Schleiereule (*Tyto alba* Scop.) als Fledermausjäger. *J. Ornithol.* **97**: 335–340.
- Bayle, P., Lasnier, J.-P., Kayser, Y. & Lucchesi, J.-L.** 1996. Le régime alimentaire de l'effraie des clochers *Tyto alba* dans une zone humide des Bouches-du-Rhône, les marais du Vigueirat. *Personnal communication*.
- Bekasinski, R., Kasprzyk, K. & Ruprecht, A.L.** 1996. Chronologiczna analiza pokarmu ptomykowi, *Tyto alba guttata* (C. L. BR.) z Równiny Rychwalskiej (Wielkopolska). *Badania fizjograficzne nad Polską zachodnią Seria C, Zoologia* **43**: 47–54.
- Benes, B.** 1985. Drobní savci v potravě sovy palene (*Tyto alba*) na severní Morave. *Cas. Slez. Muz. Opava (A)* **34**: 145–156.
- Berg, H.-M. & Ille, R.** 2002. Zur Situation der Schleiereule (*Tyto alba* Scop.) in Ostösterreich. *Egretta* **45**: 122–134.
- Bersuder, D. & Kayser, Y.** 1988. La predation des chiropters par la chouette effraie (*Tyto alba*) en Alsace et dans les contrées limitrophes. *Ciconia* **12**: 135–152.
- Bertrand, A.** 1991. Le régime alimentaire des rapaces nocturnes en Ariège. I.- La chouette effraie *Tyto alba*. *Le Pistrac, Bulletin de l'AROMP* **13**: 1–19.
- Bethge, E. & Hayo, L.** 1979. Untersuchungen an einer Population der Schleiereule *Tyto alba* in einem ländlichen Bezirk des westlichen Saarlunds. *Anz. Orn. Ges. Bayern* **18**: 161–170.
- Bielikova, J.** 1995. Spolecenstva drobných savcu v potravě sovy palené. *Diplomova prace, Ped. Fak. UP, Olomouc*.
- Binding, W. & Vauk, G.** 1966. Bemerkungen zur Kleinsäugerfauna des Stadt und Landkreises Goslar nach Untersuchungen an Schleiereulengewöllen. *Beitr. Naturk. Nieders.* **19**: 33–37.
- Blanc, M., Oppliger, J. & Schaller, J.-C.** 2003. Note à propos du régime alimentaire de l'Effraie des clochers *Tyto alba* en region jurassienne, obtenu par l'analyse de pelotes de réjection. *Nos Oiseaux* **50**: 15–20.
- Böhr, H.J.** 1962. Zur Kenntnis der Vogelwelt von Korfu. *Bonn. Zool. Beiträge* **13**: 50–114.
- Boireau, J.** 2009. Problèmes posés par l'effraie des clochers *Tyto alba* dans cinq colonies de reproduction de grand rhinolophe *Rhinolophus ferrumequinum* (Schreber, 1774) en Bretagne occidentale. *Le Rhinolophe* **18**: 43–49.
- Boisseau, J.-P. & Lahargue, J.** 1953. Les pelotes de régurgitation des rapaces nocturnes et la microfaune mammalienne de la région. *Procès-verbaux de la Société Linnéenne de Bordeaux* **95**: 84–90.
- Boldreghini, P., Casini, L. & Santolini, R.** 1982. Dati sulla predazione di *Tyto alba* (Scop.) su micromammiferi nelle Valli Bertuzzi (Delta del Po). *Boll. Zool.* **49**: 23–24.
- Borg, J. & Cachia-Zammit, R.** 1988. Avian, chiropteran and other remains in barn owl *Tyto alba* pellets from Gozo. *Il-Merill* **25**: 12–13.
- Borg, J. & Cachia-Zammit, R.** 1992–1994. Diet of the barn owl *Tyto alba* in a rural area in Gozo. *Il-Merill* **28**: 24–25.
- Bosé, M. & Guidali, F.** 2001. Seasonal and geographic differences in the diet of the barn owl in an agro-ecosystem in northern Italy. *J. Raptor Res.* **35**: 240–246.
- Bovet, J.** 1963. Etude, par l'analyse du contenu de pelotes de chouette effraie (*Tyto alba*), de fluctuations dans les populations de Micromammifères. *Rev. Suisse Zool.* **70**: 244–249.
- Bowman, N.** 1980a. The food of the barn owl (*Tyto alba*) in mid-Wales. *Nat. Wales* **17**: 84–88.
- Bowman, N.** 1980b. The food of barn owls (*Tyto alba*) at a long-used Pembrokeshire site. *Nat. Wales* **17**: 106–108.
- Brown, J.C.** 1971. Mammalian prey of the barn owl (*Tyto alba*) on Skomer Island, Pembrokeshire. *J. Zool.* **165**: 527–530.
- de Bruijn, O.** 1979. Feeding ecology of the barn owl *Tyto alba* in the Netherlands. *Limosa* **52**: 91–154.
- Bryja, J. & Rehak, Z.** 1998. Diet composition of two owl species in the Odra river floodplain (Czech Republic). *Buteo* **10**: 97–102.
- Buckley, J. & Goldsmith, J.G.** 1975. The prey of the barn owl (*Tyto alba*) in East Norfolk. *Mammal Rev.* **5**: 13–16.
- Bunn, D.S., Warburton, A.B. & Wilson, R.D.S.** 1982. *The Barn Owl*. T. & A. D. Poyser, London.
- Buser, M.** 1984. Régime alimentaire de la chouette effraie, *Tyto alba*, durant les périodes de nidification 1978 et 1979, en Ajoie (Jura suisse). *Nos Oiseaux* **37**: 392–394.
- Chaline, J., Baudvin, H., Jammot, D. & Saint Girons, M.-C.** 1974. *Les Proies des Rapaces (petits mammifères et leur environnement)*. Doin, Paris.

- Contoli, L.** 1975a. New data on the role of the barn owl in the control of mammals in central Italy. World Conference on Birds of Prey report of Proceedings, Vienna, 280–282.
- Contoli, L.** 1975b. Micro-mammals and environment in central Italy: data from *Tyto alba* (Scop.) pellets. *Boll. Zool.* **42**: 223–229.
- Contoli, L.** 1976a. Dati circa la predazione operata dal barbagianni (*Tyto alba* (Scop.)) su alcuni mammiferi nei monti della tolfa (Roma). *Supplemento alle Ricerche di Biologia della Selvaggina* **7**: 237–245.
- Contoli, L.** 1976b. Dati circa la predazione operata dal barbagianni (*Tyto alba* (Scop.)) su alcuni mammiferi nei monti della tolfa (Roma). *Ricercatore del CNR Commissione Natura Roma*, 237–245. Laboratorio di Zoologia Applicata alla Caccia, Bologna, Italy.
- Contoli, L.** 1976c. Predazione di *Tyto alba* su micromammiferi e valutazioni sullo stato dell'ambiente. Estratto dal Vol. degli Atti del VI simposio Nazionale sulla Conservazione della Natura, 229–243. Cacucci Editore, Bari, Italy.
- Contoli, L. & Sammuri, G.** 1981. Sui popolamenti di micromammiferi terragnoli della costa medio-tirrenica italiana in rapporto alla predazione operata dal barbagianni. *Accademia Nazionale dei Lincei* **7**: 237–262.
- Contoli, L., Ragonese, B. & Tizi, L.** 1978. Sul sistema trofico 'micromammiferi – *Tyto alba*' nei pantani di vendicari (Noto, Sicilia S-E). *Animalia* **5**: 79–105.
- Cooke, D., Nagle, A., Smiddy, P., Fairley, J. & Muircheartaigh, I.O.** 1996. The diet of the barn owl *Tyto alba* in county Cork in relation to land use. *Biol. Environ.: Proc R Ir Acad* **96B**: 97–111.
- Cserkés, T.** 2007. High relative frequency of *Sicista subtilis* (Dipodidae, Rodentia) in owl-pellets collected in Borsodi Mezőség (NE Hungary). *Folia Historico Naturalia Musei Matraensis* **31**: 173–177.
- Cuisin, J. & Cuisin, M.** 1979. Le régime alimentaire de la chouette effraie (*Tyto alba* (Scopoli)) dans le canton des Riceys (Aube) et ses environs immédiats. *L'Oiseau et la revue française d'ornithologie* **49**: 81–89.
- Dechambre, E.** 1938. Notes sur les petits mammifères de Camargue. *Bull. Soc. Nat. Accl. France* **85**: 464–468.
- Dietz, C., von Helversen, O. & Nill, D.** 2009. *Bats of Britain, Europe & Northwest Africa*. A & C Black Publishers Ltd., London.
- Doumeret, A. & Saint-Girons, M.-C.** 1977. Remarques sur le régime de l'Effraie, *Tyto alba*, en Charente-Maritime. *Ann. Sor. Sci. Nat. Char.-Mar.* **6**: 267–273.
- Draus, B.** 2003. Seasonal variation in the barn owl (*Tyto alba guttata*) diet in the Krakow-Czestochowa upland (South Poland). *Buteo* **13**: 21–30.
- Ekiert, T.** 2005. Sowa plomykowka *Tyto alba guttata* i badania nad nią w powiecie Ostrow Wielkopolski. Available from: <http://www.pwg.otop.org.pl/tyto.php>
- Emma, F. & Endre, F.C.** 2004. Seasonal changes in the diet of barn owl (*Tyto alba*). *Aquila* **111**: 81–87.
- Endes, M. & Harka, A.** 1988. Data concerning the small mammal fauna of the Tissa lowland based on studies of owl pellets. *A. Pusztai* **1**: 155–163.
- Erfurt, J. & Stubbe, M.** 1986. Die Aerele ausgewählter Kleinsäugerarten in der DDR. *Hercynia* **23**: 257–304.
- Fairley, J.S. & Clark, F.L.** 1972. Food of barn owls *Tyto alba* (Scopoli) over one year at a roost in Co. Galway. *Ir. Nat. J.* **17**: 219–222.
- Fairley, J.S. & Deane, C.D.** 1967. Analysis of barn owl pellets from Co. Fermanagh. *Br. Birds* **60**: 370.
- Fairley, J.S. & Smal, C.M.** 1989. Further observations on the diet of the barn owl in Ireland. *Ir. Birds* **4**: 65–68.
- Fehér, C.E.** 1996. The extraordinary great percentage of noctule (*Nyctalus noctula*) in the prey of barn owl (*Tyto alba*). *Hung. Bat Res. News* **2**: 41–42.
- Fenton, M.B.** 1990. The foraging behaviour and ecology of animal-eating bats. *Can. J. Zool.* **68**: 411–422.
- Festetics, A.** 1959. Neuere Angaben zur Ernährung der Schleiereule. *Aquila* **66**: 41–51.
- Forster, R. & Fairley, J.S.** 1974. Further data on barn owls feeding on bank voles, and a new county record of the lesser horse-shoe bat. *Ir. Nat. J.* **18**: 251–252.
- Gaborit-Loret, A.** 2010. Résultats du stage 'pelotes' et biogéographie des micromammifères dans le sud du Tarn et Garonne. *TAIS, Bulletin de liaison des Mammalogistes de Midi-Pyrénées* **4**: 43–57.
- Galan, P., Barros, A., Cerqueira, F. & Seage, R.** 2005. Datos sobre distribución de Quiropteros en el norte de Galicia. *Galemys* **17**: 71–85.
- García, L., Ona, J.A., Salas, G. & Roman, F.J.** Estudio estacional de *Tyto alba* en un medio arido casi antropogeno de Los Alrededores de Almería. Available from: http://www.google.ch/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&ved=0CCIQFjAA&url=http%3A%2F%2Fdiagonalnet.unirioja.es%2Fservlet%2Ffichero_articulo%3Fcodigo%3D2212030&ei=KbJFUMzWHeSp4gTu1oHABw&usq=AFQjCNG6xBw_h04KWJXg5r2SpXXaxK0Kog
- Georgiev, D.** 2010. Mammals (Mammalia) in "Pomorie Lake" Protected Site. Taxonomic Diversity, Distribution, and Status of Species. In Radev, R., Heebaum, G., Michev, T., & Profirov, L. (eds.) *Proceedings of the Integrated Management Plan for Protected Area "Pomorie Lake" BG0000152 and Protected Area "Pomorie" BG0000620*, 145–151. Green Balkans, Plovdiv.
- Gleinich, W. & Hummitzsch, P.** 1977. Zum Brutvorkommen der Eulen im mittleren Oberelbe-Röder-Gebiet. *Faunist. Abh. Staatl. Mus. Tierkunde Dresden* **6**: 237–262.
- Glue, D.E.** 1966. Prey taken by the barn owl in England and Wales. *Bird Study* **13**: 169–183.
- Glue, D.E.** 1971. Avian predator pellet analysis and the mammalogist. *Mammal Rev.* **1**: 53–62.
- Glue, D.E.** 1974. Food of the barn owl in Britain and Ireland. *Bird Study* **21**: 200–210.
- Glue, D.E. & Jordan, R.** 1989. Early 20th century barn owl *Tyto alba* diet in Hampshire. *Hampshire Bird Report* 1988, 79–83. Chandlers Ford, Hampshire.
- Glutz, U.N. & Bauer, K.M.** 1980. *Handbuch der Vögel Mitteleuropas*. Vol. 9, Columbiformes-Piciformes. Wiesbaden, Akademische Verlagsgesellschaft.
- Gonzalez-Kirchner, J.P.** 1992. Algunos datos de la alimentación de la lechuza (*Tyto alba*) y del carabo (*Strix aluco*) en Sierra Morena. *Oxyura* **6**: 33–39.
- Gonzalez Oreja, J.A.** 1990. Sobre la alimentación estival de la lechuza comun (*Tyto alba* Scopoli, 1769), en el Este de Zamora. *Consideraciones ecologicas. Cuad. Invest. Biol. (Bilbao)* **16**: 53–65.
- Goutner, V. & Alivizatos, H.** 2003. Diet of the barn owl (*Tyto alba*) and little owl (*Athene noctua*) in wetlands of northeastern Greece. *Belg. J. Zool.* **133**: 15–22.
- Grogan, R.A. & Whitbread, S.** 1999. The mammalian prey of the barn owl (*Tyto alba* Scopoli) on the Isle of Wight. *Proc. Isle Wight Nat. Hist. Archaeol. Soc.* **15**: 69–76.
- Guérin, G.** 1928. *Régime et croissance de l'Effraie commune en Vendée*. Thèse Faculté des Sciences de Poitiers, Paris.
- Haquart, A., Bayle, P., Cosson, E. & Rombaut, D.** 1997. Chiroptères observés dans les départements des Bouches-du Rhône et du Var. *Faune de Provence (C.E.E.P.)* **18**: 13–32.
- Helsens, B.** 1984. L'analyse des pelotes de réjection des rapaces nocturnes en Mayenne. *Biotope* **53**: 72–76.
- Herrera, C.M.** 1974. Regimen alimenticio de *Tyto alba* en Espana sudoccidental. *Ardeola* **19**: 359–396.
- Hetmanski, T. & Wolk, K.** 2007. Seasonal differences in the diet of barn owl *Tyto alba guttata* in Northern Poland. *Przeegl. Zool.* **3–4**: 169–177.
- Hoekstra, B.** 1974. Over het menu van een kerkuilenfamilie. *De Levende Natuur* **77**: 53–62.

- Hofmann, T.** 1986. Zur Verbreitung von Kleinsäugetern im Kreis Neubrandenburg. *Säugetierkd. Inf.* **2**: 319–335.
- Humphries, M.M., Thomas, D.W. & Speakman, J.R.** 2002. Climate-mediated energetic constraints on the distribution of hibernating mammals. *Nature* **418**: 313–316.
- Hurka, L.** 1981. *Kleinsäuger in der Nahrung der Schleiereule (Tyto alba guttata Brehm) in Westböhmen*. Folia Musei Rerum Naturalium Bohemiae Occidentalis. Zoologica, Vol. 15: 1–19. Západočeské muzeum Plzen.
- Indelicato, N.** 2003. Cartographie des micromammifères dans le Limousin (France) par valeurs d'un indice d'abondance. Available from: <http://perso.wanadoo.fr/nathalie.indelicato/>
- Jäckel, F.** 1867. Noch ein Wort über die Nahrung der Schleiereule. *Zool. Garten* **8**: 463–471.
- Janzekovic, F. & Ficko, M.** 2000. Diet composition of the barn owl *Tyto alba* at Goricko. *Acrocephalus* **21**: 27–29.
- Jentzsch, M.** 1987. Nachweis von Kleinsäugetern in Schleiereulengewöllen aus dem Unstrutgebiet. *Säugetierkd. Inf.* **2**: 467–474.
- Jentzsch, M.** 1992. Fledermäuse als Eulenbeute im Südharz und Helme-Unstrutgebiet. *Nyctalus* **4**: 428–431.
- Kahmann, H. & Brotzler, A.** 1956. Die Ernährung der Schleiereule (*Tyto alba*) und das Bild der Verbreitung kleiner Säugetiere auf der Insel Korsika. *Biol. Zb. Leipzig* **76**: 67–83.
- Kaspar, T. & Borovicka, J.** 1977. Kleine Säugetiere und Vögel in der Nahrung der Schleiereule (*Tyto alba guttata*) in der Umgebung von Valasské Mezirici (Nordmähren). *Zpravy Moravského Ornithologického Sdružení* **19**: 39–47.
- Kasprzyk, K. & Zalewski, A.** 1992. Small mammals of the surrounding of Torun. *Przeegl. Zool.* **36**: 211–217.
- Kasprzyk, K., Kitowski, I., Czochra, K. & Krawczyk, R.** 2004. Bats in the diet of owls from the southern part of the Lublin region (SE Poland). *Myotis* **41–42**: 75–80.
- Kayser, Y., Bayle, P., Chambouleyron, M., Disca, T., Haquart, A. & Olivier, A.** 2009. Les chauves-souris de Camargue; Synthèse et actualisation des données. *Le Vespère* **1**: 36–57.
- Kelleher, K.M., Oliver, G.A. & Sleeman, D.P.** 2010. The composition and sex of rodent prey taken by barn owls *Tyto alba* at a roost in County Cork. *Ir. Birds* **9**: 35–40.
- Kinsky, B.K.** 1942. O Potrave sovy palene (*Tyto alba guttata* Brehm) v okolí kostelce nad orlici. *Sylvia* **7**: 52–55.
- Klaas, C.** 1974. Zur Kleinsäugerbeute dreier Schleiereulenpaare (*Tyto alba*). *Natur. Wis. Museum* **104**: 241–247.
- Klys, G. & Marszalek, T.** 2008. Fauna of small mammals from Mstow (Jurajski Landscape Park) based on pellets of the barn owl, *Tyto alba* Scop. 1796. *Opole Sci. Soc. Nat. J.* **41**: 71–77.
- König, C.** 1961. Schleiereule, *Tyto a. alba* Scop., 'schlägt' fliegende Fledermäuse. *Beitr. Vogelk.* **7**: 229–233.
- Koves, E. & Schmidt, E.** 1964. Angaben zur Kenntnis der Kleinsäugetierfauna in der Umgebung von Tornyosnèmeti / nach Gewöllenuntersuchungen. *Vertebr. Hung.* **6**: 97–108.
- Kowalski, M. & Lesinski, G.** 1986. Small mammal fauna in Janowo (Warsaw voivodship) based on the analysis of barn owl (*Tyto alba* Scop.) pellets. *Przeegl. Zool.* **30**: 327–331.
- Kubic, J., Leniec, H. & Sitowski, W.** 1984. Analysis of the mammal fauna of the Lublin coal Basin based on owl pellets. *Acta Theriol.* **29**: 167–173.
- Kulczycki, A.** 1964. Study on the make up of the diet of owls from the Niski Beskid Mts. *Acta Zool. Cracov.* **9**: 529–559.
- Labes, R. & Köhler, W.** 1984. Beitrag zur Säugetierfauna des Bützow-Güstrower Beckens (Mecklenburg). *Säugetierkd. Inf.* **2**: 167–174.
- Lange, L.** 2002. Die Kleinsäugerfauna eines Marschgebietes der Elbe (Wilstermarsch, Schleswig-Holstein) – ermittelt anhand von Gewöllen der Schleiereule (*Tyto alba*). *Faun-Oekol. Mitt.* **8**: 173–177.
- Latkova, H.** 2008. Seasonal changes in food composition of the barn owl (*Tyto alba*) in the northern part of the 'Zahorie' région. *Slovak Rapt. J.* **2**: 107–112.
- Léger, F.** 1987. Chiroptères dans les pelotes d'Effraie en Lorraine. *Arvicola* **4**: 5.
- Léger, F.** 1995. Notes sur les chiroptères du département de l'Eure-et-Loir. *Soc. Amis Mus. Chartres Nat. Eure-et-Loir Bull.* **15**: 45–55.
- Lesiński, G.** 1989. Bats (Chiroptera) in the food of the barn owl, *Tyto alba* (Scop.) on the Wielun Upland. *Przeegl. Zool.* **33**: 129–135.
- Lesiński, G.** 2010. Long-term changes in abundance of bats as revealed by their frequency in tawny owls' diet. *Biologia* **65**: 749–753.
- Lesiński, G., Kasprzyk, K. & Gryz, J.** 2012. Bats taken by the tawny owl in relation to its roosting site. *North-Western J. Zool.* **8**: 247–251.
- Lopez-Gordo, J.L., Lazaro, E. & Fernandez-Jorge, A.** 1977. Comparacion de las dietas de *Strix aluco*, *Asio otus* y *Tyto alba* en un mismo biotopo de la provincia de Madrid. *Ardeola* **23**: 189–221.
- Love, R.A., Webbon, C., Glue, D.E. & Harris, S.** 2000. Changes in the food of British owls (*Tyto alba*) between 1974 and 1997. *Mammal Rev.* **30**: 107–129.
- Martelli, C.** 1980. Alimentazione del barbagianni *Tyto alba* (Scopoli) nel Parco naturale della Maremma. *Avocetta* **4**: 75–81.
- McGhie, H.** 2001. Diet of barn owls in East Ross and East Ness. *Scot. Birds* **22**: 92–103.
- Mikuska, J. & Vukovic, S.** 1980. Qualitative and quantitative analyse der Nahrung von Schleiereulen, *Tyto alba* Scop. 1769, in dem Gebiet Baranja mit Rückblick auf die Verbreitung von kleinen Säugetieren. *Larus* **31–32**: 269–288.
- Milchev, B., Boev, Z. & Kodjabashev, N.** 2006. Breeding distribution and diet composition of the barn owl *Tyto alba* (Scopoli, 1769), (Aves: Strigiformes) in the North-Western Upper Thracian plain (Bulgaria). *Acta Zool. Bulg.* **58**: 83–92.
- Miltschev, B., Boev, Z. & Georgiev, V.** 2004. Die Nahrung der Schleiereule (*Tyto alba*) in Südost-Bulgarien. *Egretta* **47**: 66–77.
- Moschetti, G., Rocco, M. & Scelba, S.** 1995. Primi dati sull'alimentazione del barbagianni *Tyto alba* in un parco urbano di Portici (NA). *Avocetta* **19**: 118.
- Mraz, L.** 1987. Small mammals in the food of owls on the territory of South Bohemia. *Lynx* **23**: 53–74.
- Nadal, J. & Palaus, X.** 1967. Micromamíferos hallados en egagropilas de *Tyto alba*. *Publ. Inst. Biol. Apl. Barcelona* **42**: 5–15.
- Niethammer, J.** 1964. Ein Beitrag zur Kenntnis der Kleinsäuger Nordspaniens. *Z. Säugetierkunde* **29**: 191–200.
- Nikodem, Z.** 1982. Materials to the bat fauna (Chiroptera) of Lublin region (south-eastern Poland). *Przeegl. Zool.* **26**: 197–205.
- Norberg, U. & Rayner, J.M.V.** 1987. Ecological morphology and flight in bats (Mammalia: Chiroptera): wing adaptations, flight performance, foraging strategy and echolocation. *Phil. Trans. R. Soc. Lond.* **B316**: 335–427.
- Nowosad, A. & Salata-Pilacinska, B.** 1987. Bats (Chiroptera) in the food of the barn owl; *Tyto alba guttata* (C. L. Brehm, 1831). *Przeegl. Zool.* **31**: 221–230.
- Obrist, M.K., Ratthey, E., Bontadina, F., Martinoli, M., Conedera, M., Christe, P. & Moretti, M.** 2011. Response of bat species to silvo-pastoral abandonment. *For. Ecol. Manage.* **261**: 789–798.
- Obuch, J.** 1992. A comparison of diet of four owl species from three regions in CSFR. *Zpravy Moravského Ornithologického Sdružení* **50**: 17–25.
- Obuch, J.** 1998. Zastupenie netopierov (Chiroptera) v potrave sov (Strigiformes) na Slovensku. *Vespertilio* **3**: 65–74.
- Ohlsen, B.** 1975. Ein Beitrag zur Kleinsäugerfauna Nordwestmecklenburgs (Kreise Grevesmühlen, Wismar, Bad Doberan) nach Gewöllfunden und Beobachtungen. *Naturschutzarbeit Mecklenburg* **18**: 34–41.

- Ohlsen, B.** 1976. Ein Beitrag zur Kleinsäugerfauna der Lewitz und des unteren Eldetals (Kreise Schwerin-Land und Ludwigslust) nach Gewölfunden. *Naturschutzarbeit Mecklenburg* **19**: 56–59.
- Pailley, M. & Pailley, P.** 2000. Le régime alimentaire de la Chouette effraie *Tyto a. alba* en Maine-et-Loire. *Crex* **5**: 41–53.
- Pénel, H., Faugier, C. & Faugier, F.** 1984. Synthèse sur les micromammifères sauvages de l'Ardèche. *Bièvre* **6**: 87–116.
- Pérez-Barbería, F.J.** 1991. Influencia de la variación latitudinal en la contribución de los murciélagos (Chiroptera) a la dieta de la lechuza común (*Tyto alba*). *Ardeola* **38**: 61–68.
- Petretti, F.** 1977. Seasonal food habits of the barn owl (*Tyto alba*) in an area of central Italy. *Le Gerfaut* **67**: 225–233.
- Pieper, H.** 1977. Fledermäuse aus Schleiereulen-Gewöllen von der Insel Kreta. *Z. Säugetierkunde* **42**: 7–12.
- Pigeon, J.** 1985. Régime alimentaire de la chouette effraie (*Tyto alba*) dans le département de la Manche. *Le Petit Lérot* **12**: 6–8.
- Prevost, O.** 2000. Cahier techniques: Chauves-souris du Poitou-Charentes. Atlas préliminaire.
- Pribbernow, M.** 1996. Nahrungsökologische Untersuchungen an Schleiereulen (*Tyto alba*, Scopoli 1769) in der Uckermark. Humboldt-Universität zu Berlin, Diplomarbeit.
- Purger, J.J.** 1992. Diet of barn owl, *Tyto alba* (Scop., 1769) using the pellet analysis in Apatin and its surroundings (western Backa, Yugoslavia). *Bull. Nat. His. Mus. Belgr* **47**: 91–99.
- Purger, J.J.** 1996. A Boronka-Melleki tájvedelmi korzet keleti határvidekenek (Somogy Megye) Kismélos faunája, Gyöngybagoly, *Tyto alba* (Scopoli, 1769) kopetek vizsgálata alapján. *Somogy Megyei Múzeumok Igazgatósága* **12**: 299–302.
- Purger, J.J.** 1997. A csokonyavísonyai halastavak (Somogy megye) környékének kismélos faunája, gyöngybagoly koetek vizsgálata alapján. *Természetvedelmi Közlemények* **5–6**: 105–109.
- Purger, J.J.** 1998. Small mammal fauna of the region of Drava river county Somogy (Hungary), obtained by barn owl, *Tyto alba* (Scopoli, 1769) pellet analysis. *Dunantuli Dolg. Term. Tud. Szorozat* **9**: 489–500.
- Purger, J.J.** 2002. Small mammal fauna of the region between Somogyoszob, Hajmas and Kalmancsa based on barn owl *Tyto alba* (Scopoli, 1769) pellet analysis. *Natura Somogyiensis* **3**: 99–110.
- Purger, J.J.** 2004. Small mammal fauna of the region between Varaszlo, Somogyvár, Iharos and Csokoly (county Somogy, Hungary), based on barn owl *Tyto alba* (Scopoli, 1769) pellet analysis. *Somogyi Múzeumok Közleményei* **16**: 409–418.
- Purger, J.J.** 2005. Small mammal fauna of Kaposvár and its surroundings (county Somogy, Hungary), based on barn owl *Tyto alba* (Scopoli, 1769) pellet analysis. *Folia Historico Naturalia Musei Matraensis* **29**: 203–215.
- Rigaux, P. & Riols, C.** 2008. Régime alimentaire de l'Effraie des clochers (*Tyto alba*) à Courpière, Val de Dore (Puy-de-Dôme). *Le Grand Duc* **72**: 27–29.
- Rodriguez, C. & Peris, S.J.** 2007. Habitat associations of small mammals in farmed landscapes: implications for agri-environmental schemes. *Anim. Biol.* **57**: 301–314.
- Rothkopf, D.** 1970. Eine Analyse von Gewöllen der Schleiereule, *Tyto alba*, aus der Eifel. *Bonn. Zool. Beitr.* **21**: 63–82.
- Roulin, A. & Dubey, S.** 2012. The occurrence of reptiles in barn owl diet in Europe. *Bird Study* **59**: 504–508.
- Roulin, A. & Dubey, S.** 2013. Amphibians in the diet of European barn owls. *Bird Study* **60**: 264–269.
- Ruprecht, A.L.** 1964. Analyse der Nahrungsbestandteile der Schleiereule *Tyto alba guttata* (C. L. Br.), vorkommend in Aleksandrow Kuj., Cieshocinek und Raciazek in den Jahren 1960–1961. *Zesz. Nauk. UMK, Biol.* **7**: 45–66.
- Ruprecht, A.L.** 1979. Bats (Chiroptera) as constituents of the food of barn owls *Tyto alba* in Poland. *Ibis* **121**: 489–494.
- Ruprecht, A.L.** 1990. Bats (Chiroptera) in the food of owls in the Nadnotecka Forest. *Przeegl. Zool.* **34**: 349–358.
- Ruprecht, A.L. & Szwagrzak, A.** 1987. Zur Ernährung der Eulen im Westteil des Bialowieza-Urwaldes. *Ökologie der Vögel (Ecol. Birds)* **9**: 89–96.
- Rutz, A.** 2009. Kleinsäuger auf den Inseln Ufnau und Lützelau. Bachelorarbeit Zürcher Hochschule für Angewandte Wissenschaften.
- Rybar, P.** 1969. Die Zweifarbige Fledermaus (*Vespertilio murinus* L.) und andere kleine Wirbeltiere in der Nahrung der Schleiereule (*Tyto alba guttata* Brehm) in Castolovice (Ostböhmen). *Zool. Listy* **18**: 239–246.
- Salata-Pilacinska, B. & Rachowiak, P.** 1990. Badania nad drobnymi ssakami (Micromammalia) Krainy Swietokrzyskiej. *Fragmenta Faunistica* **33**: 308–335.
- Scaravelli, D.** 2000. Un predatore di chiroteri in Barbagia. 1999. Lpogea '99. *Boll. Gruppo Spel. Faenza*, 36–39.
- Scaravelli, D. & Aloise, G.** 1999. La predazione sui chiroteri in Italia. Atti I° Convegno Italiano sui Chiroteri, Castell'Azzara (Grosseto), 28–29 marzo 1998, Grotte di Castro. 223–228.
- Schmidt, E.** 1964. Gyöngybagoly-Köpetvizgalatok eredmenyei. *Aquila* **69–70**: 51–55.
- Schmidt, E.** 1967. Néhány adat a Gyöngybagoly taplalkozás-ökológiájához. *Aquila* **74**: 109–116.
- Schmidt, E.** 1970–1971. Ein Versuch zur Feststellung der Populationsdichte einiger Kleinsäuger in offenen Kulturgebieten auf Grund von Schleiereulengewöllen. *Vertebr. Hung.* **12**: 79–91.
- Schmidt, E.** 1973. Die Nahrung der Schleiereule (*Tyto alba*) in Europa. *Z. Angew. Zool.* **60**: 43–70.
- Schmidt, E.** 1980. Adatok békés megye kismélosfaunájához baglyok taplalkozásvizsgálata alapján. *A Békés Megyei Múzeumok Közleményei* **6**: 179–187.
- Schmidt, E.** 2001. Nachweis der kleinäugigen Wühlmaus *Microtus subterraneus* für die Umgebung von Beeskow, LOS, und Vergleich neuer und alter Beutetierlisten der Schleiereule *Tyto alba* aus Beeskow. *Mitteilungen des LFA Säugetierkunde Brandenburg – Berlin* **9**: 18–24.
- Schmidt, E.** 2005. Nochmals Nachweise der Teichfledermaus (*Myotis dasycneme*) im Beeskower Land, O-Brandenburg, darunter der Fund in einem Gewöll der Schleiereule (*Tyto alba*). *Nyctalus* **9**: 606–610.
- Schmidt, E. & Sipos, G.Y.** 1970–1971. Kleinsäugerfaunistische Angaben aus dem Hornadbecken auf Grund der Gewölluntersuchungen der Schleiereule (*Tyto alba* Scop.). *Tiscia* **6**: 101–108.
- Schmidt, E. & Topal, G.** 1970–1971. Fledermausfunde in Eulengewöllen aus Ungarn. *Vertebr. Hung.* **12**: 93–102.
- Schnurre, O.** 1975. Ernährungsbiologische Studien an Schleiereule (*Tyto alba*) und Waldkauz (*Strix aluco*) im gleichen Lebensraum (Kloster Chorin Kr. Eberswalde). *Milu, Leipzig* **3**: 748–755.
- Sender, T. & Simon, M.** 2003. Population dynamics of the pipistrelle bat: effects of sex, age and winter weather on seasonal survival. *J. Anim. Ecol.* **72**: 308–320.
- Simeonov, S.D.** 1978. Über die Nahrung der Schleiereule (*Tyto alba* Scopoli) in einigen Gegenden Bulgariens. *BAW, Sofia* **4**: 65–71.
- Simeonov, S.D., Michev, T.M. & Simeonov, P.S.** 1981. Materials on the nesting distribution and the diet of the barn owl *Tyto alba* in Bulgaria. *Ekologiya (Sofia)* **8**: 49–54.
- Sommer, R., Zoller, H., Kock, D., Böhme, W. & Griesau, A.** 2005. Feeding of the barn owl, *Tyto alba* with first record of the European free-tailed bat, *Tadarida teniotis* on the island of Ibiza (Spain, Balearics). *Folia Zool.* **54**: 364–370.
- Sommer, R., Niederle, M., Labes, R. & Zoller, H.** 2009. Bat predation by the barn owl *Tyto alba* in a hibernation site of bats. *Folia Zool.* **58**: 98–103.

- Stastny, K.** 1973. Les petits mammifères dans les pelotes de la chouette effraye (*Tyto alba guttata* Brehm) de la Moravie septentrionale. *Lynx* **14**: 54–69.
- Stebbings, R.E.** 1988. *Conservation of European Bats*. Christopher Helm, London.
- Stubbe, M.** 1987. Gewölanalysen zur Untersuchung der Ernährungsbiologie von Eulen. *Populationsökologie Greifvogel – u. Eulenarten* **1**: 429–451.
- Thiollay, J.M.** 1968. Le régime alimentaire de nos rapaces: quelques analyses françaises. *Nos Oiseaux* **24**: 249–267.
- Torre, I., Tella, J.L. & Ballesteros, T.** 1997. Tendencias troficas de la lechuza comun (*Tyto alba*) en la depresion media del Ebro. *Hist. Anim.* **3**: 35–44.
- Torres, J.A., Raya, C., Cademas, R., Linares, N., Pérula, M.L., Rodriguez, J. & Molina, M.** 1982. Analisis de la dieta de *Tyto alba* en un medio arido antropogeno de los Alrededores de Almeria. *Donana, Acta Vert.* **9**: 397–403.
- Uhrin, M., Benda, P., Obuch, J. & Urban, M.** 2002. Mammal fauna of the Direncansky kras Karst Region surrounding areas (central Slovakia). *Lynx* **33**: 193–247.
- Uhrin, M., Benda, P., Obuch, J. & Urban, M.** 2010. Changes in abundance of hibernating bats in central Slovakia (1992–2009). *Biologia* **65**: 349–361.
- Uttendörfer, O.** 1952. *Neue Ergebnisse über die Ernährung der Greifvögel und Eulen*. Eugen Ulmer, Stuttgart/ z.Z. Ludwigsburg, Verlag für Landwirtschaft, Gartenbau und Naturwissenschaften.
- Vargas, J.M. & Antunez, A.** 1982. Sobre *Tyto alba* en la provincia de Malaga (Sur de Espana). *Mon. Trab. Zool (Malaga)* **3–4**: 63–84.
- Vauk, G.** 1963. Nahrungsökologie Untersuchungen an einer Schleiereule der nordwestdeutschen Marschlandschaft. *Beiträge zur Naturkunde Niedersachsens* **16**: 6–9.
- Veiga, J.P.** 1980. Alimentacion y relaciones troficas entre la lechuza comun (*Tyto alba*) y el buho chico (*Asio otus*) en la sierra de Guadarrama (Espana). *Ardeola* **25**: 113–142.
- Vericad, J.R., Escarre, A. & Rodriguez, E.** 1976. Datos sobre la dieta de *Tyto alba* y *Bubo bubo* en Alicante (SE de Iberia). *Mediterranea* **1**: 47–59.
- Vernier, E.** 1994. Predation of bats by barn owl (*Tyto alba*) in Italy. *Hystrix It. J. Mamm.* **5**: 105–107.
- Vicek, M. & Vondracek, J.** 1974. Potrava sovy palené (*Tyto alba guttata*, Brehm) v severozapadnich Cechach. *Biologie* **29**: 649–656.
- Vohralik, V. & Lazarova, J.** 1998. Small mammals (Insectivora, Rodentia) of the Horni Poohri region (Czech Republic) in the food of the barn owl (*Tyto alba*). *Lynx* **29**: 43–56.
- Vohralik, V. & Lazarova, J.** 2002. Small mammals (Insectivora, Rodentia) of the Jicinska pahorkatina and Vychodolabska tabule regions (Czech Republic) in the food of the barn owl (*Tyto alba*). *Lynx* **33**: 249–264.
- Vohralik, V., Hanak, V. & Andera, M.** 1972. Savci Novohradskych hor. *Lynx* **13**: 66–84.
- Von Bülow, B.** 1989. Kleine Säugetiere im Kreis Recklinghausen. Personal communication.
- Von Knorre, D.** 1961. Zur Kleinsäugetierfauna des Spreewaldes und seines südlichen Vorgeländes. *Z. Säugetierkunde* **26**: 183–187.
- Zalman, J.** 1977. Drobní savci v potravě sovy palené (*Tyto alba guttata* Brehm) v okolí Prahy. *Muzeum a Soucasnost* **1**: 125–133.
- Ziomek, J.** 1998. Small mammals (micromammalia) of the Roztocze: part II. Micromammalia of west and central Roztocze on the basis of owl pellets. *Fragmenta Faunistica (Warsaw)* **41**: 125–137.

(MS received 31 May 2013; revised MS accepted 16 September 2013)