

# SCIENTIFIC REPORTS



OPEN

## Sampling frequency affects estimates of annual nitrous oxide fluxes

Received: 20 May 2015

Accepted: 02 October 2015

Published: 02 November 2015

L. Barton<sup>1</sup>, B. Wolf<sup>2</sup>, D. Rowlings<sup>3</sup>, C. Scheer<sup>3</sup>, R. Kiese<sup>2</sup>, P. Grace<sup>3</sup>, K. Stefanova<sup>4</sup> & K. Butterbach-Bahl<sup>2,5</sup>

Quantifying nitrous oxide (N<sub>2</sub>O) fluxes, a potent greenhouse gas, from soils is necessary to improve our knowledge of terrestrial N<sub>2</sub>O losses. Developing universal sampling frequencies for calculating annual N<sub>2</sub>O fluxes is difficult, as fluxes are renowned for their high temporal variability. We demonstrate daily sampling was largely required to achieve annual N<sub>2</sub>O fluxes within 10% of the 'best' estimate for 28 annual datasets collected from three continents—Australia, Europe and Asia. Decreasing the regularity of measurements either under- or overestimated annual N<sub>2</sub>O fluxes, with a maximum overestimation of 935%. Measurement frequency was lowered using a sampling strategy based on environmental factors known to affect temporal variability, but still required sampling more than once a week. Consequently, uncertainty in current global terrestrial N<sub>2</sub>O budgets associated with the upscaling of field-based datasets can be decreased significantly using adequate sampling frequencies.

Scientists have been quantifying soil nitrous oxide (N<sub>2</sub>O) fluxes in the field for at least 60 years<sup>1</sup>. Interest in soil N<sub>2</sub>O fluxes originated from a desire to better understand the fate of soil nitrogen<sup>2,3</sup>. Efforts to measure soil N<sub>2</sub>O fluxes were further increased when N<sub>2</sub>O was recognized as a potent greenhouse gas (GHG) that also plays a significant role in the depletion of stratospheric ozone<sup>4,5</sup>. Quantifying annual soil N<sub>2</sub>O emissions, the dominating source for atmospheric N<sub>2</sub>O, has consequently become a priority for signatory countries to the UN Framework Convention on Climate Change, who are required to present national GHG inventories on an annual basis to the Conference of Parties (<http://unfccc.int>).

Quantifying annual N<sub>2</sub>O fluxes from soils is challenging. Fluxes vary spatially, and differ from day-to-day (and within the day) in response to multiple factors that regulate N<sub>2</sub>O production, consumption and emission<sup>6</sup>. Manual (static) chambers are currently the most widely used technique for quantifying soil N<sub>2</sub>O fluxes. Chamber flux measurements are short-term (e.g., hourly), repeated usually in intervals of days to weeks, and are in turn integrated across time to finally calculate an annual losses. However, fluxes estimated using static chambers can be adversely affected by inadequate mixing of the headspace air, pressure changes, and increases in headspace gas concentration in response to changes to the natural concentration gradient between the soil and atmosphere<sup>7</sup>. Despite these documented short-comings, this approach is favored for N<sub>2</sub>O measurements as fluxes can be orders of magnitude smaller than CO<sub>2</sub> fluxes and the accumulation of gas in the headspace leads to best results with respect to flux detection limit compared to, for instance, dynamic chamber or micrometeorological techniques<sup>8</sup>.

<sup>1</sup>Soil Biology and Molecular Ecology Group, School of Earth & Environment (M087), UWA Institute of Agriculture, Faculty of Sciences, The University of Western Australia, 35 Stirling Highway, Crawley, Western Australia 6009, Australia. <sup>2</sup>Karlsruhe Institute of Technology, Institute for Meteorology and Climate Research Atmospheric Environmental Research (IMK-IFU), Kreuzteckbahnstr. 19, 82467 Garmisch-Partenkirchen, Germany. <sup>3</sup>Institute for Future Environments, Queensland University of Technology, 2 George Street, Brisbane, Queensland 4000, Australia. <sup>4</sup>UWA Institute of Agriculture (M089), The University of Western Australia, 35 Stirling Highway, Crawley, Western Australia 6009, Australia. <sup>5</sup>International Livestock Research Institute (ILRI), Nairobi, Kenya. Correspondence and requests for materials should be addressed to L.B. (email: [louise.barton@uwa.edu.au](mailto:louise.barton@uwa.edu.au))

Furthermore manual chambers are simple to use, relatively inexpensive, and can be deployed in small experimental plots so the effect of multiple treatments on soil N<sub>2</sub>O fluxes can be investigated simultaneously<sup>6</sup>. Temporal coverage is typically limited to weekly, bi-weekly or monthly measurements when using manual chambers<sup>9</sup>. However, manual chambers are likely to underestimate annual N<sub>2</sub>O fluxes if the frequency of measurements does not adequately characterize N<sub>2</sub>O emissions during the year, in particular peak emissions following N fertilizer applications, irrigation, soil re-wetting or spring-thaw events, which may contribute up to 70% of the total annual flux<sup>10–12</sup>.

Guidelines for sampling frequency to estimate annual N<sub>2</sub>O fluxes using manual chambers are not well defined for all land-uses and environments. Instead the approach often taken comes down to an “educated guess” and resource availability<sup>13</sup>. A number of studies have investigated the influence of sampling frequency on cumulative N<sub>2</sub>O fluxes, however with the exception of Liu *et al.*<sup>14</sup> these studies have mainly used short-term N<sub>2</sub>O flux data sets (<1 year), and have been confined to a single study site in an agricultural setting<sup>9,15–18</sup>. These short-term studies have demonstrated that high frequency measurements should coincide with management practices likely to increase N<sub>2</sub>O fluxes, with less frequent measurements during the intervening periods. Yet, the effect of sampling frequency on annual N<sub>2</sub>O flux estimates requires investigation across a broader range of land-uses and climates.

The introduction of automated chambers has enabled researchers to better characterize temporal variation in N<sub>2</sub>O fluxes<sup>6</sup>. Although this technology is expensive and not available to all researchers, it does provide a unique opportunity to better assess how sampling frequency affects estimates of annual N<sub>2</sub>O fluxes. Such an analysis is particularly beneficial to those new to measuring *in situ* N<sub>2</sub>O fluxes from land and to those planning to investigate a previously unstudied land-use. Consequently, the objective of the following study was to investigate the effect of sampling frequency on estimates of annual soil N<sub>2</sub>O fluxes using 28 published datasets of subdaily N<sub>2</sub>O fluxes from a variety of different terrestrial ecosystems across the globe.

## Results

Annual N<sub>2</sub>O fluxes calculated from the average daily fluxes, which is used here as the reference annual flux, varied from 0.03 kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup> to 8.1 kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup> (Table 1). The smallest annual flux was recorded for a sandy soil cropped to a grain legume in a semiarid environment<sup>11</sup>, while the greatest was from a loam soil cropped to tree fruit in a subtropical climate<sup>19</sup>. Daily N<sub>2</sub>O fluxes were highly variable within each dataset, but more so for some; the coefficient of variation (CV) of the mean daily N<sub>2</sub>O flux ranged from 78% for a subtropical rainforest to 913% for a semiarid soil planted to a grain legume. The variation in daily means was not related to the magnitude of the annual N<sub>2</sub>O flux (Table 1), but instead reflected the episodic nature of the daily fluxes for a particular study site (Fig. 1). We subsequently classified the data sets as having either moderate (CV > 50–100%), high (CV > 100–200%) or extreme (CV > 200%) ‘episodicity’ based on the CV of the mean daily flux (Table 2).

Increasing the interval between sampling days increased the variance in the estimated annual N<sub>2</sub>O fluxes, and hence decreased the accuracy of the estimate (Fig. 1). As the sampling frequency decreased, the deviation from the ‘best estimate’, or expected value obtained using all daily fluxes, increased and caused annual losses to be either over- or underestimated (Fig. 2; Supplementary Table 1). Across all sites and sampling frequencies (n = 1568), 22% and 58% of annual emission values were more than 10% higher or lower, respectively, than the ‘best estimate’ annual flux. The extent that decreased sampling frequency increased the deviation from the reference annual N<sub>2</sub>O flux appeared to be largely related to the variability, or coefficient of variation, of the daily fluxes (Fig. 3). The greater the variation in daily N<sub>2</sub>O within a dataset, the greater the impact of decreasing the sampling frequency had on the accuracy of the estimated annual flux. For example, for a tropical rainforest (Bellenden Kerr) with a daily N<sub>2</sub>O flux CV of 98%, sampling every 28 days resulted in an annual N<sub>2</sub>O flux that was up to 1.2 times greater than the best estimate; whereas for cropped soil in a semiarid region with a daily N<sub>2</sub>O flux CV of 913%, sampling every 28 days resulted in an annual N<sub>2</sub>O flux that was up to 12 times greater than the best estimate (Fig. 1; Supplementary Table 1).

The minimum sampling frequency required to robustly estimate an annual N<sub>2</sub>O flux varied depending upon the ‘episodicity’ of the dataset and the required accuracy (Fig. 2; Table 2; Supplementary Table 1). Twenty, or 74%, of the datasets required daily sampling to achieve an annual N<sub>2</sub>O flux value within 10% of the best estimate (Fig. 4). In only one case (tropical rainforest, Bellenden Ker), and when the daily N<sub>2</sub>O flux CV was relatively low (98%), did weekly sampling result in annual N<sub>2</sub>O flux within 10% of the best estimate. Generally speaking, highly or extremely episodic data sets (CV > 100%) required sampling either daily or 3 days a week (Table 2). Lowering the desired accuracy decreased the required frequency of sampling, however 89% of the data sets still needed to be sampled at least weekly to achieve ±30% accuracy (Fig. 4). Lowering the level of accuracy to ±40% meant two datasets could be sampled once every 4 weeks (Fig. 4).

## Discussion

Nitrous oxide emissions need to be measured daily to accurately determine annual N<sub>2</sub>O flux in environments where data has not previously been collated. Measuring N<sub>2</sub>O fluxes on a daily basis ensured that annual N<sub>2</sub>O fluxes were estimated within 10% of the expected value for all datasets in the present study. Although a similar result could be achieved in some instances (25%, or 7 datasets) by sampling 3 days


Location <sup>†</sup> , Year	Climate	Rainfall <sup>‡</sup> (mm yr <sup>-1</sup> )	Soil C <sup>§</sup> (g kg <sup>-1</sup> )	Soil texture <sup>§</sup>	Land use	Annual datasets	Study period (days)	Annual flux (kg N <sub>2</sub> O-N ha <sup>-1</sup> yr <sup>-1</sup> )	Daily flux CV (%)	Reference
Wongan Hills, Australia. 2009–2011	Semiarid	374	10	Sand	Grain crop, rainfed, +/- lime, +/- N fertilizer	8	364–371	0.03–0.07	380–913	Barton <i>et al.</i> <sup>11</sup>
Cunderdin, Australia. 2005–2009	Semiarid	368	9.8	Sand	Grain crop, rainfed, +/- N fertilizer	8	337–379	0.08–0.16	173–428	Barton <i>et al.</i> <sup>10</sup> Li <i>et al.</i> <sup>26</sup> Barton <i>et al.</i> <sup>27</sup> Barton <i>et al.</i> <sup>28</sup>
Xilin, Inner Mongolia. 2007–2008	Semiarid, cool temperate	335	26	Sandy loam	Steppe grassland, not grazed	1	365	0.21	260	Wolf <i>et al.</i> <sup>12</sup>
Höglwald, Germany. 1996–1997	Temperate	850	22	Silty clay	Spruce and beech forest (plantation)	2	365	0.58–2.46	169–179	Papen & Butterbach-Bahl (1999) <sup>29</sup> Wu <i>et al.</i> <sup>30</sup>
Kingsthorpe, Australia. 2009–2010	Subtropical	630	15	Clay	Wheat-cotton crop, irrigated, N fertilizer	3	334	2.61–2.93	181–235	Scheer <i>et al.</i> <sup>31</sup> Scheer <i>et al.</i> <sup>32</sup>
Mooloolah Valley, Australia. 2007–2009	Subtropical	1709	28	Loam	Pasture, mowed, not grazed	2	365	1.16–2.12	155–172	Rowlings <i>et al.</i> <sup>33</sup>
Mooloolah Valley, Australia. 2008–2009	Subtropical	1709	35	Loam, silt loam	Rainforest (notophyll vine)	1	365	0.48	78	Rowlings <i>et al.</i> <sup>34</sup>
Mooloolah Valley, Australia. 2007–2009	Subtropical	1709	27	Loam	Tree crop (lychee)	2	365	1.68–8.12	93	Rowlings <i>et al.</i> <sup>19</sup>
Bellenden Ker, Australia. 2001–2002	Tropical	4360	31	Sandy loam	Rainforest (mesophyll vine)	1	365	1.16	98	Kiese <i>et al.</i> <sup>23</sup>

**Table 1. Summary of data sets used to assess the effect of sampling frequency on estimated annual N<sub>2</sub>O fluxes.** <sup>†</sup>Cunderdin, 31°36' S, 117°13' E; Wongan Hills, 30°89' S, 116°72' E; Höglwald 48°30' N, 11°10' E; Xilin 43° 33' N, 116° 42.3' E; Bellenden Ker, 17°16' S, 145°54' E; Kingsthorpe, 27°30' S, 151°46' E; Mooloolah Valley 26°75' S, 152°93' E <sup>‡</sup>Long-term average value <sup>§</sup>Surface soil (e.g., 0–15cm).

a week, this still represents a highly regularly sampling regime. Our findings are consistent with others who have investigated the effects of sampling frequency on estimates of annual N<sub>2</sub>O fluxes. For example, a relatively frequent sampling regime (once every 2 to 3 days) was required to estimate cumulative losses within 10% of the expected cumulative loss from N-fertilized crops in China and the United States of America<sup>9,14</sup>. Our findings further confirm the importance of deploying automated chamber systems when determining annual N<sub>2</sub>O fluxes in previously unstudied environments, and when the drivers of temporal variability are not well understood.

The frequency of sampling required to accurately calculate an annual N<sub>2</sub>O flux will depend on the episodic nature of the N<sub>2</sub>O flux at the study site of interest, rather than the magnitude of the annual flux. This was particularly well demonstrated by studies conducted in semiarid environments of Australia and Inner Mongolia, where relatively low annual N<sub>2</sub>O losses ( $\leq 0.21$  kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup>) resulted from a limited number of elevated daily N<sub>2</sub>O fluxes during the year<sup>11,12</sup>. For example in a cropped soil in south-western Australia, 75 to 85% of the annual fluxes were attributed to isolated, short-lived summer rainfall events<sup>11</sup>. Understanding the underlying temporal variability of daily N<sub>2</sub>O fluxes is therefore likely to improve the efficacy of sampling regimes.

Sampling efficacy for determining annual N<sub>2</sub>O fluxes may be improved, and the regularity of sampling decreased, if N<sub>2</sub>O flux responses can be anticipated. This may occur if previous research has been conducted in a similar environment, or if preliminary work is undertaken to assess the temporal variability of N<sub>2</sub>O fluxes. In either case, refining the sampling regime will require some underlying understanding of temporal variation in the N<sub>2</sub>O flux and its regulation. Using this approach, we estimated annual N<sub>2</sub>O fluxes for three of our datasets (Fig. 1) based on the authors' informed understanding of the factors driving daily losses. While we found the annual N<sub>2</sub>O fluxes estimated by the authors' did not vary statistically from the 'best' estimate calculated using all daily fluxes, the informed sampling approach still required sampling to occur every 2 to 6 days depending on the dataset (Table 3). A number of short-term studies




**Figure 1.** Daily  $N_2O$  fluxes and the influence of sampling frequency on annual  $N_2O$  fluxes. The daily  $N_2O$  flux (a,c,e) for the three data sets shown have varying coefficients of variation (CV), which influences the effect of sampling frequency on annual  $N_2O$  fluxes (b,d,f). See Table 4 for description of sampling intervals.

have also devised strategies for characterizing  $N_2O$  fluxes in response to N inputs to cropping and grazed soils setting<sup>9,15–18</sup>. Notably, various authors used automated chambers to develop a sampling regime for measuring  $N_2O$  emissions from temperate grasslands in response to ruminant urine deposition<sup>16</sup> in New Zealand, N fertilized potato fields in Europe<sup>18</sup>, and rainfed cereal crops in subtropical Australia<sup>17</sup>; agricultural land use not captured in the present study. Interestingly, some of these authors recommended weekly sampling (with a higher frequency following anticipated  $N_2O$  events), which is less frequent than our analysis would recommend for agricultural and non-agricultural study sites in the present study.

The uncertainty of current global  $N_2O$  estimates maybe partly attributed to the sampling frequency of the datasets selected for inclusion in the analysis. Modelling of global soil  $N_2O$  emissions has been largely derived from manual chambers measurements covering more than 300 days in a year<sup>20</sup>. However, less than a third of the 464 studies included in the metadata analysis by Stehfest and Bouwman<sup>20</sup> measured  $N_2O$  on at least a daily basis, with 50% of the data used collected 3 days a week, or less than weekly. Given the influence of sampling frequency on annual  $N_2O$  fluxes in the present paper, it is likely that current global  $N_2O$  values have not been accurately captured. Instead, we recommend that revision of global estimates using high frequency measurements (at least daily) or an ‘informed’ sampling approach for at least a year.

Location	Episodicity <sup>†</sup>	Minimum sampling frequency	
		10% accuracy	0% accuracy
Wongan Hills, Australia.	Extreme	Daily	Daily to 3 days a week
Cunderdin, Australia.	High to Extreme	Daily or 3 days a week	Daily to weekly
Xilin, Inner Mongolia.	Extreme	3 days a week	Weekly
Höglwald, Germany.	High	3 days a week	Weekly to bi-weekly
Kingsthorpe, Australia.	High to Extreme	Daily	Daily
Mooloolah Valley, (Pasture) Australia.	High	Daily to 3 days a week	3 days a week to weekly
Mooloolah Valley, (Rainforest) Australia.	Moderate	3 days a week	Bi-weekly
Mooloolah Valley, (Tree crop) Australia.	Moderate	3 days a week	Weekly
Bellenden Ker, Australia.	Moderate	Weekly	Bi-weekly

**Table 2. The relationship between the ‘episodicity’ of each study location and the minimum sampling frequency needed to meet a given accuracy.** <sup>†</sup>Episodicity determined using coefficient of the mean daily flux (Table 1). Moderate, CV > 50–100%; high, CV > 100–200%; extreme, CV > 200%.




**Figure 2. The proportion (%) of the ‘best estimate’ annual N<sub>2</sub>O flux estimated by each sampling frequency.** For each dataset (28), the average annual flux estimate (calculated from replicate chambers) for each sampling frequency (and each permutation, Table 4) was compared to the ‘best estimate’ flux calculated from the average daily fluxes (expressed as a %). The ‘best estimate’ was calculated using all daily fluxes. For each sampling frequency, the datasets are presented in the same order (from left to right in the above Figure) as that listed in Supplementary Table 1. Specific values for each dataset are listed in Supplementary Table 1.

Finally, we recommend data from automated chambers should be continuously used to build on existing guidelines for use of manual chambers<sup>21</sup>. While the present study included and discussed a large number of datasets from a variety of climates, soils and land uses, there were a number of environments not represented. For example, grazed soils outside temperate climate, a broader range of horticultural soils, and non-agricultural soils in semiarid environments. We therefore encourage researchers utilizing automated chamber systems to determine annual N<sub>2</sub>O fluxes from soils, to in turn also utilize the data to investigate the impacts of sampling frequency on these losses.


## Methods

**Study sites.** The meta-analysis included datasets from published research studies, and where N<sub>2</sub>O fluxes had been measured on a subdaily basis for approximately one year using automated chambers. Annual data sets were sourced from measurements in Australia, Germany, and Inner Mongolia, representing a variety of climates, soil types and land uses (Table 1). Climates ranged from semiarid (including a Mediterranean-type climate) to tropical, soil textures varied from sands to heavy clays, while land use included various agricultural production and forest systems. A number of study sites also included different treatments (Table 1). Consequently our meta-analysis included 28 sub-daily N<sub>2</sub>O datasets.

**Automated chamber system.** Nitrous oxide fluxes were measured at each study location using soil chambers connected to a fully automated system that enabled *in situ* determination of N<sub>2</sub>O fluxes.



**Figure 3.** Relationship between the coefficient of variation of the daily  $N_2O$  flux and the deviation (range) from the ‘best estimate’ annual  $N_2O$  flux. For each dataset (28; represented as single point in the above Figure), the range in deviation was determined after comparing the annual  $N_2O$  fluxes calculated from a sample interval of 4-weekly (every 28 days) with the ‘best estimate’ for each permutation (Table 4). The ‘best estimate’ was calculated using all daily fluxes. Specific values for each dataset are listed in Supplementary Table 1.



**Figure 4.** Impact of sampling interval on estimating annual  $N_2O$  fluxes at a given accuracy. The number of datasets obtaining annual  $N_2O$  fluxes at a given accuracy are listed as a function of sampling interval. Specific values for each datasets are listed in Supplementary Table 1.

Location <sup>†</sup>	Best estimate	Informed sampling regime <sup>‡</sup>	
	Annual flux (kg $N_2O-N$ ha <sup>-1</sup> yr <sup>-1</sup> )	Annual flux (kg $N_2O-N$ ha <sup>-1</sup> yr <sup>-1</sup> )	No. measurements
Wongan Hills, Australia.	0.04 (0.0) <sup>a</sup>	0.03 (0.0) <sup>a</sup>	60
Höglwald, Germany.	0.58 (0.1) <sup>a</sup>	0.64 (0.1) <sup>a</sup>	83
Bellenden Ker, Australia.	1.16 (0.1) <sup>a</sup>	1.35 (0.2) <sup>a</sup>	156

**Table 3.** Annual  $N_2O$  fluxes for three contrasting study sites estimated using either an informed sampling strategy or from daily measurements. Values represent means (and standard errors) of three to five replicates depending on the study site. Values followed by the same letter in the same row are not statistically different ( $P < 0.05$ ). <sup>†</sup>For further details see Table 1. Corresponds to data set numbers 2 (Wongan Hills), 19 (Höglwald) and 28 (Bellenden Ker) in Supplementary Table 1. <sup>‡</sup>Informed sampling strategy for each location: Wongan Hills,  $N_2O$  fluxes measured daily for five consecutive days when daily rainfall  $> 5$  mm rainfall (December–June) or  $> 10$  mm rainfall (July–November); Höglwald,  $N_2O$  fluxes measured daily for seven days consecutive day when air temperature between  $-0.7$  and  $0.7^\circ C$  or if daily rainfall  $> 15$  mm; Bellenden Ker,  $N_2O$  fluxes measured daily for six days consecutive day when daily rainfall  $> 15$  mm.

Sampling frequency	Sampling interval (days)	Permutations	Examples
Daily	1	1	Sunday
			Monday
			Tuesday <i>etc</i>
3 days a week	2	7	Sunday-Tuesday-Thursday
			Monday-Wednesday-Friday
			Tuesday-Thursday-Saturday <i>etc</i>
1 day a week (weekly)	7	7	Every Sunday
			Every Monday
			Every Tuesday <i>etc</i>
1 day every 2 weeks (bi-weekly)	14	14	Every Sunday (week 1 of 2)
			Every Sunday (week 2 of 2)
			Every Monday (week 1 of 2)
			Every Monday (week 2 of 2) <i>etc</i>
1 day every 4 weeks (4-weekly)	28	28	Every Sunday (week 1 of 4)
			Every Sunday (week 2 of 4)
			Every Sunday (week 3 of 4)
			Every Sunday (week 4 of 4) <i>etc</i>

**Table 4.** Description of sampling frequencies.

Details of the design and operation of the automated gas sampling systems have been described by Breuer *et al.*<sup>22</sup> and Kiese *et al.*<sup>23</sup>. Briefly, the various systems consisted of a gas chromatograph (e.g., Texas Instruments, SRI 8610C) equipped with an electron capture detector (ECD) for N<sub>2</sub>O analysis, an automated sampling unit for collecting and distributing gas samples, and a series of chambers (three to five replicates depending on the study site). Chambers (0.5 m × 0.5 m or 0.7 m × 0.7 m) were placed on metal bases inserted into the ground (0.05–0.1 m), and fitted with a top (0.15 m or 0.3 m in height) that could be automatically opened and closed by means of pneumatic actuators. The height of the chambers was progressively increased to accommodate crop growth at some study sites, with a maximum height of 0.95 m. Furthermore, in some instances the chambers were programmed to open if the air temperature in the chamber exceeded a set value, or if rain fell while the chambers were closed. The automated gas sampling unit enabled N<sub>2</sub>O to be monitored continuously, providing up to eight (hourly) emission rates per day. Specific N<sub>2</sub>O measurement details for each study site are described in the associated published papers (Table 1).

**Evaluating sample frequency effects.** The effect of sampling frequency on estimates of annual N<sub>2</sub>O-N fluxes was assessed using a modified jackknife technique<sup>24,25</sup>. Average daily flux measurements were calculated for each replicate chamber in each dataset from the sub-daily flux measurements as we did not consistently observe diurnal flux variations at each location. Each site's daily flux population was subsequently subsampled daily, three times per week, weekly, bi-weekly and 4-weekly, and for each permutation of the time interval, for each dataset (Table 4). There were 7 to 28 jackknifed populations depending on the sampling frequency (Table 4). Estimates of annual N<sub>2</sub>O-N flux for a given chamber, site and frequency permutation were then calculated by linear interpolation and integration of daily fluxes with time. Missing daily N<sub>2</sub>O flux data was not replaced. The average annual flux estimate (calculated from replicate chambers) from each sampling frequency, and for each of the dataset, was then compared to the 'best estimate' annual flux calculated from the average daily fluxes (expressed as a %) so as to assess the accuracy of each of the sampling frequencies. An annual flux determined using an informed sampling regime (based on the authors' understanding of the factors driving daily N<sub>2</sub>O fluxes) was also compared with the 'best estimate' annual flux using a general analysis of variance (Genstat for Windows, 14<sup>th</sup> Edition, VSN International).

## References

1. Arnold, P. Losses of nitrous oxide from soil *Journal of Soil Science* **5**, 116–128 (1954).
2. Denmead, O. Chamber systems for measuring nitrous oxide emission from soils in the field. *Soil Sc. Soc. Am. J.* **43**, 89–95 (1979).
3. Hutchinson, G. & Mosier, A. Improved Soil cover method for field measurement of nitrous oxide fluxes. *Soil Sc. Soc. Am. J.* **45**, 311–316 (1981).
4. Ravishankara, A., Daniel, J. & Portmann, R. Nitrous oxide (N<sub>2</sub>O): the dominant ozone-depleting substance emitted in the 21st century. *Science* **326**, 123–125 (2009).

5. Crutzen, P. J. in *Denitrification, nitrification, and atmospheric nitrous oxide* (ed C C Delwiche) 17–44 (John Wiley & Sons, 1981).
6. Butterbach-Bahl, K., Baggs, E., Dannenmann, M., Kiese, R. & Zechmeister-Boltenstern, S. Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Phil Trans R Soc B* **368**, 20130122 (2013).
7. Pihlatie, M. *et al.* Comparison of static chambers to measure CH<sub>4</sub> emissions from soils. *Agr. Forest. Meteorol.* **171–172**, 124–136 (2013).
8. Denmead, O. T. Approaches to measuring fluxes of methane and nitrous oxide between landscapes and the atmosphere. *Plant Soil* **309**, 5–24 (2008).
9. Parkin, T. B. Effect of sampling frequency on estimates of cumulative nitrous oxide emissions. *J. Environ. Qual.* **37**, 1390–1395 (2008).
10. Barton, L. *et al.* Nitrous oxide emissions from a cropped soil in a semiarid climate. *Glob. Change Biol.* **14**, 177–192 (2008).
11. Barton, L., Murphy, D. & Butterbach-Bahl, K. Influence of crop rotation and liming on greenhouse gas emissions from a semiarid soil. *Agric. Ecosyst. Environ.* **167**, 23–32 (2013).
12. Wolf, B. *et al.* Grazing-induced reduction of natural nitrous oxide release from continental steppe. *Nature* **464**, 881–884 (2010).
13. Morris, S., Kimber, S., Grace, P. & Van Zwielen, L. Improving the statistical preparation for measuring soil N<sub>2</sub>O flux by closed chamber. *Sci. Total Environ.* **465**, 166–172 (2013).
14. Liu, C. *et al.* Nitrous oxide and nitric oxide emissions from an irrigated cotton field in Northern China. *Plant Soil* **332**, 123–134 (2010).
15. Smith, K. A. & Dobbie, K. E. The impact of sampling frequency and sampling times on chamber-based measurements of N<sub>2</sub>O emissions from fertilized soils. *Global Change Biol.* **7**, 933–945 (2001).
16. van der Weeden, T., Clough, T. & Styles, T. Using near-continuous measurements of N<sub>2</sub>O emission from urine-affected soil to guide manual gas sampling regimes. *New Zeal. J. Agr. Res.* **56**, 60–76 (2013).
17. Reeves, S. & Wang, W. Optimum sampling time and frequency for measuring N<sub>2</sub>O emissions from a rain-fed cereal cropping system. *Science of The Total Environment* **530–531**, 219–226, doi: <http://dx.doi.org/10.1016/j.scitotenv.2015.05.117> (2015).
18. Flessa, H. *et al.* N<sub>2</sub>O and CH<sub>4</sub> fluxes in potato fields: automated measurement, management effects and temporal variation. *Geoderma* **105**, 307–325 (2002).
19. Rowlings, D., Grace, P., Scheer, C. & Kiese, R. Influence of nitrogen fertiliser application and timing on greenhouse gas emissions from a lychee (*Litchi chinensis*) orchard in humid subtropical Australia. *Agric. Ecosyst. Environ.* **179**, 168–178 (2013).
20. Stehfest, E. & Bouwman, L. N<sub>2</sub>O and NO emission from agricultural fields and soils under natural vegetation: summarizing available measurement data and modeling of global annual emissions. *Nutr. Cycl. Agroecosystem* **74**, 207–228 (2006).
21. de Klein, C. & Harvey, M. *Nitrous oxide chamber methodology guidelines*. (Ministry for Primary Industries, 2012).
22. Breuer, L., Papen, H. & Butterbach-Bahl, K. N<sub>2</sub>O emission from tropical forest soils of Australia. *J. Geophys. Res.* **105**, 26, 353–326, 367 (2000).
23. Kiese, R., Hewett, B., Graham, A. & Butterbach-Bahl, K. Seasonal variability of N<sub>2</sub>O emissions and CH<sub>4</sub> uptake by tropical rainforest soils of Queensland, Australia. *Global Biogeochem. Cycles* **17**, 1043 (2003).
24. Efron, B. Computers and the theory of statistics: Thinking the unthinkable. *SIAM Review* **21**, 460–480 (1979).
25. Efron, B. & Gong, G. A leisurely look at the bootstrap, the jackknife, and cross-validation. *Am. Stat.* **37**, 36–48 (1983).
26. Li, Y., Barton, L. & Chen, D. Simulating response of N<sub>2</sub>O emissions to fertiliser N application and climatic variability from a rain-fed and wheat-cropped soil in Western Australia. *J. Sci. Food Agric.* **92**, 1130–1143 (2012).
27. Barton, L., Murphy, D. V., Kiese, R. & Butterbach-Bahl, K. Soil nitrous oxide and methane fluxes are low from a bioenergy crop (canola) grown in a semiarid climate. *Glob. Change Biol. Bioenergy* **2**, 1–15 (2010).
28. Barton, L., Butterbach-Bahl, K., Kiese, R. & Murphy, D. Nitrous oxide fluxes from a grain-legume crop (narrow-leaved lupin) grown in a semiarid climate. *Glob. Change Biol.* **17**, 1153–1166 (2011).
29. Papen, H. & Butterbach-Bahl, K. A 3-year continuous record of nitrogen trace gas fluxes from untreated and limed soil of a N-saturated spruce and beech forest ecosystem in Germany: 1. N<sub>2</sub>O emissions. *J. Geophys. Res.* **104**, 18, 487–418, 503 (1999).
30. Wu, X. *et al.* Environmental controls over soil-atmosphere exchange of N<sub>2</sub>O, NO and CO<sub>2</sub> in a temperate Norway spruce forest. *Global Biogeochem. Cycles* **24**, 773–787 (2010).
31. Scheer, C., Grace, P., Rowlings, D. & Payero, J. Nitrous oxide emissions from irrigated wheat in Australia: impact of irrigation management. *Plant Soil* **359**, 351–362 (2012).
32. Scheer, C., Grace, P., Rowlings, D. & Payero, J. Soil N<sub>2</sub>O and CO<sub>2</sub> emissions from cotton in Australia under varying irrigation management. *Nutr. Cycl. Agroecosystem* **95**, 43–56 (2013).
33. Rowlings, D., Grace, P., Scheer, C. & Liu, S. Rainfall variability drives interannual variation in N<sub>2</sub>O emissions from a humid, subtropical pasture. *Sci. Total Environ.* **512–513**, 8–18 (2015).
34. Rowlings, D., Grace, P., Kiese, R. & Weier, K. Environmental factors controlling temporal and spatial variability in the soil-atmosphere exchange of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O from an Australian subtropical rainforest. *Glob. Change Biol.* **18**, 726–738 (2012).

## Acknowledgements

The research was funded by the Australian Government's Climate Change Research Program, Australian Research Council (DP0559791), Department of Agriculture & Food Western Australia, Grains & Research Development Corporation (GRDC), EU Integrating Activity (IA) project InGOS, the Climate Change, Agriculture and Food Security (CCFAS) of CGIAR institutes and the German Science Foundation.

## Author Contributions

L.B. had the main responsibility for the design and writing of the manuscript together with B.W. All co-authors participated, commented and provided contributions relevant to their own data and area of expertise.

## Additional Information

**Supplementary information** accompanies this paper at <http://www.nature.com/srep>

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Barton, L. *et al.* Sampling frequency affects estimates of annual nitrous oxide fluxes. *Sci. Rep.* **5**, 15912; doi: 10.1038/srep15912 (2015).





This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>