

RESEARCH

Open Access



The convergence analysis and error estimation for unique solution of a p -Laplacian fractional differential equation with singular decreasing nonlinearity

Jing Wu¹, Xinguang Zhang^{2,3*} , Lishan Liu^{4,3}, Yonghong Wu³ and Yujun Cui⁵

*Correspondence:

zxg123242@163.com

²School of Mathematical and Informational Sciences, Yantai University, Yantai, China³Department of Mathematics and Statistics, Curtin University of Technology, Perth, Australia
Full list of author information is available at the end of the article

Abstract

In this paper, we focus on the convergence analysis and error estimation for the unique solution of a p -Laplacian fractional differential equation with singular decreasing nonlinearity. By introducing a double iterative technique, in the case of the nonlinearity with singularity at time and space variables, the unique positive solution to the problem is established. Then, from the developed iterative technique, the sequences converging uniformly to the unique solution are formulated, and the estimates of the error and the convergence rate are derived.

Keywords: Convergence analysis; Error estimation; p -Laplacian fractional differential equation; Double iterative technique; Riemann–Stieltjes integral conditions

1 Introduction

This paper is motivated by the following singular nonlocal fractional differential equation:

$$\begin{cases} -\mathcal{D}_x^{\frac{3}{2}}(\varphi_{\frac{3}{2}}(-\mathcal{D}_x^{\frac{4}{3}}z))(x) = x^{-1}z^{-\frac{1}{2}}, & 0 < x < 1, \\ z(0) = 0, \quad \mathcal{D}_x^{\frac{4}{3}}z(0) = \mathcal{D}_x^{\frac{4}{3}}z(1) = 0, & z(1) = \int_0^1 z(x) d\chi(x), \end{cases} \quad (1.1)$$

where χ is a function of bounded variation satisfying $\chi(x) = 0$, $x \in [0, \frac{1}{3})$, $\chi(x) = \frac{1}{2}$, $x \in [\frac{1}{3}, \frac{2}{3})$, $\chi(x) = 1$, $x \in [\frac{2}{3}, 1]$, which exhibits a blow-up behaviour at $x = 0$ and $z = 0$. These types of singular behaviours [1–11] as well as impulsive phenomena [12–21] often exhibit some blow-up properties [22, 23] which occur in many complex physical processes, for example, in mechanics process [1], the stress near the crack tip in elastic fracture exhibits a singularity of $r^{-0.5}$, where r is the distance measured from the crack tip.

Inspired by the above problem, this paper presents the convergence analysis and error estimation for the unique solution of the general fractional differential equation with singular decreasing nonlinearity and a p -Laplacian operator

$$\begin{cases} -\mathcal{D}_x^\alpha(\varphi_p(-\mathcal{D}_x^\gamma z))(x) = f(x, z(x)), & 0 < x < 1, \\ z(0) = 0, \quad \mathcal{D}_x^\gamma z(0) = \mathcal{D}_x^\gamma z(1) = 0, & z(1) = \int_0^1 z(x) d\chi(x), \end{cases} \quad (1.2)$$

where $\mathcal{D}_x^\alpha, \mathcal{D}_x^\gamma$ are the standard Riemann–Liouville derivatives with $\gamma, \alpha \in (1, 2]$, $\int_0^1 z(x) d\chi(x)$ is a Riemann–Stieltjes integral and χ is a function of bounded variation, $\varphi_p(x) = |x|^{p-2}x$, $p > 1$ is the p -Laplacian operator with conjugate index $q > 1$ satisfying $\frac{1}{p} + \frac{1}{q} = 1$.

Fractional calculus is a new research area of analytical mathematics which provides many useful tools for modelling various complex physical and biological processes with long memory [24–31]. For example, in fluid dynamics, laboratory data [24] and numerical experiment [25] show that solutes moving through a highly heterogeneous aquifer do not abide by Fick’s first law, and thus in order to improve the accuracy of the model, one can adopt fractional order advection–dispersion equation to describe the convection–diffusion process in a highly heterogeneous aquifer, see [24, 32–40]. In biomedicine, Arafa et al. [41] introduced a fractional-order HIV-1 infection of CD4+ T cells dynamics model and then used the generalised Euler method to find a numerical solution of the HIV-1 infection fractional order model. Subsequently, by analytical techniques, Wang et al. [42] and Zhang et al. [43] studied the existence of positive solution for some abstract fractional dynamic systems for bioprocess, respectively.

On the other hand, the p -Laplacian equation is a second order quasilinear differential operator with the ability of modelling various fundamental nonlinear phenomena in non-Newtonian fluids, nonlinear elasticity, torsional creep problem, radiation of heat, etc. [44–56]. Thus fractional order differential equations with p -Laplacian operator not only can describe the nonlinear phenomena in non-Newtonian fluids but also can model complex processes with long memory. For example, by using the monotone iterative technique, Wu et al. [57] investigated the existence of twin iterative solutions for a fractional differential turbulent flow model

$$\begin{cases} -\mathcal{D}_x^\alpha(\varphi_p(-\mathcal{D}_x^\gamma z))(x) = g(x)h(z), & 0 < t < 1, \\ z(0) = 0, \quad \mathcal{D}_x^\gamma z(0) = \mathcal{D}_x^\gamma z(1) = 0, \quad z(1) = \int_0^1 z(x) d\chi(x), \end{cases}$$

where $\mathcal{D}_x^\gamma, \mathcal{D}_x^\alpha$ are the standard Riemann–Liouville derivatives such that $1 < \alpha, \gamma \leq 2$, and $h : [0, +\infty) \rightarrow [0, \infty)$ is a continuous and increasing function in the variable. The above work (also see [58–67]) shows that the monotone iterative technique is an effective analysis tool for obtaining iterative solutions and numerical solutions of the relative differential equations. However, to the best of our knowledge, in the application of iterative techniques, almost all works require the nonlinear term to be increasing in space variables and not to have singularity at space variables. So, even for the simplest case as Eq. (1.1), iterative solutions are difficult to construct by using classical iterative techniques. Thus in this paper, by introducing a double iterative technique, we study the convergence analysis and error estimation of the unique solution for the case where the nonlinearity in the equation is decreasing in space variables and is allowed to be singular at some time and space variables.

This paper is organised as follows. In Sect. 2, we firstly recall the definitions and properties of the Riemann–Liouville fractional derivative and integral, and then give some lemmas which will be used in the rest of this paper. In Sect. 3, we introduce a double iterative technique and establish the condition for which Eq. (1.2) has a unique positive solution, then from the developed iterative technique, the sequences converging uniformly to the unique positive solution are formulated, and the estimates of the approximation error and the convergence rate are derived.

2 Preliminaries and lemmas

In this section, we firstly recall the definitions and properties of the Riemann–Liouville fractional derivative and integral, and then give some useful lemmas.

Definition 2.1 ([68]) The Riemann–Liouville fractional integral of order $\gamma > 0$ of a function $z : (0, +\infty) \rightarrow \mathbb{R}$ is given by

$$I^\gamma z(x) = \frac{1}{\Gamma(\gamma)} \int_0^x (x - y)^{\gamma-1} z(y) dy$$

provided that the right-hand side is pointwise defined on $(0, +\infty)$.

Definition 2.2 ([68]) The Riemann–Liouville fractional derivative of order $\gamma > 0$ of a function $z : (0, +\infty) \rightarrow \mathbb{R}$ is given by

$$\mathcal{D}_x^\gamma z(x) = \frac{1}{\Gamma(n - \gamma)} \left(\frac{d}{dx} \right)^n \int_0^x (x - y)^{n-\gamma-1} z(y) dy,$$

where $n = [\gamma] + 1$, $[\gamma]$ denotes the integer part of number γ , provided that the right-hand side is pointwise defined on $(0, +\infty)$.

Property 2.1 ([68])

(1) If $z \in L^1(0, 1)$, $\gamma > \alpha > 0$, then

$$I^\gamma I^\alpha z(x) = I^{\gamma+\alpha} z(x), \quad \mathcal{D}_x^\alpha I^\gamma z(x) = I^{\gamma-\alpha} z(x), \quad \mathcal{D}_x^\alpha I^\alpha z(x) = z(x).$$

(2) If $\gamma > 0$, $\alpha > 0$, then

$$\mathcal{D}_x^\gamma x^{\alpha-1} = \frac{\Gamma(\alpha)}{\Gamma(\alpha - \gamma)} x^{\alpha-\gamma-1}.$$

(3) Let $\gamma > 0$, and $z(x)$ is integrable, then

$$I^\gamma \mathcal{D}_x^\gamma z(x) = z(x) + c_1 x^{\gamma-1} + c_2 x^{\gamma-2} + \dots + c_n x^{\gamma-n},$$

where $c_i \in \mathbb{R}$ ($i = 1, 2, \dots, n$), n is the smallest integer greater than or equal to γ .

According to the definitions and properties of the Riemann–Liouville fractional derivative and integral and discussion in [34], we have the following lemma.

Lemma 2.1 Given $h \in L^1(0, 1)$, the following boundary value problem

$$\begin{cases} -\mathcal{D}_x^\alpha z(x) = h(x), & 0 < x < 1, \\ z(0) = z(1) = 0, \end{cases} \tag{2.1}$$

has the unique solution

$$z(x) = \int_0^1 K_\alpha(x, y) h(y) dy,$$

where

$$K_\alpha(x, y) = \frac{1}{\Gamma(\alpha)} \begin{cases} [x(1-y)]^{\alpha-1}, & 0 \leq x \leq y \leq 1, \\ [x(1-y)]^{\alpha-1} - (x-y)^{\alpha-1}, & 0 \leq y \leq x \leq 1, \end{cases} \tag{2.2}$$

with an index α .

On the other hand, by using Property 2.1(3), we get that the unique solution of the equation

$$\begin{cases} -\mathcal{D}_x^\gamma z(x) = 0, & 0 < x < 1, \\ z(0) = 0, & z(1) = 1, \end{cases} \tag{2.3}$$

is $x^{\gamma-1}$. Thus let

$$\mathcal{L} = \int_0^1 x^{\gamma-1} d\chi(x), \quad \mathcal{K}_\chi(y) = \int_0^1 K_\gamma(x, y) d\chi(x),$$

and according to the strategy of [45], we have the following lemma.

Lemma 2.2 *Suppose $1 < \gamma \leq 2$ and $h \in L^1(0, 1)$, then the following nonlocal boundary value problem*

$$\begin{cases} -\mathcal{D}_x^\gamma z(x) = h(x), & x \in (0, 1), \\ z(0) = 0, & z(1) = \int_0^1 z(x) d\chi(x), \end{cases} \tag{2.4}$$

has the unique solution

$$z(x) = \int_0^1 H(x, y)h(y) dy, \tag{2.5}$$

where

$$H(x, y) = \frac{x^{\gamma-1}}{1-\mathcal{L}}\mathcal{K}_\chi(y) + K_\gamma(x, y). \tag{2.6}$$

Lemma 2.3 ([69]) *Let $0 \leq \mathcal{L} < 1$ and $\mathcal{K}_\chi(y) \geq 0$ for $y \in [0, 1]$, then $K_\alpha(x, y)$ and $H(x, y)$ have the following properties:*

- (1) $K_\alpha(x, y)$ and $H(x, y)$ are nonnegative and continuous for $(x, y) \in [0, 1] \times [0, 1]$.
- (2) $K_\alpha(x, y)$ satisfies

$$\frac{x^{\alpha-1}(1-x)y(1-y)^{\alpha-1}}{\Gamma(\alpha)} \leq K_\alpha(x, y) \leq \frac{\alpha-1}{\Gamma(\alpha)}y(1-y)^{\alpha-1}, \quad \text{for } x, y \in [0, 1]. \tag{2.7}$$

- (3) *There exist two constants a, b such that*

$$ax^{\gamma-1}\mathcal{K}_\chi(y) \leq H(x, y) \leq bx^{\gamma-1}, \quad y, x \in [0, 1]. \tag{2.8}$$

Let q be the conjugate index of p , and consider the following associated linear nonlocal boundary value problem:

$$\begin{cases} -\mathcal{D}_x^\alpha (\varphi_p(-\mathcal{D}_x^\gamma z))(x) = h(x), & x \in (0, 1), \\ z(0) = 0, \quad \mathcal{D}_x^\gamma z(0) = \mathcal{D}_x^\gamma z(1) = 0, & z(1) = \int_0^1 z(x) d\chi(x), \end{cases} \tag{2.9}$$

for $h \in L^1(0, 1)$ and $h \geq 0$. We have the following result.

Lemma 2.4 *The associated linear nonlocal boundary value problem (2.9) has a unique positive solution with the form*

$$z(x) = \int_0^1 H(x, y) \left(\int_0^1 K_\alpha(y, \tau) h(\tau) d\tau \right)^{q-1} dy.$$

Proof Let $w = -\mathcal{D}_x^\gamma z$, $v = \varphi_p(w) = \varphi_p(-\mathcal{D}_x^\gamma z)$, then we have

$$v(0) = \varphi_p(-\mathcal{D}_x^\gamma z(0)) = 0, \quad v(1) = \varphi_p(-\mathcal{D}_x^\gamma z(1)) = 0, \quad -\mathcal{D}_x^\gamma z = \varphi_p^{-1}(v). \tag{2.10}$$

Now consider the fractional Dirichlet boundary value problem

$$\begin{cases} -\mathcal{D}_x^\alpha v(x) = h(x), & t \in (0, 1), \\ v(0) = v(1) = 0. \end{cases}$$

It follows from Lemma 2.1 that

$$v(x) = \int_0^1 K_\alpha(x, y) h(y) dy, \quad x \in [0, 1]. \tag{2.11}$$

Thus by (2.9)–(2.11), one gets that the solution of (2.9) satisfies

$$\begin{cases} -\mathcal{D}_x^\gamma z(x) = \varphi_p^{-1} \left(\int_0^1 K_\alpha(x, y) h(y) dy \right), & x \in (0, 1), \\ z(0) = 0, \quad z(1) = \int_0^1 z(x) d\chi(x). \end{cases}$$

Hence, according to Lemma 2.2, the solution of the boundary value problem (2.9) can be written by

$$z(x) = \int_0^1 H(x, y) \varphi_p^{-1} \left(\int_0^1 K_\alpha(y, \tau) h(\tau) d\tau \right) dy, \quad x \in [0, 1].$$

As $h(y) \geq 0, y \in [0, 1]$, the solution of Eq. (2.9) is also positive. □

3 Main results

In this section, we firstly list some assumptions and then give the proof of our main results.

(K_0) χ is a function of bounded variation satisfying $\mathcal{K}_\chi(y) \geq 0$ for $y \in [0, 1]$ and $0 \leq \mathcal{L} < 1$.

(F_1) $f \in C((0, 1) \times (0, +\infty), [0, +\infty))$, and $f(x, z)$ is decreasing in z and for any $r \in (0, 1)$, there exists a constant $0 < \mu < \frac{1}{p-1}$ such that, for any $(x, z) \in (0, 1) \times (0, +\infty)$,

$$f(x, rz) \leq r^{-\mu} f(x, z). \tag{3.1}$$

Remark 3.1 Obviously, if $p = \frac{3}{2}$, then $f(x, z) = x^{-1}z^{-\frac{1}{2}}$ satisfies the assumption (F_1) which implies that f can be allowed to have singularity at $x = 0$ and $z = 0$.

Remark 3.2 If (F_1) holds, from (3.1), for any $r \geq 1$, one has the following equivalent statement:

$$f(x, rz) \geq r^{-\mu}f(x, z) \quad \text{for any } (x, z) \in (0, 1) \times (0, +\infty). \tag{3.2}$$

In this paper, our work space is a Banach space $E = C[0, 1]$ with the norm $\|z\| = \max_{x \in [0, 1]} |z(x)|$ for any $z \in E$. Let $P = \{z \in C[0, 1] : z(x) \geq 0, x \in [0, 1]\}$, then P is a normal cone of E with normality constant 1. Now define a subset of P and a nonlinear integral operator $T: E \rightarrow E$ by

$Q = \{z(x) \in P : \text{there exists a positive number } 0 < l_z < 1 \text{ such that}$

$$l_z x^{\gamma-1} \leq z(x) \leq l_z^{-1} x^{\gamma-1}, x \in [0, 1]\},$$

and

$$(Tz)(x) = \int_0^1 H(x, y) \left(\int_0^1 K_\alpha(y, \tau) f(\tau, z(\tau)) d\tau \right)^{q-1} dy, \quad x \in [0, 1]. \tag{3.3}$$

It follows from Lemma 2.4 that $z \in C[0, 1]$ is a solution of the p -Laplacian fractional differential Eq. (1.2) if and only if $z \in C[0, 1]$ is a fixed point of the nonlinear operator T .

Theorem 3.1 *Suppose (K_0) and (F_1) hold. If*

$$0 < \int_0^1 x(1-x)^{\alpha-1} f(x, x^{\gamma-1}) dx < +\infty, \tag{3.4}$$

then

- (i) *the p -Laplacian fractional differential Eq. (1.2) has a unique positive solution $z^* \in C[0, 1]$;*
- (ii) *for any initial value $z_0 \in Q$, the sequence of functions $\{z_n\}_{n \geq 1}$ defined by*

$$z_n = \int_0^1 H(x, y) \left(\int_0^1 K_\alpha(y, \tau) f(\tau, z_{n-1}(\tau)) d\tau \right)^{q-1} dy, \quad n = 1, 2, 3, \dots, \tag{3.5}$$

converge uniformly to the unique positive solution z^ of Eq. (1.2) on $[0, 1]$;*

- (iii) *the error between the iterative value z_n and the exact solution z^* can be estimated by*

$$\|z_n - z^*\| \leq (1 - \epsilon^{[\mu(q-1)]^{2n}}) \epsilon^{-\frac{1}{2}},$$

with an exact convergence rate

$$\|z_n - z^*\| = o(1 - \epsilon^{[\mu(q-1)]^{2n}}),$$

where $0 < \epsilon < 1$ is a positive constant.

(iv) *there exists a constant $0 < l < 1$ such that the exact solution z^* of Eq. (1.2) intervenes between two known curves $lx^{\gamma-1}$ and $l^{-1}x^{\gamma-1}$, i.e.,*

$$lx^{\gamma-1} \leq z^*(x) \leq l^{-1}x^{\gamma-1}, \quad x \in [0, 1].$$

Proof Step 1. We show that $T: Q \rightarrow Q$ is a compact operator.

In fact, for any $z \in Q$, it follows from the definition of the set Q that there exists a constant $0 < l_z < 1$ such that

$$l_z x^{\gamma-1} \leq z(x) \leq l_z^{-1} x^{\gamma-1}, \quad x \in [0, 1]. \tag{3.6}$$

Notice that $f(x, z)$ is decreasing in z , by Lemma 2.3, (3.1), (3.4) and (3.6), one has

$$\begin{aligned} (Tz)(x) &= \int_0^1 H(x, y) \left(\int_0^1 K_\alpha(y, \tau) f(\tau, z(\tau)) d\tau \right)^{q-1} dy \\ &\leq \int_0^1 bx^{\gamma-1} \left(\int_0^1 \frac{\alpha-1}{\Gamma(\alpha)} \tau(1-\tau)^{\alpha-1} f(\tau, l_z \tau^{\gamma-1}) d\tau \right)^{q-1} dy \\ &\leq bl_z^{-\mu(q-1)} \left(\frac{\alpha-1}{\Gamma(\alpha)} \right)^{q-1} \left(\int_0^1 \tau(1-\tau)^{\alpha-1} f(\tau, \tau^{\gamma-1}) d\tau \right)^{q-1} \\ &< +\infty. \end{aligned}$$

So T is well defined and uniformly bounded.

On the other hand, since $H(x, y)$ is uniformly continuous on $[0, 1] \times [0, 1]$, let $0 \leq x_1 < x_2 \leq 1$, for all $z \in Q$, one has

$$\begin{aligned} &|(Tz)(x_1) - (Tz)(x_2)| \\ &\leq \int_0^1 |H(x_1, y) - H(x_2, y)| \left(\int_0^1 K_\alpha(y, \tau) f(\tau, z(\tau)) d\tau \right)^{q-1} dy \\ &\leq \int_0^1 |H(x_1, y) - H(x_2, y)| \left(\int_0^1 K_\alpha(y, \tau) f(\tau, l_z \tau^{\gamma-1}) d\tau \right)^{q-1} dy \\ &\leq l_z^{-\mu(q-1)} \left(\frac{\alpha-1}{\Gamma(\alpha)} \right)^{q-1} \left(\int_0^1 \tau(1-\tau)^{\alpha-1} f(\tau, \tau^{\gamma-1}) d\tau \right)^{q-1} \int_0^1 |H(x_1, y) - H(x_2, y)| dy, \end{aligned}$$

which implies that $T(Q)$ is equicontinuous, and then T is a compact operator in Q .

In the following, we shall show that $T(Q) \subset Q$. In fact, by (2.7), (2.8), (3.6) and (3.1), for any $z \in Q$, we have

$$\begin{aligned} (Tz)(x) &= \int_0^1 H(x, y) \left(\int_0^1 K_\alpha(y, \tau) f(\tau, z(\tau)) d\tau \right)^{q-1} dy \\ &\leq \frac{bx^{\gamma-1}}{\Gamma^{q-1}(\alpha)} \left(\int_0^1 \tau(1-\tau)^{\alpha-1} f(\tau, l_z \tau^{\gamma-1}) d\tau \right)^{q-1} \\ &\leq l_z^{-\mu(q-1)} \left(\frac{\alpha-1}{\Gamma(\alpha)} \right)^{q-1} x^{\gamma-1} \left(\int_0^1 \tau(1-\tau)^{\alpha-1} f(\tau, \tau^{\gamma-1}) d\tau \right)^{q-1} \\ &\leq \tilde{l}_{Tz}^{-1} x^{\gamma-1}, \end{aligned} \tag{3.7}$$

and

$$\begin{aligned}
 (Tz)(x) &= \int_0^1 H(x, y) \left(\int_0^1 K_\alpha(y, \tau) f(\tau, z(\tau)) d\tau \right)^{q-1} dy \\
 &\geq ax^{\gamma-1} \int_0^1 \mathcal{K}_\chi(y) \left(\int_0^1 K_\alpha(y, \tau) f(\tau, l_z^{-1} \tau^{\gamma-1}) d\tau \right)^{q-1} dy \\
 &\geq ax^{\gamma-1} l_z^{\mu(q-1)} \int_0^1 \mathcal{K}_\chi(y) \left(\int_0^1 K_\alpha(y, \tau) f(\tau, \tau^{\gamma-1}) d\tau \right)^{q-1} dy \\
 &\geq ax^{\gamma-1} \left(\frac{l_z^\mu}{\Gamma(\alpha)} \right)^{q-1} \int_0^1 \mathcal{K}_\chi(y) y^{(\alpha-1)(q-1)} (1-y)^{q-1} dy \\
 &\quad \times \left(\int_0^1 \tau(1-\tau)^{\alpha-1} f(\tau, \tau^{\gamma-1}) d\tau \right)^{q-1} \\
 &\geq \tilde{l}_{T_z} x^{\gamma-1},
 \end{aligned} \tag{3.8}$$

where \tilde{l}_{T_z} satisfies

$$\begin{aligned}
 0 &< \tilde{l}_{T_z} \\
 &< \min \left\{ \frac{1}{2}, \left\{ l_z^{-\mu(q-1)} \left(\frac{\alpha-1}{\Gamma(\alpha)} \right)^{q-1} \left(\int_0^1 \tau(1-\tau)^{\alpha-1} f(\tau, \tau^{\gamma-1}) d\tau \right)^{q-1} \right\}^{-1}, \right. \\
 &\quad a \left(\frac{l_z^\mu}{\Gamma(\alpha)} \right)^{q-1} \int_0^1 \mathcal{K}_\chi(y) y^{(\alpha-1)(q-1)} (1-y)^{q-1} dy \\
 &\quad \left. \times \left(\int_0^1 \tau(1-\tau)^{\alpha-1} f(\tau, \tau^{\gamma-1}) d\tau \right)^{q-1} \right\}.
 \end{aligned} \tag{3.9}$$

Hence we have $T(Q) \subset Q$.

Step 2. In this step, we prove that Eq. (1.2) has a unique positive solution $z^* \in C[0, 1]$.

In fact, let $\eta(x) = x^{\gamma-1}$, then $\eta \in Q$. By Step 1, we have $T\eta \in Q$. Thus there exists a constant l_{T_η} such that $0 < l_{T_\eta} < 1$ and

$$l_{T_\eta} \eta(x) \leq T\eta(x) \leq l_{T_\eta}^{-1} \eta(x), \tag{3.10}$$

where l_{T_η} can be chosen as in (3.9). Notice that $0 < \mu(q-1) < 1$, for some $\kappa \in (0, 1)$, we can choose a sufficiently large positive constant σ such that

$$[\kappa^{(-\mu(q-1)+1)}]^\sigma \leq l_{T_\eta}. \tag{3.11}$$

Now fix the initial value $z_0 = \kappa^\sigma \eta(x)$ and let

$$z_n = Tz_{n-1}, \quad n = 1, 2, \dots \tag{3.12}$$

We firstly show

$$z_0 \leq z_2 \leq \dots \leq z_{2n} \leq \dots \leq z_{2n+1} \leq \dots \leq z_3 \leq z_1. \tag{3.13}$$

In fact, since T is a decreasing operator in z , it follows from (3.10)–(3.12) that

$$\begin{aligned} z_0(x) &\leq \eta(x), \\ z_1 = Tz_0 &\geq T\eta \geq l_{T\eta} \eta(x) \geq (\kappa^{-\mu(q-1)+1})^\sigma \eta(x) = (\kappa^{\mu(q-1)})^{-\sigma} \kappa^\sigma \eta(x) \\ &= (\kappa^{\mu(q-1)})^{-\sigma} z_0 \geq z_0, \end{aligned} \tag{3.14}$$

and then

$$z_2 = Tz_1(x) \leq Tz_0(x) = z_1. \tag{3.15}$$

On the other hand, it follows from (3.1) and (3.10) that

$$\begin{aligned} z_1 = T(z_0) &= \int_0^1 H(x, y) \left(\int_0^1 K_\alpha(y, \tau) f(\tau, \kappa^\sigma \eta(\tau)) d\tau \right)^{q-1} dy \\ &\leq \kappa^{-\mu\sigma(q-1)} \int_0^1 H(x, y) \left(\int_0^1 K_\alpha(y, \tau) f(\tau, \eta(\tau)) d\tau \right)^{q-1} dy \\ &= \kappa^{-\mu\sigma(q-1)} T\eta \leq \kappa^{-\mu\sigma(q-1)} l_{T\eta}^{-1} \eta(x) \leq \kappa^{-\sigma} \eta(x), \end{aligned} \tag{3.16}$$

and then by (3.2), (3.10), (3.16) and the monotonicity of T , one gets

$$\begin{aligned} z_2 = Tz_1(x) &\geq T(\kappa^{-\sigma} \eta(x)) = \int_0^1 H(x, y) \left(\int_0^1 K_\alpha(y, \tau) f(\tau, \kappa^{-\sigma} \eta(\tau)) d\tau \right)^{q-1} dy \\ &\geq \kappa^{\sigma\mu(q-1)} T\eta(x) \geq \kappa^{\sigma\mu(q-1)} l_{T\eta} \eta(x) \geq \kappa^\sigma \eta(x) = z_0. \end{aligned} \tag{3.17}$$

Equation (3.14), (3.15) and (3.17) yield

$$z_0 \leq z_2 \leq z_1. \tag{3.18}$$

Consequently, by applying induction for (3.18), we obtain (3.13).

Now, for any $c \in (0, 1)$, from (3.1) and (3.3) we have

$$T^2(cz) \geq c^{\mu^2(q-1)^2} T^2 z. \tag{3.19}$$

Noticing that T^2 is a nondecreasing operator with respect to z , by using (3.19) repeatedly, we obtain

$$\begin{aligned} z_{2n} = Tz_{2n-1}(x) &= T^{2n} z_0 = T^{2n} (\kappa^\sigma \eta(x)) = T^{2n} (\kappa^{2\sigma} \kappa^{-\sigma} \eta(x)) \\ &\geq T^{2n-2} (T^2 (\kappa^{2\sigma} z_1(x))) \geq T^{2n-2} ((\kappa^{2\sigma})^{\mu^2(q-1)^2} T^2 z_1(x)) \\ &= T^{2n-4} T^2 ((\kappa^{2\sigma})^{\mu^2(q-1)^2} T^2 z_1(x)) \geq T^{2n-4} ((\kappa^{2\sigma})^{\mu^4(q-1)^4} T^4 z_1(x)) \\ &\geq \dots \geq (\kappa^{2\sigma})^{\mu^{2n}(q-1)^{2n}} T^{2n} z_1(x) = (\kappa^{2\sigma})^{\mu^{2n}(q-1)^{2n}} T^{2n+1} z_0(x) \\ &= (\kappa^{2\sigma})^{\mu^{2n}(q-1)^{2n}} z_{2n+1}, \end{aligned} \tag{3.20}$$

that is,

$$(\kappa^{2\sigma})^{\mu^{2n}(q-1)^{2n}} z_{2n+1} \leq z_{2n} \leq z_{2n+1}. \tag{3.21}$$

Consequently, for all natural numbers n and p , one has

$$\begin{aligned} 0 \leq z_{2(n+p)}(x) - z_{2n}(x) &\leq z_{2n+1}(x) - z_{2n}(x) \leq (1 - (\kappa^{2\sigma})^{\mu^{2n}(q-1)^{2n}}) z_{2n+1} \\ &\leq (1 - (\kappa^{2\sigma})^{\mu^{2n}(q-1)^{2n}}) z_1 \leq (1 - (\kappa^{2\sigma})^{\mu^{2n}(q-1)^{2n}}) \kappa^{-\sigma} \eta(x), \end{aligned} \tag{3.22}$$

and

$$0 \leq z_{2n+1}(x) - z_{2(n+p)+1}(x) \leq z_{2n+1}(x) - z_{2n}(x) \leq (1 - (\kappa^{2\sigma})^{\mu^{2n}(q-1)^{2n}}) \kappa^{-\sigma} \eta(x). \tag{3.23}$$

It follows from (3.22), (3.23) and the fact that P is a normal cone with normality constant 1 that

$$\|z_{n+p} - z_n\| \leq (1 - (\kappa^{2\sigma})^{\mu^{2n}(q-1)^{2n}}) \kappa^{-\sigma} \rightarrow 0, \quad n \rightarrow +\infty. \tag{3.24}$$

Since $\{z_n\} \in Q$ and $T(Q) \subset Q$ is compact, $\{z_n\}$ is a Cauchy sequence of compact set, and then $\{z_n\}$ converges to some $z^* \in Q$ with

$$z_{2n} \leq z^* \leq z_{2n+1}.$$

So

$$z_{2n+2} = Tz_{2n+1} \leq Tz^* \leq Tz_{2n} = z_{2n+1}. \tag{3.25}$$

Let $n \rightarrow \infty$ in (3.25), we get $z^*(x) = Tz^*(x)$, which implies that z^* is a positive solution of Eq. (1.2).

Now we prove $z^* \in Q$ is unique. Let \tilde{z} be another positive solution of Eq. (1.2). Take $r_1 = \sup\{r > 0 \mid \tilde{z} \geq rz^*\}$. Obviously, $0 < r_1 < +\infty$. We assert $r_1 \geq 1$. If not, we have $0 < r_1 < 1$, which leads to

$$\tilde{z} = T\tilde{z} = T^2\tilde{z} \geq T^2(r_1 z^*) \geq r_1^{\mu^2(q-1)^2} T^2 z^* = r_1^{\mu^2(q-1)^2} z^*.$$

Since $r_1^{\mu^2(q-1)^2} > r_1$, this contradicts with the definition of r_1 . Hence $r_1 \geq 1$ and $\tilde{z} \geq z^*$. Similarly, we also have $\tilde{z} \leq z^*$. Therefore $\tilde{z} = z^*$, which implies that the positive solution of Eq. (1.2) is unique.

Step 3. At the end, we give the convergence analysis and error estimation for the unique solution of Eq. (1.2).

For any initial value $\omega_0 \in Q$, there exists a constant $l_{\omega_0} \in (0, 1)$ such that

$$l_{\omega_0} \eta(x) \leq \omega_0(x) \leq l_{\omega_0}^{-1} \eta(x), \quad x \in [0, 1].$$

Since $T(Q) \subset Q$, there still exists a constant $l_{\omega_1} \in (0, 1)$ such that

$$l_{\omega_1} \eta(x) \leq \omega_1 = T\omega_0 \leq l_{\omega_1}^{-1} \eta(x), \quad x \in [0, 1].$$

Choose sufficiently large $\tilde{\sigma} > 2\sigma$ such that

$$\kappa^{\tilde{\sigma}-\sigma} \leq \min\{l_{\omega_0}, l_{\omega_1}\},$$

where $\kappa \in (0, 1)$ and $\sigma > 0$ are defined by (3.11). Thus

$$z_0 = \kappa^\sigma \eta(x) \leq \kappa^{\tilde{\sigma}-\sigma} \eta(x) \leq l_{\omega_0} \eta(x) \leq \omega_0, \quad z_0 = \kappa^\sigma \eta(x) \leq \kappa^{\tilde{\sigma}-\sigma} \eta(x) \leq l_{\omega_1} \eta(x) \leq \omega_1,$$

which implies that $\omega_1 = T\omega_0 \leq Tz_0 = z_1$, and then

$$z_0 \leq \omega_1 \leq z_1. \tag{3.26}$$

Let

$$\omega_n = \int_0^1 H(x, y) \left(\int_0^1 K_\alpha(y, \tau) f(\tau, \omega_{n-1}(\tau)) d\tau \right)^{q-1} dy, \quad n = 1, 2, 3, \dots, \tag{3.27}$$

it follows from (3.26) and (3.27) that

$$z_{2n}(x) \leq \omega_{2n}(x) \leq z_{2n+1}(x), \quad z_{2n+2}(x) \leq \omega_{2n+1}(x) \leq z_{2n+1}(x). \tag{3.28}$$

Letting $n \rightarrow \infty$ in (3.28) and using (3.25), we get that ω_n uniformly converges to the unique positive solution z^* of Eq. (1.2).

Moreover, by (3.23) and (3.28), we have the following estimate of error:

$$\|\omega_n - z^*\| \leq (1 - (\kappa^{2\sigma})^{\mu^{2n}(q-1)^{2n}}) \kappa^{-\sigma} = (1 - \epsilon^{\mu^{2n}(q-1)^{2n}}) \epsilon^{-\frac{1}{2}}, \tag{3.29}$$

with an exact rate of convergence

$$\|\omega_n - z^*\| = o(1 - (\kappa^{2\sigma})^{\mu^{2n}(q-1)^{2n}}) = o(1 - \epsilon^{\mu^{2n}(q-1)^{2n}}),$$

where $0 < \epsilon = \kappa^{2\sigma} < 1$ is a positive constant which is determined by $z_0 = \kappa^\sigma \eta(x)$, that is, it is independent of the initial value ω_0 .

At the end, it follows from $z^* \in Q$ that there exists a constant $0 < l_1 < 1$ such that

$$l_1 x^{\gamma-1} \leq z^*(x) \leq l_1^{-1} x^{\gamma-1}.$$

The proof is completed. □

4 Example

Now we recall the singular nonlocal fractional differential Eq. (1.1). By simple computation, we get that Eq. (1.1) is equivalent to the following 4-point boundary value problem:

$$\begin{cases} -\mathcal{D}_x^{\frac{3}{2}} (\varphi_{\frac{3}{2}} (-\mathcal{D}_x^{\frac{4}{3}} z))(x) = x^{-1} z^{-\frac{1}{2}}, & 0 < x < 1, \\ z(0) = 0, \quad \mathcal{D}_x^{\frac{4}{3}} z(0) = \mathcal{D}_x^{\frac{4}{3}} z(1) = 0, & z(1) = \frac{1}{2} z(\frac{1}{3}) + \frac{1}{2} z(\frac{2}{3}). \end{cases} \tag{4.1}$$

In the following, we shall verify that Eq. (1.1) satisfies all conditions of Theorem 3.1. Let $\alpha = \frac{3}{2}, \gamma = \frac{4}{3}, p = \frac{3}{2}$ and

$$f(x, z) = x^{-1}z^{-\frac{1}{2}},$$

then $f \in C((0, 1) \times [0, \infty), [0, +\infty))$, and for any fixed $x \in (0, 1), f(x, z)$ is nondecreasing in z .

Take $\mu = \frac{2}{3}$, then $0 < \mu < \frac{1}{p-1} = 2$. For any $r \in (0, 1)$ and $(x, z) \in (0, 1) \times (0, +\infty)$, we have

$$f(x, rz) = r^{-\frac{1}{2}}x^{-1}z^{-\frac{1}{2}} \leq r^{-\frac{2}{3}}x^{-1}z^{-\frac{1}{2}} = r^{-\frac{2}{3}}f(x, z). \tag{4.2}$$

Thus condition (F_1) is satisfied.

Next we verify condition (K_0) . In fact, since

$$K_{\frac{4}{3}}(x, y) = \frac{1}{\Gamma(\frac{4}{3})} \begin{cases} [x(1-y)]^{\frac{1}{3}}, & 0 \leq x \leq y \leq 1, \\ [x(1-y)]^{\frac{1}{3}} - (x-y)^{\frac{1}{3}}, & 0 \leq y \leq x \leq 1, \end{cases} \tag{4.3}$$

we have

$$K_{\chi}(y) = \int_0^1 K_{\frac{4}{3}}(x, y) d\chi(x) = \frac{1}{2}K_{\frac{4}{3}}\left(\frac{1}{3}, y\right) + \frac{1}{2}K_{\frac{4}{3}}\left(\frac{2}{3}, y\right) \geq 0, \quad \text{for all } y \in [0, 1],$$

and

$$\mathcal{L} = \int_0^1 x^{\gamma-1} d\chi(x) = \frac{1}{2}\left(\frac{2}{3}\right)^{\frac{1}{3}} + \frac{1}{2}\left(\frac{1}{3}\right)^{\frac{1}{3}} = 0.7835 < 1.$$

So condition (K_0) is also satisfied.

Now we check condition (3.1). In fact, substituting $f(x, z) = x^{-1}z^{-\frac{1}{2}}$ into (3.1), we get

$$0 < \int_0^1 x(1-x)^{\alpha-1} f(x, x^{\gamma-1}) dx = \int_0^1 x^{-\frac{1}{6}}(1-x)^{\frac{1}{2}} dx < +\infty,$$

which implies that (3.1) holds. Thus, according to Theorem 3.1, we have the following conclusions:

- (i) the p -Laplacian fractional differential Eq. (1.1) has a unique positive solution $z^* \in C[0, 1]$;
- (ii) for any initial value $z_0 \in Q$, the sequence of functions $\{z_n\}_{n \geq 1}$ defined by

$$\begin{aligned} z_n &= \int_0^1 \left[\frac{x^{\frac{1}{3}}}{0.2635} \left(\frac{1}{2}K_{\frac{4}{3}}\left(\frac{1}{3}, y\right) + \frac{1}{2}K_{\frac{4}{3}}\left(\frac{2}{3}, y\right) \right) + K_{\frac{4}{3}}(x, y) \right] \\ &\quad \times \left(\int_0^1 K_{\frac{3}{2}}(y, \tau) \tau^{-1} z_n^{-\frac{1}{2}}(\tau) d\tau \right)^2 dy, \\ &n = 1, 2, 3, \dots, \end{aligned}$$

converges uniformly to the unique positive solution z^* of Eq. (1.1) on $[0, 1]$;

(iii) the error between the iterative value z_n and the exact solution z^* can be estimated by

$$\|z_n - z^*\| \leq (1 - \epsilon^{[\frac{4}{3}]^{2n}})\epsilon^{-\frac{1}{2}},$$

and the convergence rate can be formulated by

$$\|z_n - z^*\| = o(1 - \epsilon^{[\frac{4}{3}]^{2n}}),$$

where $0 < \epsilon < 1$ is a positive constant which is determined by the fixed function $\kappa^\sigma x^{\frac{4}{3}}$;

(iv) there exists a constant $0 < l < 1$ such that the exact solution z^* of Eq. (1.1) intervenes between two known curves $lx^{\frac{2}{3}}$ and $l^{-1}x^{\frac{2}{3}}$, i.e.,

$$lx^{\frac{2}{3}} \leq z^*(x) \leq l^{-1}x^{\frac{2}{3}}, \quad x \in [0, 1].$$

5 Conclusion

In this paper, by introducing a double iterative technique, we established the convergence analysis and error estimation for the unique solution of a p -Laplacian fractional differential equation with singular decreasing nonlinearity. The equation we studied in the present paper exhibits a blow-up behaviour at time and space variables, which occurs in many complex physical processes, such as mechanics processes, the convection-diffusion process and the bioprocess with long memory. The developed double iterative technique can be applied for solving the case where the nonlinear term is decreasing and has singularity at time and space variables.

Acknowledgements

We are thankful to the editor and the anonymous reviewers for many valuable suggestions to improve this paper.

Funding

The authors are supported financially by the National Natural Science Foundation of China (11571296).

Availability of data and materials

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

The study was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Author details

¹School of Economic Mathematics, Southwestern University of Finance and Economics, Chengdu, China. ²School of Mathematical and Informational Sciences, Yantai University, Yantai, China. ³Department of Mathematics and Statistics, Curtin University of Technology, Perth, Australia. ⁴School of Mathematical Sciences, Qufu Normal University, Qufu, China. ⁵Department of Mathematics, Shandong University of Science and Technology, Qingdao, China.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 11 April 2018 Accepted: 18 May 2018 Published online: 25 May 2018

References

- Borberg, K.B.: *Cracks and Fracture*. Academic Press, San Diego (1999)
- Sun, F., Liu, L., Zhang, X., Wu, Y.: Spectral analysis for a singular differential system with integral boundary conditions. *Mediterr. J. Math.* **13**, 4763–4782 (2016)

3. Hao, X., Liu, L., Wu, Y.: On positive solutions of m -point nonhomogeneous singular boundary value problem. *Nonlinear Anal.* **73**(8), 2532–2540 (2010)
4. Zhang, X., Liu, L.: A necessary and sufficient condition of positive solutions for nonlinear singular differential systems with four-point boundary conditions. *Appl. Math. Comput.* **215**, 3501–3508 (2010)
5. Liu, L., Hao, X., Wu, Y.: Positive solutions for singular second order differential equations with integral boundary conditions. *Math. Comput. Model.* **57**, 836–847 (2013)
6. Liu, L., Sun, F., Zhang, X., Wu, Y.: Bifurcation analysis for a singular differential system with two parameters via topological degree theory. *Nonlinear Anal., Model. Control* **22**(1), 31–50 (2017)
7. Liu, J., Zhao, Z.: Existence of positive solutions to a singular boundary-value problem using variational methods. *Electron. J. Differ. Equ.* **2014**, 135 (2014)
8. Wu, J., Zhang, X., Liu, L., Wu, Y.: Positive solution of singular fractional differential system with nonlocal boundary conditions. *Adv. Differ. Equ.* **2014**, 323 (2014)
9. Jiang, J., Liu, L., Wu, Y.: Symmetric positive solutions to singular system with multi-point coupled boundary conditions. *Appl. Math. Comput.* **220**, 536–548 (2013)
10. Jiang, J., Liu, L., Wu, Y.: Positive solutions to singular fractional differential system with coupled boundary conditions. *Commun. Nonlinear Sci. Numer. Simul.* **18**, 3061–3074 (2013)
11. Jiang, J., Liu, L., Wu, Y.: Second-order nonlinear singular Sturm–Liouville problems with integral boundary conditions. *Appl. Math. Comput.* **215**, 1573–1582 (2009)
12. Hao, X., Liu, L.: Mild solution of semilinear impulsive integro-differential evolution equation in Banach spaces. *Math. Methods Appl. Sci.* **40**(13), 4832–4841 (2017)
13. Xu, Y., Zhang, H.: Positive solutions of an infinite boundary value problem for n th-order nonlinear impulsive singular integro-differential equations in Banach spaces. *Appl. Math. Comput.* **218**, 5806–5818 (2012)
14. Liu, J., Zhao, Z.: Multiple solutions for impulsive problems with non-autonomous perturbations. *Appl. Math. Lett.* **64**, 143–149 (2017)
15. Guan, Y., Zhao, Z., Lin, X.: On the existence of solutions for impulsive fractional differential equations. *Adv. Math. Phys.* **2017**, Article ID 1207456 (2017)
16. Wang, Y., Zhao, Z.: Existence and multiplicity of solutions for a second-order impulsive differential equation via variational methods. *Adv. Differ. Equ.* **2017**, 46 (2017)
17. Zhang, H., Liu, L., Wu, Y.: Positive solutions for n th-order nonlinear impulsive singular integro-differential equations on infinite intervals in Banach spaces. *Nonlinear Anal.* **70**, 772–787 (2009)
18. Gao, L., Wang, D., Wang, G.: Further results on exponential stability for impulsive switched nonlinear time-delay systems with delayed impulse effects. *Appl. Math. Comput.* **268**, 186–200 (2015)
19. Hao, X., Zuo, M., Liu, L.: Multiple positive solutions for a system of impulsive integral boundary value problems with sign-changing nonlinearities. *Appl. Math. Lett.* **82**, 24–31 (2018)
20. Hao, X., Liu, L., Wu, Y.: Positive solutions for second order impulsive differential equations with integral boundary conditions. *Commun. Nonlinear Sci. Numer. Simul.* **16**, 101–111 (2011)
21. Liu, J., Zhao, Z.: An application of variational methods to second-order impulsive differential equation with derivative dependence. *Electron. J. Differ. Equ.* **2014**, 62 (2014)
22. Sun, F., Liu, L., Wu, Y.: Finite time blow-up for a class of parabolic or pseudo-parabolic equations. *Comput. Math. Appl.* **75**, 3685–3701 (2018)
23. Sun, F., Liu, L., Wu, Y.: Finite time blow-up for a thin-film equation with initial data at arbitrary energy level. *J. Math. Anal. Appl.* **458**, 9–20 (2018)
24. Benson, D., Wheatcraft, S., Meerschaert, M.: Application of a fractional advection dispersion equation. *Water Resour. Res.* **36**, 1403–1412 (2000)
25. Eggleston, J., Rojstaczer, S.: Identification of large-scale hydraulic conductivity trends and the influence of trends on contaminant transport. *Water Resour. Res.* **34**(9), 2155–2168 (1998)
26. Zhang, X., Liu, L., Wu, Y., Wiwatanapataphee, B.: The spectral analysis for a singular fractional differential equation with a signed measure. *Appl. Math. Comput.* **257**, 252–263 (2015)
27. Zhang, X., Liu, L., Wu, Y.: The eigenvalue problem for a singular higher fractional differential equation involving fractional derivatives. *Appl. Math. Comput.* **218**(17), 8526–8536 (2012)
28. Zuo, M., Hao, X., Liu, L., Cui, Y.: Existence results for impulsive fractional integro-differential equation of mixed type with constant coefficient and antiperiodic boundary conditions. *Bound. Value Probl.* **2017**, 161 (2017)
29. Zhang, X., Liu, L., Wu, Y.: Existence results for multiple positive solutions of nonlinear higher order perturbed fractional differential equations with derivatives. *Appl. Math. Comput.* **219**(4), 1420–1433 (2012)
30. Zhang, X., Liu, L.: Multiple positive solutions of a singular fractional differential equation with negatively perturbed term. *Math. Comput. Model.* **55**, 1263–1274 (2012)
31. Guo, L., Liu, L., Wu, Y.: Existence of positive solutions for singular fractional differential equations with infinite-point boundary conditions. *Nonlinear Anal., Model. Control* **21**(5), 635–650 (2016)
32. Zhang, X., Liu, L., Wu, Y., Wiwatanapataphee, B.: Nontrivial solutions for a fractional advection dispersion equation in anomalous diffusion. *Appl. Math. Lett.* **66**, 1–8 (2017)
33. Zhang, X., Liu, L., Wu, Y.: Variational structure and multiple solutions for a fractional advection–dispersion equation. *Comput. Math. Appl.* **68**, 1794–1805 (2014)
34. Zhang, X., Liu, L., Wu, Y.: The uniqueness of positive solution for a fractional order model of turbulent flow in a porous medium. *Appl. Math. Lett.* **37**, 26–33 (2014)
35. Wang, Y., Liu, L., Wu, Y.: Positive solutions for a nonlocal fractional differential equation. *Nonlinear Anal.* **74**, 3599–3605 (2011)
36. Qin, H., Zuo, X., Liu, J., Liu, L.: Approximate controllability and optimal controls of fractional dynamical systems of order $1 < q < 2$ in Banach spaces. *Adv. Differ. Equ.* **2015**, 73 (2015)
37. Zhang, X., Liu, L., Wu, Y., Cui, Y.: New result on the critical exponent for solution of an ordinary fractional differential problem. *J. Funct. Spaces* **2017**, Article ID 3976469 (2017)
38. Zhu, B., Liu, L., Wu, Y.: Local and global existence of mild solutions for a class of semilinear fractional integro-differential equations. *Fract. Calc. Appl. Anal.* **20**(6), 1338–1355 (2017)

39. Wang, Y, Liu, L., Wu, Y: Positive solutions for a class of fractional boundary value problem with changing sign nonlinearity. *Nonlinear Anal.* **74**, 6434–6441 (2011)
40. Zhu, B., Liu, L., Wu, Y: Local and global existence of mild solutions for a class of nonlinear fractional reaction-diffusion equation with delay. *Appl. Math. Lett.* **61**, 73–79 (2016)
41. Arafat, A.A.M., Rida, S.Z., Khalil, M.: Fractional modeling dynamics of HIV and CD4+ T-cells during primary infection. *Nonlinear Biomed. Phys.* **6**, 1 (2012)
42. Wang, Y, Liu, L., Zhang, X., Wu, Y: Positive solutions of an abstract fractional semipositone differential system model for bioprocesses of HIV infection. *Appl. Math. Comput.* **258**, 312–324 (2015)
43. Zhang, X., Mao, C., Liu, L., Wu, Y: Exact iterative solution for an abstract fractional dynamic system model for bioprocess. *Qual. Theory Dyn. Syst.* **16**, 205–222 (2017)
44. Kawohl, B.: On a family of torsional creep problems. *J. Reine Angew. Math.* **410**, 1–22 (1990)
45. Zhang, X., Liu, L., Wiwatanapataphee, B., Wu, Y: The eigenvalue for a class of singular p -Laplacian fractional differential equations involving the Riemann–Stieltjes integral boundary condition. *Appl. Math. Comput.* **235**, 412–422 (2014)
46. Zhang, X., Liu, L., Wu, Y., Cui, Y.: Entire blow-up solutions for a quasilinear p -Laplacian Schrödinger equation with a non-square diffusion term. *Appl. Math. Lett.* **74**, 85–93 (2017)
47. Wang, Y, Jiang, J.: Existence and nonexistence of positive solutions for the fractional coupled system involving generalized p -Laplacian. *Adv. Differ. Equ.* **2017**, 337 (2017)
48. Guo, L., Liu, L., Wu, Y: Iterative unique positive solutions for singular p -Laplacian fractional differential equation system with several parameters. *Nonlinear Anal., Model. Control* **23**(2), 182–203 (2018)
49. Zhao, Z.: Fixed points of $\tau - \varphi$ -convex operators and applications. *Appl. Math. Lett.* **23**(5), 561–566 (2010)
50. Hao, X., Wang, H., Liu, L., Cui, Y.: Positive solutions for a system of nonlinear fractional nonlocal boundary value problems with parameters and p -Laplacian operator. *Bound. Value Probl.* **2017**, 182 (2017)
51. Wang, Y, Liu, L., Wu, Y: Extremal solutions for p -Laplacian fractional integro-differential equation with integral conditions on infinite intervals via iterative computation. *Adv. Differ. Equ.* **2015**, 24 (2015)
52. Jiang, J., Liu, L., Wu, Y: Positive solutions for p -Laplacian fourth-order differential system with integral boundary conditions. *Discrete Dyn. Nat. Soc.* **2012**, Article ID 293734 (2012)
53. Zhang, X., Liu, L., Wu, Y., Caccetta, L.: Entire large solutions for a class of Schrödinger systems with a nonlinear random operator. *J. Math. Anal. Appl.* **423**, 1650–1659 (2015)
54. Kong, D., Liu, L., Wu, Y: Triple positive solutions of a boundary value problem for nonlinear singular second order differential equations of mixed type with p -Laplacian. *Comput. Math. Appl.* **58**(7), 1425–1432 (2009)
55. Xu, F., Liu, L., Wu, Y: Multiple positive solutions of four-point nonlinear boundary value problems for higher-order p -Laplacian operator with all derivatives. *Nonlinear Anal.* **71**(9), 4309–4319 (2009)
56. Sun, F., Liu, L., Wu, Y: Infinitely many sign-changing solutions for a class of biharmonic equation with p -Laplacian and Neumann boundary condition. *Appl. Math. Lett.* **73**, 128–135 (2017)
57. Wu, J., Zhang, X., Liu, L., Wu, Y: Twin iterative solutions for a fractional differential turbulent flow model. *Bound. Value Probl.* **2016**, 98 (2016)
58. Zhang, X., Liu, L., Wu, Y., Lu, Y.: The iterative solutions of nonlinear fractional differential equations. *Appl. Math. Comput.* **219**, 4680–4691 (2013)
59. Zhang, X., Liu, L.: The existence and nonexistence of entire positive solutions of semilinear elliptic systems with gradient term. *J. Math. Anal. Appl.* **371**(1), 300–308 (2010)
60. Zhao, Z.: Existence and uniqueness of fixed points for some mixed monotone operators. *Nonlinear Anal.* **73**(6), 1481–1490 (2010)
61. Lin, X., Zhao, Z.: Iterative technique for a third-order differential equation with three-point nonlinear boundary value conditions. *Electron. J. Qual. Theory Differ. Equ.* **12**, 1 (2016)
62. Lin, X., Zhao, Z.: Existence and uniqueness of symmetric positive solutions of $2n$ -order nonlinear singular boundary value problems. *Appl. Math. Lett.* **26**(7), 692–698 (2013)
63. Zhang, X., Liu, L., Wu, Y: The uniqueness of positive solution for a singular fractional differential system involving derivatives. *Commun. Nonlinear Sci. Numer. Simul.* **18**, 1400–1409 (2013)
64. Zhang, X., Liu, L., Wu, Y: The entire large solutions for a quasilinear Schrödinger elliptic equation by the dual approach. *Appl. Math. Lett.* **55**, 1–9 (2016)
65. Zhang, X., Liu, L., Wu, Y., Cui, Y.: The existence and nonexistence of entire large solutions for a quasilinear Schrödinger elliptic system by dual approach. *J. Math. Anal. Appl.* (2018). <https://doi.org/10.1016/j.jmaa.2018.04.040>
66. Sun, Y., Liu, L., Wu, Y: The existence and uniqueness of positive monotone solutions for a class of nonlinear Schrödinger equations on infinite domains. *J. Comput. Appl. Math.* **321**, 478–486 (2017)
67. Mao, J., Zhao, Z.: The existence and uniqueness of positive solutions for integral boundary value problems. *Bull. Malays. Math. Sci. Soc.* **34**, 153–164 (2011)
68. Podlubny, I.: *Fractional Differential Equations*. Mathematics in Science and Engineering. Academic Press, New York (1999)
69. Zhang, X., Han, Y.: Existence and uniqueness of positive solutions for higher order nonlocal fractional differential equations. *Appl. Math. Lett.* **25**, 555–560 (2012)