



**Regenerating drylands in
response to a changing climate**



With support from IDRC and CGIAR global research program Climate Change, Agriculture, and Food Security (CAAFS), IIRR and its local NGO partners is implementing climate smart villages (CSV) to demonstrate community-based adaptation in agriculture in different agroecological zones in Myanmar.

This primer is based on IIRR baseline studies as well as desk research that IIRR commissioned to develop profiles of each CSV in the project. The purpose of this primer is to provide the reader background information as to the agriculture, livelihoods, nutrition, gender and climate change context of each CSV.



Climate change and natural disasters in Myanmar have adversely affected lives, livelihoods and natural resources. Food security and the nutritional well being of poor households are most at risk. Climate risks will continue to threaten agriculture, nutrition and enterprises of farming households and the landless workers.



Water is scarce in the dry zone and the priority always is for humans and livestock use. Very efficient use of water is practiced.



Preparatory anticipatory measures are needed to develop a portfolio of technological and social interventions and/or innovations to address these risks. Just as important is the need to strengthen community level capacities to innovate and adapt to climate change.



Climate smart villages (CSVs) provide local platforms to test, develop and outscale climate smart agriculture technologies and social processes. Because the manifestations of climate change differ from site to site, these CSVs help generate potential solutions for each unique agro ecological, socio cultural and market situation. Participatory action research is an important element of this effort. Adaptation capacities are built in a gradual manner: *in order to identify solutions of local relevance.*



These social and institutional processes are usually referred to as community-based adaptation(CBA). These processes include action research, strengthening of groups, leadership development, and farmer to farmer extension. Location-relevant approaches help facilitate horizontal, local and spontaneous outscaling. Farmer-to-farmer diffusion is nurtured.



These local processes are best undertaken in local townships or municipalities. These CSVs serve as platforms for generating evidence as basis for learning, sharing, and advocacy. Wherever feasible, local funding is leveraged in an effort to build local ownership and demonstrate the potential for scalability. With time CSVs emerge as models for other local governments, civil society and investors.



Regenerating drylands in response to a changing climate

The central dry zone of Myanmar is characterized by low and unpredictable rainfall, degrading soils with low fertility and an environment that is rapidly desertifying. Water is scarce and so are trees.




NyaungOo township in the Central Dry Zone (where the CSV is located) often has the highest temperature of all the regions of Myanmar with a maximum temperature of 33 to 35°C. At times the temperatures in summer can cross 40°C



Rainfall in Nyaung Oo is less than 40 inches a year and is generally erratic, variable and unevenly distributed. This complicates the crop planning decision-making processes.

Figure 2: Rainfall Trend in Nyaung Oo Township from 2007 to 2017.



Source: Department of Meteorology and Hydrology, Nyaung Oo Township

Farming communities in the central dry zone of Myanmar, over the centuries have coped with limited rainfall, poor fertility and unpredictability. To reduce risks, they rely on a range of “climate smart” crops. Most of the sesame, peanuts, sunflower, pigeon pea and chick pea in Myanmar is grown in the dry zone.



Farmers in the central dry zone already practice many elements of what we now refer to as Climate Smart Agriculture (CSA). They take full advantage of the limited rainfall and soil moisture. Farmers undertake very early land preparation in order to “harvest” water from early rain showers.



They use varieties that are known to perform under limited fertility conditions (eg. legumes which fix nitrogen by themselves) and they rely very little on chemical fertilizers relying on micro-nutrient rich animal manure and compost. However, low dosage with chemical fertilizers used at the right boosting yields and biomass.



Traditional farms and farming systems in the central dry zone are nutritionally sensitive. There are very few farming communities in Southeast Asia where there is such a heavy emphasis on protein rich legumes and mineral rich millets (mung beans, pigeon peas, peanut, dolichos, sesame, millets, and sesame).



Local food culture features greens, legumes, and sesame and peanut oil which provide minerals, vitamins, and carbohydrates. Snack foods are also nutritionally important. These attributes of local food systems (the food culture associated with it) need to be protected, encouraged and conserved.



The rich agro biodiversity of central drylands of Myanmar can be characterized as climate smart and nutritional relevant. Their conservation through productivity enhancement and their sustainable use is therefore a high priority for proponents of CSA.




Farmers conserve valuable livestock agrobiodiversity in the central dry zone. Cattle provide tillage, manure, transportation and other ecosystem services. These animals are raised via low carbon footprint and rely on locally sourced feed and residues. Livestock rely on palm trees and palm nuts to provide protein and mineral rich supplements during the long dry season.



These farm animals are economic assets serving local communities well in case of crop failure. CSA can provide pathways to further enhance productivity, improve animal health care and improve feed resources through agroforestry interventions. The Bagan goat is a climate handy and valuable livestock biodiversity resource.



Livestock is an important asset building approach for Myanmar small farmers. These are important coping mechanisms in case of crop failure. Pigs and cattle are economic assets which enhance resilience building are sold for cash in difficult times (IIRR baseline study) .



Livestock are raised for meat. Most of the meat consumed by households is farm grown (ie., not purchased). Similarly eggs and poultry though consumed on fairly regular basis, is rarely purchased from outside. (IIRR baseline studies)





The dry zone of Myanmar would benefit from CSA that supports fodder trees and grasses grown in community forests and in homesteads and school areas. Fodder banks in schools can help farmers with emergency sources of fodder during the dry season.



The restoration of community forests and other common property areas are a critical approach to transforming the environment and landscapes around villages and farms. They serve as fodder and fuel banks for communities during emergencies.



Traditional knowledge and capacities for local innovation have characterized these communities. This capacity to innovate can be directed towards the anticipated challenges of climate variability. Here farmers store peanut seed for over six month via very innovative humidity and temperature management methods.

NOTE: Peanut seed storage is a complicated process as seeds cannot normally be stored for more than 3-4 months due to high oil content and rancidity that affects seed germination. Farmers in NyaungOo have overcome this problem through indigenous seed storage methods they have developed.



Continuous crop cultivation and the over-extraction of vegetation and feeding of all crop residues to livestock has degraded the soils in the dryzone. Low soil fertility levels, low organic matter and enhanced soil erosion reduces productivity from year to year. This “soil mining” year after year further makes farms vulnerable to climate change.



Green leafy vegetables, roots and tubers, and legumes are eaten fairly regularly. These are farm grown or collected from the wild. Cereals grains, oil, fish, exotic vegetables are purchased from the markets. Overall farm households surprisingly have a moderate level of dietary diversity in the Htee Puu CSV (IIRR baseline studies). These are good practices that should be conserved.



CSA programs should consider distributing diversity kits of planting materials (intra species, varietal diversity of beans, green leafy vegetables, roots and tubers and millets) as part of an effort to restore or strengthen local agro biodiversity. Planting materials can even be sourced from markets.



Without rebuilding soil organic matter and reducing degradation, farmers will not be able to take full advantage of CSA and new crops and varieties, and other CSA options. In the CSV village in NyaungOo farmers are planting *Cassia Siamea*, a non-browsable tree that is planted on farm boundaries primarily to rebuild soil organic matter and reduce the effects of winds.



Regenerating drylands in response to a changing climate

Building on the local farming systems (proven to be relatively resilient), CSA programs can also help introduce new cultivars and varieties from research stations for trial and observation in CSVs. Here pigeon pea and green gram from the NyaungOo research station are being tested in a climate smart village.



In the NyaungOo CSV, efforts are also made to re-introduce traditional crops which have been lost. Dolichos lab lab and sorghum from other parts of Myanmar (with analogous climates) were sourced and introduced for small scale testing and promotion.



Regenerating drylands in response to a changing climate

Sorghum used to be an important crop in Nyaung Oo, but with the market opportunities, farmers shifted to peanut (a crop that has proven to be more risky). A renewed role for sorghum without totally replacing peanuts should be considered.

Scaling-out Community based Adaptation (CBA) via Climate Smart and Nutrition Villages as Platforms to Address Food Insecurity in Myanmar

Update Crop Field Based On Mango

မတောင်သွားမည်	- နှိုင်းစာဝင်
ပိုင်ပိုင်ပိုင်	- (၅၀) ခု
ပိုင်ပိုင်ပိုင်	- သရက်၊ မြေညှိ၊ စားနပ်ရိက္ခာ၊ ပဲနီ၊ ပဲနီ၊ မယ်တင်
မသွားမည်	- ဆီ၊ သရက်
မြို့နယ်တစ်ခု	- မတောင်ပိုင်



Sorghum is also featured as a green fodder source for livestock. Sorghum grains are rich in protein, B1, B2, and B3. It is being re-introduced as feed for chickens and as concentrate for pigs, goats and cattle.



Some farmers in dry zone already understand the need to prepare feed from sorghum residues (they mix green and dry sorghum stalks) by chopping fine and mixing with grains and peanut waste. These are low carbon footprint feedings system that need wider promotion.



There is a role for new ideas. In NyaungOo dryland horticulture is being introduced. Twenty-five farmers in the climate smart village are growing mango trees (known as a drought tolerant crops once established for 2 years). The spaces in between are used for annual crop cultivation.



There are various methods to get trees started in dry areas. These include shading, protection from grazing animals and deep pit planting for conserving water (in sandy loam soils) and with mulch around each trees.



Aside from farms, homesteads (which currently only have livestock components) can also be used for CSA interventions. Small-scale fruit production, vegetable production, small livestock, and value addition processing are examples of CSA for homestead spaces.



Homesteads and schools offer “new” spaces for using CSA to empower the poor, tenants, women and the landless. Homesteads and schools in Myanmar are endowed with land that is often under utilized. This can include bio-intensive or backyard gardens.



Landlessness is a major concern in the dryland areas. Livestock can be a special CSA for this group of people provided fodder and water needs are considered. Improved livestock productivity rely on better feed and water management practices.



**At the start
(year one)
every CSV has
to undertake
a baseline
survey,
conduct
participatory
vulnerability
studies, and
identify
climate, food
security and
livelihood
risks.**



Participatory trials help to build capacities to adapt to climate change and to address poverty in an incremental fashion. With time, a portfolio of CSA options and associated social processes are identified for each CSV.



Location specific Participatory Varietal Selection trials (PVS) provide opportunities for determining which varieties are best suited to the area. The Dryland Research Station in NyaungOo provides proven varieties to farmers for on-farm testing.




 Scaling-out Community based Adaptation (CBA) via Climate Smart and Nutrition Villages as Platforms to Address Food Insecurity in Myanmar
 တနင်္သာရီတက္ကသိုလ်မှ ရှိနေသော သက်တောင့်သက်သာရှိသည့် စေ့စပ်စေ့စပ်သည့် စောင့်ရှောက်မှုများ ဆောင်ရွက်ခြင်း၊ လုပ်ငန်းစဉ်များကို အခြေခံပြီး လူထုအခြေပြုလက်ကားနှင့် ဆက်သွယ်သုံးစွဲခြင်းဆိုင်ရာစီမံခန့်ခွဲမှု
Upland Crop Field Based On Mango
 သစ်တောအခြေခံသော မာသီနီစိုက်ခင်း
 တောင်သူအမည် - ဦးညိုစိုင်း
 စိုက်ပျိုးရေးက - (သေ) ဇော
 စိုက်ပျိုးသီးနှံ - သရက်၊ မြေပဲ၊ တောဝတ်ပဲ၊ နံနားပြောင်း၊ မယ်ဝလီ၊
 ကော့ရှားအမည် - ထီးယုကော့ရှား
 ပြုစုပေးအမည် - ဧညာင်ဦးမြို့နယ်၊



Women farmers and female headed households provide new opportunities for CSA introduction especially for small livestock and fruit trees and intensive vegetable production. With proper targetting of households, CSA can be a tool for the economic empowerment of women.



When a critical mass of farmers become engaged in a CSV in addressing climate change, nutritional and poverty risks, these villages emerges as focal points for learning, sharing, and outscaling to other villages in the township. Local farmers play a big role in this process, with research, government, and civil society playing primarily a facilitating role.



Resiliency to climate disasters is closely associated with higher levels of farm biodiversity (polyculture, agroforestry, and small livestock). Accumulated farmer experience resulting from their interaction with their local environment provides a strong foundation for introducing improvements (ie. building on what people already know best).



Regenerating drylands in response to a changing climate

When local communities become engaged in finding solutions to their climate risk and livelihood priorities, our chances for outscaling and sustainability are greatly enhanced. CSVs provide this opportunity for demonstrating wider uptake of CSA.



Implementing/Support Team

IIRR

Wilson John Barbon
Julian Gonsalves
Chan Myae
Yinn Minn Latt

Radanar Ayar

Naing Lin Tun
San Oo

KMF

Kyaw Soe
Ye Win Aung

CDA

Pyae Phyo Hein
Moe Myint Sandar Khaing

Nyo Mar Htwe - ACARE, YAU
Nang Ei Mon Thae - Dep't. of
Agricultural Economics, YAU
Tin Moe Khaing - Food Security
Working Group (FSWG)

KMSS Hakha

Salai Philip Ngun Khar
Van Buai Lian

Research/Compilation

Julian Gonsalves
Wilson John Barbon
Chan Myae
Yinn Minn Latt

Photo Credits

Julian Gonsalves
Chan Myae

Publication Design/ Layout

Celso Amutan

*(Permission is granted for reproduction and wider use.
However, please acknowledge IIRR and its partners.)*



INTERNATIONAL INSTITUTE OF RURAL RECONSTRUCTION

114, Dana Theikdi Lane, Gandamar St, Gabaraye Pagoda Road
8 Ward, Mayangone Township, Yangon

Website: www.iirr.org



RESEARCH PROGRAM ON
Climate Change,
Agriculture and
Food Security



IDRC | CRDI

International Development Research Centre

Centre de recherches pour le développement international



RADANAR AYAR
for the Rural People

