

```

gap> g:= SymmetricGroup( 4 );
Sym( [ 1 .. 4 ] )
gap> tbl:= CharacterTable( g ); HasIrr( tbl );
i5 : betti(t,Weights=>{1,0})
false
0 1 2 3 4
o5 = total: 1 4 13 14 4
0: 1 . . .
1: . 2 2 4 2
2: . 2 5 6 .
3: . . 4 . 2
4: . . . 4 .
5: . . 2 . .
gap> tblmod2:= CharacterTable( tbl, 2 );
BrauerTable( Sym( [ 1 .. 4 ] ), 2 )
gap> tblmod2 = CharacterTable( tbl, 2 );
true
gap> tblmod2 = BrauerTable( tbl, 2 );
true
o5 : BrauerTable
i6 : betti(t,Weights=>{0,1})
0 1 2 3 4
o6 = total: 1 4 13 14 4
0: 1 . . .
1: . 2 . . .
2: . 2 . . .
3: . . 4 . 2
4: . . . 4 .
5: . . 2 . .
gap> libtbl:= CharacterTable( "M" );
CharacterTable( "M" )
gap> CharacterTableRegular( libtbl, 2 );
BrauerTable( "M", 2 )
gap> BrauerTable( libtbl, 2 );
fail
o6 : BettiTally
i7 : t1 = betti(t,Weights=>{1,1})
0 1 2 3 4
o7 = total: 1 4 13 14 4
0: 1 . . .
1: . . . .
2: . . . .
3: . 2 . .
4: . . . .
5: . 2 . .
6: . . 1 .
7: . . 8 6 .
8: . . 4 8 4
gap> CharacterTable( "Symmetric", 4 );
CharacterTable( "Sym(4)" )
gap> ComputedBrauerTables( tbl );
[ , BrauerTable( Sym( [ 1 .. 4 ] ), 2 ) ]
ring r1 = 32003,(x,y,z),ds;
int a,b,c,t=11,5,3,0;
poly f = x^a*y^b+z^(3*c)+x^(c+2)*y^(c-1)+x^(c-2)*y^c*(y^2+t*x)^2;
option(noprot);
timer=1;
ring r2 = 32003,(x,y,z),dp;
poly f=imap(r1,f);
ideal j=jacob(f);
vdim(std(j));
==> 536
vdim(std(j+f));
==> 195
timer=0; // reset timer
o8 = BettiTally{0, {0, 0}, 0} => 1 }
(1, {2, 2}, 4) => 2
(1, {3, 3}, 6) => 2
(2, {3, 7}, 10) => 2
(2, {4, 4}, 8) => 1
(2, {4, 5}, 9) => 4
(2, {5, 4}, 9) => 4
(2, {7, 3}, 10) => 2
(3, {4, 7}, 11) => 4
(3, {7, 4}, 11) => 4
(4, {5, 7}, 12) => 2
(4, {7, 5}, 12) => 2

```

Journal of Software for Algebra and Geometry

Strongly stable ideals and Hilbert polynomials

DAVIDE ALBERELLI AND PAOLO LELLA

Strongly stable ideals and Hilbert polynomials

DAVIDE ALBERELLI AND PAOLO LELLA

ABSTRACT: The `StronglyStableIdeals.m2` package for Macaulay2 provides a method to compute all saturated strongly stable ideals in a given polynomial ring with a fixed Hilbert polynomial. A description of the main method and auxiliary tools is given.

INTRODUCTION. Strongly stable ideals are a key tool in commutative algebra and algebraic geometry. These ideals have nice combinatorial properties that make them well suited for both theoretical and computational applications. In the case of polynomial rings with coefficients in a field of characteristic zero, the notion of strongly stable ideals coincides with the notion of Borel-fixed ideals. Such ideals are fixed by the action of the Borel subgroup of triangular matrices and play a special role in the theory of Gröbner bases because initial ideals in generic coordinates are of this type [Galligo 1974].

In the context of parameter spaces of algebraic varieties, Galligo’s theorem says that each component and each intersection of components of a Hilbert scheme contains at least one point corresponding to a scheme defined by a Borel-fixed ideal. Hence, these ideals are distributed throughout the Hilbert scheme and can be used to study its local structure. To this end, in recent years several authors [Lella and Roggero 2011; 2016; Cioffi and Roggero 2011; Bertone et al. 2013a; 2017a; 2017b] developed algorithmic methods based on the use of strongly stable ideals to construct flat families corresponding to special loci of the Hilbert scheme. In particular, a new open cover of the Hilbert scheme has been defined using strongly stable ideals and the action of the projective linear group [Bertone et al. 2013b; Brachat et al. 2016]. In this construction, the list of all points corresponding to Borel-fixed ideals in a given Hilbert scheme is needed. The main feature of the package `StronglyStableIdeals.m2` is a method to compute this set of points, i.e., the list of all saturated strongly stable ideals in a polynomial ring with a given

The second author is a member of GNSAGA .

MSC2010: primary 13P10; secondary 13P99.

Keywords: strongly stable ideal, Borel-fixed ideal, Hilbert polynomial, Gotzmann number, Hilbert scheme.

`StronglyStableIdeals.m2` version 1.1

Hilbert polynomial. The method has been theoretically introduced in [Cioffi et al. 2011] and improved in [Lella 2012]. Several other tools are developed and presented in the current paper.

1. STRONGLY STABLE IDEALS. Let us denote by $\mathbb{K}[\mathbf{x}]$ the polynomial ring in $n + 1$ variables $\mathbb{K}[x_0, \dots, x_n]$ with coefficients in a field \mathbb{K} . We assume that $x_0 > x_1 > \dots > x_n$. We use the multi-index notation to describe monomials, i.e., $\mathbf{x}^\alpha := x_0^{\alpha_0} \cdots x_n^{\alpha_n}$ for every $\alpha = (\alpha_0, \dots, \alpha_n) \in \mathbb{Z}_{\geq 0}^{n+1}$, and we denote by $\mathbb{T}_{n,s}$ the set of monomials of $\mathbb{K}[\mathbf{x}]$ of degree s . For any monomial \mathbf{x}^α , we denote by $\min \mathbf{x}^\alpha$ and $\max \mathbf{x}^\alpha$ the indices of the minimal and maximal variable dividing \mathbf{x}^α .

Following [Green 2010], *increasing* and *decreasing elementary moves* are defined as the multiplications

$$e_i^+(\mathbf{x}^\alpha) := \frac{x_{i-1}}{x_i} \cdot \mathbf{x}^\alpha, \quad i > 0, \quad \text{and} \quad e_j^-(\mathbf{x}^\alpha) := \frac{x_{j+1}}{x_j} \cdot \mathbf{x}^\alpha, \quad j < n.$$

We say that an elementary move $e_i^{+/-}$ is *admissible* for a monomial \mathbf{x}^α if $\alpha_i > 0$, that is, $e_i^{+/-}(\mathbf{x}^\alpha)$ is a monomial of $\mathbb{K}[\mathbf{x}]$.

Definition 1.1. An ideal $I \subset \mathbb{K}[\mathbf{x}]$ is called *strongly stable* if

- (i) I is a monomial ideal;
- (ii) for every $\mathbf{x}^\alpha \in I$ and for every admissible increasing move e_i^+ , the monomial $e_i^+(\mathbf{x}^\alpha)$ is contained in I .

We recall that a strongly stable ideal is a Borel-fixed ideal. We now summarize some properties holding in general for Borel-fixed ideals and useful in this context.

Proposition 1.2 [Green 2010, Section 2]. *Let $I \subset \mathbb{K}[\mathbf{x}]$ be a strongly stable ideal.*

- (i) *The regularity of I is equal to the maximal degree of a generator.*
- (ii) *Let \mathfrak{m} be the irrelevant ideal of $\mathbb{K}[\mathbf{x}]$. Then, $(I : \mathfrak{m}) = (I : x_n)$, so that the ideal I is saturated if no generator involves the last variable x_n .*
- (iii) *The last variable x_n is a regular element for I , i.e., the multiplication by x_n induces the short exact sequence*

$$0 \longrightarrow \frac{\mathbb{K}[\mathbf{x}]}{I}(t-1) \xrightarrow{\cdot x_n} \frac{\mathbb{K}[\mathbf{x}]}{I}(t) \longrightarrow \frac{\mathbb{K}[\mathbf{x}]}{(x_n, I)}(t) \longrightarrow 0.$$

2. HILBERT POLYNOMIALS. The Hilbert polynomial $p(t)$ of a homogeneous ideal $I \subset \mathbb{K}[\mathbf{x}]$ is the numerical polynomial such that for s sufficiently large

$$\dim_{\mathbb{K}} \left(\frac{\mathbb{K}[\mathbf{x}]}{I} \right)_s = \dim_{\mathbb{K}} \left(\frac{\mathbb{K}[\mathbf{x}]_s}{I_s} \right) = \binom{n+s}{n} - \dim_{\mathbb{K}} I_s = p(s).$$

Obviously, not every numerical polynomial is a Hilbert polynomial of some homogeneous ideal. Those being Hilbert polynomials have been completely described by Gotzmann [1978].

Gotzmann's decomposition. *A numerical polynomial $p(t) \in \mathbb{Q}[t]$ is a Hilbert polynomial if, and only if, it can be written as*

$$p(t) = \binom{n+a_1}{a_1} + \binom{n+a_2-1}{a_2} + \cdots + \binom{n+a_r-(r-1)}{a_r}, \quad a_1 \geq \cdots \geq a_r \geq 0. \quad (1)$$

This decomposition is strictly related to Macaulay's decomposition

$$p(t) = \sum_{k=0}^d \left[\binom{t+k}{k+1} - \binom{t+k-m_k}{k+1} \right],$$

where $d = \deg p(t)$. For all $n \geq d+1$ the saturated lexicographic ideal within $\mathbb{K}[x_0, \dots, x_n]$ with Hilbert polynomial $p(t)$ is

$$(x_0, \dots, x_{n-d-2}, x_{n-d-1}^{b_d+1}, x_{n-d-1}^{b_d} x_{n-d}^{b_{d-1}+1}, \dots, x_{n-d-1}^{b_d} x_{n-d}^{b_{d-1}} \cdots x_{n-1}^{b_0}),$$

where

$$b_d = \#\{a_j \mid a_j = d\} = m_d \quad \text{and} \quad b_k = \#\{a_j \mid a_j = k\} = m_k - m_{k+1}, \quad 0 \leq k < d.$$

The description of the lexicographic ideal in terms of Gotzmann's decomposition gives an insight to the following theorem.

Gotzmann's regularity theorem. *The regularity of a saturated ideal $I \subset \mathbb{K}[\mathbf{x}]$ with Hilbert polynomial $p(t)$ is at most r , where r is the number of terms in the decomposition (1) and it is called the **Gotzmann number** of $p(t)$.*

Example 2.1. The package `StronglyStableIdeals.m2` provides the method `isHilbertPolynomial` to determine if a numerical polynomial is a Hilbert polynomial.

```
Macaulay2, version 1.11
with packages: ConwayPolynomials, Elimination, IntegralClosure,
               InverseSystems, LLLBases, PrimaryDecomposition,
               ReesAlgebra, TangentCone

i1 : loadPackage "StronglyStableIdeals";
i2 : QQ[t];
i3 : isHilbertPolynomial (4*t)
o3 = true
i4 : isHilbertPolynomial (5*t-6)
o4 = false
```

Gotzmann's and Macaulay's decompositions of a Hilbert polynomial can be computed using `gotzmannDecomposition` and `macaulayDecomposition`. These

methods return the list of terms in the decompositions. The summand $\binom{t+e}{c}$ is constructed with the command `projectiveHilbertPolynomial(c, c-e)`.

```
i5 : gotzmannDecomposition (4*t)
o5 = {P1, - P0 + P1, - 2*P0 + P1, - 3*P0 + P1, P0, P0}
o5 : List
i6 : macaulayDecomposition (4*t)
o6 = {- P0 + P1, 7*P0 - P1, - P1 + P2, - 10*P0 + 5*P1 - P2}
o6 : List
```

Finally, the saturated lexicographic ideal L with Hilbert polynomial $p(t)$ in the polynomial ring $\mathbb{K}[\mathbf{x}]$ can be computed with the method `lexIdeal` and its regularity is equal to the Gotzmann number of $p(t)$.

```
i7 : L = lexIdeal (4*t, QQ[x,y,z,w])
o7 = ideal (x, y5, y4z2)
o7 : Ideal of QQ[x, y, z, w]
i8 : regularity L == gotzmannNumber (4*t)
o8 = true
```

3. THE MAIN ALGORITHM. In this section, we outline the strategy of the main algorithm. This algorithm was firstly described in [Cioffi et al. 2011] and then optimized in [Lella 2012]. The same problem has been previously discussed in [Reeves 1992] and an alternative algorithm was later presented in [Moore and Nagel 2014].

We need to relate the properties of a strongly stable ideal with its Hilbert polynomial. If I is a strongly stable ideal, for each $s \in \mathbb{N}$ the monomial basis of the homogeneous piece I_s of the ideal is a subset of $\mathbb{T}_{n,s}$ closed by increasing elementary moves. We call *Borel sets* such subsets of $\mathbb{T}_{n,s}$ (see Figure 1 for an example). Proposition 1.2(i) implies that the monomial basis of I_s for a saturated strongly stable ideal $I \subset \mathbb{K}[\mathbf{x}]$ with Hilbert polynomial $p(t)$ and regularity at most s is a Borel set with $q(s) := \binom{n+s}{s} - p(s)$ elements. Thus, we consider the map

$$\left\{ \begin{array}{l} \text{saturated strongly stable ideals in } \mathbb{K}[\mathbf{x}] \text{ with} \\ \text{Hilbert polynomial } p(t) \text{ and regularity } \leq s \end{array} \right\} \leftrightarrow \left\{ \begin{array}{l} \text{Borel sets of } \mathbb{T}_{n,s} \\ \text{with } q(s) \text{ elements} \end{array} \right\}. \quad (2)$$

Moreover, Gotzmann's regularity theorem suggests considering s equal to the Gotzmann number of $p(t)$ to determine all saturated strongly stable ideals with Hilbert polynomial $p(t)$. Obviously, there are many Borel sets in $\mathbb{T}_{n,s}$ with $q(s)$ elements not corresponding to an ideal with Hilbert polynomial $p(t)$. To identify the image of the previous map, we recall a definition and a proposition by Mall.




Figure 1. The Borel sets defined in $\mathbb{T}_{2,3}$ and $\mathbb{T}_{2,4}$ by the ideal $(x_0^2, x_0 x_1, x_1^4) \subset \mathbb{K}[x_0, x_1, x_2]$.

Definition 3.1 [Mall 1997, Definition 2.7]. Let $B \subset \mathbb{T}_{n,s}$ be a Borel set. The set $B^{(i)} := \{\mathbf{x}^\alpha \in B \mid \min \mathbf{x}^\alpha = n - i\}$ is called the i -growth class of B . The sequence $\text{gv}(B) := (|B^{(0)}|, \dots, |B^{(n)}|)$ is called the growth vector of B .

Proposition 3.2 [Mall 1997, Proposition 3.2]. Let $I \subset \mathbb{K}[\mathbf{x}]$ be a strongly stable ideal generated by the monomials of a Borel set $B \subset \mathbb{T}_{n,s}$ and let $p(t)$ be its Hilbert polynomial. Then,

$$p(t) = \binom{n+t}{n} - \sum_{k=0}^n |B^{(k)}| \binom{k+t-s}{k}, \quad \text{for all } t \geq s. \quad (3)$$

We can use this result to determine the growth vector of a Borel set $B \subset \mathbb{T}_{n,s}$ starting from the Hilbert polynomial. The i -th difference polynomial of $p(t)$ is

$$(\Delta^i p)(t) = (\Delta^{i-1} p)(t) - (\Delta^{i-1} p)(t-1) = \binom{n+t-i}{n-i} - \sum_{k=i}^n |B^{(k)}| \binom{k+t-s-i}{k-i}.$$

Evaluating these identities at $t = s$, we obtain the linear system

$$\begin{cases} \sum_{k=0}^n |B^{(k)}| = \binom{n+s}{n} - p(s), \\ \vdots \\ \sum_{k=i}^n |B^{(k)}| = \binom{n+s-i}{n-i} - (\Delta^i p)(s), \\ \vdots \\ |B^{(n)}| = \binom{s}{0} - (\Delta^n p)(s), \end{cases} \quad (4)$$

whose solution is

$$|B^{(i)}| = \sum_{k=i}^n |B^{(k)}| - \sum_{k=i+1}^n |B^{(k)}| = \binom{n+s-i-1}{n-i} - (\Delta^i p)(s) + (\Delta^{i+1} p)(s), \quad i < n,$$

and $|B^{(n)}| = 1$ (recall that $(\Delta^i p)(t) \equiv 0$ for $i > \deg p(t)$ and $\deg p(t) < n$). Let us call the growth vector of $p(t)$ in degree s the solution of the linear system (4) and let us denote it by $\text{gv}_s(p(t))$.




Figure 2. Borel sets in $\mathbb{T}_{2,5}$ corresponding to the saturated strongly stable ideals $(x_0^3, x_0^2 x_1, x_0 x_1^4)$ (on the left) and $(x_0^3, x_0^2 x_1^2, x_0 x_1^3)$ (on the right) in 3 variables with Hilbert polynomial $t + 6$ and regularity at most 5.

Proposition 3.3 (cf. [Lella 2012, Theorem 3.3]). *Let $p(t)$ be a Hilbert polynomial. There is a bijective map*

$$\left\{ \begin{array}{l} \text{saturated strongly stable ideals} \\ \text{in } \mathbb{K}[\mathbf{x}] \text{ with Hilbert polynomial} \\ p(t) \text{ and regularity } \leq s \end{array} \right\} \xleftrightarrow{1:1} \left\{ \begin{array}{l} \text{Borel sets of } \mathbb{T}_{n,s} \\ \text{with } q(s) \text{ elements and} \\ \text{growth vector } \text{gv}_s(p(t)) \end{array} \right\}, \quad (5)$$

$$I \longrightarrow \text{monomial basis of } I_s,$$

$$\text{saturation of } (B) \longleftarrow B.$$

In order to determine the Borel sets of Proposition 3.3, we use a recursive algorithm based on Proposition 1.2(iii). Indeed, if $I \subset \mathbb{K}[x_0, \dots, x_n]$ is a strongly stable ideal with Hilbert polynomial $p(t)$ and B is the associated Borel set in $\mathbb{T}_{n,s}$, then the subset $B' = \{\mathbf{x}^\alpha \in B \mid \min \mathbf{x}^\alpha > n\} \subset B$ is a Borel set in $\mathbb{T}_{n-1,s}$ corresponding to the strongly stable ideal $I' = (x_n, I) \cap \mathbb{K}[x_0, \dots, x_{n-1}] \subset \mathbb{K}[x_0, \dots, x_{n-1}]$ with Hilbert polynomial $(\Delta p)(t)$.

Example 3.4. We want to determine the set of strongly stable ideals in the polynomial ring $\mathbb{K}[x_0, x_1, x_2]$ with regularity at most 5 defining schemes with Hilbert polynomial $p(t) = t + 6$. The Gotzmann number of $p(t)$ is 6 and its growth vector in degree 5 is $\text{gv}_5(t + 6) = (5, 4, 1)$. We start considering the set of strongly stable ideals in $\mathbb{K}[x_0, x_1]$ with Hilbert polynomial $\Delta p(t) = 1$ and regularity at most 5 corresponding to Borel sets with growth vector $\text{gv}_5(\Delta p(t)) = (4, 1)$. There is a unique Borel set

$$B' = \{x_0^5, x_0^4 x_1, x_0^3 x_1^2, x_0^2 x_1^3, x_0 x_1^4\}.$$

Since x_1^5 is not contained in B' , a Borel set $B \subset \mathbb{T}_{2,5}$ with growth vector $(5, 4, 1)$ does not contain monomials obtained from x_1^5 by applying decreasing elementary moves, i.e., $x_1^4 x_2$, $x_1^3 x_2^2$, $x_1^2 x_2^3$, $x_1 x_2^4$ and x_2^5 . Hence, we need to select five monomials divisible by both x_0 and x_2 producing a set closed by increasing elementary moves (see Figure 2).

Our package provides the method `stronglyStableIdeals` to compute the set of strongly stable ideals of a given polynomial ring with fixed Hilbert polynomial and bounded regularity.

```
i9 : stronglyStableIdeals (4*t, QQ[x,y,z,w])
o9 = {ideal (x^5, y^4, z^2), ideal (x^2*z, x*y, x^2, y^4, z^5, y^5),
      ideal (x*y, x^2, x*z^2, y^4), ideal (x*y, x^2, y^3)}
o9 : List
i10 : stronglyStableIdeals (4*t, QQ[x,y,z,w], MaxRegularity => 4)
o10 = {ideal (x*y, x^2, x*z^2, y^4), ideal (x*y, x^2, y^3)}
o10 : List
```

4. SEGMENT IDEALS. The transitive closure of the order relation

$$\mathbf{x}^\alpha >_B \mathbf{x}^\beta \iff \mathbf{x}^\beta = e_i^-(\mathbf{x}^\alpha) \quad (6)$$

induces a partial order on the set of monomials of any degree called the *Borel order*. Every graded term ordering is a refinement of this partial order. Since a Borel set B is closed with respect to the Borel order, i.e., $\mathbf{x}^\alpha >_B \mathbf{x}^\beta$, $\mathbf{x}^\beta \in B \Rightarrow \mathbf{x}^\alpha \in B$, it is natural to ask whether there exists a term ordering $<$ with the same property. For instance, for the lexicographic ideal, the graded lexicographic order separates, in each degree, monomials contained in the ideal from those outside. In [Cioffi et al. 2011], several notions of segment ideals are introduced.

Definition 4.1 [Cioffi et al. 2011, Definitions 3.1 and 3.7]. A Borel set $B \subset \mathbb{T}_{n,s}$ is called a *segment* if there exists a term ordering $<$ such that $\mathbf{x}^\alpha \succ \mathbf{x}^\beta$, for all $\mathbf{x}^\alpha \in B$ and $\mathbf{x}^\beta \in \mathbb{T}_{n,s} \setminus B$.

Let $I \subset \mathbb{K}[\mathbf{x}]$ be a saturated strongly stable ideal.

- (i) I is called a *hilb-segment* if the Borel set $I \cap \mathbb{T}_{n,r}$ is a segment, where r is the Gotzmann number of the Hilbert polynomial of I .
- (ii) I is called a *reg-segment* if the Borel set $I \cap \mathbb{T}_{n,m}$ is a segment, where m is the regularity of I .
- (iii) I is called a *gen-segment* if there exists a term ordering $<$ such that $\mathbf{x}^\alpha \succ \mathbf{x}^\beta$ for each minimal generator \mathbf{x}^α of degree s of I and for all $\mathbf{x}^\beta \in \mathbb{T}_{n,s} \setminus I_s$.

These notions are very important in the construction of flat families based on properties of Gröbner bases and in general for the study of the Hilbert scheme. The `StronglyStableIdeals.m2` package provides three methods for determining whether a strongly stable ideal may be some type of segment (and, in case, gives the term ordering). These methods use tools of the package `gfAnInterface.m2` and the term ordering is given as a weight vector.

```

i11 : sevenPointsP2 = stronglyStableIdeals (7, 3, MaxRegularity => 5)
o11 = {ideal (x2, x x, x5), ideal (x2, x4, x3),
        ideal (x x, x x, x3, x4)}
o11 : List
i12 : for J in sevenPointsP2 list isHilbSegment J
o12 = {(true, {7, 3, 1}), (false, ), (true, {4, 3, 1})}
o12 : List
i13 : for J in sevenPointsP2 list isRegSegment J
o13 = {(true, {7, 3, 1}), (false, ), (true, {4, 3, 1})}
o13 : List
i14 : for J in sevenPointsP2 list isGenSegment J
o14 = {(true, {6, 3, 1}), (true, {4, 3, 1}), (true, {4, 3, 1})}
o14 : List

```

SUPPLEMENT. Version 1.1 of `StronglyStableIdeals.m2` is contained in the online supplement.

REFERENCES.

- [Bertone et al. 2013a] C. Bertone, F. Cioffi, P. Lella, and M. Roggero, “Upgraded methods for the effective computation of marked schemes on a strongly stable ideal”, *J. Symbolic Comput.* **50** (2013), 263–290. MR Zbl
- [Bertone et al. 2013b] C. Bertone, P. Lella, and M. Roggero, “A Borel open cover of the Hilbert scheme”, *J. Symbolic Comput.* **53** (2013), 119–135. MR Zbl
- [Bertone et al. 2017a] C. Bertone, F. Cioffi, and M. Roggero, “Double-generic initial ideal and Hilbert scheme”, *Ann. Mat. Pura Appl.* **196**:1 (2017), 19–41. MR Zbl
- [Bertone et al. 2017b] C. Bertone, F. Cioffi, and M. Roggero, “Macaulay-like marked bases”, *J. Algebra Appl.* **16**:5 (2017), 1750100, 36. MR Zbl
- [Brachat et al. 2016] J. Brachat, P. Lella, B. Mourrain, and M. Roggero, “Extensors and the Hilbert scheme”, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* **16**:1 (2016), 65–96. MR Zbl
- [Cioffi and Roggero 2011] F. Cioffi and M. Roggero, “Flat families by strongly stable ideals and a generalization of Gröbner bases”, *J. Symbolic Comput.* **46**:9 (2011), 1070–1084. MR Zbl
- [Cioffi et al. 2011] F. Cioffi, P. Lella, M. G. Marinari, and M. Roggero, “Segments and Hilbert schemes of points”, *Discrete Math.* **311**:20 (2011), 2238–2252. MR Zbl
- [Galligo 1974] A. Galligo, “À propos du théorème de-préparation de Weierstrass”, pp. 543–579 in *Fonctions de plusieurs variables complexes*, edited by F. Norguet, Lecture Notes in Math. **409**, Springer, 1974. MR Zbl
- [Gotzmann 1978] G. Gotzmann, “Eine Bedingung für die Flachheit und das Hilbertpolynom eines graduierten Ringes”, *Math. Z.* **158**:1 (1978), 61–70. MR Zbl
- [Green 2010] M. L. Green, “Generic initial ideals”, pp. 119–186 in *Six lectures on commutative algebra*, edited by J. Elias et al., Birkhäuser Verlag, Basel, 2010. MR

- [Lella 2012] P. Lella, “An efficient implementation of the algorithm computing the Borel-fixed points of a Hilbert scheme”, pp. 242–248 in *ISSAC 2012—Proceedings of the 37th International Symposium on Symbolic and Algebraic Computation*, edited by J. van der Hoeven and M. van Hoeij, ACM, New York, 2012. MR Zbl
- [Lella and Roggero 2011] P. Lella and M. Roggero, “Rational components of Hilbert schemes”, *Rend. Semin. Mat. Univ. Padova* **126** (2011), 11–45. MR Zbl
- [Lella and Roggero 2016] P. Lella and M. Roggero, “On the functoriality of marked families”, *J. Commut. Algebra* **8**:3 (2016), 367–410. MR Zbl
- [Mall 1997] D. Mall, “Betti numbers, Castelnuovo Mumford regularity, and generalisations of Macaulay’s theorem”, *Comm. Algebra* **25**:12 (1997), 3841–3852. MR Zbl
- [Moore and Nagel 2014] D. Moore and U. Nagel, “Algorithms for strongly stable ideals”, *Math. Comp.* **83**:289 (2014), 2527–2552. MR Zbl
- [Reeves 1992] A. A. Reeves, *Combinatorial structure on the Hilbert scheme*, Ph.D. thesis, Cornell University, Ann Arbor, MI, 1992, Available at <https://search.proquest.com/docview/303996863>. MR

RECEIVED: 26 Jun 2014

REVISED: 22 Jun 2018

ACCEPTED: 4 Nov 2018

DAVIDE ALBERELLI:

davide.alberelli@gmail.com

PAOLO LELLA:

paolo.lella@polimi.it

Dipartimento di Matematica, Politecnico di Milano, Milano, Italy

<i>Strongly stable ideals and Hilbert polynomials</i>	1
Davide Alberelli and Paolo Lella	
<i>DiffAlg: a Differential algebra package</i>	11
Manuel Dubinsky, César Massri, Ariel Molinuevo and Federico Quallbrunn	
<i>Matroids: a Macaulay2 package</i>	19
Justin Chen	

