

dipartimento
economia
Marco Biagi



DEMB Working Paper Series

N. 131

On the financial connectedness of the commodity market: a replication of the Diebold and Yilmaz (2012) study

Francesco Caloia*, Andrea Cipollini*, Silvia Muzzioli***

September 2018

* Cà Foscari University of Venice, Department of Economics
Email: francesco.caloia@unive.it

** University of Palermo, Department of Economics, Accounting and Statistics
CEFIN, Centro Studi Banca e Finanza and RECent, Center for Economic Research - Unimore
Email: andrea.cipollini@unipa.it

*** University of Modena and Reggio Emilia
CEFIN, Centro Studi Banca e Finanza
Address: Viale Berengario 51, 41121, Modena, Italy
Email: silvia.muzzioli@unimore.it

ISSN: 2281-440X online



UNIMORE
UNIVERSITÀ DEGLI STUDI DI
MODENA E REGGIO EMILIA



Dipartimento di Economia Marco Biagi
Università degli studi di Modena e Reggio Emilia
Via Berengario 51 | 41121 Modena
tel. 059 2056711 | fax. 059 2056937
info.economia@unimore.it | www.economia.unimore.it

On the financial connectedness of the commodity market: a replication of the Diebold and Yilmaz (2012) study

Francesco Giuseppe Caloia^a
Department of Economics
Cà Foscari University of Venice

Andrea Cipollini^b
Department of Economics, Accounting and Statistics
University of Palermo

Silvia Muzzioli^c
Department of Economics
University of Modena and Reggio Emilia

Abstract:

In this paper we replicate the Diebold and Yilmaz (2012) study on the connectedness of the Commodity market and three other financial markets: the stock market, the bond market, and the FX market. We show that both the row and the column normalization schemes of the Generalized Forecast Error Variance Decomposition, suggested by the authors, lead to inaccurate measures of net contribution to risk transmission, in terms of ranking and sign. We show that, considering data generating processes characterized by different degrees of comovement and persistence, a scalar based normalization of the Generalized Forecast Error Variance Decomposition yields consistent (free of sign and ranking errors) net spillovers.

Keywords: causality; normalization schemes; generalized forecast error variance decomposition; spillover; simulation; Vector Autoregression Models.

JEL classification: C15; C53; C58; G17.

^a Department of Economics, Cà Foscari University of Venice, 873 Cannaregio, Fondamenta San Giobbe, 30121 Venezia. E-mail: francesco.caloia@unive.it.

^b Department of Economics, Accounting and Statistics, University of Palermo, V.le delle Scienze, 90128 Palermo. E-mail: andrea.cipollini@unipa.it, *CEFIN*, *RECent* Modena;

^c Corresponding author, Department of Economics, University of Modena and Reggio Emilia; *CEFIN*; Viale J. Berengario, 51, 41121 Modena. E-mail: silvia.muzzioli@unimore.it

1 Introduction

A normalization scheme is a set of one or more constraints to be imposed on a matrix such that the resulting scaled version will satisfy certain conditions. Equilibration, i.e. scaling a matrix such that its rows or columns sum to one is one of the most common normalization schemes. A normalization scheme is adopted either for estimation purposes or simply for interpretative purposes.

The aim of this paper is threefold. First we provide a review of the most common normalization schemes used in different financial applications, with a particular focus on forecast error variance decomposition. In fact, the implementation of the generalized forecast error variance decomposition yields a variance decomposition table that has to be normalized for interpretative purposes.

Second, we suggest a scalar based normalization scheme overcoming the limits of the traditional row-normalization scheme, used in Diebold and Yilmaz (2012)¹. The advantages and disadvantages of the normalization schemes are assessed through simulation, using data characterized by different degrees of comovement and persistence.

Third, we replicate the analysis in Diebold and Yilmaz (2012) in order to show how results change (in terms of net spillovers) as the normalization scheme changes. The results of the paper are intended to be useful not only for deriving spillovers measures, but also in any other field where a matrix normalization scheme is adopted, such as network analysis or spatial econometrics.

The structure of the paper is as follows. In Section 2 we provide an overview of the methodology suggested by Diebold and Yilmaz (2012) to construct a network graph and, in particular, indices of connectedness. In Section 3 we review the most common normalization schemes used in various fields. Section 4 highlights how persistence and comovement among the series affect the results of the spillover analysis based on different normalization schemes. Section 5 replicates the study by Diebold and Yilmaz (2012). The final section concludes.

2. Networks and connectedness

Networks are usually represented in graphs, where nodes and edges are graphically displayed. A weighted network is a network that allows for weights on the edges in order to represent stronger or weaker connections between nodes, while direct networks are networks that allow for asymmetries.

One example of a weighted and direct network (also varying across forecast horizons) is the forecast error variance decomposition (FEVD).

¹ Diebold and Yilmaz (2009) is the first study to compute a total spillover index based on the forecast error variance decomposition (FEVD) by using the Cholesky decomposition of the VAR residuals covariance matrix. It is well known that the Cholesky decomposition is sensitive to the variable ordering. Klobner and Wagner (2013) provide an algorithm for swiftly calculating the spillover index's maximum and minimum over all renumerations.

Forecast error variance decomposition is a standard econometric tool used in multivariate time series analysis to assess the contribution in terms of forecast error variance of each variable due to a shock to any of the other variables. Consider a covariance stationary VAR(p) with k endogenous variables:

$$x_t = A_1 x_{t-1} + \dots + A_p x_{t-p} + \varepsilon_t \quad (1)$$

where ε_t are i.i.d. disturbances with contemporaneous covariance matrix Σ . In order to derive the moving average representation of the VAR(p), we rewrite (1) as a first order system:

$$\begin{bmatrix} x_t \\ x_{t-1} \\ \vdots \\ \vdots \\ x_{t-p+1} \end{bmatrix} = \begin{bmatrix} A_1 & A_2 & & A_p \\ I & 0 & & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & & 0 \\ 0 & 0 & & I \end{bmatrix} \begin{bmatrix} x_{t-1} \\ x_{t-2} \\ \vdots \\ \vdots \\ x_{t-p} \end{bmatrix} + \begin{bmatrix} \varepsilon_t \\ 0 \\ \vdots \\ \vdots \\ 0 \end{bmatrix} \quad (2)$$

which can be written in compact form:

$$\bar{x}_t = \bar{A} \bar{x}_{t-1} + \bar{\varepsilon}_t \quad (3)$$

where \bar{A} is the $kp \times kp$ companion matrix. Then, using the selection matrix $\bar{e} = \begin{bmatrix} I & \bar{0} & \dots & \bar{0} \end{bmatrix}$ which has k rows and kp columns (and the first k rows and columns are given by the identify matrix I),

we obtain the $k \times k$ moving average coefficient matrix Ψ_h :

$$\Psi_h = \bar{e} \bar{A}^h \bar{e}' \quad (4)$$

Given a set of k endogenous variables, Diebold and Yilmaz (2012; 2014) use the Pesaran and Shin (1998) "generalized" approach (GFEVD) that allows shocks to be correlated (which is insensitive to variable ordering,) to obtain the element in row i and column j of the connectedness matrix proxied by FEVD. More specifically, the contribution of the j -th shock to the h step ahead forecast error variance of the i -th endogenous series is computed as follows:

$$\theta_{ij}^g = \frac{\sigma_{jj}^{-1} \sum_{l=0}^{h-1} (e_i' \Psi_l \Sigma e_j)^2}{\sum_{l=0}^{h-1} (e_i' \Psi_l \Sigma \Psi_l' e_i)} \quad (5)$$

where σ_{jj} the standard deviation of the disturbance of the j th equation, and e_i is the selection vector with one as the i th element and zeros otherwise.

Due to the non-orthogonality of shocks, the sum of the contributions to the forecast error variance (i.e. the row sum) is not equal to one². The authors, therefore, propose a row-normalization of the values of the variance decomposition in equation (5), in order to interpret its elements as variance shares:

$$\bar{\theta}_{ij}^g = \frac{\theta_{ij}^g}{\sum_{j=1}^k \theta_{ij}^g} \quad (6)$$

They also mention the equivalency between this row-normalization scheme and the alternative column-normalization scheme (see Section 3).

Diebold and Yilmaz (2014) rely on the absolute value of net pairwise spillovers when graphing the network, which is obtained from generalized forecast error variance decomposition. Therefore in this case, the network graph depends on the chosen normalization scheme for the GFEVD. Moreover, the summary descriptive statistics of the network are given by the directional connectedness indices. In particular, the directional spillover received by each market from all the other markets (FROM others) is computed as the off-diagonal row sum; the spillover transmitted by each market to all the other markets (TO others) is computed as the off-diagonal column sum. A measure of net contribution (NET) of each market is obtained as the difference between the directional spillovers TO others and FROM others. In this way we are able to distinguish markets that are net donors from those that are net receivers in terms of risk transmission.

Consequently, not only the network graph, but also the summary statistics described by the directional connectedness indices depend on the chosen normalization scheme for the GFEVD. Given the critical role of the normalization scheme, in the next section we review the different normalization schemes proposed in the literature.

3. Normalization schemes

In this section, we review the most used normalization schemes in GFEVD and spatial regression models. While GFEVD, allowing for the construction of weighed directed network, relies mainly on row normalization, spatial regression models, mainly used to obtain weighed networks, allowing only for symmetric linkages, rely on other normalization schemes. More specifically, consider a standard generalised spatial autoregressive model of order p , or simply SAR(p) model:

$$u = \sum_{h=1}^p \phi_h W_h u + \varepsilon \quad (7)$$

² The moving average coefficients necessary to computed the spillover indices are obtained through the estimation of a traditional VAR in Diebold and Yilmaz (2012). Cipollini et al. (2017) and Fengler et al. (2018) obtain the moving average coefficients through the estimation of a fractionally integrated VAR and of a multivariate GARCH, respectively.

Where $W = (w_{i,j})$ for $i, j = 1, \dots, k$ is the spatial weight matrix and ϕ_h are autoregressive parameters. Equivalently, we can rewrite equation (7) as follows:

$$u = \left(I_N - \sum_{h=1}^p \phi_h W_h \right)^{-1} \varepsilon \quad (8)$$

We recall, in the following, the different normalization schemes of the spatial weight matrix W used in the literature to make $(I_N - \sum_{h=1}^p \phi_h W_h)$ non-singular. These schemes can be applied also to GFEVD.

3.1 Row normalization

Given a $(k \times k)$ unscaled matrix $W^* = (w_{ij}^*)$, we can obtain the corresponding row-stochastic matrix $W = (w_{ij})$ by row-normalizing W^* such that:

$$w_{ij} = \frac{w_{ij}^*}{\sum_{j=1}^k w_{ij}^*} \quad (9)$$

The resulting matrix W has row sums equal to one. However, row normalization is not a restrictive task since the same result can be achieved by constraining the parameter space of the autoregressive parameters ϕ_h (Caporin and Paruolo (2015)); as a result, the normalization task would be absorbed by the AR parameter through scaling.

Moreover, this normalization is useful in interpreting spatial weight matrices, whose elements can be thought of as a fraction of all spatial influence. This interpretative advantage also applies for a forecast error variance decomposition that does not rely on Cholesky factorization (or any other identifying scheme of structural VAR models) so that the matrix coefficients can be interpreted as variance shares. This is the normalization scheme proposed by Diebold and Yilmaz (2012) when using the generalized forecast error variance decomposition. However, this scheme also has certain drawbacks: by scaling the elements of each row by the corresponding row sum, the order of magnitude is preserved only by row.

3.2 Column normalization

This scheme is specular to the row-normalization scheme described above. The only difference is that the normalization is done by column: in this case only the columns sum to one. The critical issues concerning the row-normalization scheme apply also in this case. Note that for the variance decomposition Diebold and Yilmaz (2012) suggest this normalization scheme as an alternative to row normalization.

3.3 Max row normalization

In this normalization scheme, the normalization factor is a scalar equal to the maximum row sum of the unscaled matrix W^* , then the scaled matrix is obtained as $W = W^*/k$ where:

$$k = \max(r_1, \dots, r_k) \quad (10)$$

and:

$$r_i = \sum_{j=1}^k w_{ij}^* \quad (11)$$

where w_{ij}^* is the element in row i and column j of the unscaled matrix W^* . This scheme is characterized by a single normalization factor instead of the k factors of the row normalization scheme (one for each row). As a result, it preserves the magnitude relation among the elements of rows and columns and column and row values can therefore be safely compared. Moreover, it allows for a comparison between different rows and column sums, making it possible to distinguish between stronger or weaker influences. As argued by Billio et al. (2016) it is also possible to normalize by the maximum row sum over time in order to compare spatial weight matrices in different time periods while preserving a reasonable magnitude of autoregressive parameters.

3.4 Max column normalization

This scheme is specular to the max row normalization described above: the only difference is that the scalar is equal to the maximum column sum of the unscaled matrix W^* . The same advantages of the max row normalization apply.

3.5 Spectral radius normalization

Let W^* be the $(k \times k)$ positive unscaled matrix and let $\{\lambda_1, \dots, \lambda_k\}$ be the eigenvalues of W^* . The spectral radius is the maximum eigenvalue (in modul), formally:

$$\tau = \max\{|\lambda_1|, |\lambda_2|, \dots, |\lambda_k|\} \quad (12)$$

The scalar normalization factor is set equal to the spectral radius and the scaled matrix W is therefore obtained as follows:

$$W = W^*/\tau \quad (13)$$

Under the Perron and Frobenius theorem, the spectral radius satisfies the following inequalities:

$$\min_i \sum_{j=1}^N w_{ij} \leq \tau \leq \max_i \sum_{j=1}^N w_{ij} \quad (14)$$

As a result, some row sums and column sums exceed unity, while others can be less than one. This normalization scheme therefore has one main drawback: the elements can no longer be interpreted as fractions of the overall influence (e.g. the sum by row and by column).

Nevertheless, this normalization scheme is widely used in spatial econometrics: in fact, following LeSage and Pace (2010) a matrix W^* can be transformed to have maximum eigenvalue equal to one using $W = W^* \max(\lambda_{W^*})$, and this is a desirable property because it constrains the autoregressive parameter to

have maximum possible value equal to one. In particular, Kelejian and Prucha (2010) show that $(I_N - \sum_{h=1}^p \phi_h W_h)$ is non-singular for all the values of the parameter space in the interval $(-1 ; 1)$.

4 Comparison of the normalization schemes in GFEVD

In this section, we show how the normalization schemes reviewed in section 3 affect the GFEVD, by using data generating processes characterized by different degrees of comovement and persistence. The review of different normalization schemes in section 3 shows that row normalization has interpretative limits and, in this framework, leads to misspecified spillover measures. In particular:

- If the normalization is carried out by row, the column sum is not necessarily equal to one. As a result, while FROM directional spillovers can be interpreted as a fraction of the total variance received via spillovers, TO directional spillovers lack this kind of interpretation (some column sums are above unity, while some others are beyond unity).
- Normalization by row implies that the order of magnitude of the entries of the variance decomposition table is preserved only by row. As a result, NET spillovers are obtained as the difference between two values incomparable in magnitude.

We now show that the values of the spillover measures are sensitive to this normalization choice, leading to misspecified measures of net contribution (NET). We consider four cases: a) LL (Low Persistence; Low Comovement); b) LH (Low Persistence; High Comovement); c) HL (High Persistence; Low Comovement); d) HH (High Persistence; High Comovement), according to the different setup of the VAR model described in eq. (1). We set the number of endogenous variables, k , equal to five. The model configurations differ for the coefficient matrices in the lag operator $A(L)$ and of the covariance matrix $\Sigma = P P'$. In particular, the Low Comovement case is defined by using a lower triangular matrix P set as follows:

$$P = \begin{bmatrix} 0.10 & 0 & 0 & 0 & 0 \\ 0.15 & 0.15 & 0 & 0 & 0 \\ 0.20 & 0.20 & 0.20 & 0 & 0 \\ 0.25 & 0.25 & 0.25 & 0.25 & 0 \\ 0.30 & 0.30 & 0.30 & 0.30 & 0.30 \end{bmatrix} \quad (15)$$

while the High Comovement case is defined by using the following lower triangular matrix P :

$$P = \begin{bmatrix} 0.40 & 0 & 0 & 0 & 0 \\ 0.45 & 0.45 & 0 & 0 & 0 \\ 0.50 & 0.50 & 0.50 & 0 & 0 \\ 0.55 & 0.55 & 0.55 & 0.55 & 0 \\ 0.60 & 0.60 & 0.60 & 0.60 & 0.60 \end{bmatrix} \quad (16)$$

where a scalar equal to 0.3 has been added to all nonzero entries of matrix P in equation (15). In this way, the resulting variance-covariance matrix has higher variance and covariance entries and, thus, higher comovements, while maintaining the underlying correlation between the endogenous variables the same.

To ensure a stationary VAR(p) (e.g. with roots of the characteristic polynomial A(L) outside the unit circle) characterised by Low Persistence, we consider a VAR(2) with coefficient matrices A₁ and A₂ with values equal to 0.05. A stationary VAR(p) characterised by High Persistence is a restricted VAR(22) given by the parsimonious Vector HAR representation with coefficient matrices A^(d), A^(w), A^(m) described as follows: A^(d) with values equal to 0.05, A^(w) with values equal to -0.02 and A^(m) with values equal to 0.01³.

Consequently, we compute the generalized forecast error variance decomposition as defined by equation (5) and we obtain the measures of NET contribution. Formally, the non-normalized NET spillovers for the forecast horizon *h*, which are taken as benchmark, are obtained as follows:

$$NET_i(h) = DS_{\bullet \rightarrow}^g(h) - DS_{\rightarrow \bullet}^g(h) \quad (17)$$

where:

$$DS_{\bullet \rightarrow}^g(h) = \sum_{\substack{i=1 \\ i \neq j}}^K \theta_{ij}^g \quad ; \quad DS_{\rightarrow \bullet}^g(h) = \sum_{\substack{j=1 \\ j \neq i}}^K \theta_{ij}^g \quad (18)$$

where $DS_{\bullet \rightarrow}^g$ denotes the non-normalized directional spillover transmitted by the market *i* to all other markets *j* (named TO others), while $DS_{\rightarrow \bullet}^g$ denotes the non-normalized directional spillover received by market *i* from all the other markets *j* (named FROM others). Second, we compute the \overline{NET} spillovers obtained from the forecast error variance decomposition normalized by the different schemes:

$$\overline{NET}_i(h) = \overline{DS}_{\bullet \rightarrow}^g(h) - \overline{DS}_{\rightarrow \bullet}^g(h) \quad (19)$$

where the over bar denotes the normalized spillovers. These normalized measures are compared to the benchmark spillovers in equation (5). The comparison is intended to assess the reliability of the different normalization schemes both in terms of order of ranking (to assess which market is the largest net contributor to the total connectedness) and in terms of sign (to distinguish net donors from net receivers).

³ In the Vector HAR model the matrices A^(d), A^(w) and A^(m) are coefficient matrices associated with the three terms of daily, weekly and monthly partial volatility components, respectively. In particular, the Vector HAR model can be written as follows:

$$x_t^{(d)} = c + \phi^{(d)} x_{t-1}^{(d)} + \phi^{(w)} x_{t-1}^{(w)} + \phi^{(m)} x_{t-1}^{(m)} + \varepsilon_t$$

where x_t are daily volatilities, while the terms representing the weekly and monthly volatilities are obtained as the arithmetic average of the daily volatilities recorded in the last week and the last month, respectively.

4.1 Results based on population parameters

In this section we cast light on how the choice of the normalization scheme can affect the ranking and the sign of the NET spillovers, by means of an introductory example. Moreover, in order to show how the spillover tables change for different forecast horizons, two different horizons are reported: the two-day horizon is reported in the upper panel of every Table, while the lower panel contains the ten-day forecast horizon.

For this introductory example we report the results based on the population parameters for the “High Persistence, High Comovement” scenario, which is the most illuminating one. Table 1 shows the spillover table based on the non-normalized forecast error variance decomposition which is taken as a benchmark. Tables 2 to 6 show the same spillover table after applying the different normalization schemes outlined in Section 3 (Table 2 for row normalization, Table 3 for column normalization, Table 4 for normalization by spectral radius, Table 5 for normalization by maximum row sum, Table 6 for normalization by maximum column sum). These Tables show the directional spillover received from others (FROM others), the directional spillover received from others including own (FROM others including own), the directional spillover transmitted to others (TO others), the directional spillover transmitted to others including own (TO others including own), and the net contribution (NET) defined as the difference between the directional spillover transmitted TO others and the directional spillover received FROM others for each variable V_i for $i = 1, \dots, 5$. The tables also show the sign of the NET spillover (NET sign): negative if the market is the net receiver and positive if the market is the net donor, and the ranking of the NET spillover from the highest to the lowest (NET ranking).

In Table 2 we show the standard row-normalization scheme proposed by Diebold and Yilmaz (2012) which has the interpretative advantage that the directional spillovers received FROM others including own sum to one, and as a result each element of the forecast error variance decomposition matrix can be interpreted as a variance share (by row). For example, in the upper panel variable 2 receives the most from variable 3 (0.219), and the least from variable 5 (0.140). Moreover, variable 1 represents the market least affected by the others (FROM others=0.653), while variable 3 represents the market most affected by the others (FROM others=0.705).

On the contrary in Table 3 all the columns (TO others including own) sum to one: each element of the forecast error variance decomposition matrix can be interpreted as the fraction of total variance transmitted. For example, in the upper panel of Table 3 variable 2 gives the least to variable 5 (0.131), and the most to variable 3 (0.216). Moreover, variable 3 represents the market that transmits the most to the others (TO others=0.714), while variable 1 represents the market that transmits the least to others (TO others=0.592). In the case of the column normalization, the focus is on how much one variable (market or country) affects the system. Despite the neat interpretation, the row normalization or column normalization schemes affect the NET spillovers, which may have the opposite sign and the wrong ranking if compared to the non-normalized ones. In fact, the first variable in the column

normalization scheme (Table 3) is misconceived as the net donor, while it is a net receiver in the non-normalized case (Table 1), whereas variables 3 and 4 are mistakenly considered as net receivers instead of net donors, as apparent in Table 1. The same happens in the row normalization scheme (Table 2): for example, variable 2 is misconceived as net donor in the two-day forecast horizon, while it is a net receiver in the non-normalized case (Table 1). As a result, also there is a change in the ranking of the variables (ranging from the one giving the most to the system, that is the major net donor and has rank 1, to the variable receiving the most from the system, which is the major net receiver and has rank 5). For example in the non-normalized case the variable transmitting the most to the system is variable 5 (for both forecast horizons), but in the row-normalized case it emerges that the variable transmitting the most to the system is variable 3 for the two-day forecast horizon and variable 4 for the ten-day horizon.

Tables 4 to 6 show the scalar-normalization cases. The scalar factors applied are: the spectral radius (Table 4), the maximum row sum (Table 5) and the maximum column sum (Table 6). In the spectral radius normalization, it is not possible to interpret each element of the forecast error variance decomposition matrix as variance shares by column, or by row. In fact, the sum by row and by column (FROM others including own and TO others including own) can attain values higher or lower than 1, given the mathematical property of the maximum eigenvalue described in eq. (16). Despite the lack of interpretability in terms of variance shares, all the net spillovers maintain the correct sign after normalization and the correct ranking as in the non-normalized case.

It may be noted that in the maximum row sum normalization scheme in Table 5 and in the maximum column sum normalization scheme in Table 6, the only values which sum to one are those in the row with the maximum sum (the third row in both Panels of Table 5) and those in the column with the maximum sum (column 3 for Panel A and column 4 for Panel B of in Table 6), respectively. Only for these values is it possible to give a percentage interpretation: in Table 5 it may be seen that for $h=2$ variable 3 receives 70.5% FROM others, while variable 3 in Table 6 Panel A transmits 71.4% TO others.

In conclusion, it may be stated that the max row sum and max col sum normalization are slightly better than the spectral radius since they can preserve the ranking and the sign of the spillovers and, at least for one variable, they can preserve their interpretation as variance share.

As Tables 2 to 6 focus on the normalization issue for only the high correlated and high persistence scenario, in Table 7 and Table 8 we show the results based on population parameters for all the other scenarios: by looking at the sign of the net spillovers (Table 7) it is clear that the row-normalization scheme performs fairly well with no errors in sign for the horizon $H=2$ and only one error in sign in each of the high-correlated scenarios: on the contrary in each scenario the column normalization produces from 1 to 3 errors in sign. By looking at the ranking errors in Table 8, what emerges is that both the row normalization and the column normalization scheme affect the ranking in most cases. On the other hand, any scalar normalization scheme does not affect the ranking.

4.2 Results based on simulation

In order to account for the role played by parameter estimation on the rank and sign of net spillovers, we simulate a multivariate dynamic system, using the DGP given by eq. (1). The shocks ε_t are given by $P \eta_t$, where η_t are iid Gaussian and orthogonal innovations with unit variance. In order to assess the reliability of the different normalization schemes in preserving the order of magnitude and the sign of net contributions (NET spillovers) obtained from the generalized forecast error variance decomposition, the simulation experiment involves the following steps:

1) Five artificial data series (where the time series dimension is equal to 500) are obtained by simulating either the VAR(2) (in the case of Low Persistence) or the restricted VAR(22) (in the case of High Persistence) with Gaussian innovations. The coefficient matrices for the lags and the lower triangular matrices P aiming at capturing the different degrees of contemporaneous comovement are those used in section 5.

2) For each replication, we estimate the model parameters by OLS, obtaining the impulse-responses for the forecast horizons $h = 2$, $h = 10$ and computing the corresponding generalized forecast error variance decomposition as defined in eq. (5).

After obtaining the simulated datasets, we compare the non-normalized matrix W^* (e.g. the non-normalized variance decomposition table for a given forecast horizon) and the five normalized matrices W (e.g. the normalized variance decomposition table for a given forecast horizon) in terms of sign and ranking errors.

First, we measure the number of errors in the sign of the net spillovers. Errors are counted when the net spillover obtained from the normalized matrix has a sign opposite to that of the net spillover obtained from the non-normalized matrix. The total number of possible errors is 5000 for each scenario (5 variables times 1000 simulations for each scenario).

Second, we measure the errors in the ranking. Errors are counted when the ranking of the net spillovers obtained from the normalized matrix is different from that of the non-normalized matrix. The total number of possible errors is 1000 for each scenario (one ranking times 1000 simulations for each scenario). Results are shown in Table 9 for sign errors and in Table 10 for ranking errors.

Table 9 shows that over a total number of 5000 possible errors for each scenario (5 variables times 1000 simulations for each scenario), the row normalization performs much better than the column normalization for each scenario: in fact, for $H=2$ ($H=10$) the average number of errors in sign is about 354 (169) for the row-normalization scheme and about 2525 (1997) for the column normalization scheme. This result is surprising since the row normalization and column normalization schemes should theoretically be equal. In both normalization schemes, the number of errors increases with the degree of comovement. On the contrary, the sum of errors in sign in the low persistence scenarios is slightly higher than the sum of the same errors in the high persistence scenarios for both forecast horizons.

Moreover, as shown in Table 10, the row-normalization proposed by Diebold and Yilmaz (2012) and the alternative column normalization schemes affect the ranking of the spillovers more than 850 times out of 1000 for $H=2$ and more than 950 times out of 1000 for $H=10$ (with the sole exception of the row normalization scheme in the high persistence scenarios).

To conclude, even if the row normalization scheme and the column normalization scheme allow for a better interpretation of the values of the generalized forecast error variance decomposition, there is a need to be cautious in interpreting the resulting net spillovers that should discriminate markets which are net donors from those which are net receivers. On the contrary, any scalar normalization scheme (by maximum eigenvalue, maximum row sum or maximum column sum) will outperform the traditional normalization schemes, preserving the ranking and the sign of the NET spillovers. As a result, we suggest using a scalar normalization scheme to derive the correct measures of net contribution. Among the scalar normalization schemes, the maximum row sum or the maximum column sum are preferred to the spectral radius since they allow for a better interpretation of how much one variable receives or transmits in terms of percentage values.

5 Replication of Diebold and Yilmaz (2012)

In order to shed further light on the normalization issue in obtaining reliable spillover measures we replicate the original paper of Diebold and Yilmaz (2012) to see whether results would have changed if a scalar normalization were applied. Therefore we compute the 10-days-ahead generalized forecast error variance decomposition, on a VAR of order 4, by using the same daily data as in Diebold and Yilmaz (2012). The data consists of range based volatilities of the S&P 500 (stock market), the 10 years Treasury bond yield (bond market), the New York Board of Trade US dollar index futures (FX market) and the Dow Jones UBS Commodity index (commodity market), recorded from January 25, 1999 to January 29, 2010. Full sample results are presented in Table 11.

Panel I corresponds to Table 2 in the Diebold and Yilmaz (2012) paper (variance decomposition by using row normalization) while Panel II displays the variance decomposition by using scalar normalization (the max row sum). By comparing the NET spillovers in Panels I and II, we can see that the stock market turns out to be a net donor while all the others variables are net receivers of volatility spillovers during the entire period. As a result there are no sign errors in Diebold and Yilmaz (2012), even if the row-normalization scheme is used in place of the max row sum one. However, we can detect some ranking errors. If we concentrate on the bond market, it turns out to have received less volatility spillovers than the commodity market if the row-normalization scheme is used, as in Diebold and Yilmaz (2012). However, if we use scalar normalization, as in Panel II, we can see that the result is the opposite: the bond market turns out to have received more volatility spillovers than the commodity market. As a result we have a ranking error if the row-normalization scheme is used. This result is not surprising: since Diebold and Yilmaz (2012) consider four distinct asset classes and use a short-order VAR(p)

model, the results of their application can be “classified” in our “low-persistence, low-comovement” case that, in fact, recorded the lowest number of sign errors. On the contrary, from our simulation results, it is evident that ranking errors are frequent in data of any degree of comovement and persistence.

6 Concluding remarks

In this study we replicate the full sample results of Diebold and Yilmaz (2012) applied to range based volatilities of the Dow Jones UBS Commodity index and of three other financial markets: the S&P 500 (stock market), the 10 years Treasury bond yield (bond market), the New York Board of Trade US dollar index futures (FX market). Moreover, we show that the net spillover indices used to assess the net contribution of one market to systemic risk are sensitive to the normalization scheme. In particular, the row normalization scheme of the Generalized Forecast Error Variance Decomposition suggested by Diebold and Yilmaz (2012), although allowing for a better interpretation (as variance shares) of the results, may fail to establish whether the market is a net risk transmitter or net risk receiver. Moreover, it is also unable to assess the degree to which a single market influences all the others in net absolute terms. The column normalization scheme, suggested by Diebold and Yilmaz (2012), suffers from the same drawbacks. As a result, we suggest using a scalar normalization scheme (as in Caloia et al. 2018) to avoid the misspecification of results. Among the scalar normalization schemes, the maximum row sum or the maximum column sum schemes are preferable to the spectral radius since they allow for a better interpretation of how much one variable receives or transmits in terms of percentage values.

Acknowledgements.

The authors gratefully acknowledge financial support from FAR2017 project “ The role of Asymmetry and Kolmogorov equations in financial Risk Modelling (ARM)”, from Fondazione Cassa di Risparmio di Modena, for the project 2015.0333 “Volatility and higher order moments: new measures and indices of financial connectedness” and from the FAR2015 project “A SKEWness index for Europe (EU-SKEW)”. The usual disclaimer applies.

References:

- Billio, M., Caporin, M., Frattarolo L. & Pelizzon, L. (2016). Networks in risk spillovers, a multivariate GARCH perspective. *Working Paper*, Department of Economics, Cà Foscari University of Venice.
- Caporin, M. & Paruolo, P. (2015). Proximity structured multivariate volatility models. *Econometric Reviews*, 34(5) p. 559-593 .
- Caloia, F.G., Cipollini, A. & Muzzioli, S. (2018). Symmetric semi-volatility spillover effects in EMU stock markets. *International Review of Financial Analysis*, 57, p. 221-230.
- Diebold, F. X., & Yilmaz, K. (2009). Measuring financial asset return and volatility spillovers, with application to global equity markets. *Economic Journal*, 119(534), p. 158–171.
- Diebold, F. X., & Yilmaz, K. (2012). Better to Give than to Receive: Predictive Directional Measurements of Volatility Spillovers. *International Journal of Forecasting*, 28 (1), p. 57-66.
- Diebold, F. X., & Yilmaz, K. (2014). On the Network Topology of Variance Decompositions: Measuring the Connectedness of Financial Firms. *Journal of Econometrics*, 182(1), p. 119-134.
- Fengler, M.R., & Herwartz, H. (2018). Measuring Spot Variance Spillovers when (Co)variances are Time-varying – The Case of Multivariate GARCH Models. *Oxford Bulletin of Economics and Statistics*, 80(1), p. 135-159.
- Kelejian, H. H. & Prucha, I. R. (2010). Specification and estimation of spatial autoregressive model with autoregressive and heteroskedastic disturbances. *Journal of Econometrics*, 157(1), p. 53-67.
- Keiler, S. & Eder, A. (2013). CDS spreads and systemic risk: a spatial econometric approach. *Discussion Paper 01/2013, Deutsche Bundesbank*.
- Klobner, S. & Wagner, S. (2013). Exploring all VAR orderings for calculating spillovers? Yes, we can!-A note on Diebold and Yilmaz (2009). *Journal of Applied Econometrics*, 29, p. 172-179.
- LeSage, J. & Pace, R. K. (2010). *Introduction to Spatial Econometrics*. Chapman and Hall/CRC.
- Pesaran, H. & Shin, Y. (1998). Generalized Impulse Response in Linear Multivariate Models. *Economic Letters*, 58, p. 17-29.

Table 1. Spillover Table based on the non-normalized variance decomposition table (VDT).

Panel A: h=2							
	V1	V2	V3	V4	V5	FROM others including own	FROM others
V1	0.889	0.539	0.435	0.376	0.324	2.561	1.672
V2	0.493	0.975	0.683	0.535	0.438	3.124	2.149
V3	0.337	0.668	0.994	0.758	0.612	3.369	2.375
V4	0.255	0.506	0.753	0.997	0.801	3.313	2.316
V5	0.204	0.406	0.605	0.802	0.997	3.015	2.018
TO others including own	2.178	3.094	3.470	3.467	3.172		
TO others	1.289	2.119	2.477	2.470	2.175		
NET	-0.383	-0.030	0.102	0.154	0.157		
NET sign	-	-	+	+	+		
NET ranking	5	4	3	2	1		
Panel B: h=10							
	V1	V2	V3	V4	V5	FROM others including own	FROM others
V1	0.773	0.579	0.540	0.506	0.453	2.850	2.077
V2	0.483	0.940	0.706	0.582	0.490	3.202	2.262
V3	0.343	0.671	0.984	0.769	0.630	3.397	2.413
V4	0.263	0.516	0.759	0.993	0.804	3.334	2.341
V5	0.211	0.416	0.614	0.806	0.992	3.040	2.048
TO others including own	2.073	3.122	3.603	3.656	3.368		
TO others	1.300	2.181	2.619	2.663	2.377		
NET	-0.777	-0.080	0.206	0.322	0.329		
NET sign	-	-	+	+	+		
NET ranking	5	4	3	2	1		

Note. This figure shows the spillover Table based on the non-normalized generalized forecast error variance decomposition, which is displayed in the central frame. Results refer to the HH scenario (high-persistent and high-correlated series) and to the forecast horizon $h=2$ (Panel A) and $h=10$ (Panel B). The Table shows the directional spillover received from others (FROM others), the directional spillover received from others including own (FROM others (including own)) the directional spillover transmitted to others (TO others), the directional spillover transmitted to others including own (TO others (including own)) and the net contribution (NET) defined as the difference between the directional spillover transmitted TO others and the directional spillover received FROM others for variable V_i , $i=1, \dots, 5$. The bottom lines show for each variable the sign of the NET spillover (NET sign): negative if the variable is a net receiver and positive if the variable is a net donor, and the ranking based on the value of the NET spillover from the highest to the lowest (NET ranking).

Table 2. Spillover Table based on the row-normalized variance decomposition table (VDT).

Panel A: h=2								
	V1	V2	V3	V4	V5	FROM others including own	FROM others	
V1	0.347	0.210	0.170	0.147	0.126	1	0.653	
V2	0.158	0.312	0.219	0.171	0.140	1	0.688	
V3	0.100	0.198	0.295	0.225	0.182	1	0.705	
V4	0.077	0.153	0.227	0.301	0.242	1	0.699	
V5	0.068	0.135	0.201	0.266	0.331	1	0.669	
TO others including own	0.750	1.008	1.112	1.110	1.021			
TO others	0.403	0.696	0.817	0.809	0.690			
NET	-0.250	0.008	0.112	0.110	0.021			
NET sign	-	+	+	+	+			
NET ranking	5	4	1	2	3			
Panel B: h=10								
	V1	V2	V3	V4	V5	FROM others including own	FROM others	
V1	0.271	0.203	0.189	0.178	0.159	1	0.729	
V2	0.151	0.294	0.221	0.182	0.153	1	0.706	
V3	0.101	0.198	0.290	0.226	0.185	1	0.710	
V4	0.079	0.155	0.228	0.298	0.241	1	0.702	
V5	0.070	0.137	0.202	0.265	0.326	1	0.674	
TO others including own	0.671	0.986	1.129	1.149	1.065			
TO others	0.400	0.692	0.840	0.851	0.738			
NET	-0.329	-0.014	0.129	0.149	0.065			
NET sign	-	-	+	+	+			
NET ranking	5	4	2	1	3			

Note. This figure shows the spillover Table based on the row-normalized generalized forecast error variance decomposition, which is displayed in the central frame. Results refer to the HH scenario (high-persistent and high-correlated series) and to the forecast horizon $h=2$ (Panel A) and $h=10$ (Panel B). The Table reports the directional spillover received from others (FROM others), the directional spillover received from others including own (FROM others (including own)) the directional spillover transmitted to others (TO others), the directional spillover transmitted to others including own (TO others (including own)) and the net contribution (NET) defined as the difference between the directional spillover transmitted TO others and the directional spillover received FROM others for variable V_i , $i=1, \dots, 5$. The bottom lines show for each variable the sign of the NET spillover (NET sign): negative if the variable is a net receiver and positive if the variable is a net donor, and the ranking based on the value of the NET spillover from the highest to the lowest (NET ranking).

Table 3: Spillover Table based on the column-normalized variance decomposition table (VDT).

Panel A: h=2							
	V1	V2	V3	V4	V5	FROM others including own	FROM others
V1	0.408	0.174	0.125	0.108	0.102	0.918	0.510
V2	0.226	0.315	0.197	0.154	0.138	1.031	0.716
V3	0.155	0.216	0.286	0.218	0.193	1.069	0.782
V4	0.117	0.164	0.217	0.288	0.253	1.038	0.750
V5	0.094	0.131	0.174	0.231	0.314	0.945	0.631
TO others including own	1	1	1	1	1		
TO others	0.592	0.685	0.714	0.712	0.686		
NET	0.082	-0.031	-0.069	-0.038	0.055		
NET sign	+	-	-	-	+		
NET ranking	1	3	5	4	2		
Panel B: h=10							
	V1	V2	V3	V4	V5	FROM others including own	FROM others
V1	0.373	0.185	0.150	0.138	0.134	0.981	0.608
V2	0.233	0.301	0.196	0.159	0.146	1.035	0.734
V3	0.166	0.215	0.273	0.210	0.187	1.051	0.778
V4	0.127	0.165	0.211	0.272	0.239	1.013	0.741
V5	0.102	0.133	0.170	0.220	0.294	0.921	0.626
TO others including own	1	1	1	1	1		
TO others	0.627	0.699	0.727	0.728	0.706		
NET	0.019	-0.035	-0.051	-0.013	0.079		
NET sign	+	-	-	-	+		
NET ranking	2	4	5	3	1		

Note. This figure shows the spillover Table based on the column-normalized generalized forecast error variance decomposition, which is displayed in the central frame. Results refer to the HH scenario (high-persistent and high-correlated series) and to the forecast horizon $h=2$ (Panel A) and $h=10$ (Panel B). The Table shows the directional spillover received from others (FROM others), the directional spillover received from others including own (FROM others (including own)) the directional spillover transmitted to others (TO others), the directional spillover transmitted to others including own (TO others (including own)) and the net contribution (NET) defined as the difference between the directional spillover transmitted TO others and the directional spillover received FROM others for variable V_i , $i=1, \dots, 5$. The bottom lines report for each variable the sign of the NET spillover (NET sign): negative if the variable is a net receiver and positive if the variable is a net donor, and the ranking based on the value of the NET spillover from the highest to the lowest (NET ranking).

Table 4: Spillover Table based on the variance decomposition table (VDT) normalized by the spectral radius.

Panel A: h=2							
	V1	V2	V3	V4	V5	FROM others including own	FROM others
V1	0.284	0.172	0.139	0.120	0.103	0.818	0.534
V2	0.157	0.311	0.218	0.171	0.140	0.998	0.686
V3	0.108	0.213	0.317	0.242	0.195	1.076	0.758
V4	0.081	0.162	0.241	0.318	0.256	1.058	0.740
V5	0.065	0.130	0.193	0.256	0.318	0.963	0.644
TO others including own	0.695	0.988	1.108	1.107	1.013		
TO others	0.412	0.677	0.791	0.789	0.695		
NET	-0.122	-0.010	0.032	0.049	0.050		
NET sign	-	-	+	+	+		
NET ranking	5	4	3	2	1		
Panel B: h=10							
	V1	V2	V3	V4	V5	FROM others including own	FROM others
V1	0.241	0.181	0.169	0.158	0.141	0.890	0.649
V2	0.151	0.294	0.221	0.182	0.153	1.000	0.706
V3	0.107	0.210	0.307	0.240	0.197	1.061	0.754
V4	0.082	0.161	0.237	0.310	0.251	1.041	0.731
V5	0.066	0.130	0.192	0.252	0.310	0.949	0.639
TO others including own	0.647	0.975	1.125	1.142	1.052		
TO others	0.406	0.681	0.818	0.832	0.742		
NET	-0.243	-0.025	0.064	0.100	0.103		
NET sign	-	-	+	+	+		
NET ranking	5	4	3	2	1		

Note. This figure shows the spillover Table based on the generalized forecast error variance decomposition normalized by the spectral radius, which is displayed in the central frame. Results refer to the HH scenario (high-persistent and high-correlated series) and to the forecast horizon $h=2$ (Panel A) and $h=10$ (Panel B). The Table shows the directional spillover received from others (FROM others), the directional spillover received from others including own (FROM others (including own)) the directional spillover transmitted to others (TO others), the directional spillover transmitted to others including own (TO others (including own)) and the net contribution (NET) defined as the difference between the directional spillover transmitted TO others and the directional spillover received FROM others for variable V_i , $i=1, \dots, 5$. The bottom lines show for each variable the sign of the NET spillover (NET sign): negative if the variable is a net receiver and positive if the variable is a net donor, and the ranking based on the value of the NET spillover from the highest to the lowest (NET ranking).

Table 5: Spillover Table based on the variance decomposition table (VDT) normalized by the maximum row sum.

Panel A: h=2								
	V1	V2	V3	V4	V5	FROM others including own	FROM others	
V1	0.264	0.160	0.129	0.111	0.096	0.760	0.496	
V2	0.146	0.289	0.203	0.159	0.130	0.927	0.638	
V3	0.100	0.198	0.295	0.225	0.182	1	0.705	
V4	0.076	0.150	0.224	0.296	0.238	0.984	0.688	
V5	0.061	0.121	0.180	0.238	0.296	0.895	0.599	
TO others including own	0.647	0.918	1.030	1.029	0.942			
TO others	0.383	0.629	0.735	0.733	0.646			
NET	-0.114	-0.009	0.030	0.046	0.047			
NET sign	-	-	+	+	+			
NET ranking	5	4	3	2	1			
Panel B: h=10								
	V1	V2	V3	V4	V5	FROM others including own	FROM others	
V1	0.227	0.170	0.159	0.149	0.133	0.839	0.612	
V2	0.142	0.277	0.208	0.171	0.144	0.943	0.666	
V3	0.101	0.198	0.290	0.226	0.185	1	0.710	
V4	0.077	0.152	0.223	0.292	0.237	0.982	0.689	
V5	0.062	0.123	0.181	0.237	0.292	0.895	0.603	
TO others including own	0.610	0.919	1.061	1.076	0.992			
TO others	0.383	0.642	0.771	0.784	0.700			
NET	-0.229	-0.024	0.061	0.095	0.097			
NET sign	-	-	+	+	+			
NET ranking	5	4	3	2	1			

Note. This figure shows the spillover Table based on the generalized forecast error variance decomposition normalized by the maximum row sum, which is displayed in the central frame. Results refer to the HH scenario (high-persistent and high-correlated series) and to the forecast horizon $h=2$ (Panel A) and $h=10$ (Panel B). The Table shows the directional spillover received from others (FROM others), the directional spillover received from others including own (FROM others (including own)) the directional spillover transmitted to others (TO others), the directional spillover transmitted to others including own (TO others (including own)) and the net contribution (NET) defined as the difference between the directional spillover transmitted TO others and the directional spillover received FROM others for variable V_i , $i=1, \dots, 5$. The bottom lines show for each variable the sign of the NET spillover (NET sign): negative if the variable is a net receiver and positive if the variable is a net donor, and the ranking based on the value of the NET spillover from the highest to the lowest (NET ranking).

Table 6: Spillover Table based on the variance decomposition table (VDT) normalized by the maximum column sum.

Panel A: h=2								
	V1	V2	V3	V4	V5	FROM others including own	FROM others	
V1	0.256	0.155	0.125	0.108	0.093	0.738	0.482	
V2	0.142	0.281	0.197	0.154	0.126	0.900	0.619	
V3	0.097	0.193	0.286	0.218	0.176	0.971	0.684	
V4	0.073	0.146	0.217	0.287	0.231	0.955	0.667	
V5	0.059	0.117	0.174	0.231	0.287	0.869	0.582	
TO others including own	0.628	0.892	1	0.999	0.914			
TO others	0.372	0.611	0.714	0.712	0.627			
NET	-0.110	-0.009	0.029	0.044	0.045			
NET sign	-	-	+	+	+			
NET ranking	5	4	3	2	1			
Panel B: h=10								
	V1	V2	V3	V4	V5	FROM others including own	FROM others	
V1	0.211	0.158	0.148	0.138	0.124	0.779	0.568	
V2	0.132	0.257	0.193	0.159	0.134	0.876	0.619	
V3	0.094	0.184	0.269	0.210	0.172	0.929	0.660	
V4	0.072	0.141	0.208	0.272	0.220	0.912	0.640	
V5	0.058	0.114	0.168	0.220	0.271	0.831	0.560	
TO others including own	0.567	0.854	0.986	1	0.921			
TO others	0.356	0.597	0.716	0.728	0.650			
NET	-0.212	-0.022	0.056	0.088	0.090			
NET sign	-	-	+	+	+			
NET ranking	5	4	3	2	1			

Note. This figure shows the spillover Table based on the generalized forecast error variance decomposition normalized by the maximum column sum, which is displayed in the central frame. Results refer to the HH scenario (high-persistent and high-correlated series) and to the forecast horizon $h=2$ (Panel A) and $h=10$ (Panel B). The Table shows the directional spillover received from others (FROM others), the directional spillover received from others including own (FROM others (including own)) the directional spillover transmitted to others (TO others), the directional spillover transmitted to others including own (TO others (including own)) and the net contribution (NET) defined as the difference between the directional spillover transmitted TO others and the directional spillover received FROM others for variable V_i , $i=1, \dots, 5$. The bottom lines report for each variable the sign of the NET spillover (NET sign): negative if the variable is a net receiver and positive if the variable is a net donor, and the ranking based on the value of the NET spillover from the highest to the lowest (NET ranking).

Table 7: Errors in sign (using population parameters).

Panel A: h=2				
	L. L.	L.H.	H.L.	H.H.
normalization by row	0	1	0	1
normalization by column	3	3	3	3
normalization by spectral radius	0	0	0	0
normalization by max row sum	0	0	0	0
normalization by max col sum	0	0	0	0
Panel B: h=10				
	L. L.	L.H.	H.L.	H.H.
normalization by row	0	0	0	0
normalization by column	1	3	1	3
normalization by spectral radius	0	0	0	0
normalization by max row sum	0	0	0	0
normalization by max col sum	0	0	0	0

Note. The Table shows the number of errors in sign for each scenario (L.L. , L.H. , H.L. , H.H. , where L.L. = low persistence low comovement, L.H.=low persistence high comovement, H.L.=high persistence low comovement, H.H.=high persistence high comovement). Results refer to the forecast horizon h=2 (panel A) and h=10 (Panel B). Errors are counted when the net spillover obtained from the normalized matrix has a sign opposite to the one of the net spillover obtained from the non-normalized matrix. The total number of possible errors is 5 for each scenario.

Table 8: Errors in ranking (using population parameters).

Panel A: h=2				
	L. L.	L.H.	H.L.	H.H.
normalization by row	1	1	1	1
normalization by column	1	1	1	1
normalization by spectral radius	0	0	0	0
normalization by max row sum	0	0	0	0
normalization by max col sum	0	0	0	0
Panel B: h=10				
	L. L.	L.H.	H.L.	H.H.
normalization by row	1	1	1	1
normalization by column	0	1	0	1
normalization by spectral radius	0	0	0	0
normalization by max row sum	0	0	0	0
normalization by max col sum	0	0	0	0

Note. The Table shows the number of errors in ranking for each scenario (L.L. , L.H. , H.L. , H.H. , where L.L. = low persistence low comovement, L.H.=low persistence high comovement, H.L.=high persistence low comovement, H.H.=high persistence high comovement). Results refer to the forecast horizon h=2 (panel A) and h=10 (Panel B). Errors are counted when the ranking of the net spillovers obtained from the normalized matrix is different from that of the non-normalized matrix. The total number of possible errors is 1 for each scenario.

Table 9: Errors in sign (using simulations).

Panel A: h=2				
	L. L.	L.H.	H.L.	H.H.
normalization by row	118	607	198	492
normalization by column	2802	3226	1746	2324
normalization by spectral radius	0	0	0	0
normalization by max row sum	0	0	0	0
normalization by max col sum	0	0	0	0
Panel B: h=10				
	L. L.	L.H.	H.L.	H.H.
normalization by row	28	293	111	243
normalization by column	1368	3143	1644	1833
normalization by spectral radius	0	0	0	0
normalization by max row sum	0	0	0	0
normalization by max col sum	0	0	0	0

Note. The Table shows the number of errors in sign for each scenario (L.L. , L.H. , H.L. , H.H. , where L.L. = low persistence low comovement, L.H.=low persistence high comovement, H.L.=high persistence low comovement, H.H.=high persistence high comovement). Results refer to the forecast horizon H=2 (panel A) and H=10 (Panel B). Errors are counted when the net spillover obtained from the normalized matrix has a sign opposite to the one of the net spillover obtained from the non-normalized matrix. The total number of possible errors is 5000 for each scenario (5 variables times 1000 simulations for each scenario).

Table 10: Errors in ranking (using simulations).

Panel A: h=2				
	L. L.	L.H.	H.L.	H.H.
normalization by row	1000	990	870	989
normalization by column	1000	1000	934	949
normalization by spectral radius	0	0	0	0
normalization by max row sum	0	0	0	0
normalization by max col sum	0	0	0	0
Panel B: h=10				
	L. L.	L.H.	H.L.	H.H.
normalization by row	1000	991	481	743
normalization by column	966	1000	981	977
normalization by spectral radius	0	0	0	0
normalization by max row sum	0	0	0	0
normalization by max col sum	0	0	0	0

Note. The Table shows the number of errors in ranking for each scenario (L.L. , L.H. , H.L. , H.H. , where L.L. = low persistence low comovement, L.H.=low persistence high comovement, H.L.=high persistence low comovement, H.H.=high persistence high comovement). Results refer to the forecast horizon H=2 (panel A) and H=10 (Panel B). Errors are counted when the ranking of the net spillovers obtained from the normalized matrix is different from the one of the non-normalized matrix. The total number of possible errors is 1000 for each scenario (one ranking times 1000 simulations for each scenario).

Table 11: Comparison with the results in Diebold and Yilmaz (2012)

Panel I: Row Normalization						
	Stocks	Bonds	Commodities	FX	FROM others including own	FROM others
Stocks	0.8876	0.0729	0.0035	0.0361	1	0.112
Bonds	0.1021	0.8145	0.0273	0.0561	1	0.186
Commodities	0.0047	0.0370	0.9369	0.0214	1	0.063
FX	0.0569	0.0703	0.0155	0.8573	1	0.143
TO others including own	1.051	0.995	0.983	0.971		
TO others	0.164	0.180	0.046	0.114		TOTAL
NET	0.051	-0.005	-0.017	-0.029		0.126
Panel II: Max row Normalization						
	Stocks	Bonds	Commodities	FX	FROM others including own	FROM others
Stocks	0.8444	0.0694	0.0033	0.0343	0.951	0.107
Bonds	0.1021	0.8145	0.0273	0.0561	1	0.186
Commodities	0.0041	0.0323	0.8194	0.0187	0.875	0.055
FX	0.0547	0.0675	0.0149	0.8238	0.961	0.137
TO others including own	1.005	0.984	0.865	0.933		
TO others	0.161	0.169	0.045	0.109		TOTAL
NET	0.054	-0.016	-0.010	-0.028		0.121

Note. This Table shows the spillover based on the generalized forecast error variance decomposition normalized by row (Panel A), as in Diebold and Yilmaz (2012), and normalized by the maximum row sum (Panel B). Results refer to the 10-days-ahead forecast horizon.