

 EXTRACTA MATHEMATICAE Vol. 31, Núm. 1, 109–117 (2016)

A Note on Rational Approximation with Respect to Metrizable Compactifications of the Plane

M. FRAGOULOPOULOU, V. NESTORIDIS

*Department of Mathematics, University of Athens
Panepistimiopolis, Athens 157 84, Greece
fragoulop@math.uoa.gr, vnestor@math.uoa.gr*

Presented by Manuel Maestre

Received February 10, 2015

Abstract: In the present note we examine possible extensions of Runge, Mergelyan and Arakelian Theorems, when the uniform approximation is meant with respect to the metric ϱ of a metrizable compactification (S, ϱ) of the complex plane \mathbb{C} .

Key words: compactification, Arakelian's theorem, Mergelyan's theorem, Runge's theorem, uniform approximation in the complex domain.

AMS *Subject Class.* (2010): 30E10.

1. INTRODUCTION

It is well known that the class of uniform limits of polynomials in $\overline{D} = \{z \in \mathbb{C} : |z| \leq 1\}$ coincides with the disc algebra $A(D)$. A function $f : \overline{D} \rightarrow \mathbb{C}$ belongs to $A(D)$ if and only if it is continuous on \overline{D} and holomorphic in the open unit disc D . It is less known (see [3, 7]) what is the corresponding class when the uniform convergence is not meant with respect to the usual Euclidean metric on \mathbb{C} , but it is meant with respect to the chordal metric χ on $\mathbb{C} \cup \{\infty\}$. The class of χ -uniform limits of polynomials on \overline{D} is denoted by $\tilde{A}(D)$ and contains $A(D)$. A function $f : \overline{D} \rightarrow \mathbb{C} \cup \{\infty\}$ belongs to $\tilde{A}(D)$ if and only if $f \equiv \infty$, or it is continuous on \overline{D} , $f(D) \subset \mathbb{C}$ and $f|_D$ is holomorphic. The function $f(z) = \frac{1}{1-z}$, $z \in D$, belongs to $\tilde{A}(D)$, but not to $A(D)$; thus, it cannot be uniformly approximated on D , by polynomials with respect to the usual Euclidean metric on \mathbb{C} , but it can be uniformly approximated by polynomials with respect to the chordal metric χ .

More generally, if $K \subset \mathbb{C}$ is a compact set with connected complement, then according to Mergelyan's theorem [10] polynomials are dense in $A(K)$ with respect to the usual Euclidean metric on \mathbb{C} . We recall that a function $f : K \rightarrow \mathbb{C}$ belongs to $A(K)$ if and only if it is continuous on K and holomorphic in the interior K° of K .

An open problem is to characterize the class $\tilde{A}(K)$ of χ -uniform limits of polynomials on K .

CONJECTURE. ([1, 6]) Let $K \subset \mathbb{C}$ be a compact set with connected complement K^c . A function $f : K \rightarrow \mathbb{C} \cup \{\infty\}$ belongs to $\tilde{A}(K)$ if and only if it is continuous on K and for each component V of K° , either $f(V) \subset \mathbb{C}$ and $f|_V$ is holomorphic, or $f|_V \equiv \infty$.

Extensions of this result have been obtained in [5] when K^c has a finite number of components and K is bounded by a finite set of disjoint Jordan curves. In this case, the χ -uniform approximation is achieved using rational functions with poles out of K instead of polynomials. Furthermore, extensions of Runge's theorem are also proved in [5]. Finally a first result has been obtained in [5] concerning an extension of the approximation theorem of Arakelian ([2]).

Instead of considering the one point compactification $\mathbb{C} \cup \{\infty\}$ of the complex plane \mathbb{C} , we can consider an arbitrary metrizable compactification (S, ϱ) of \mathbb{C} and investigate the analogues of all previous results. This is the content of the present paper.

2. PRELIMINARIES

We say that (S, ϱ) is a *metrizable compactification of the plane* \mathbb{C} , if ϱ is a metric on S , S is compact, $S \supset \mathbb{C}$ and \mathbb{C} is an open dense subset of S . Obviously, $S \setminus \mathbb{C}$ is a closed subset of S . We say that the points in $S \setminus \mathbb{C}$ are the points at infinity.

Let (S, ϱ) be a metrizable compactification of \mathbb{C} with metric ϱ . Many such compactifications can be found in [1]. The one point compactification $\mathbb{C} \cup \{\infty\}$ with the chordal metric χ is a distinct one of them. We note that in this case, the continuous function $\pi : S \rightarrow \mathbb{C} \cup \{\infty\}$, such that $\pi(c) = c$, for every $c \in \mathbb{C}$ and $\pi(x) = \infty$, for every $x \in S \setminus \mathbb{C}$, is useful.

Another metrizable compactification is the one defined in [8] and constructed as follows: consider the map

$$\begin{aligned} \phi : \mathbb{C} &\longrightarrow D = \{\lambda \in \mathbb{C} : |\lambda| < 1\} \\ z &\longmapsto \frac{z}{1 + |z|} \end{aligned} ,$$

which is a homeomorphism. A compactification of the image D of ϕ is \overline{D} , the closure of D , with the usual metric. This leads to the following compactifica-

tion of \mathbb{C}

$$(2.1) \quad S_1 := \mathbb{C} \cup \{\infty e^{i\vartheta} : 0 \leq \vartheta \leq 2\pi\},$$

with metric d given by

$$(2.2) \quad \begin{aligned} d(z, w) &= \left| \frac{z}{1+|z|} - \frac{w}{1+|w|} \right| && \text{if } z, w \in \mathbb{C}, \\ d(z, \infty e^{i\vartheta}) &= \left| \frac{z}{1+|z|} - e^{i\vartheta} \right| && \text{if } z \in \mathbb{C}, \vartheta \in \mathbb{R}, \\ d(\infty e^{i\vartheta}, \infty e^{i\varphi}) &= \left| e^{i\vartheta} - e^{i\varphi} \right| && \text{if } \vartheta, \varphi \in \mathbb{R}. \end{aligned}$$

In what follows, with a compactification (S, ϱ) of \mathbb{C} , we shall always mean a metrizable compactification.

An important question for a given compactification of \mathbb{C} is, whether for $c \in \mathbb{C}$ and $x \in S \setminus \mathbb{C}$, the addition $c + x$ is well defined. In other words, having two convergent sequences $\{z_n\}, \{w_n\}$ in \mathbb{C} , such that $z_n \rightarrow c$ and $w_n \rightarrow x$ does the sequence $\{z_n + w_n\}$ have a limit in S ?

If the answer is positive for any such sequences $\{z_n\}, \{w_n\}$ in \mathbb{C} , then the limit $y \in S$ of the sequence $\{z_n + w_n\}$ is uniquely determined and we write $c + x = y = x + c$. We are interested in compactifications (S, ϱ) , where $c + x$ is well defined for any $c \in \mathbb{C}$ and $x \in S$ (it suffices to take $x \in S \setminus \mathbb{C}$). In this case, the map $\mathbb{C} \times S \rightarrow S$, $(c, x) \mapsto c + x$, is automatically continuous.

Indeed, let $x \in S \setminus \mathbb{C}$, $y \in \mathbb{C}$ and $w = x + y \in S \setminus \mathbb{C}$. Let $\{z_n\}$ in S and $\{y_n\}$ in \mathbb{C} , such that $z_n \rightarrow x$ and $y_n \rightarrow y$. If all but finitely many z_n belong to \mathbb{C} , then by our assumption $z_n + y_n \rightarrow x + y$. Suppose that infinitely many z_n belong to $S \setminus \mathbb{C}$. Without loss of generality we may assume that all z_n belong to $S \setminus \mathbb{C}$ and by compactness we can assume that $z_n + y_n \rightarrow l \neq w = x + y$.

Let $d = \varrho(l, w) > 0$. Then there exists $n_0 \in \mathbb{N}$, such that

$$\varrho(z_n + y_n, l) < \frac{d}{2} \quad \text{for all } n \geq n_0.$$

Fix $n \geq n_0$. Since, $z_n + y_n$ is well defined, there exists $z'_n \in \mathbb{C}$, such that

$$\varrho(z_n, z'_n) < \frac{1}{n} \quad \text{and} \quad \varrho(z_n + y_n, z'_n + y_n) < \frac{1}{n}.$$

It follows that

$$\varrho(z'_n, x) \leq \varrho(z'_n, z_n) + \varrho(z_n, x) < \frac{1}{n} + \varrho(z_n, x) \rightarrow 0.$$

Hence, $z'_n \rightarrow x$, $y_n \rightarrow y$ and $z'_n, y_n \in \mathbb{C}$. By our assumption, it follows that $z'_n + y_n \rightarrow x + y = w$. But

$$\begin{aligned} \varrho(z'_n + y_n, l) &\leq \varrho(z'_n + y_n, z_n + y_n) + \varrho(z_n + y_n, l) \\ &\leq \frac{1}{n} + \varrho(z_n + y_n, l) < \frac{1}{n} + \frac{d}{2} \rightarrow \frac{d}{2}. \end{aligned}$$

Thus, for all n large enough we have

$$\varrho(z'_n + y_n, l) \leq \frac{3d}{4} < d = \varrho(l, w).$$

It follows that $\varrho(z'_n + y_n, w) \geq \frac{d}{4}$, for all n large enough. Therefore, we cannot have $z'_n + y_n \rightarrow w$.

Consequently, one concludes that the addition map is continuous at every (x, y) with $x \in S \setminus \mathbb{C}$ and $y \in \mathbb{C}$. Obviously, it is also continuous at every (x, y) with x and y in \mathbb{C} . Thus, addition is continuous on $S \times \mathbb{C}$. Furthermore, the following holds:

Let $K \subset \mathbb{C}$ be compact. Obviously, the map $K \times S \rightarrow S$, $(c, x) \mapsto c + x$, is uniformly continuous.

Remark 1. The preceding certainly holds for the compactification (S_1, d) (see (2.1)), since

$$c + \infty e^{i\vartheta} = \infty e^{i\vartheta} \quad \text{for all } c \in \mathbb{C} \text{ and } \vartheta \in \mathbb{R},$$

and we have continuity.

Remark 2. If we identify \mathbb{R} with the interval $(-1, 1)$, up to a homeomorphism, then $\mathbb{C} \cong \mathbb{R}^2$ is identified with the square $(-1, 1) \times (-1, 1)$. An obvious compactification of \mathbb{C} is then the closed square with the usual metric. The points at infinity are those on the boundary of the square, for instance, those points on the side $\{1\} \times [-1, 1]$. If $x \in \{1\} \times (-1, 1)$ and $c \in \mathbb{C}$, then $c + x$ is a point in the same side; if $\text{Im } c \neq 0$, then $c + x \neq x$. If $x = (1, 1)$ and $c \in \mathbb{C}$, then $x + c = x$. If $\text{Im } c > 0$, then $c + x$ lies higher than x in the side $\{1\} \times (-1, 1)$.

In this example, the addition is well defined and continuous, but the points at infinity are not stabilized as in Remark 1.

QUESTION. Is there a metrizable compactification of \mathbb{C} such that the addition $c + x$ is not well defined for some $c \in \mathbb{C}$ and $x \in S \setminus \mathbb{C}$?

The answer is “yes”. An example comes from the previous square in Remark 2, if we identify all the points of $\{1\} \times [-\frac{1}{2}, \frac{1}{2}]$ and make them just one point.

3. RUNGE AND MERGELYAN TYPE THEOREMS

In this section using a compactification of \mathbb{C} satisfying all properties discussed in the Preliminaries, we obtain the following theorem, that extends [5, Theorem 3.3].

THEOREM 3.1. *Let $\Omega \subset \mathbb{C}$ be a bounded domain, whose boundary consists of a finite set of pairwise disjoint Jordan curves. Let $K = \overline{\Omega}$ and A a set containing one point from each component of $(\mathbb{C} \cup \{\infty\}) \setminus K$. Let (S, ϱ) be a compactification of \mathbb{C} , such that the addition $+: \mathbb{C} \times S \rightarrow S$ is well defined. Let $f: K \rightarrow S$ be a continuous function, such that $f(\Omega) \subset \mathbb{C}$ and $f|_{\Omega}$ is holomorphic. Let $\varepsilon > 0$. Then, there exists a rational function R with poles only in A and such that $\varrho(f(z), R(z)) < \varepsilon$, for all $z \in K$.*

Proof. If Ω is a disk, the proof has been given in [1]. If Ω is the interior of a Jordan curve, the proof is given again in [1], but also in [6]. In the general case, we imitate the proof of [5, Theorem 3.3]. Namely, we consider the Laurent decomposition of f , given by $f = f_0 + f_1 + \dots + f_N$ (see [4]). The function f_0 is defined on a simply connected domain, bounded by a Jordan curve, and it can be uniformly approximated by a polynomial or a rational function R_0 with pole in the unbounded component. Similarly, f_1 is approximated by a rational function R_1 with pole in A and so on. Thus, the function $R_0 + R_1 + \dots + R_N$ approximates, with respect to ϱ , the function $f = f_0 + f_1 + \dots + f_N$. This is due to the fact that at every point z all the f_i 's, $i = 1, 2, \dots, N$, except maybe one, take values in \mathbb{C} and the one, maybe has as a value, an infinity point in $S \setminus \mathbb{C}$. In this way, the addition map $\mathbb{C} \times S \rightarrow S$, $(c, x) \mapsto c + x$, is well defined and uniformly continuous on compact sets and so we are done. ■

Another Runge-type theorem is the following, where we do not need any assumption for the compactification S , or the addition map $+: \mathbb{C} \times S \rightarrow S$.

THEOREM 3.2. *Let $\Omega \subset \mathbb{C}$ be open, $f: \Omega \rightarrow \mathbb{C}$ be holomorphic and (S, ϱ) a compactification of \mathbb{C} . Let A be a set containing one point from each component of $(\mathbb{C} \cup \infty) \setminus \Omega$. Let $\varepsilon > 0$ and $L \subset \Omega$ compact. Then, there*

exists a rational function R with poles in A , such that $\varrho(f(z), R(z)) < \varepsilon$ for all $z \in L$.

Proof. Clearly the subset $f(L)$ of \mathbb{C} is compact. Then, from the classical theorem of Runge, there exist rational functions $\{R_n\}$, with poles only in A , converging uniformly to f on L , with respect to the Euclidean metric $|\cdot|$. Hence, there is a positive integer n_0 and a compact K , such that

$$f(L) \subset K \subset \mathbb{C} \quad \text{and} \quad R_n(L) \subset K \quad \text{for all } n \geq n_0.$$

But on K the metrics $|\cdot|$ and ϱ are uniformly equivalent. Therefore, $R_n \rightarrow f$ uniformly on L , with respect to ϱ . To conclude the proof, it suffices to put $R = R_n$, for n large enough. ■

Theorem 3.2 easily yields the following

COROLLARY 3.3. *Under the assumptions of Theorem 3.2 there exists a sequence $\{R_n\}$ of rational functions with poles in A , such that $R_n \rightarrow f$, ϱ -uniformly, on each compact subset of Ω .*

Remark. According to Corollary 3.3, some of the ϱ -uniform limits, on compacta, of rational functions with poles in A , are the holomorphic functions $f : \Omega \rightarrow \mathbb{C}$. Those are limits of the finite type. The other limits of sequences $\{R_n\}$ as above may be functions $f : \Omega \rightarrow S \setminus \mathbb{C}$ of infinite type, continuous (but maybe not all of them, as the Example (S_1, d) shows; cf. [8]).

QUESTION. Is a characterization possible for such limits $f : \Omega \rightarrow S_1 \setminus \mathbb{C}$?

An imitation of the arguments in [8, p. 1007] gives that f must be of the form $f(z) = \infty e^{i\vartheta(z)}$, $z \in \Omega$, where ϑ is a multivalued harmonic function.

The following extends [5, Section 5].

THEOREM 3.4. *Let $\Omega \subset \mathbb{C}$ be open and f a meromorphic function on Ω . Let B denote the set of poles of f . Let (S, ϱ) be a compactification of \mathbb{C} , such that the addition $+$: $\mathbb{C} \times S \rightarrow S$ is well defined. Let $\varepsilon > 0$ and $K \subset \Omega$ be a compact set. Then, there is a rational function g , such that $\varrho(f(z), g(z)) < \varepsilon$, for every $z \in K \setminus B$.*

Proof. Since $B \cap K$ is a finite set, the function f decomposes to $f = h + w$, where h is a rational function with poles in $B \cap K$ and w is holomorphic on an open set containing K . By Runge's theorem there exists a rational function R

with poles off K , such that $|w(z) - R(z)| < \varepsilon'$ on K . Since $w(K)$ is a compact subset of \mathbb{C} and the addition $+: \mathbb{C} \times S \rightarrow S$ is well defined, a suitable choice of ε' gives

$$\varrho([h(z) + w(z)], [h(z) + R(z)]) < \varepsilon \quad \text{on } K \setminus B.$$

We set $g = h + R$ and the result follows. ■

4. ARAKELIAN SETS

A closed set $F \subset \mathbb{C}$ is said a *set of approximation* if every function $f : F \rightarrow \mathbb{C}$ continuous on F and holomorphic in F° can be approximated by entire functions, uniformly on the whole F . This is equivalent to the fact that F is an Arakelian set (see [2]), that is $(\mathbb{C} \cup \{\infty\}) \setminus F$ is connected and locally connected (at ∞).

We can now ask about an extension of the Arakelian theorem in the context of metrizable compactifications. A result in this direction is the following

PROPOSITION 4.1. *Let $F \subset \mathbb{C}$ be a closed Arakelian set with empty interior, i.e., $F^\circ = \emptyset$. We consider the compactification (S_1, d) of \mathbb{C} (see (2.1) and (2.2)) and let $f : F \rightarrow S_1$ be a continuous function. Let $\varepsilon > 0$. Then, there is an entire function g such that $d(f(z), g(z)) < \varepsilon$, for every $z \in F$.*

Proof. According to (1.1), the compactification S_1 is homeomorphic to $\overline{D} = \{z \in \mathbb{C} : |z| \leq 1\}$. For each $0 < R < 1$ let us define

$$\begin{aligned} \phi_R : \overline{D} &\longrightarrow \{z \in \mathbb{C} : |z| \leq R\} \subset \overline{D} \\ z &\longmapsto \begin{cases} z, & \text{if } |z| \leq R, \\ \frac{Rz}{|z|}, & \text{if } R \leq |z| \leq 1. \end{cases} \end{aligned}$$

In other words, the whole line segment $[Re^{i\vartheta}, e^{i\vartheta}]$ is mapped at the end point $Re^{i\vartheta}$. The function ϕ_R is continuous and induces a continuous function $\tilde{\phi}_R : S_1 \rightarrow S_1$. It suffices to take $\tilde{\phi}_R := T^{-1} \circ \phi_R \circ T$, where $T : S_1 \rightarrow \{w \in \mathbb{C} : |w| \leq 1\}$ is defined as follows

$$\begin{aligned} T(z) &:= \frac{z}{1 + |z|} && \text{for } z \in \mathbb{C} \subset S_1, \\ T(\infty e^{i\vartheta}) &:= e^{i\vartheta} && \text{for } \vartheta \in \mathbb{R}. \end{aligned}$$

If $\varepsilon > 0$ is given, then there exists $R_\varepsilon < 1$, such that for $R_\varepsilon \leq R < 1$ and $z \in S_1$, we have $d(z, \tilde{\phi}_R(z)) < \frac{\varepsilon}{2}$.

Let now f be as in the statement of the Proposition 4.1. Then,

$$d\left(f(z), (\tilde{\phi}_R \circ f)(z)\right) < \frac{\varepsilon}{2} \quad \text{for all } z \in F.$$

Moreover, the function $\tilde{\phi}_R \circ f : F \rightarrow \mathbb{C}$ is continuous. Since F is a closed Arakelian set, with empty interior, and $(\tilde{\phi}_R \circ f)(F) \subset K$, is included in a compact subset K of \mathbb{C} , there exists g entire, such that

$$\left|(\tilde{\phi}_R \circ f)(z) - g(z)\right| < \varepsilon' \quad \text{for all } z \in F.$$

Since $(\tilde{\phi}_R \circ f)(F)$ is contained in a compact subset K of \mathbb{C} , for a suitable choice of ε' , it follows that

$$d\left((\tilde{\phi}_R \circ f)(z), g(z)\right) < \frac{\varepsilon}{2} \quad \text{for all } z \in F.$$

The triangle inequality completes the proof. ■

An analogue of Proposition 4.1 for the one point compactification $\mathbb{C} \cup \{\infty\}$ of \mathbb{C} has been established in [5].

REFERENCES

- [1] I. ANDROULIDAKIS, V. NESTORIDIS, Extension of the disc algebra and of Mergelyan's theorem, *C.R. Math. Acad. Sci. Paris* **349**(13–14) (2011), 745–748.
- [2] N.U. ARAKELIAN, Uniform approximation on closed sets by entire functions, *Izv. Akad. Nauk SSSR Ser. Mat.* **28** (1964), 1187–1206 (Russian).
- [3] L. BROWN, P.M. GAUTHIER, W. HENGARTNER, Continuous boundary behaviour for functions defined in the open unit disc, *Nagoya Math. J.* **57** (1975), 49–58.
- [4] G. COSTAKIS, V. NESTORIDIS, I. PAPADOPERAKIS, Universal Laurent series, *Proc. Edinb. Math. Soc. (2)* **48**(3) (2005), 571–583.
- [5] M. FRAGOULOPOULOU, V. NESTORIDIS, I. PAPADOPERAKIS, Some results on spherical approximation, *Bull. Lond. Math. Soc.* **45**(6) (2013), 1171–1180.
- [6] V. NESTORIDIS, Compactifications of the plane and extensions of the disc algebra, in “Complex Analysis and Potential Theory”, CRM Proc. Lecture Notes, 55, Amer. Math. Soc., Providence, RI, 2012, 61–75.
- [7] V. NESTORIDIS, An extension of the disc algebra, I, *Bull. Lond. Math. Soc.* **44**(4) (2012), 775–788.

- [8] V. NESTORIDIS, N. PAPADATOS, An extension of the disc algebra, II, *Complex Var. Elliptic Equ.* **59** (7) (2014), 1003–1015.
- [9] V. NESTORIDIS, I. PAPADOPERAKIS, A remark on two extensions of the disc algebra and Mergelian's theorem, *preprint* 2011, arxiv: 1104.0833.
- [10] W. RUDIN, "Real and Complex Analysis", McGraw-Hill Book Co., New York-Toronto, Ont.-London, 1966.