

Rend. Lincei Mat. Appl. 29 (2018), 607–617
DOI 10.4171/RLM/824



Probability Theory — *Connectedness of random set attractors*, by MICHAEL SCHEUTZOW and ISABELL VORKASTNER, communicated on April 20, 2018.¹

ABSTRACT. — We examine the question whether random set attractors for continuous-time random dynamical systems on a connected state space are connected. In the deterministic case, these attractors are known to be connected. In the probabilistic setup, however, connectedness has only been shown under stronger connectedness assumptions on the state space. Under a weak continuity condition on the random dynamical system we prove connectedness of the pullback attractor on a connected space. Additionally, we provide an example of a weak random set attractor of a random dynamical system with even more restrictive continuity assumptions on an even path-connected space which even attracts all bounded sets and which is not connected. On the way to proving connectedness of a pullback attractor we prove a lemma which may be of independent interest and which holds without the assumption that the state space is connected. It states that even though pullback convergence to the attractor allows for exceptional nullsets which may depend on the compact set, these nullsets can be chosen independently of the compact set (which is clear for σ -compact spaces but not at all clear for spaces which are not σ -compact).

KEY WORDS: Random dynamical system, pullback attractor, weak attractor, pullback continuity, measurable selection

MATHEMATICS SUBJECT CLASSIFICATION: 37H99, 37B25, 37C70, 28B20

1. INTRODUCTION

While attractors for (deterministic) dynamical systems have been studied for a long time, attractors for *random dynamical systems* were only introduced and studied in the nineties of the last century. The question of connectedness of a random pullback attractor was first addressed in the seminal paper [4]. Proposition 3.13 of that paper states that if a random dynamical system in discrete or continuous time taking values in a connected Polish space admits a pullback attractor A (in the sense that A attracts every bounded set in the pullback sense almost surely) then A is almost surely connected. Later, a gap was found in the proof of that proposition and an example in [6] shows that the claim does not even hold true in the deterministic case when time is discrete. Positive results (in discrete and continuous time) have been found in [2] under the additional condition that any compact set in the state space can be covered by a connected compact set (a property which clearly does not hold in the example in [6]).

¹Presented by Prof. G. Da Prato

The aim of this paper is to examine the question whether random set attractors of continuous-time random dynamical systems on a connected state space are connected.

In this paper, we distinguish between two kinds of random set attractors, pullback and weak attractors (precise definitions will be provided in the next section). By *set* attractor we mean an attractor which either attracts every deterministic compact set or every deterministic bounded set (we will state explicitly in each case if we want the attractor to attract every compact or even every bounded set). Pullback and weak attractors differ in the type of convergence of compact (or bounded) sets under the action of the random dynamical system to the attractor. *Pullback* stands for almost sure convergence and *weak* for convergence in probability. Both of these set attractors are known to be (almost surely) unique, see [5, Lemma 1.3].

In Section 3, we consider pullback attractors for continuous-time random dynamical systems taking values in a connected Polish space. Under a rather weak continuity assumption on the random dynamical system which we call *pullback continuity* we show that the pullback attractor (if it exists) is almost surely connected (even if it is only required to attract all compact sets). The first lemma in that section may be of independent interest. It states that even though pullback convergence to the attractor allows for exceptional nullsets which may depend on the compact set, these nullsets can be chosen independently of the compact set (even if the space is not σ -compact). This lemma does not assume the state space to be connected. The result allows us to argue pathwise (for fixed ω) in the proof of the main result.

In Section 4 we provide an example of a random dynamical system on a path-connected state space where the weak attractor is not connected. In that example the random dynamical system enjoys even stronger continuity properties than in the previous section and the attractor even attracts all bounded and not just compact sets. The state space in that example is the same as that in [6] but the random dynamical system on that space is more sophisticated.

Apart from set attractors for continuous-time system other types of random attractors such as *random point attractors* or *random Hausdorff-Delta-attractors* have been studied in the literature either in the pullback or weak sense ([2], [9]). These are generally not connected even if the ambient space is connected and the attractors are chosen to be *minimal* (unlike set attractors they are generally not unique). As an example for a disconnected minimal point attractor consider the scalar differential equation $dx = (x - x^3) dt$ on the interval $[0, 1]$. Each trajectory converges to $\{0\}$ or $\{1\}$. Hence, $\{0\} \cup \{1\}$ is the minimal (pullback or weak) point attractor (while the set attractor is the whole interval $[0, 1]$).

2. NOTATION AND PRELIMINARIES

Let (X, d) be a Polish (i.e. separable complete metric) space with Borel σ -algebra $\mathcal{B}(X)$ and $(\Omega, \mathcal{F}, \mathbb{P}, \theta)$ be a metric dynamical system, i.e. $(\Omega, \mathcal{F}, \mathbb{P})$ is a probability space and $(\theta_t)_{t \in \mathbb{R}}$ a group of jointly measurable maps on $(\Omega, \mathcal{F}, \mathbb{P})$ such that

$\theta_0 = \text{id}$ with invariant measure \mathbb{P} . Denote by $\overline{\mathcal{F}}$ the completion of \mathcal{F} with respect to \mathbb{P} . We further denote by $\overline{\mathbb{P}}$ the (unique) extension of \mathbb{P} to $\overline{\mathcal{F}}$.

Let $\varphi : \mathbb{R}_+ \times \Omega \times X \rightarrow X$ be jointly measurable, $\varphi_0(\omega, x) = x$, $\varphi_{s+t}(\omega, x) = \varphi_t(\theta_s\omega, \varphi_s(\omega, x))$ for all $x \in X$, and $x \mapsto \varphi_t(\omega, x)$ continuous, $s, t \in \mathbb{R}_+$ and $\omega \in \Omega$. Then, φ is called a *cocycle* and the collection $(\Omega, \mathcal{F}, \mathbb{P}, \theta, \varphi)$ is called a *random dynamical system (RDS)*, see [1] for a comprehensive treatment. We call an RDS *pullback continuous* if $t \mapsto \varphi_t(\theta_{-t}\omega, x)$ is continuous for each $\omega \in \Omega$ and $x \in X$.

A *semi-flow* $\phi : \{-\infty < s \leq t < \infty\} \times \Omega \times X \rightarrow X$ satisfies $\phi_{s,u}(\omega, x) = \phi_{t,u}(\omega, \cdot) \circ \phi_{s,t}(\omega, x)$, $\phi_{s,t}(\omega, x) = \phi_{s+h,t+h}(\theta_h\omega, x)$ and $\phi_{s,s}(\omega, x) = x$ for $\omega \in \Omega$, $x \in X$, $h \in \mathbb{R}$ and $-\infty < s \leq t \leq u < \infty$. There is a one-to-one relation between cocycles and semi-flows. One can either define a semi-flow by $\phi_{s,t}(\omega, x) := \varphi_{t-s}(\theta_s\omega, x)$ or a cocycle by $\varphi_t(\omega, x) := \phi_{0,t}(\omega, x)$. We say a semi-flow respectively RDS is *jointly continuous* if $(s, t, x) \mapsto \phi_{s,t}(\omega, x)$ respectively $(s, t, x) \mapsto \varphi_{t-s}(\theta_s\omega, x)$ is continuous. Note that a jointly continuous RDS is pullback continuous but the converse does not necessarily hold true.

For a set $A \subset X$ we denote

$$A^\varepsilon := \left\{ x \in X : d(x, A) := \inf_{a \in A} d(x, a) < \varepsilon \right\}.$$

DEFINITION 2.1. A family $\{A(\omega)\}_{\omega \in \Omega}$ of non-empty subsets of X is called

- (i) a *random compact set* if it is \mathbb{P} -almost surely a compact set and $\omega \mapsto d(x, A(\omega))$ is $\overline{\mathcal{F}}$ -measurable for each $x \in X$.
- (ii) φ -invariant if $\varphi_t(\omega, A(\omega)) = A(\theta_t\omega)$ for almost all $\omega \in \Omega$, $t \in \mathbb{R}_+$.

DEFINITION 2.2. Let $(\Omega, \mathcal{F}, \mathbb{P}, \theta, \varphi)$ be a random dynamical system. A random compact set A is called a *pullback attractor* if it satisfies the following properties

- (i) A is φ -invariant
- (ii) for every compact set $B \subset X$

$$\limsup_{t \rightarrow \infty} \sup_{x \in B} d(\varphi_t(\theta_{-t}\omega, x), A(\omega)) = 0 \quad \mathbb{P}\text{-almost surely.}$$

If the convergence in (ii) is merely in probability, then A is called a *weak attractor*.

3. PULLBACK ATTRACTOR

In this section, we show that the pullback attractor of a pullback continuous RDS on a connected space is connected. The pullback attractor attracts any compact set almost surely. We prove that the nullsets where it may not converge can be chosen independently of the compact set. This allows us to analyze the RDS pathwise and to use similar arguments as in the deterministic proof of [6, Theorem 3.1].

LEMMA 3.1. *Let A be the pullback attractor of the pullback continuous RDS φ . Then, there exists some $\hat{\Omega} \in \mathcal{F}$ with $\mathbb{P}(\hat{\Omega}) = 1$ such that for any $\omega \in \hat{\Omega}$ and compact set $K \subset X$,*

$$\limsup_{t \rightarrow \infty} \sup_{x \in K} d(\varphi_t(\theta_{-t}\omega, x), A(\omega)) = 0.$$

PROOF. First, we consider convergent sequences in X . Let

$$\hat{c} := \left\{ (x_\infty, x_1, x_2, x_3, \dots) \in X^{\mathbb{N}} : d(x_n, x_\infty) \leq \frac{1}{n} \text{ for all } n \in \mathbb{N} \right\}$$

which is closed in the Polish space $X^{\mathbb{N}}$ and hence itself a Polish space. Further, let

$$M(\omega) := \bigcup_{n \in \mathbb{N}} \bigcap_{m \in \mathbb{N}} \bigcup_{q \in \mathbb{Q}, q \geq m} \bigcup_{k \in \mathbb{N} \cup \{\infty\}} \{ (x_\infty, x_1, x_2, \dots) \in \hat{c} : \varphi_q(\theta_{-q}\omega, x_k) \in A(\omega)^{\frac{1}{n}} \}^c$$

be the set of sequences of \hat{c} that are not uniformly attracted. By measurability of φ and A , the graph of M is measurable.

Assume there is a subset $\tilde{\Omega} \in \mathcal{F}$ with $\mathbb{P}(\tilde{\Omega}) > 0$ such that $M(\omega) \neq \emptyset$ for all $\omega \in \tilde{\Omega}$. Define

$$\tilde{M}(\omega) := \begin{cases} M(\omega) & \text{if } \omega \in \tilde{\Omega} \\ \hat{c} & \text{else.} \end{cases}$$

Then the graph of M is in $\mathcal{F} \times \mathcal{B}(X)$ and hence in $\overline{\mathcal{F}} \times \mathcal{B}(X)$. Note that $\overline{\mathcal{F}}$ is closed under the Souslin operation (see [10, Example 3.5.20 and Theorem 3.5.22]). Hence, [8, Corollary of Theorem 7] (see also the survey by Wagner [11, Theorem 3.4]) implies the existence of a $\overline{\mathcal{F}}$ -measurable selection $x(\omega) = (x_\infty(\omega), x_1(\omega), x_2(\omega), \dots) \in \tilde{M}(\omega)$. The set $\bigcup_{k \in \mathbb{N} \cup \{\infty\}} \{x_k(\omega)\}$ is sequentially compact for each $\omega \in \tilde{\Omega}$. By the same arguments as in [3, Proposition 2.15], there exists some deterministic compact set $\tilde{K} \subset X$ such that

$$\overline{\mathbb{P}}(x_k(\omega) \in \tilde{K} \text{ for all } k \in \mathbb{N} \cup \{\infty\}) > 1 - \mathbb{P}(\tilde{\Omega}).$$

Using the definition of $\tilde{\Omega}$ and \tilde{M} it follows that

$$\overline{\mathbb{P}}(x(\omega) \in M(\omega) \text{ and } x_k(\omega) \in \tilde{K} \text{ for all } k \in \mathbb{N} \cup \{\infty\}) > 0.$$

This contradicts the fact that the pullback attractor attracts \tilde{K} almost surely. Hence, $M(\omega) = \emptyset$ almost surely. Using pullback continuity of φ , it follows that there exists some $\hat{\Omega} \in \mathcal{F}$ with $\mathbb{P}(\hat{\Omega}) = 1$ such that for any $\omega \in \hat{\Omega}$ and $(x_\infty, x_1, x_2, \dots) \in \hat{c}$,

$$(1) \quad \limsup_{t \rightarrow \infty} \sup_{k \in \mathbb{N} \cup \{\infty\}} d(\varphi_t(\theta_{-t}\omega, x_k), A(\omega)) = 0.$$

Now, assume there exists some compact set K , $\varepsilon > 0$, $\omega \in \hat{\Omega}$ and sequence t_m going to infinity such that $\varphi_{t_m}(\theta_{-t_m}\omega, K) \not\subset A(\omega)^\varepsilon$ for all $m \in \mathbb{N}$. Hence, there are $y_m \in K$ such that $\varphi_{t_m}(\theta_{-t_m}\omega, y_m) \notin A(\omega)^\varepsilon$ for all $m \in \mathbb{N}$. Since K is compact, there is a convergent subsequence y_{m_k} with $y_\infty := \lim_{k \rightarrow \infty} y_{m_k}$ and $(y_\infty, y_{m_1}, y_{m_2}, \dots) \in \hat{c}$ which is a contradiction to (1). \square

REMARK 3.2. The statement of Lemma 3.1 remains true for pullback attractors of RDS in discrete time.

LEMMA 3.3. *Let A be the pullback attractor of the RDS φ . For $\delta > 0$ there exist compact sets $K_n \subset X$ and $t_n \geq 0$, $n \in \mathbb{N}$ such that*

$$\mathbb{P}(\varphi_{t_n}(\theta_{-t_n}\omega, K_n) \supset A(\omega) \text{ and } \varphi_t(\theta_{-t}\omega, K_n) \subset A(\omega)^{\frac{1}{n}} \text{ for all } t \geq t_n, n \in \mathbb{N}) \geq 1 - \delta.$$

PROOF. Let $n \in \mathbb{N}$. By [3, Proposition 2.15] there exists some compact set $K_n \subset X$ such that

$$(2) \quad \mathbb{P}(A(\omega) \subset K_n) \geq 1 - \frac{\delta}{2^{n+1}}.$$

The definition of the pullback attractor implies that there exists some $t_n > 0$ such that

$$(3) \quad \mathbb{P}(\varphi_t(\theta_{-t}\omega, K_n) \subset A(\omega)^{\frac{1}{n}} \text{ for all } t \geq t_n) \geq 1 - \frac{\delta}{2^{n+1}}.$$

By φ -invariance of A , θ -invariance of \mathbb{P} and (2) it follows that

$$\mathbb{P}(\varphi_{t_n}(\theta_{-t_n}\omega, K_n) \supset A(\omega)) \geq 1 - \frac{\delta}{2^{n+1}}.$$

Combining this estimate and (3), we conclude

$$\mathbb{P}(\varphi_{t_n}(\theta_{-t_n}\omega, K_n) \supset A(\omega) \text{ and } \varphi_t(\theta_{-t}\omega, K_n) \subset A(\omega)^{\frac{1}{n}} \text{ for all } t \geq t_n) \geq 1 - \frac{\delta}{2^n}$$

which implies the claim. \square

THEOREM 3.4. *Let X be a connected Polish space and φ be a pullback continuous RDS. If there exists a pullback attractor A , then A is almost surely connected.*

PROOF. Assume A is not connected with positive probability. By Lemma 3.1 and 3.3 we can choose $\tilde{\Omega} \in \mathcal{F}$ with $\mathbb{P}(\tilde{\Omega}) > 0$, compact sets $K_n \subset X$ and a sequence t_n such that for any $\omega \in \tilde{\Omega}$, $n \in \mathbb{N}$ and compact set $K \subset X$ it holds that

- $A(\omega)$ is not connected,
- $\lim_{t \rightarrow \infty} \sup_{x \in K} d(\varphi_t(\theta_{-t}\omega, x), A(\omega)) = 0$,
- $\varphi_{t_n}(\theta_{-t_n}\omega, K_n) \supset A(\omega)$ and $\varphi_t(\theta_{-t}\omega, K_n) \subset A(\omega)^{\frac{1}{n}}$ for all $t \geq t_n$.

Fix $\omega \in \tilde{\Omega}$. For this fixed ω we will follow the idea of the proof in the deterministic case (see [6, Theorem 3.1]). Note however that Step 3 below requires some extra argument in our case.

Step 1: Let $A(\omega) = A_1 \cup A_2$, where A_1 and A_2 are nonempty, disjoint, compact sets. There exists some $\varepsilon > 0$ such that $A_1^\varepsilon \cap A_2^\varepsilon = \emptyset$. Define

$$\begin{aligned} X_1 &:= \{x \in X : \text{there exists } t \text{ such that } \varphi_s(\theta_{-s}\omega, x) \in A_1^\varepsilon \text{ for all } s \geq t\} \\ X_2 &:= \{x \in X : \text{there exists } t \text{ such that } \varphi_s(\theta_{-s}\omega, x) \in A_2^\varepsilon \text{ for all } s \geq t\}. \end{aligned}$$

If we show that X_1 and X_2 are disjoint nonempty open sets satisfying $X_1 \cup X_2 = X$, then we found a contradiction to X being connected. Obviously, $X_1 \cap X_2 = \emptyset$.

Step 2: We show that $X_1 \cup X_2 = X$.

Let $x \in X$. By definition of $\tilde{\Omega}$, there exists some $t > 0$ such that $\varphi_s(\theta_{-s}\omega, x) \in A(\omega)^\varepsilon$ for all $s \geq t$. Define

$$S_t := \{\varphi_s(\theta_{-s}\omega, x) : s \geq t\}.$$

Then, $S_t \subset A(\omega)^\varepsilon$ and S_t is connected by pullback continuity. Therefore, S_t is either totally contained in A_1^ε or totally contained in A_2^ε .

Step 3: We show that $X_i \neq \emptyset$ for $i = 1, 2$.

Let $n \in \mathbb{N}$ with $\frac{1}{n} \leq \varepsilon$. By definition of $\tilde{\Omega}$, $\varphi_{t_n}(\theta_{-t_n}\omega, K_n) \supset A(\omega)$ and $\varphi_t(\theta_{-t}\omega, K_n) \subset A(\omega)^\varepsilon$ for all $t \geq t_n$ for some $n \in \mathbb{N}$. Hence, there exists $x \in K_n \subset X$ such that $\varphi_{t_n}(\theta_{-t_n}\omega, x) \in A_i$. By continuity in time, $\varphi_t(\theta_{-t}\omega, x) \in A_i^\varepsilon$ for all $t \geq t_n$.

Step 4: We show that X_i is open for $i = 1, 2$.

Assume that X_i is not open. Then, there exist an $x \in X_i$, a sequence x_k converging to x and a sequence s_k converging to infinity such that $\varphi_{s_k}(\theta_{-s_k}\omega, x_k) \notin A_i^\varepsilon$ for all $k \in \mathbb{N}$. By definition of $\tilde{\Omega}$, there exists some $s > 0$ such that $\varphi_t(\theta_{-t}\omega, x_k) \in A(\omega)^\varepsilon$ for all $k \in \mathbb{N}$ and $t \geq s$. Since $x \in X_i$, x_k is converging to x and φ is continuous in the state space, there exists some k^* such that $\varphi_s(\theta_{-s}\omega, x_k) \in A_i^\varepsilon$ for $k \geq k^*$. Using pullback continuity, it follows that $\varphi_t(\theta_{-t}\omega, x_k) \in A_i^\varepsilon$ for $t \geq s$ and $k \geq k^*$ which is a contradiction to the definition of x_k . \square

4. WEAK ATTRACTOR

The question arises whether the result in the previous section can be extended to weak attractors. In contrast to pullback attractors, convergence to weak attractors is merely in probability. We give an example of an RDS where the weak attractor

is not connected. In addition to the assumption on the RDS and state space of Section 3, this example has a jointly continuous RDS, a path-connected state space and every bounded set converges to the attractor.

EXAMPLE 4.1. *Step 1: The metric space.* We choose the same metric space as in [6, Remark 5.2]. Set $s_n = \sum_{i=0}^n 2^{-i}$ for $n \in \mathbb{N}_0$. Let us consider the following sets in \mathbb{R}^2 :

$$\begin{aligned}
 P_{-\infty} &:= (-1, 0), & P_{\infty} &:= (2, 0), \\
 P_n &:= (s_{n-1}, 0), & P_{-n} &:= (1 - s_n, 0), \\
 X_n^L &:= \{(x, y) \in \mathbb{R}^2 : x = s_{n-1} + \lambda 2^{-n-1} \text{ and} \\
 & \quad y = \lambda 2^{-n} \text{ for some } \lambda \in [0, 1]\}, \\
 X_n^R &:= \{(x, y) \in \mathbb{R}^2 : x = s_{n-1} + (2 - \lambda)2^{-n-1} \text{ and} \\
 & \quad y = \lambda 2^{-n} \text{ for some } \lambda \in [0, 1]\}, \\
 X_{-n}^L &:= \{(x, y) \in \mathbb{R}^2 : x = 1 - s_n + \lambda 2^{-n-1} \text{ and} \\
 & \quad y = \lambda 2^{-n} \text{ for some } \lambda \in [0, 1]\}, \\
 X_{-n}^R &:= \{(x, y) \in \mathbb{R}^2 : x = 1 - s_n + (2 - \lambda)2^{-n-1} \text{ and} \\
 & \quad y = \lambda 2^{-n} \text{ for some } \lambda \in [0, 1]\}, \\
 X_{-\infty} &:= \{(-1, y) \in \mathbb{R}^2 : y \geq 0\}, \\
 Y &:= \{(x, y) \in \mathbb{R}^2 : y \leq 0, (x - 0.5)^2 + y^2 = 2.25\}
 \end{aligned}$$

and


$$X_z := X_z^L \cup X_z^R$$

for $n \in \mathbb{N}_0$ and $z \in \mathbb{Z}$. The sets X_z are the two equal sides of isosceles triangles in the halfplane with base $P_z P_{z+1}$ and height 2^{-z} . The left- respectively right-hand side of X_z is denoted by X_z^L respectively X_z^R . Finally we define the complete metric space

$$X := \bigcup_{z \in \mathbb{Z}} X_z \cup X_{-\infty} \cup Y$$

with the metric induced by \mathbb{R}^2 .

Step 2: The dynamics. We characterize the dynamics by phases of length one. To each phase there corresponds a random variable ξ_m where $(\xi_m)_{m \in \mathbb{Z}}$ is a sequence of independent identically distributed random variables with $\mathbb{P}(\xi_0 = k) = 2^{-k}$ for $k \in \mathbb{N}$. In a phase with corresponding $\xi_m = k$ all points to the right of $P_{-(k+1)!+1}$ get pushed $k!$ triangles to the right and all points on the lower half of the triangles to the left of $P_{-(k+1)!}$ decrease their height. We describe the dynamics during a phase by a function $f : \{0 \leq s \leq t \leq 1\} \times \mathbb{N} \times X \mapsto X$. Let f be such that

Figure 1. Bounded subset of X

- $P \mapsto f_{0,t}(k, P)$ is bijective
- $f_{s,t} = f_{0,t} \circ f_{0,s}^{-1}$
- $(s, t) \mapsto f_{s,t}(k, P)$ is continuous
- if $z \geq -(k+1)!$ and $P = (x, y) \in X_z^R$, then $f_{0,1}(k, P) \in \{(\tilde{x}, \tilde{y}) \in X_{z+k!}^R : \tilde{y} = 2^{-k!}y\}$
- if $z \geq -(k+1)! + 1$ and $P = (x, y) \in X_z^L$, then $f_{0,1}(k, P) \in \{(\tilde{x}, \tilde{y}) \in X_{z+k!}^L : \tilde{y} = 2^{-k!}y\}$
- if $z \geq -(k+1)! + 1$ and $P \in X_{z-1}^R \cup X_z^L$, then $|f_{0,t}(k, P) - f_{0,t}(k, P_z)| \leq |P - P_z|$
- if $z \leq -(k+1)!$ and $P = (x, y) \in X_z^L$ with $y \leq 2^{-z-1}$, then $f_{0,t}(k, P) \in \{(\tilde{x}, \tilde{y}) \in X_z^L : \tilde{y} = 2^{-t}y\}$
- if $z \leq -(k+1)! - 1$ and $P = (x, y) \in X_z^R$ with $y \leq 2^{-z-1}$, then $f_{0,t}(k, P) \in \{(\tilde{x}, \tilde{y}) \in X_z^R : \tilde{y} = 2^{-t}y\}$
- if $P, Q \in X_z^L$ or $P, Q \in X_z^R$ for $z \in \mathbb{Z}$, then $|f_{s,t}(k, P) - f_{s,t}(k, Q)| \leq 4(k! + 1)|P - Q|$
- if $P \in X_{-\infty}$ and $P = (-1, y)$, then $f_{0,t}(k, P) = (-1, 2^{-t}y)$
- if $P \in Y$, then $f_{0,t}(k, P) = P$.

Then, $t \mapsto f_{s,t}(\zeta_m, P)$ describes the dynamics of the system started in a point P at time s in a phase with corresponding random variable ζ_m . Since $(s, t) \mapsto f_{s,t}(k, P)$ is continuous and $P \mapsto f_{s,t}(k, P)$ is Lipschitz continuous with Lipschitz constant depending on k , the map $(s, t, P) \mapsto f_{s,t}(k, P)$ is continuous.

In the following steps we show that the weak attractor of this system exists and is not connected.

Step 3: Attractor of discrete-time system. Let $r \in \mathbb{N}$ be arbitrary. Define the bounded set $K_r := \{(x, y) \in X : y \leq 2^r\}$ and the neighborhood $U_r = \{(x, y) \in X : y \leq 2^{-r}\}$ of $\bigcup_{z \in \mathbb{Z}} P_z \cup Y$. Consider the discrete-time system generated by the iterated functions $(f_{0,1}(\zeta_m, \cdot))_{m \in \mathbb{Z}}$. If $\zeta_m \geq k$ for some phase with $k! \geq 2r$,

then the process started in $\bigcup_{z=-r}^{\infty} X_z \cap K_r$ stays in U_r after this phase. Running $2r$ phases, all points in $K_r \cap (\bigcup_{i=r+1}^{\infty} X_{-i} \cup X_{-\infty})$ decrease their height and reach U_r . Therefore, after $2r$ phases where at least one corresponding $\xi_m \geq k$ with $k! \geq 2r$ the discrete-time process started in K_r is in U_r . In contrast to the continuous-time process, the discrete-time process cannot leave U_r afterwards. By [2, Theorem 3.4], there exists a pullback attractor of the discrete-time process and this attractor is a subset of $\bigcup_{z \in \mathbb{Z}} P_z \cup Y$. For $n \in \mathbb{N}$ define

$$F_n(\xi_{-1}, \xi_{-2}, \dots, \xi_{-n}) := f_{0,1}(\xi_{-1}, \cdot) \circ f_{0,1}(\xi_{-2}, \cdot) \circ \dots \circ f_{0,1}\left(\xi_{-n}, \bigcup_{z \in \mathbb{Z}} P_z\right) \\ \subset \bigcup_{z \in \mathbb{Z}} P_z.$$

By definition of the pullback attractor, $F_n(\xi_{-1}, \xi_{-2}, \dots, \xi_{-n})$ converges to the pullback attractor as n goes to infinity \mathbb{P} -almost surely. Therefore, $P_0 \in F_n$ for large enough n implies that P_0 is in the attractor as well. The point P_0 is not in F_n iff there exist $k \in \mathbb{N}$ and times $-n \leq t_0 < t_1 < \dots < t_k < 0$ such that $\xi_{t_i} = k$ for all $0 \leq i \leq k$ and $\xi_s \leq k$ for all $t_0 \leq s < 0$. Then,

$$\mathbb{P}(P_0 \text{ is in the attractor}) = \lim_{n \rightarrow \infty} \mathbb{P}(P_0 \in F_n(\xi_{-1}, \xi_{-2}, \dots, \xi_{-n})) \\ \geq 1 - \sum_{k \in \mathbb{N}} \mathbb{P}(\xi_0 = k \mid \xi_0 \geq k)^{k+1} = \frac{1}{2}$$

which implies that the pullback attractor is not connected with positive probability. More generally, the attractor is not connected if there exists an $m \geq 0$ such that for all $n \in \mathbb{N}$ the point $P_0 \in F_n(\xi_{-m-1}, \xi_{-m-2}, \dots, \xi_{-m-n})$. This event is in the terminal sigma algebra. By Kolmogorov's zero-one law, the pullback attractor of the discrete-time system is almost surely not connected.

Step 4: Attractor of continuous-time system. When we consider the continuous-time system we need to add a random phase shift which is uniformly distributed on $[0, 1)$. For $0 \leq s, t < 1$ and $n \in \mathbb{N}$, the system started in a point P at time s of a phase is described by

$$\varphi_{-s+n+t}(\omega, P) = f_{0,t}(\xi_n, \cdot) \circ f_{0,1}(\xi_{n-1}, \cdot) \circ \dots \circ f_{0,1}(\xi_1, \cdot) \circ f_{s,1}(\xi_0, P).$$

with $\omega = (s, (\xi_m)_{m \in \mathbb{Z}}) \in [0, 1) \times \mathbb{N}^{\mathbb{Z}} =: \Omega$ and canonical shift on Ω and the basic probability measure on Ω is the product of Lebesgue measure on $[0, 1)$ and the laws of $(\xi_m)_{m \in \mathbb{Z}}$. Then, φ is a jointly continuous RDS as a composition of jointly continuous maps.

Let $r \geq 2$. If we start in a set K_r as in Step 3 in an incomplete phase with corresponding $\xi_m \leq r$, then at the end of this phase the process is still in K_r . The pullback attractor of the discrete-time system attracts this bounded set. Hence, there exists a time $n_r \in \mathbb{N}$ such that the discrete process started in K_r stays in a ball around the discrete-time attractor with radius $2^{-(r+1)!}$ after time n_r with probability $1 - 2^{-r}$.

We extend the discrete-time attractor to continuous time in such a way that the so constructed random set stays strictly invariant under the given dynamics. If one starts the end phase in a ball around the discrete-time attractor with radius $2^{-(r+1)!}$, one can leave the ball around the invariantly extended random set with radius $2^{-(k+1)!}$ only during a phase with corresponding $\xi_m \geq r$.

Combining these three parts, the continuous-time process started in K_r at time $t \geq n_r + 1$ is in a ball around the discrete-time attractor with radius $2^{-(r+1)!}$ with probability $1 - 2^{-r+1}$.

This probability tends to one as r goes to infinity. Therefore, the continuous-time extension of the discrete-time attractor is the weak attractor of the continuous-time system. By construction, the weak attractor of the continuous system is almost surely not connected. Note that the weak attractor will not almost surely be contained in the set $\bigcup_{z \in \mathbb{Z}} P_z \cup Y$.

REMARK 4.2. If every compact set in X can be covered by a connected compact set, then the weak attractor is connected. This follows by the same arguments as in [2, Proposition 3.7] where this result was stated for the pullback attractor. Here, one does not need to assume continuity in time.

This assumption is in particular satisfied for an attractor that attracts bounded sets in probability on a connected and locally connected Polish space. By local connectedness, a compact set can be covered by finitely many open connected sets. Since a connected and locally connected Polish space is also path-connected (see Mazurkiewicz-Moore-Menger theorem in [7, p. 254, Theorem 1 and p. 253, Theorem 2]), one can connect these sets by paths.

REFERENCES

- [1] L. ARNOLD, *Random Dynamical Systems*, Springer Monographs in Mathematics. Springer-Verlag, Berlin, 1998.
- [2] H. CRAUEL, *Random point attractors versus random set attractors*, J. London Math. Soc. (2), 63(2):413–427, 2001.
- [3] H. CRAUEL, *Random Probability Measures on Polish Spaces*, Stochastics Monographs. CRC Press, 2003.
- [4] H. CRAUEL - F. FLANDOLI, *Attractors for random dynamical systems*, Probab. Theory Related Fields, 100(3):365–393, 1994.
- [5] F. FLANDOLI - B. GESS - M. SCHEUTZOW, *Synchronization by noise*, Probab. Theory Related Fields, 168(3–4):511–556, 2017.
- [6] M. GOBBINO - M. SARDELLA, *On the connectedness of attractors for dynamical systems*, J. Differential Equations, 133(1):1–14, 1997.
- [7] K. KURATOWSKI, *Topology. Vol. II*, Academic Press, New York–London, 1968.
- [8] S. J. LEESE, *Multifunctions of Souslin type*, Bull. Austral. Math. Soc., 11:395–411, 1974.
- [9] M. SCHEUTZOW - M. WILKE-BERENQUER, *Random Delta-Hausdorff-attractors*, Discrete Contin. Dyn. Syst. Ser. B, 23(3):1199–1217, 2018.
- [10] S. M. SRIVASTAVA, *A course on Borel sets*, volume 180 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1998.

- [11] D. H. WAGNER, *Survey of measurable selection theorems: an update*, in Measure theory, Oberwolfach 1979 (Proc. Conf., Oberwolfach, 1979), volume 794 of Lecture Notes in Math., pages 176–219. Springer, Berlin–New York, 1980.

Received 15 December 2017,
and in revised form 4 April 2018.

Michael Scheutzow
Institut für Mathematik
MA 7-5, Fakultät II
Technische Universität Berlin
Straße des 17. Juni 136
10623 Berlin, Germany
ms@math.tu-berlin.de

Isabell Vorkastner
Institut für Mathematik
MA 7-5, Fakultät II
Technische Universität Berlin
Straße des 17. Juni 136
10623 Berlin, Germany
vorkastn@math.tu-berlin.de

