

1 Red mud a byproduct of Aluminum production contains soluble vanadium that causes
2 genotoxic and cytotoxic effects in higher plants

3
4 3
5
6
7 4 Miroslav Mišík¹, Ian T. Burke², Matthias Reismüller¹, Clemens Pichler¹, Bernhard Reiner¹,
8
9 5 Katarina Mišíková³, William M. Mayes⁴, Siegfried Knasmueller^{1*}

10
11 6
12
13
14 7
15
16 8 ¹Institute of Cancer Research, Department of Medicine I, Comprehensive Cancer Center,
17
18 9 Medical University of Vienna, Austria

19
20
21 10 ²Earth Surface Science Institute, School of Earth and Environment, University of Leeds,
22
23 11 Leeds LS2 9JT, UK

24
25
26 12 ³Department of Botany, Faculty of Natural Sciences, Comenius University Bratislava,
27
28 13 Slovakia

29
30
31 14 ⁴Centre for Environmental and Marine Sciences, University of Hull, Scarborough, YO11
32
33 15 3AZ, UK

34
35
36 16
37
38
39 17
40
41 18
42
43 19 Corresponding author:

44
45 20 Siegfried Knasmueller; Institute of Cancer Research, Borschkegasse 8a, A-1090 Vienna
46
47
48 21 Austria; Phone: ++43-1-40160 57561, Fax ++43 1 40160 957500; mail:
49
50 22 siegfried.knasmueller@meduniwien.ac.at

23 ABSTRACT

1
2 24 Red mud (RM) is a byproduct of aluminum production, worldwide between 70 and 120
3
4 25 million tons are produced annually. We analyzed RM which was released in the course of the
5
6
7 26 Kolonatar disaster in Hungary into the environment in acute and genotoxicity experiments
8
9 27 with plants which are widely used for environmental monitoring. We detected induction of
10
11 28 micronuclei which reflect chromosomal damage in tetrads of Tradescantia and in root cells of
12
13
14 29 Allium as well as retardation of root growth with contaminated soils and leachates. Chemical
15
16
17 30 analyses showed that RM contains metals, in particular high concentrations of vanadium.
18
19 31 Follow up experiments indicated that vanadate causes the effects in the plants. This
20
21 32 compound causes also in humans DNA damage and positive results were obtained in
22
23
24 33 carcinogenicity studies. Since it was found also in RM from other production sites our
25
26 34 findings indicate that its release in the environment is a global problem which should be
27
28
29 35 studied in more detail.

30
31
32 36

33
34
35 37

36
37 38

38
39
40 39

41
42 40 **Capsule abstract:** Our findings indicate that the red mud causes genotoxic effect in plants
43
44 41 probably due to the presence of vanadate which is contained at high concentrations in the
45
46 42 residue.

47
48
49 43

50
51
52 44

53
54 45

55
56 46 Key words: red mud, Tradescantia, micronuclei, Allium, vanadium

57
58
59 47

1. Introduction

Red mud is a by-product of aluminum production with the Bayer process. Its global production is in the range between of 70 to 120 million tons per year (Mayes et al., 2011b). The material consists mainly of iron-, aluminum- and titanium- oxides and hydroxides (Burke et al., 2012; Mayes et al., 2011b). Chemical analyses showed that its contains also radionuclides (e.g. ^{226}Ra , ^{230}Th and ^{40}K), as well as heavy metals including As, Cr, Co, Cd, Ni and V (Mayes et al., 2011a; Ruyters et al., 2011).

On Oct 4th 2010, approximately one million m³ of the residue was released into the environment from the aluminum plant Ajkai Timfoldgyar Zrt in Western Hungary. According to the Hungarian Ministry of Interior the “Kolontar disaster” is the biggest environmental catastrophe which ever happened in this country (Ádám et al., 2011). Hundreds of houses were destroyed, 265 individuals were injured and ten died (Gundy et al., 2013).

After the accident, attempts were made to investigate the impact of the release of the material into the environment and to assess the health consequences in humans. Ecotoxicological studies were conducted with different plant species and bacteria concerning acute toxic effects (Klebercz et al., 2012; Ruyters et al., 2011); the motility and the concentrations of toxic trace elements were studied in physico-chemical measurements (Burke et al., 2012). Furthermore, studies were conducted to assess the consequences of inhalation of dust particles in humans and rodents (Czovek et al., 2012; Gelencser et al., 2011).

Radionuclides as well as certain heavy metals (found in red mud) cause damage of the genetic material (Knasmuller et al., 1998; Minouflet et al., 2005) which may lead to destabilization of ecosystems (Sarkar et al., 2006; Zvereva et al., 2008) and also causes adverse effects in humans such as cancer, ageing, infertility and birth defects in the offspring (Aitken and De Iuliis, 2007; Assem and Levy, 2009). Therefore it is of particular interest if

73 the release of this material into the environment induces chromosomal damage. This question
1
2 74 has been addressed in a human study (Gundy et al., 2013) and in bacterial tests (Gelencser et
3
4 75 al., 2011) but no firm conclusions can be drawn of these studies (for details see discussion).
5
6

7 76 The primary aim of the present study was the investigation of the genotoxic properties
8
9 77 of red mud in two plant bioassays, namely in the micronucleus (MN) test with tetrads of
10
11 78 *Tradescantia* (Trad-MN assay) and with root tip cells of *Allium cepa* (A-MN assay). The
12
13 79 experiments were conducted with tetrads as they reflect damage in meiotic germ cells. Root
14
15 80 tip cells were used as they enable the detection of effects in mitotic cells. It was postulated
16
17 81 that differences exist in regard to sensitivity of these cell types towards DNA reactive
18
19 82 compounds (Rodrigues et al., 1997). MN are formed as a consequence of chromosomal
20
21 83 breakage or aneuploidy and can be monitored in a variety of organisms (Heddle et al., 2011).
22
23 84 In addition acute toxic effects were studied in root cells of *A. cepa* by measuring the impact of
24
25 85 the material on the root growth and by calculating the division rates of the cells. These
26
27 86 bioassays are at present the most widely used genotoxicity tests with higher plants and have
28
29 87 been employed in more than 300 investigations for the detection of DNA damaging properties
30
31 88 of chemicals and complex mixtures (for reviews see Leme and Marin-Morales, 2009; Misik et
32
33 89 al., 2011). We used these test systems since they provide information on environmental
34
35 90 effects and it is known that they are, in contrast to other genotoxicity assays with bacterial
36
37 91 indicators and mammalian cells (which are also used for environmental monitoring), highly
38
39 92 sensitive towards radiation (Ma and Davies, 2009; Misik et al., 2011) and heavy metals
40
41 93 (Knasmuller et al., 1998; Majer et al., 2002; Steinkellner et al., 1998).
42
43
44
45
46
47
48
49

50
51 94 We investigated the acute cytotoxic and genotoxic activities of red mud and of soils
52
53 95 which were contaminated with the material collected from fields and gardens which were
54
55 96 used for cultivation of crops and vegetables. Furthermore, we studied also the effects of
56
57 97 waters leached from red mud and affected soils. In additional experiments, attempts were
58
59
60
61
62
63
64
65

98 made to identify the compound(s) which cause genotoxic effects. We determined the
1
2 99 concentrations of trace elements in solid samples and in leachates and conducted additional
3
4 100 Trad-MN assays with sodium metavanadate (NaVO_3) since vanadium was found to be the
5
6
7 101 most abundant heavy metal in the samples.
8
9

102 103 **2. Materials and methods**

104 105 2.1 Sampling

106 Soil samples were collected from the Torna catchment on December 15th 2010. Table
107 1 contains a description of the sampling sites and specifications of their GPS locations. From
108 each site, samples were collected from the 0-5; 5-15 and 15-25 cm horizons to assess if
109 potentially toxic constituents in red mud contaminated the soils.

110 Total organic carbon (TOC) was determined in soil samples using a LECO SC-144DR
111 elemental analyzer after removal of the inorganic carbon fraction using 20% HCl.

112
113 Table 1

114 Figure 1

115 2.2 Preparation of the leachates.

116 Soils were used as sampled (field moist). 50 g of each sample were suspended in 100
117 mL of deionised water in a 250 mL glass beaker for 2 hours using a magnetic stirrer. The
118 extracts were filtered with filter paper before they were tested in stem absorption experiments.
119 The pH values of the leachates were measured with a pH Meter 526 (WTW, Weilheim,
120 Germany) and are listed in Table 1.

122

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

123 2.3 Measurements of the trace elements in the soil samples and leachates.

124 The detection of the trace elements is described in detail in a recent paper of Renforth
125 et al. (2012). Prior to analysis by XRF the samples was prepared as follows. For major
126 element analysis, samples were prepared as fused beads (after loss on ignition at 1050 °C)
127 with lithium metaborate/tetraborate flux (Johnson - Matthey Spectroflux JM100B) (0.6g
128 sample; 3 g Flux). For minor / trace element analysis approximately 10 g dried sample was
129 prepared as a pressed pellet using ~10-20 drops of 6.6% w/v polyvinyl alcohol in a 1:6 mix of
130 methanol and distilled deionised water as a binder (Moviol 88 solution). Elemental analysis of
131 the soil composition was achieved using a PANalytical Axios Advanced X-ray Fluorescence
132 (XRF) spectrometer (data corrected for loss on ignition; % weight loss after furnace treatment
133 at 1050°C). The aqueous phase produced during water extractions was separated from solids
134 by centrifugation (10 min; 2000 g) followed by membrane filtration (0.45 µm); the filtered
135 samples were then acidified by addition of 2% v/v HNO₃. Aqueous elemental concentrations
136 were then determined using a Perkin Elmer Optima 5300 DV ion-coupled plasma, optical
137 emission spectrometer (ICP-OES).

138

139 2.4 Tradescantia micronucleus assays

140 Tradescantia clone #4430 was cultivated according to the protocol of Ma et al. (1994)
141 For stem absorption experiments, 15 young inflorescences were treated in each group. The
142 stems were cut to a length of 10 - 15 cm, transferred to plastic breakers (250 mL) and exposed
143 to aqueous leachates (soil:water – 1:2) of the contaminated soils for 24 h. Subsequently, they
144 were transferred to water for a 24 h recovery period. The flower buds were then collected and

145 fixed in acetic acid and ethanol (1/3, v/v). After 24 h, they were transferred to 70% ethanol.

146 Maleic hydrazide (20 mg L⁻¹; MH, Sigma-Aldrich, SL, US) was used as a positive control.

147 The protocol for root absorption assays is described in an article of Steinkellner et al.
148 (1998). The soil which was used in the control cultures was biological, pesticide free (from
149 Composana, Wien, Austria). Intact plants were removed from hydroponic culture,
150 subsequently the roots were rinsed and individual plants with at least 15 inflorescences placed
151 into plastic pots (250 ml, diameter 12 cm) which were filled with the different soil samples.
152 The plants were exposed under standard conditions for 48 h (Ma et al., 1994). Subsequently,
153 15 inflorescences were collected from each soil sample in all experiments and fixed in a
154 solution of acetic acid and ethanol (1/3, v/v). After 24 h, they were transferred to 70%
155 ethanol.

156 To find out if vanadium accounts for the effects which were found in the soils and in
157 the leachates, additional experiments were carried out with sodium metavanadate. The salt
158 was dissolved in 0.1M NaOH (pH 13) at 1000 ppm (colorless solution) since earlier findings
159 indicate that vanadium is present as V⁵⁺ in the soil (Burke et al., 2012). The stock solution
160 was diluted with tap water and different concentrations (0.5-10.0 ppm) were used in
161 experiments with *Tradescantia*.

162 Slides were prepared and evaluated as described in the protocol of Ma et al. (1994)
163 The tetrads were stained with a 2% aceto-carmin solution. Early tetrads were analyzed under
164 400-fold magnification. For each experimental point a minimum of five buds with early
165 tetrads was evaluated (300 tetrads were evaluated per slide, 1500 per sample).

166 2.5 Micronucleus assay with *Allium cepa*

167 The experiments were carried out according to the standard protocol published by Ma
168 et al. (1995) with slight modifications. Young onion bulbs (diameter 12-21 mm, Schneeball

169 weis, Austroaat, Wien, Austria) were placed in 13 ml glass tubes filled with tap water for
1
2 170 24 h in the dark. Subsequently, the roots (lengths ca. 0.5-1 cm) were exposed directly in the
3
4
5 171 soils in the dark for 24 h and then transferred to fresh tap water for another 24 h. At the end of
6
7 172 the recovery phase, the maximal root length of each onion was measured and the material was
8
9 173 fixed in a mix of acetic acid and ethanol (1/3, v/v) for 24 h and stored in 70% ethanol. MH
10
11
12 174 (10 mg L⁻¹; Sigma-Aldrich, SL, US) was used in all experiments as a positive control.
13

14 175 The root tips were hydrolyzed in a 1:1 mix of HCl (5.0 N) and ethanol (99%) for
15
16 176 3 min and washed in tap water before they were stained with 2% acetocarmine. MN were
17
18
19 177 scored according to the criteria described by Ma et al. (1995). For each experimental point,
20
21 178 the MN frequencies were determined in five plants. From each bulb, two slides were made
22
23
24 179 and 500 cells were evaluated per slide (5000 cells per dose). Furthermore, also the mitotic
25
26 180 indices (MIs) were determined in 1000 cells (100 cells/root) per experimental point. The
27
28
29 181 microscopic evaluation was carried out under a light microscope (Nikon YS200, Japan) with
30
31 182 400-fold magnification.
32

33
34 183

35 36 37 184 2.6 Statistical evaluation 38

39
40 185 The results of the MN experiments were analyzed with one-way ANOVA followed by
41
42 186 Dunnett's multiple comparison test. The results of the experiments concerning the mitotic
43
44
45 187 indices (MI) were analyzed with the Kruskal-Wallis test followed by Dunn's comparison test.
46
47 188 P-values ≤ 0.05 were considered as significant.
48

49
50 189

51
52 190

53
54 191

55
56 192

57
58

59
60

61
62

63
64
65

193 **3. Results**

195 3.1 TOC and pH values of the samples.

196 It can be seen in Table 1 that values between 8.0 and 9.3 were found in the soil
197 leachates which were contaminated with red mud. Most of the samples had a total organic
198 carbon content between 1.0-1.5%, which is typical for agricultural soils in the Ajka area
199 (Lehoux et al., 2013). The sample from the upper horizon of the site 3 had a substantially
200 lower TOC content (0.22%) and a higher pH (9.33) which is typical for red mud (Lehoux et
201 al., 2013).

203 3.2 Micronucleus assays with Tradescantia (experiments with soils and leachates)

204 The results which were obtained in Trad-MN assays with red mud contaminated soil
205 and with aqueous leachates from these samples are summarized in Fig. 2A and 2B. It can be
206 seen that direct exposure of intact plants in material from the contaminated sites 2 (S2) and 3
207 (S3) caused a significant increase of the MN frequencies. In the case of S3, the strongest
208 effect was seen with material from the upper layer, samples from deeper horizons caused only
209 a statistically not significant increase. With the two samples from S4, clear cut negative
210 results were obtained. As described above (Table 1), this material was collected after
211 remediation (i.e. after removal of the upper layer).

212 Also with some leachates, positive results were obtained (Fig. 2B). Again the strongest
213 effects were seen with extracts prepared with material from the upper layers (S2 and S3); soils
214 from deeper horizons did not cause significant effects, also the sample which was collected
215 from the cleanup site was devoid of activity.

216
217 Figures 2A and B

218

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

219 3.3 Micronucleus experiments with mitotic root tip cells of *Allium cepa*

220 The results of assays with *A. cepa* are shown in Figures 3A – C. The roots were
221 exposed in these experiments directly in the samples. Since division delays may decrease the
222 formation of MN (Fenech and Morley, 1985), we determined in all experiments also the
223 mitotic indices (MI).

224 It can be seen in Figure 3A that a significant increase of the MN rates was observed
225 with some samples. The most pronounced effect was detected with material collected from
226 the middle layer of S2 while red mud enriched sample from surface of the same location did
227 not cause a significant increase of the MN rates. This finding can be explained by inhibition
228 of cell division, which was most pronounced with material collected from the upper horizon
229 of this site (Figure 3B). As in the *Tradescantia* experiments, no increase was seen with
230 samples from the remediation site (S4).

231 The impact of the material on root growth is shown in Figure 3C. It can be seen, that
232 clear effects were obtained with samples from the surface of sites S2 and S3.

233
234 Figures 3A – C

235

236 3.4 Chemical analyses of the soil samples and of the leachates

237 The chemical composition of samples which were collected from S3 and S1 (Table
238 2A-B) and of the corresponding leachates (Table 2C) are shown in Table 2.

239

240 Table 2

241

242 The chemical composition of the topmost soil sample (S3 0-5 cm) is similar to that of
1
2 243 other samples of red mud which were collected after the Ajka spill (Burke et al., 2012;
3
4 244 Ruyters et al., 2011). Several potentially toxic metals including As, Cr, Ni, Pb and V were
5
6
7 245 present at elevated concentrations.
8

9 246 The chemical composition of the control sample (S1) is consistent with unaffected
10
11
12 247 soils found in the region (see Table 2c) (Mayes et al., 2011b).
13

14 248 The corresponding leachates from S3, contained elevated concentrations of potentially
15
16
17 249 toxic elements, such as V, As, Cu, Al and Cr, notably, V concentrations were an order of
18
19 250 magnitude higher than the levels of the other metals (except Al). Also chromium levels were
20
21
22 251 relatively high.
23

24 252

27 253 3.5 Tradescantia micronucleus assays with vanadate

28
29
30 254 We found in earlier experiments with Tradescantia that the concentrations of Ni, Cr,
31
32
33 255 Cd, Pb and As, which cause MN induction in tetrads are in the range of hundred to thousand
34
35 256 ppm (Knasmuller et al., 1998; Steinkellner et al., 1998). We hypothesized that V, which was
36
37
38 257 detected in higher levels than other metals, may account for the induction of MN, which was
39
40 258 seen with the soils and the leachates. Therefore, we tested different concentrations of an
41
42
43 259 aqueous vanadate solution in subsequent Trad MN assays. The results are summarized in
44
45 260 Figure 4. It can be seen that exposure of the cuttings to concentrations ≥ 1.0 mg/L caused
46
47 261 significant induction of the MN frequencies i.e. an approximately 6-fold increase over the
48
49
50 262 background.
51

52 263

54 264 Figure 4

57 265

266 3.6 Impact of pH on MN formation in Tradescantia

267 Since the pH values of the soil extracts and of the vanadate solutions were quite high,
268 i.e. in the range between 8.2 and 9.3 (for details see Table 1) we conducted an additional
269 experiment in which plant cuttings were exposed in aqueous solutions (without metals) to
270 different pH values. No indication of pH depended induction of MN by the pH itself was
271 found in this experiment. The numbers of MN per 100 tetrads were 1.7 ± 0.6 ; 2.1 ± 0.6 and
272 1.6 ± 0.6 for pH 8, 8.5 and 9.3 respectively (numbers are means \pm SD of results obtained from
273 1500 tetrads).

274 **4. Discussion**

275
276 The results of this study show that red mud, which was released in large amounts into
277 the environment, causes damage of the genetic material and also acute toxic effects in higher
278 plants.

279
280 4.1 Acute toxic and genotoxic effects of red mud

281 Several ecotoxicological studies have been conducted with red mud from Ajka
282 (Klebercz et al., 2012; Ruyters et al., 2011) and adverse effects were seen in experiments with
283 higher plants; i.e. the material was found to inhibit the root of *Sinapis alba* and affect the
284 shoot yield in barley (Klebercz et al., 2012; Ruyters et al., 2011). In the latter study, the
285 authors monitored also the levels of trace elements which are contained in red mud such as
286 Cu, Cr, Fe and Ni in exposed plants. They detected these metals in the shoots but stress that
287 their levels did not exceed the toxic limits and hypothesize that Na is the prime cause which
288 affected the growth of the indicator plants. Another factor which was made responsible for the
289 toxic effects in *Sinapis* and in experiments with the Ostracod *Heterocypris incongruens* is

290 gypsum which may limit the nutrient supply and increase the availability of contaminants
1
2 291 (Klebercz et al., 2012). The genotoxic effect of red mud has not been investigated in the
3
4 292 invertebrates so far according to our knowledge. However, it is possible that several species
5
6
7 293 may be affected due to the high metal concentration of the material. For example, it is know
8
9 294 that earthworms are highly sensitive towards specific contaminants and specific metals.
10
11
12 295 Genotoxic effects were detected with chromium, nickel and cadmium (Bigorgne et al., 2010;
13
14 296 Bonnard et al., 2010; Manerikar et al., 2008).

16
17 297 The results of the present study show clearly that red mud inhibits the division and
18
19 298 growth of root tip cells in *A. cepa* (Figures 3B and 3C) and causes induction of MN in this
20
21 299 species. Furthermore, we showed that also aqueous leachates of red mud contaminated soils
22
23
24 300 cause induction of MN in both plant bioassays. As shown in Figure 3 a stronger effect was
25
26 301 obtained with sample S2 (5-15 cm) as with the sample from upper layer from the same site.
27
28
29 302 This effect can be explained by reduction of MN formation due to inhibition of cell division.
30
31 303 Furthermore, it is notable that the effects which were detected with soil exposure are more
32
33
34 304 pronounced as those which were found with the leachates. This observation is in agreement
35
36 305 with the results obtained with heavy metals contaminated soils and can be explained by more
37
38 306 efficient uptake via the roots by active transport mechanism (for detail see Steinkellner et al.,
39
40
41 307 1998).

42
43 308 The only genotoxicity assay conducted so far with red mud is the SOS chromotest
44
45 309 which is based on the detection of SOS responses in bacterial indicator cells (Gelencser et al.,
46
47
48 310 2011). Consistently negative results were obtained with samples, which were prepared from
49
50
51 311 fugitive dust from the area where the accident had happened. However, it is notable that it is
52
53 312 known from earlier investigations that bacterial assays are insensitive towards heavy metals
54
55 313 (Majer et al., 2002).

56 314
57
58
59
60
61
62
63
64
65

315 4.2 Identification of vanadate as the active principle of red mud

1
2 316 The results of the chemical analyses show that red mud contaminated soils and their
3
4
5 317 aqueous leachates contain elevated concentrations of V (Table 2C). This observation is in
6
7 318 agreement with the results of earlier investigations with material from Ajka (Mayes et al.,
8
9
10 319 2011a; Mayes et al., 2011b; Renforth et al., 2012). In this context, it is notable that vanadium
11
12 320 was also detected in red mud from several other aluminium production sites (Fontanier et al.,
13
14
15 321 2012; Gawu et al., 2012; Rajeev et al., 1999; Samal et al., 2012).

16
17 322 Spectroscopic evidence showed that vanadium is present primarily as V^{5+} phases
18
19 323 (Burke et al., 2012). Under alkaline conditions (high pH is induced in red mud leachate by
20
21
22 324 dissolution of the NaOH present), V^{5+} is predicted to become readily solubilized in water as
23
24 325 vanadate (Peacock and Sherman, 2004).

26
27 326 Vanadium is more abundant in red mud than other alkali-soluble trace metals (e.g.
28
29 327 As), therefore its environmental mobility is enhanced relative to those of other carcinogenic
30
31
32 328 heavy metals which are present in the material. Therefore, we conducted Trad-MN assays
33
34 329 with sodium metavanadate solutions and compared the concentrations which cause positive
35
36 330 results with those contained in the aqueous leachates. The extracts prepared from the upper
37
38
39 331 layer of S3 contained 1.1 mg/L V (Table 2) and the results which were obtained with aqueous
40
41 332 vanadate solutions (Figure 4) show, that levels ≥ 1.0 mg/L cause significant induction of the
42
43
44 333 MN frequencies in Tradescantia indicating that vanadate accounts for the genotoxic effects
45
46 334 which we detected in the leachate.

48
49 335 The solubility of vanadate in the environment is largely controlled by sorption
50
51 336 reactions with mineral surfaces, with low solution concentrations expected at circumneutral
52
53
54 337 pH (Peacock and Sherman, 2004). If the pH of red mud affected soils can be controlled to pH
55
56 338 < 9 , high aqueous V concentrations and their accompanying genotoxic effects may be avoided
57
58 339 (Lehoux et al., 2013). In this context, it is notable that, several technologies have been

340 developed to recover vanadium from sludge (for review see Rajeev et al., 1999). As
341 mentioned above, we found no impact of the pH on MN frequencies in experiments with
342 waters to which NaOH was added in absence of metal ions. However, the situation may be
343 different in soils which are contaminated with vanadium compounds.

344

4.3 Vanadate is an extremely potent mutagen in the Trad MN assay

346 As shown in Table 2A-C, a number of other genotoxic and carcinogenic heavy metals
347 such as Ni, Cd, Cr and As were detected in the aqueous extracts. Results of earlier Trad-MN
348 assays show that the concentrations of these metals, which are required to cause significant
349 effects, are 10^2 - to 10^3 -fold higher as the levels which were detected in the leachates. The
350 strongest activity was seen in these experiments with $\text{Cr}^{(\text{VI})}\text{O}_3$ (129) followed by As_2O_3 (98.5),
351 $\text{Ni}^{(\text{II})}\text{Cl}_2$ (645), CdCl_2 (1006) and $\text{Pb}(\text{NO}_3)_2$ (1820); numbers in parenthesis indicate the
352 LOELs in mg/L (Knasmueller et al., 2003; Knasmuller et al., 1998; Steinkellner et al., 1998).
353 Furthermore, the results of the chemical analyses demonstrate that, these effects can be
354 attributed to the presence of soluble vanadium. As mentioned above, we found in the present
355 study significant induction of MN by vanadate in *Tradescantia* already at levels ≥ 1 mg/L.
356 Comparisons with the effects seen with other metals indicate that vanadate is more potent in
357 regard to induction of chromosomal damage. According to our knowledge, it has never been
358 tested before in genotoxicity experiments with higher plants, but earlier findings with *Allium*
359 roots showed that 25 mg/L cause stickiness of the chromosomes and a decrease of the MI
360 (Marcano et al., 2006).

361 Results which were obtained in MN assays and other genotoxicity tests with
362 mammalian cells and rodents with vanadium compounds are summarized in an IARC
363 monograph (IARC, 2006) and in a review by Assem and Levy (2009) and Taylor et al.

364 (2012). The available data show that positive results were obtained in most in vitro
1
2 365 experiments with mammalian cells and also in a number of in vivo studies with rodents.
3

4
5 366

6 7 8 367 4.4 Possible consequences of the release of DNA damaging toxins in red mud to the 9 10 368 environment

11
12
13 369 It has been postulated that genotoxins may have an impact on the stability of ecosystems.

14
15
16 370 In this context it is notable, that we found in an earlier study concerning urban air pollution,

17
18 371 that the induction of MN in Tradescantia is paralleled by a decrease of the fertility of wildlife

19
20 372 plants (Misik et al., 2007). It was also shown by other groups that DNA damaging toxins in

21
22 373 the environment cause mutations which lead to transgenerational genomic instability in the

23
24
25 374 offspring (Glen and Dubrova, 2012). The fact that we detected pronounced induction with

26
27 375 leachates of red mud indicates that the compounds which induce genetic damage (presumably

28
29
30 376 V^{5+}) contaminate surface and/or ground waters. Therefore, it is likely that they cause

31
32 377 environmental damage.
33
34

35 378 In regard to potential effects in humans it is known that DNA damage is causally

36
37 379 related to diseases such as cancer, infertility and heritable diseases (Shaugnessy and

38
39
40 380 DeMarini, 2009) and leads to accelerated ageing. We found in an earlier occupational study

41
42 381 that exposure of workers to V^{5+} causes induction of MN in peripheral lymphocytes (Ehrlich et

43
44 382 al., 2008) which is a reliable biomarker for increased cancer risks (Bonassi et al., 2011).

45
46
47 383 Furthermore, it was shown in a long term carcinogenicity study that inhalative exposure to

48
49 384 this compound causes an increase of the tumor rates in the lungs of rats (Assem and Levy,

50
51
52 385 2009), as a consequence of those results V_2O_5 was classified as a possible (group 2B) human

53
54 386 carcinogen by the IARC in 2006 (IARC, 2006).
55
56
57
58
59
60
61
62
63
64
65

387 It can be not excluded that humans living in the area where the accident happened are
1
2 388 exposed to the genotoxic components which are contained in red mud. Gundy et al. (2013)
3
4 389 analyzed chromosomal aberrations (CA) in lymphocytes of individuals who were either
5
6
7 390 injured by the accident or exposed during cleanup activities. The authors conclude that the
8
9 391 exposure does not pose an immediate short term genotoxic hazard but recommend further
10
11 392 studies. In fact, no firm conclusions can be drawn from their investigation. The CA
12
13 393 frequencies of the exposed group were almost 30% higher as those in an urban control group
14
15 394 while no differences were seen with controls from a rural village. Furthermore, it is notable
16
17 395 that the exposure periods of the individuals in the study group were quite heterogeneous (i.e.
18
19 396 between 9 and 336 hrs) which enhardens the detection of effects.
20
21
22
23

24 397 Taken together, the results of the present investigation show for the first time that
25
26 398 DNA damaging toxins were released in the course of the Kolontar disaster and that red mud
27
28 399 contains vanadate which causes damage of the genetic material at low concentrations. As
29
30 400 mentioned above, the metal was detected also in samples from other production sites and
31
32 401 environmental pollution has been reported from other countries, for example from Brazil and
33
34 402 India (2011; Gawu et al., 2012; Lima et al., 2009). The release of genotoxins into the
35
36 403 environment may affect the stability of ecosystems and have a negative impact on human and
37
38 404 environmental health; due to the worldwide production of the residue, the contamination of
39
40 405 the material with DNA reactive and potentially carcinogenic vanadium compounds is a global
41
42 406 problem which should be studied in more detail in the future.
43
44
45
46
47

48 407

49 50 51 408 **Acknowledgement**

52
53
54
55 409 We should like to thank to Thomas Trenker and Herwig Schuster for help in regard to
56
57 410 the collection of the samples.
58
59

60 411

412 References

- 1
2 413 2011. Vedanta Red Mud - Vedanta Resources [www.vedantaresources.com/media/161444/vedanta-](http://www.vedantaresources.com/media/161444/vedanta-red-mud.pdf)
3 414 [red-mud.pdf](http://www.vedantaresources.com/media/161444/vedanta-red-mud.pdf).
- 4 415 Ádám, J., Bánvölgyi, G., Dura, G., Grenerczy, G., Gubek, N., Gutper, I., Simon, G., Szegfalvi, Z., Székács,
5 416 A., Szépvölgyi, J., Ujlaky, E., 2011. The Kolontár Report - Causes and lessons learned from the
6 417 red mud disaster, in: Jávör, B., Hargitai, M. (Eds.), Budapest.
- 8 418 Aitken, R.J., De Iulius, G.N., 2007. Origins and consequences of DNA damage in male germ cells.
9 419 *Reprod Biomed Online* 14, 727-733.
- 10 420 Assem, F.L., Levy, L.S., 2009. A review of current toxicological concerns on vanadium pentoxide and
11 421 other vanadium compounds: gaps in knowledge and directions for future research. *J Toxicol*
12 422 *Environ Health B Crit Rev* 12, 289-306.
- 14 423 Bigorgne, E., Cossu-Leguille, C., Bonnard, M., Nahmani, J., 2010. Genotoxic effects of nickel, trivalent
15 424 and hexavalent chromium on the *Eisenia fetida* earthworm. *Chemosphere* 80, 1109-1112.
- 16 425 Bonassi, S., El-Zein, R., Bolognesi, C., Fenech, M., 2011. Micronuclei frequency in peripheral blood
17 426 lymphocytes and cancer risk: evidence from human studies. *Mutagenesis* 26, 93-100.
- 18 427 Bonnard, M., Devin, S., Leyval, C., Morel, J.L., Vasseur, P., 2010. The influence of thermal desorption
20 428 on genotoxicity of multipolluted soil. *Ecotoxicol Environ Saf* 73, 955-960.
- 21 429 Burke, I.T., Mayes, W.M., Peacock, C.L., Brown, A.P., Jarvis, A.P., Gruiz, K., 2012. Speciation of arsenic,
22 430 chromium, and vanadium in red mud samples from the Ajka spill site, Hungary. *Environ Sci*
23 431 *Technol* 46, 3085-3092.
- 24 432 Czovek, D., Novak, Z., Somlai, C., Asztalos, T., Tizslavicz, L., Bozoki, Z., Ajtai, T., Utry, N., Filep, A., Bari,
26 433 F., Petak, F., 2012. Respiratory consequences of red sludge dust inhalation in rats. *Toxicol Lett*
27 434 209, 113-120.
- 28 435 Ehrlich, V.A., Nersesyan, A.K., Atefie, K., Hoelzl, C., Ferk, F., Bichler, J., Valic, E., Schaffer, A., Schulte-
29 436 Hermann, R., Fenech, M., Wagner, K.H., Knasmuller, S., 2008. Inhalative exposure to vanadium
30 437 pentoxide causes DNA damage in workers: results of a multiple end point study. *Environ*
31 438 *Health Perspect* 116, 1689-1693.
- 33 439 Fenech, M., Morley, A.A., 1985. Measurement of micronuclei in lymphocytes. *Mutat Res* 147, 29-36.
- 34 440 Fontanier, C., Fabri, M.C., Buscail, R., Biscara, L., Koho, K., Reichart, G.J., Cossa, D., Galaup, S.,
35 441 Chabaud, G., Pigot, L., 2012. Deep-sea foraminifera from the Cassidaigne Canyon (NW
36 442 Mediterranean): assessing the environmental impact of bauxite red mud disposal. *Mar Pollut*
37 443 *Bull* 64, 1895-1910.
- 39 444 Gawu, S.K.Y., Amisshah, E.E., Kuma, J.S., 2012. The proposed alumina industry and how to mitigate
40 445 against the red mud footprint in Ghana. *Journal of Urban and Environmental Engineering* 6, 48-
41 446 56.
- 42 447 Gelencser, A., Kovats, N., Turoczy, B., Rostasi, A., Hoffer, A., Imre, K., Nyiro-Kosa, I., Csakberenyi-
43 448 Malasics, D., Toth, A., Czitrovsky, A., Nagy, A., Nagy, S., Acs, A., Kovacs, A., Ferincz, A.,
44 449 Hartyani, Z., Posfai, M., 2011. The red mud accident in Ajka (Hungary): characterization and
46 450 potential health effects of fugitive dust. *Environ Sci Technol* 45, 1608-1615.
- 47 451 Glen, C.D., Dubrova, Y.E., 2012. Exposure to anticancer drugs can result in transgenerational genomic
48 452 instability in mice. *Proc Natl Acad Sci U S A* 109, 2984-2988.
- 49 453 Gundy, S., Farkas, G., Szekely, G., Kasler, M., 2013. No short-term cytogenetic consequences of
51 454 Hungarian red mud catastrophe. *Mutagenesis* 28, 1-5.
- 52 455 Heddle, J.A., Fenech, M., Hayashi, M., MacGregor, J.T., 2011. Reflections on the development of
53 456 micronucleus assays. *Mutagenesis* 26, 3-10.
- 54 457 IARC, 2006. Vanadium pentoxide, IARC (International Agency for Research on Cancer) Monogr Eval
55 458 Carcinog Risks Hum, pp. 227-292.
- 57 459 Klebercz, O., Mayes, W.M., Anton, A.D., Feigl, V., Jarvis, A.P., Gruiz, K., 2012. Ecotoxicity of fluvial
58 460 sediments downstream of the Ajka red mud spill, Hungary. *J Environ Monit* 14, 2063-2071.

461 Knasmueller, S., Uhl, M., Gottmann, E., Hoelzl, C., Majer, B.J., 2003. The *Tradescantia* micronucleus
1 462 bioassay, in: Maluszynska, J., Plewa, M. (Eds.), *Bioassays in plant cells for improvement of*
2 463 *ecosystem and human health*. Wydawnictwo Uniwersytetu Śląskiego, Katowice, pp. 95-106.

3 464 Knasmuller, S., Gottmann, E., Steinkellner, H., Fomin, A., Pickl, C., Paschke, A., God, R., Kundi, M.,
4 465 1998. Detection of genotoxic effects of heavy metal contaminated soils with plant bioassays.
5 466 *Mutat Res* 420, 37-48.

7 467 Lehoux, A.P., Lockwood, C.L., Mayes, W.M., Stewart, D.I., Mortimer, R.J.G., Gruiz, K., Burke, I.T., 2013.
8 468 Gypsum addition to soils contaminated by red mud: Implications for aluminium, arsenic,
9 469 molybdenum and vanadium solubility. *Environmental Geochemistry and Health* 35, 643-656.

10 470 Leme, D.M., Marin-Morales, M.A., 2009. *Allium cepa* test in environmental monitoring: a review on
11 471 its application. *Mutat Res* 682, 71-81.

12 472 Lima, M., CARNEIRO, B., COSTA, V., 2009. Caracterização preliminar dos impactos ambientais, danos
13 473 ao ecossistema e riscos a saúde decorrentes do lançamento no rio Murucupi de efluentes do
14 474 processo de beneficiamento de bauxita, Barcarena-Pará.
15 475 2009. Disponível em:<http://iah.iec.pa.gov.br/iah/fulltext/pc/relatorios/barcarena2009.pdf>>.

17 475 2009. Disponível em:<http://iah.iec.pa.gov.br/iah/fulltext/pc/relatorios/barcarena2009.pdf>>.

18 476 Ma, T.H., Cabrera, G.L., Chen, R., Gill, B.S., Sandhu, S.S., Vandenberg, A.L., Salamone, M.F., 1994.
19 477 *Tradescantia* micronucleus bioassay. *Mutat Res* 310, 221-230.

20 478 Ma, T.H., Davies, B.M., 2009. Effects of extreme low dose radiation in wet condition detected by
21 479 Trad-MCN assay, Genomics in the environmental century. 40th Annual meeting EMS. EMS, St.
22 480 louis Missouri.

24 481 Ma, T.H., Xu, Z., Xu, C., McConnell, H., Rabago, E.V., Arreola, G.A., Zhang, H., 1995. The improved
25 482 *Allium/Vicia* root tip micronucleus assay for clastogenicity of environmental pollutants. *Mutat*
26 483 *Res* 334, 185-195.

27 484 Majer, B.J., Tschерko, D., Paschke, A., Wennrich, R., Kundi, M., Kandeler, E., Knasmuller, S., 2002.
28 485 Effects of heavy metal contamination of soils on micronucleus induction in *Tradescantia* and
29 486 on microbial enzyme activities: a comparative investigation. *Mutat Res* 515, 111-124.

31 487 Manerikar, R.S., Apte, A.A., Ghole, V.S., 2008. In vitro and in vivo genotoxicity assessment of Cr(VI)
32 488 using comet assay in earthworm coelomocytes. *Environ Toxicol Pharmacol* 25, 63-68.

33 489 Marcano, L., Carruyo, I., Fernandez, Y., Montiel, X., Torrealba, Z., 2006. Determination of vanadium
34 490 accumulation in onion root cells (*Allium cepa* L.) and its correlation with toxicity. *Biocell* 30,
35 491 259-267.

37 492 Mayes, W., Jarvis, A., Burke, I., Walton, M., Gruiz, K., 2011a. Trace and rare earth element dispersal
38 493 downstream of the Ajka red mud spill, Hungary in: Rűde, R.T., Freund, A., Wolkersdorfer, C.
39 494 (Eds.), *Mine Water - Managing the challenges*. International Mine Water Association, pp. 29 -
40 495 34

42 496 Mayes, W.M., Jarvis, A.P., Burke, I.T., Walton, M., Feigl, V., Klebercz, O., Gruiz, K., 2011b. Dispersal
43 497 and attenuation of trace contaminants downstream of the Ajka bauxite residue (red mud)
44 498 depository failure, Hungary. *Environ Sci Technol* 45, 5147-5155.

45 499 Minouflet, M., Ayrault, S., Badot, P.M., Cotelle, S., Ferard, J.F., 2005. Assessment of the genotoxicity
46 500 of ¹³⁷Cs radiation using *Vicia*-micronucleus, *Tradescantia*-micronucleus and *Tradescantia*-
47 501 *stamen-hair* mutation bioassays. *J Environ Radioact* 81, 143-153.

49 502 Misik, M., Ma, T.H., Nersesyan, A., Monarca, S., Kim, J.K., Knasmueller, S., 2011. Micronucleus assays
50 503 with *Tradescantia* pollen tetrads: an update. *Mutagenesis* 26, 215-221.

51 504 Misik, M., Micieta, K., Solenska, M., Misikova, K., Pisarcikova, H., Knasmuller, S., 2007. In situ
52 505 biomonitoring of the genotoxic effects of mixed industrial emissions using the *Tradescantia*
53 506 micronucleus and pollen abortion tests with wild life plants: demonstration of the efficacy of
54 507 emission controls in an eastern European city. *Environ Pollut* 145, 459-466.

56 508 Peacock, C.L., Sherman, D.M., 2004. Vanadium(V) adsorption onto goethite (alpha-FeOOH) at pH 1.5
57 509 to 12: A surface complexation model based on ab initio molecular geometries and EXAFS
58 510 spectroscopy. *Geochimica Et Cosmochimica Acta* 68, 1723-1733.

511 Rajeev, Pradhan, J., Das, S.N., Thakur, R.S., 1999. Vanadium sludge - An useful byproduct of alumina
1 512 plant. *Journal of Scientific & Industrial Research* 58, 948-953.

2 513 Renforth, P., Mayes, W.M., Jarvis, A.P., Burke, I.T., Manning, D.A., Gruiz, K., 2012. Contaminant
3 514 mobility and carbon sequestration downstream of the Ajka (Hungary) red mud spill: The
4 515 effects of gypsum dosing. *Sci Total Environ* 421-422, 253-259.

5 516 Rodrigues, G.S., Ma, T.-H., Pimentel, D., Weinstein, L.H., 1997. *Tradescantia* Bioassays as Monitoring
6 517 Systems for Environmental Mutagenesis: A review. *Crit. Rev. Plant Sci.* 16, 325-359.

8 518 Ruyters, S., Mertens, J., Vassilieva, E., Dehandschutter, B., Poffijn, A., Smolders, E., 2011. The red
9 519 mud accident in Ajka (Hungary): plant toxicity and trace metal bioavailability in red mud
10 520 contaminated soil. *Environ Sci Technol* 45, 1616-1622.

11 521 Samal, S., Ray, A.K., Bandopadhyay, A., 2012. Proposal for resources, utilization and processes of red
12 522 mud in India - A review. *International Journal of Mineral Processing*.

14 523 Sarkar, A., Ray, D., Shrivastava, A.N., Sarker, S., 2006. Molecular Biomarkers: their significance and
15 524 application in marine pollution monitoring. *Ecotoxicology* 15, 333-340.

16 525 Shaughnessy, D.T., DeMarini, D.M., 2009. Types and consequence of DNA damage in: Knasmueller, S.,
17 526 DeMarini, D.M., Johnson, I., Gerhauser, C. (Eds.), *Chemoprevention of Cancer and DNA*
18 527 *Damage by Dietary Factors*. Wiley-VCH, Weinheim, pp. 21-34.

20 528 Steinkellner, H., Mun-Sik, K., Helma, C., Ecker, S., Ma, T.H., Horak, O., Kundi, M., Knasmuller, S., 1998.
21 529 Genotoxic effects of heavy metals: comparative investigation with plant bioassays. *Environ Mol*
22 530 *Mutagen* 31, 183-191.

23 531 Taylor, J., Keith, S., Cseh, L., Ingerman, L., Chappell, L., Rhoades, J., Hueber, A., 2012. Toxicological
24 532 Profile for Vanadium. Agency for Toxic Substances and Disease Registry US.
25 533 <http://www.atsdr.cdc.gov/toxprofiles/tp58.pdf>, Atlanta, p. 255.

27 534 Zvereva, E., Toivonen, E., Kozlov, M.V., 2008. Changes in species richness of vascular plants under the
28 535 impact of air pollution: a global perspective. *Global Ecology and Biogeography* 17, 305-319.

29
30 536
31
32
33 537
34
35 538
36
37
38 539
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 1 Red mud a byproduct of Aluminum production contains soluble vanadium that causes
4
5 2 genotoxic and cytotoxic effects in higher plants
6
7 3

8
9 4 Miroslav Mišík¹, Ian T. Burke², Matthias Reismüller¹, Clemens Pichler¹, Bernhard Reiner¹,
10
11 5 Katarina Mišíková³, William M. Mayes⁴, Siegfried Knasmueller^{1*}
12
13 6
14
15 7

16
17 8 ¹Institute of Cancer Research, Department of Medicine I, Comprehensive Cancer Center,
18
19 9 Medical University of Vienna, Austria
20

21 10 ²Earth Surface Science Institute, School of Earth and Environment, University of Leeds,
22
23 11 Leeds LS2 9JT, UK
24

25 12 ³Department of Botany, Faculty of Natural Sciences, Comenius University Bratislava,
26
27 13 Slovakia
28

29 14 ⁴Centre for Environmental and Marine Sciences, University of Hull, Scarborough, YO11
30
31 15 3AZ, UK
32
33 16
34
35 17
36
37 18
38

39 19 Corresponding author:

40
41 20 Siegfried Knasmueller; Institute of Cancer Research, Borschkegasse 8a, A-1090 Vienna
42
43 21 Austria; Phone: ++43-1-40160 57561, Fax ++43 1 40160 957500; mail:
44
45 22 siegfried.knasmueller@meduniwien.ac.at
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

23 ABSTRACT

24 Red mud (RM) is a byproduct of aluminum production, worldwide between 70 and 120
25 million tons are produced annually. We analyzed RM which was released in the course of the
26 Kolonatar disaster in Hungary into the environment in acute and genotoxicity experiments
27 with plants which are widely used for environmental monitoring. We detected induction of
28 micronuclei which reflect chromosomal damage in tetrads of Tradescantia and in root cells of
29 Allium as well as retardation of root growth with contaminated soils and leachates. Chemical
30 analyses showed that RM contains metals, in particular high concentrations of vanadium.
31 Follow up experiments indicated that vanadate causes the effects in the plants. This
32 compound causes also in humans DNA damage and positive results were obtained in
33 carcinogenicity studies. Since it was found also in RM from other production sites our
34 findings indicate that its release in the environment is a global problem which should be
35 studied in more detail.

40 **Capsule abstract:** Our findings indicate that the red mud causes genotoxic effect in plants
41 probably due to the presence of vanadate which is contained at high concentrations in the
42 residue.

46 Key words: red mud, Tradescantia, micronuclei, Allium, vanadium

1
2
3
4 48 **1. Introduction**

5 49 Red mud is a by-product of aluminum production with the Bayer process. Its global
6
7 50 production is in the range between of 70 to 120 million tons per year (Mayes et al., 2011b).
8
9 51 The material consists mainly of iron-, aluminum- and titanium- oxides and hydroxides (Burke
10
11 52 et al., 2012; Mayes et al., 2011b). Chemical analyses showed that its contains also
12
13 53 radionuclides (e.g. ²²⁶RA, ²³⁰Th and ⁴⁰K), as well as heavy metals including As, Cr, Co, Cd,
14
15 54 Ni and V (Mayes et al., 2011a; Ruyters et al., 2011).

16
17 55 On Oct 4th 2010, approximately one million m³ of the residue was released into the
18
19 56 environment from the aluminum plant Ajkai Timfoldgyar Zrt in Western Hungary. According
20
21 57 to the Hungarian Ministry of Interior the “Kolontar disaster” is the biggest environmental
22
23 58 catastrophe which ever happened in this country (Ádám et al., 2011). Hundreds of houses
24
25 59 were destroyed, 265 individuals were injured and ten died (Gundy et al., 2013).

26
27 60 After the accident, attempts were made to investigate the impact of the release of the
28
29 61 material into the environment and to assess the health consequences in humans.
30
31 62 Ecotoxicological studies were conducted with different plant species and bacteria concerning
32
33 63 acute toxic effects (Klebercz et al., 2012; Ruyters et al., 2011); the motility and the
34
35 64 concentrations of toxic trace elements were studied in physico-chemical measurements
36
37 65 (Burke et al., 2012). Furthermore, studies were conducted to assess the consequences of
38
39 66 inhalation of dust particles in humans and rodents (Czovek et al., 2012; Gelencser et al.,
40
41 67 2011).

42
43 68 Radionuclides as well as certain heavy metals (found in red mud) cause damage of the
44
45 69 genetic material (Knasmuller et al., 1998; Minouflet et al., 2005) which may lead to
46
47 70 destabilization of ecosystems (Sarkar et al., 2006; Zvereva et al., 2008) and also causes
48
49 71 adverse effects in humans such as cancer, ageing, infertility and birth defects in the offspring
50
51 72 (Aitken and De Iuliis, 2007; Assem and Levy, 2009). Therefore it is of particular interest if

1
2
3 73 the release of this material into the environment induces chromosomal damage. This question
4
5 74 has been addressed in a human study (Gundy et al., 2013) and in bacterial tests (Gelencser et
6
7 75 al., 2011) but no firm conclusions can be drawn of these studies (for details see discussion).

8
9 76 The primary aim of the present study was the investigation of the genotoxic properties
10
11 77 of red mud in two plant bioassays, namely in the micronucleus (MN) test with tetrads of
12
13 78 *Tradescantia* (Trad-MN assay) and with root tip cells of *Allium cepa* (A-MN assay). The
14
15 79 experiments were conducted with tetrads as they reflect damage in meiotic germ cells. Root
16
17 80 tip cells were used as they enable the detection of effects in mitotic cells. It was postulated
18
19 81 that differences exist in regard to sensitivity of these cell types towards DNA reactive
20
21 82 compounds (Rodrigues et al., 1997). MN are formed as a consequence of chromosomal
22
23 83 breakage or aneuploidy and can be monitored in a variety of organisms (Heddle et al., 2011).
24
25 84 In addition acute toxic effects were studied in root cells of *A. cepa* by measuring the impact of
26
27 85 the material on the root growth and by calculating the division rates of the cells. These
28
29 86 bioassays are at present the most widely used genotoxicity tests with higher plants and have
30
31 87 been employed in more ~~the~~than 300 investigations for the detection of DNA damaging
32
33 88 properties of chemicals and complex mixtures (for reviews see Leme and Marin-Morales,
34
35 89 2009; Misik et al., 2011). We used these test systems since they provide information on
36
37 90 environmental effects and it is known that they are, in contrast to other genotoxicity assays
38
39 91 with bacterial indicators and mammalian cells (which are also used for environmental
40
41 92 monitoring), highly sensitive towards radiation (Ma and Davies, 2009; Misik et al., 2011) and
42
43 93 heavy metals (Knasmuller et al., 1998; Majer et al., 2002; Steinkellner et al., 1998).

44
45 94 We investigated the acute cytotoxic and genotoxic activities of red mud and of soils
46
47 95 which were contaminated with the material collected from fields and gardens which were
48
49 96 used for cultivation of crops and vegetables. Furthermore, we studied also the effects of
50
51 97 waters leached from red mud and affected soils. In additional experiments, attempts were

1
2
3 98 made to identify the compound(s) which cause genotoxic effects. We determined the
4
5 99 concentrations of trace elements in solid samples and in leachates and conducted additional
6
7 100 Trad-MN assays with sodium metavanadate (NaVO_3) since vanadium was found to be the
8
9 101 most abundant heavy metal in the samples.

14 103 **2. Materials and methods**

17 105 2.1 Sampling

19 106 Soil samples were collected from the Torna catchment on December 15th 2010. Table
20
21 107 1 contains a description of the sampling sites and specifications of their GPS locations. From
22
23 108 each site, samples were collected from the 0-5; 5-15 and 15-25 cm horizons to assess if
24
25 109 potentially toxic constituents in red mud contaminated the soils.

27 110 Total organic carbon (TOC) was determined in soil samples using a LECO SC-144DR
28
29 111 elemental analyzer after removal of the inorganic carbon fraction using 20% HCl.

33 113 Table 1

35 114 Figure 1

38 115 2.2 Preparation of the leachates.

40 116 Soils were used as sampled (field moist). 50 g of each sample were suspended in 100
41
42 117 mL of deionised water in a 250 mL glass beaker for 2 hours using a magnetic stirrer. The
43
44 118 extracts were filtered with filter paper before they were tested in stem absorption experiments.
45
46 119 The pH values of the leachates were measured with a pH Meter 526 (WTW, Weilheim,
47
48 120 Germany) and are listed in Table 1.

1
2
3 122
4
5
6 123
7
8 124
9
10 125
11
12 126
13
14 127
15
16 128
17
18 129
19
20 130
21
22 131
23
24 132
25
26 133
27
28 134
29
30 135
31
32 136
33
34 137
35
36 138
37
38
39 139
40
41 140
42
43 141
44
45 142
46
47 143
48
49 144
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2.3 Measurements of the trace elements in the soil samples and leachates.

The detection of the trace elements is described in detail in a recent paper of Renforth et al. (2012). Prior to analysis by XRF the samples was prepared as follows. For major element analysis, samples were prepared as fused beads (after loss on ignition at 1050 °C) with lithium metaborate/tetraborate flux (Johnson - Matthey Spectroflux JM100B) (0.6g sample; 3 g Flux). For minor / trace element analysis approximately 10 g dried sample was prepared as a pressed pellet using ~10-20 drops of 6.6%w/v polyvinyl alcohol in a 1:6 mix of methanol and distilled deionised water as a binder (Moviol 88 solution). Elemental analysis of the soil composition was achieved using a PANalytical Axios Advanced X-ray Fluorescence (XRF) spectrometer (data corrected for loss on ignition; % weight loss after furnace treatment at 1050°C). The aqueous phase produced during water extractions was separated from solids by centrifugation (10 min; 2000 g) followed by membrane filtration (0.45 µm); the filtered samples were then acidified by addition of 2% v/v eHNO₃. Aqueous elemental concentrations were then determined using a Perkin Elmer Optima 5300 DV ion-coupled plasma, optical emission spectrometer (ICP-OES).

2.4 Tradescantia micronucleus assays

Tradescantia clone #4430 was cultivated according to the protocol of Ma et al. (1994) For stem absorption experiments, 15 young inflorescences were treated in each group. The stems were cut to a length of 10 - 15 cm, transferred to plastic breakers (250 mL) and exposed to aqueous leachates (soil:water – 1:2) of the contaminated soils for 24 h. Subsequently, they were transferred to water for a 24 h recovery period. The flower buds were then collected and

1
2
3 145 fixed in acetic acid and ethanol (1/3, v/v). After 24 h, they were transferred to 70% ethanol.

4
5 146 Maleic hydrazide (20 mg L⁻¹; MH, Sigma-Aldrich, SL, US) was used as a positive control.

6
7 147 The protocol for root absorption assays is described in an article of Steinkellner et al.
8
9 148 (1998). The soil which was used in the control cultures was biological, pesticide free (from
10
11 149 Composana, Wien, Austria). Intact plants were removed from hydroponic culture,
12
13 150 subsequently the roots were rinsed and individual plants with at least 15 inflorescences placed
14
15 151 in-to plastic pots (250 ml, diameter 12 cm) which were filled with the different soil samples.
16
17 152 The plants were exposed under standard conditions for 48 h (Ma et al., 1994). Subsequently,
18
19 153 15 inflorescences were collected from each soil sample in all experiments and fixed in a
20
21 154 solution of acetic acid and ethanol (1/3, v/v). After 24 h, they were transferred to 70%
22
23 155 ethanol.

24
25 156 To find out if vanadium accounts for the effects which were found in the soils and in
26
27 157 the leachetes, additional experiments were carried out with sodium metavanadate. The salt
28
29 158 was dissolved in 0.1M NaOH (pH 13) at 1000 ppm (colorless solution) since earlier findings
30
31 159 indicate that vanadium is present as V⁵⁺ in the soil (Burke et al., 2012). The stock solution
32
33 160 was diluted with tap water and different concentrations (0.5-10.0 ppm) were used in
34
35 161 experiments with Tradescantia.

36
37 162 Slides were prepared and evaluated as described in the protocol of Ma et al. (1994)
38
39 163 The tetrads were stained with a 2% aceto-carmin solution. Early tetrads were analyzed under
40
41 164 400-fold magnification. For each experimental point a minimum of five buds with early
42
43 165 tetrads was evaluated (300 tetrads were evaluated per slide, 1500 per sample).

44 45 46 166 2.5 Micronucleus assay with *Allium cepa*

47
48 167 The experiments were carried out according to the standard protocol published by Ma
49
50 168 et al. (1995) with slight modifications. Young onion bulbs (diameter 12-21 mm, Schneeball

1
2
3 169 weis, Austrosaat, Wien, Austria) were placed in 13 ml glass tubes filled with tap water for
4
5 170 24 h in the dark. Subsequently, the roots (lengths ca. 0.5-1 cm) were exposed directly in the
6
7 171 soils in the dark for 24 h and then transferred to fresh tap water for another 24 h. At the end of
8
9 172 the recovery phase, the maximal root length of each onion was measured and the material was
10
11 173 fixed in a mix of acetic acid and ethanol (1/3, v/v) for 24 h and stored in 70% ethanol. MH
12
13 174 (10 mg L⁻¹; Sigma-Aldrich, SL, US) was used in all experiments as a positive control.
14

15 175 The root tips were hydrolyzed in a 1:1 mix of HCl (5.0 N) and ethanol (99%) for
16
17 176 3 min and washed in tap water before they were stained with 2% acetocarmine. MN were
18
19 177 scored according to the criteria described by Ma et al. (1995). For each experimental point,
20
21 178 the MN frequencies were determined in five plants. From each bulb, two slides were made
22
23 179 and 500 cells were evaluated per slide (5000 cells per dose). Furthermore, also the mitotic
24
25 180 indices (MIs) were determined in 1000 cells (100 cells/root) per experimental point. The
26
27 181 microscopic evaluation was carried out under a light microscope (Nikon YS200, Japan) with
28
29 182 400-fold magnification.
30

31 183

32

33

34 184 2.6 Statistical evaluation

35

36 185 The results of the MN experiments were analyzed with one-way ANOVA followed by
37
38 186 Dunnett's multiple comparison test. The results of the experiments concerning the mitotic
39
40 187 indices (MI) were analyzed with the Kruskal-Wallis test followed by Dunn's comparison test.
41
42 188 P-values ≤ 0.05 were considered as significant.
43

44 189

45

46 190

47

48 191

49

50 192

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

1
2
3 193 **3. Results**
4 194
5

6 195 3.1 TOC and pH values of the samples.
7

8
9 196 It can be seen in Table 1 that values between 8.0 and 9.3 were found in the soil
10
11 197 leachates which were contaminated with red mud. Most of the samples had a total organic
12
13 198 carbon content between 1.0-1.5%, which is typical for agricultural soils in the Ajka area
14
15 199 (Lehoux et al., 2013). The sample from the upper horizon of the site 3 had a substantially
16
17 200 lower TOC content (0.22%) and a higher pH (9.33) which is typical for red mud (Lehoux et
18
19 201 al., 2013).
20
21 202
22

23 203 3.2 Micronucleus assays with Tradescantia (experiments with soils and leachates)
24

25
26 204 The results which were obtained in Trad-MN assays with red mud contaminated soil
27
28 205 and with aqueous leachates from these samples are summarized in Fig. 2A and 2B. It can be
29
30 206 seen that direct exposure of intact plants in material from the contaminated sites 2 (S2) and 3
31
32 207 (S3) caused a significant increase of the MN frequencies. In the case of S3, the strongest
33
34 208 effect was seen with material from the upper layer, samples from deeper horizons caused only
35
36 209 a statistically not significant increase. With the two samples from S4, clear cut negative
37
38 210 results were obtained. As described above (Table 1), this material was collected after
39
40 211 remediation (i.e. after removal of the upper layer).
41

42 212 Also with some leachates, positive results were obtained (Fig. 2B). Again the strongest
43
44 213 effects were seen with extracts prepared with material from the upper layers (S2 and S3); soils
45
46 214 from deeper horizons did not cause significant effects, also the sample which was collected
47
48 215 from the cleanup site was devoid of activity.
49

50 216
51 217 Figures 2A and B
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 218
4
5
6 219
7
8 220
9
10 221
11
12 222
13
14 223
15
16 224
17
18 225
19
20 226
21
22 227
23
24 228
25
26 229
27
28 230
29
30 231
31
32 232
33
34 233
35
36 234
37
38 235
39
40
41 236
42
43 237
44
45 238
46
47 239
48
49 240
50
51 241
52
53
54
55
56
57
58
59
60
61
62
63
64
65

3.3 Micronucleus experiments with mitotic root tip cells of *Allium cepa*

The results of assays with *A. cepa* are shown in Figures 3A – C. The roots were exposed in these experiments directly in the samples. Since division delays may decrease the formation of MN (Fenech and Morley, 1985), we determined in all experiments also the mitotic indices (MI).

It can be seen in Figure 3A that a significant increase of the MN rates was observed with some samples. The most pronounced effect was detected with material collected from the middle layer of S2 while red mud enriched sample from surface of the same location did not cause a significant increase of the MN rates. This finding can be explained by inhibition of cell division, which was most pronounced with material collected from the upper horizon of this site (Figure 3B). As in the *Tradescantia* experiments, no increase was seen with samples from the remediation site (S4).

The impact of the material on root growth is shown in Figure 3C. It can be seen, that clear effects were obtained with samples from the surface of sites S2 and S3.

Figures 3A – C

3.4 Chemical analyses of the soil samples and of the leachates

The chemical composition of samples which were collected from S3 and S1 (Table 2A-B) and of the corresponding leachates (Table 2C) are shown in Table 2.

Table 2

1
2
3 242 The chemical composition of the topmost soil sample (S3 0-5 cm) is similar to that of
4
5 243 other samples of red mud which were collected after the Ajka spill (Burke et al., 2012;
6
7 244 Ruyters et al., 2011). Several potentially toxic metals including As, Cr, Ni, Pb and V were
8
9 245 present at elevated concentrations.

10
11 246 The chemical composition of the control sample (S1) is consistent with unaffected
12
13 247 soils found in the region (see Table 2c) (Mayes et al., 2011b).

14
15 248 The corresponding leachates from S3, contained elevated concentrations of potentially
16
17 249 toxic elements, such as V, As, Cu, Al and Cr, notably, V concentrations were an order of
18
19 250 magnitude higher than the levels of the other metals (except Al). Also chromium levels were
20
21 251 relatively high.

22
23 252
24
25
26 253 3.5 Tradescantia micronucleus assays with vanadate

27
28 254 We found in earlier experiments with Tradescantia that the concentrations of Ni, Cr,
29
30 255 Cd, Pb and As, which cause MN induction in tetrads are in the range of hundred to thousand
31
32 256 ppm (Knasmuller et al., 1998; Steinkellner et al., 1998). We hypothesized that V, which was
33
34 257 detected in higher levels than other metals, may account for the induction of MN, which was
35
36 258 seen with the soils and the leachates. Therefore, we tested different concentrations of an
37
38 259 aqueous vanadate solution in subsequent Trad MN assays. The results are summarized in
39
40 260 Figure 4. It can be seen that exposure of the cuttings to concentrations ≥ 1.0 ppm-mg/L caused
41
42 261 significant induction of the MN frequencies i.e. an approximately 6-fold increase over the
43
44 262 background.

45
46 263
47
48 264 Figure 4

49
50 265
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 266 3.6 Impact of pH on MN formation in Tradescantia
4

5 267 Since the pH values of the soil extracts and of the vanadate solutions were quite high,
6
7 268 i.e. in the range between 8.2 and 9.3 (for details see Table 1) we conducted an additional
8
9 269 experiment in which plant cuttings were exposed in aqueous solutions (without metals) to
10
11 270 different pH values. No indication of pH depended induction of MN by the pH itself was
12
13 271 found in this experiment. The numbers of MN per 100 tetrads were 1.7 ± 0.6 ; 2.1 ± 0.6 and
14
15 272 1.6 ± 0.6 for pH 8, 8.5 and 9.3 respectively (numbers are means \pm SD of results obtained from
16
17 273 1500 tetrads).
18
19

20 274 **4. Discussion**
21 275
22

23 276 The results of this study show that red mud, which was released in large amounts into
24
25 277 the environment, causes damage of the genetic material and also acute toxic effects in higher
26
27 278 plants.
28
29 279
30
31

32 280 4.1 Acute toxic and genotoxic effects of red mud
33

34 281 Several ecotoxicological studies have been conducted with red mud from Ajka
35
36 282 (Klebercz et al., 2012; Ruyters et al., 2011) and adverse effects were seen in experiments with
37
38 283 higher plants; i.e. the material was found to inhibit the root of *Sinapis alba* and affect the
39
40 284 shoot yield in barley (Klebercz et al., 2012; Ruyters et al., 2011). In the latter study, the
41
42 285 authors monitored also the levels of trace elements which are contained in red mud such as
43
44 286 Cu, Cr, Fe and Ni in exposed plants. They detected these metals in the shoots but stress that
45
46 287 their levels did not exceed the toxic limits and hypothesize that Na is the prime cause which
47
48 288 affected the growth of the indicator plants. Another factor which was made responsible for the
49
50 289 toxic effects in *Sinapis* and in experiments with the Ostracod *Heterocypris incongruens* is
51

1
2
3 290 gypsum which may limit the nutrient supply and increase the availability of contaminants
4
5 291 (Klebercz et al., 2012). The genotoxic effect of red mud has not been investigated in the
6
7 292 invertebrates so far according to our knowledge. However, it is possible that several species
8
9 293 may be affected due to the high metal concentration of the material. For example, it is know
10
11 294 that earthworms are highly sensitive towards specific contaminants and specific metals.
12
13 295 Genotoxic effects were detected with chromium, nickel and cadmium (Bigorgne et al., 2010;
14
15 296 Bonnard et al., 2010; Manerikar et al., 2008).

16
17 297 The results of the present study show clearly that red mud inhibits the division and
18
19 298 growth of root tip cells in *A. cepa* (Figures 3B and 3C) and causes induction of MN in this
20
21 299 species. Furthermore, we showed that also aqueous leachates of red mud contaminated soils
22
23 300 cause induction of MN in both plant bioassays. As shown in Figure 3 a stronger effect was
24
25 301 obtained with sample S2 (5-15 cm) as with the sample from upper layer from the same site.
26
27 302 This effect can be explained by reduction of MN formation due to inhibition of cell division.
28
29 303 Furthermore, it is notable that the effects which were detected with soil exposure are more
30
31 304 pronounced as those which were found with the leachetes. This observation is in agreement
32
33 305 with the results obtained with heavy metals contaminated soils and can be explained by more
34
35 306 efficient uptake via the roots by active transport mechanism (for detail see Steinkellner et al.,
36
37 307 1998).

38
39 308 The only genotoxicity assay conducted so far with red mud is the SOS chromotest
40
41 309 which is based on the detection of SOS responses in bacterial indicator cells (Gelencser et al.,
42
43 310 2011). Consistently negative results were obtained with samples, which were prepared from
44
45 311 fugitive dust from the area where the accident had happened. However, it is notable that it is
46
47 312 known from earlier investigations that bacterial assays are insensitive towards heavy metals
48
49 313 (Majer et al., 2002).

50
51 314
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 315 4.2 Identification of vanadate as the active principle of red mud
4

5 316 The results of the chemical analyses show that red mud contaminated soils and their
6
7 317 aqueous leachates contain elevated concentrations of V (Table 2C). This observation is in
8
9 318 agreement with the results of earlier investigations with material from Ajka (Mayes et al.,
10
11 319 2011a; Mayes et al., 2011b; Renforth et al., 2012). In this context, it is notable that vanadium
12
13 320 was also detected in red mud from several other aluminium production sites (Fontanier et al.,
14
15 321 2012; Gawu et al., 2012; Rajeev et al., 1999; Samal et al., 2012).
16

17 322 Spectroscopic evidence showed that vanadium is present primarily as V^{5+} phases
18
19 323 (Burke et al., 2012). Under alkaline conditions (high pH is induced in red mud leachate by
20
21 324 dissolution of the NaOH present), V^{5+} is predicted to become readily solubilized in water as
22
23 325 vanadate (Peacock and Sherman, 2004).
24

25 326 Vanadium is more abundant in red mud than other alkali-soluble trace metals (e.g.
26
27 327 As), therefore its environmental mobility is enhanced relative to those of other carcinogenic
28
29 328 heavy metals which are present in the material. Therefore, we conducted Trad-MN assays
30
31 329 with sodium metavanadate solutions and compared the concentrations which cause positive
32
33 330 results with those contained in the aqueous leachates. The extracts prepared from the upper
34
35 331 layer of S3 contained 1.1 mg/L V (Table 2) and the results which were obtained with aqueous
36
37 332 vanadate solutions (Figure 4) show, that levels ≥ 1.0 mg/L cause significant induction of the
38
39 333 MN frequencies in Tradescantia indicating that vanadate accounts for the genotoxic effects
40
41 334 which we detected in the leachate.
42

43 335 The solubility of vanadate in the environment is largely controlled by sorption
44
45 336 reactions with mineral surfaces, with low solution concentrations expected at circumneutral
46
47 337 pH (Peacock and Sherman, 2004). ~~Therefore, if~~ the pH of red mud affected soils can be
48
49 338 controlled to $pH < 9$, high aqueous V concentrations and their accompanying genotoxic
50
51 339 effects may be avoided (Lehoux et al., 2013). In this context, it is notable that, several
52

1
2
3 340 technologies have been developed to recover vanadium from sludge (for review see Rajeev et
4
5 341 al., 1999). As mentioned above, we found no impact of the pH on MN frequencies in
6
7 342 experiments with waters to which NaOH was added in absence of metal ions. However, the
8
9 343 situation may be different in soils which are contaminated with vanadium compounds. In this
10
11 344 context, it is notable that, several technologies have been developed to recover vanadium from
12
13 345 sludge (for review see Rajeev et al., 1999).
14

15 346 16 17 18 347 4.3 Vanadate is an extremely potent mutagen in the Trad MN assay 19

20 348 As shown in Table 2A-C, a number of other genotoxic and carcinogenic heavy metals
21
22 349 such as Ni, Cd, Cr and As were detected in the aqueous extracts. Results of earlier Trad-MN
23
24 350 assays show that the concentrations of these metals, which are required to cause significant
25
26 351 effects, are 10^2 - to 10^3 -fold higher as the levels which were detected in the leachates. The
27
28 352 strongest activity was seen in these experiments with $\text{Cr}^{(VI)}\text{O}_3$ (129) followed by As_2O_3 (98.5),
29
30 353 $\text{Ni}^{(II)}\text{Cl}_2$ (645), CdCl_2 (1006) and $\text{Pb}(\text{NO}_3)_2$ (1820); numbers in parenthesis indicate the
31
32 354 LOELs in mg/L (Knasmueller et al., 2003; Knasmuller et al., 1998; Steinkellner et al., 1998).
33
34 355 Furthermore, the results of the chemical analyses demonstrate that, these effects can be
35
36 356 attributed to the presence of soluble vanadium. As mentioned above, we found in the present
37
38 357 study significant induction of MN by vanadate in Tradescantia already at levels ≥ 1 mg/L.
39
40 358 Comparisons with the effects seen with other metals indicate that vanadate is more potent in
41
42 359 regard to induction of chromosomal damage. According to our knowledge, it has never been
43
44 360 tested before in genotoxicity experiments with higher plants, but earlier findings with Allium
45
46 361 roots showed that 25 mg/L cause stickiness of the chromosomes and a decrease of the MI
47
48 362 (Marcano et al., 2006).
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 363 Results which were obtained in MN assays and other genotoxicity tests with
4
5 364 mammalian cells and rodents with vanadium compounds are summarized in an IARC
6
7 365 monograph (IARC, 2006) and in a review by Assem ~~and Levy et al.~~ (2009) and Taylor et al.
8
9 366 (2012). The available data show that positive results were obtained in most in vitro
10
11 367 experiments with mammalian cells and also in a number of in vivo studies with rodents.
12

13 368 14 15 16 369 4.4 Possible consequences of the release of DNA damaging toxins in red mud to the 17 18 370 environment

19
20 371 It has been postulated that genotoxins may have an impact on the stability of ecosystems.
21
22 372 In this context it is notable, that we found in an earlier study concerning urban air pollution,
23
24 373 that the induction of MN in Tradescantia is paralleled by a decrease of the fertility of wildlife
25
26 374 plants (Misik et al., 2007). It was also shown by other groups that DNA damaging toxins in
27
28 375 the environment cause mutations which lead to transgenerational genomic instability in the
29
30 376 offspring (Glen and Dubrova, 2012). The fact that we detected pronounced induction with
31
32 377 leachates of red mud indicates that the compounds which induce genetic damage (presumably
33
34 378 V⁵⁺) contaminate surface and/or ground waters. Therefore, it is likely that they cause
35
36 379 environmental damage.

37
38 380 In regard to potential effects in humans it is known that DNA damage is causally
39
40 381 related to diseases such as cancer, infertility and heritable diseases (Shaugnessy and
41
42 382 DeMarini, 2009) and leads to accelerated ageing. We found in an earlier occupational study
43
44 383 that exposure of workers to V⁵⁺ causes induction of MN in peripheral lymphocytes (Ehrlich et
45
46 384 al., 2008) which is a reliable biomarker for increased cancer risks (Bonassi et al., 2011).
47
48 385 Furthermore, it was shown in a long term carcinogenicity study that inhalative exposure to
49
50 386 this compound causes an increase of the tumor rates in the lungs of rats (Assem and Levy,
51

1
2
3 387 2009), as a consequence of those results V₂O₅ was classified as a possible (group 2B) human
4
5 388 carcinogen by the IARC in 2006 (IARC, 2006).
6

7 389 It can be not excluded that humans living in the area where the accident happened are
8
9 390 exposed to the genotoxic components which are contained in red mud. Gundy et al. (2013)
10
11 391 analyzed chromosomal aberrations (CA) in lymphocytes of individuals who were either
12
13 392 injured by the accident or exposed during cleanup activities. The authors conclude that the
14
15 393 exposure does not pose an immediate short term genotoxic hazard but recommend further
16
17 394 studies. In fact, no firm conclusions can be drawn from their investigation. The CA
18
19 395 frequencies of the exposed group were almost 30% higher as those in an urban control group
20
21 396 while no differences were seen with controls from a rural village. Furthermore, it is notable
22
23 397 that the exposure periods of the individuals in the study group were quite heterogeneous (i.e.
24
25 398 between 9 and 336 hrs) which enhardens the detection of effects.
26

27 399 Taken together, the results of the present investigation show for the first time that
28
29 400 DNA damaging toxins were released in the course of the Kolontar disaster and that red mud
30
31 401 contains vanadate which causes damage of the genetic material at low concentrations. As
32
33 402 mentioned above, the metal was detected also in samples from other production sites and
34
35 403 environmental pollution has been reported from other countries, for example from Brazil and
36
37 404 India (2011; Gawu et al., 2012; Lima et al., 2009). The release of genotoxins into the
38
39 405 environment may affect the stability of ecosystems and have a negative impact on human and
40
41 406 environmental health; due to the worldwide production of the residue, the contamination of
42
43 407 the material with DNA reactive and potentially carcinogenic vanadium compounds is a global
44
45 408 problem which should be studied in more detail in the future.
46

47 409
48
49 410 **Acknowledgement**
50

51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 411
4
5 412
6
7
8 413
9
10 414
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

We should like to thank to Thomas Trenker and Herwig Schuster for help in regard to
the collection of the samples.

References

2011. Vedanta Red Mud - Vedanta Resources www.vedantaresources.com/media/161444/vedanta-red-mud.pdf.
- Ádám, J., Bánvölgyi, G., Dura, G., Grenerczy, G., Gubek, N., Gutper, I., Simon, G., Szegfalvi, Z., Székács, A., Szépvölgyi, J., Ujlaky, E., 2011. The Kolontár Report - Causes and lessons learned from the red mud disaster, in: Jávör, B., Hargitai, M. (Eds.), Budapest.
- Aitken, R.J., De luliis, G.N., 2007. Origins and consequences of DNA damage in male germ cells. *Reprod Biomed Online* 14, 727-733.
- Assem, F.L., Levy, L.S., 2009. A review of current toxicological concerns on vanadium pentoxide and other vanadium compounds: gaps in knowledge and directions for future research. *J Toxicol Environ Health B Crit Rev* 12, 289-306.
- Bigorgne, E., Cossu-Leguille, C., Bonnard, M., Nahmani, J., 2010. Genotoxic effects of nickel, trivalent and hexavalent chromium on the *Eisenia fetida* earthworm. *Chemosphere* 80, 1109-1112.
- Bonassi, S., El-Zein, R., Bolognesi, C., Fenech, M., 2011. Micronuclei frequency in peripheral blood lymphocytes and cancer risk: evidence from human studies. *Mutagenesis* 26, 93-100.
- Bonnard, M., Devin, S., Leyval, C., Morel, J.L., Vasseur, P., 2010. The influence of thermal desorption on genotoxicity of multipolluted soil. *Ecotoxicol Environ Saf* 73, 955-960.
- Burke, I.T., Mayes, W.M., Peacock, C.L., Brown, A.P., Jarvis, A.P., Gruiz, K., 2012. Speciation of arsenic, chromium, and vanadium in red mud samples from the Ajka spill site, Hungary. *Environ Sci Technol* 46, 3085-3092.
- Czovek, D., Novak, Z., Somlai, C., Asztalos, T., Tiszlavicz, L., Bozoki, Z., Ajtai, T., Utry, N., Filep, A., Bari, F., Petak, F., 2012. Respiratory consequences of red sludge dust inhalation in rats. *Toxicol Lett* 209, 113-120.
- Ehrlich, V.A., Nersesyan, A.K., Atefie, K., Hoelzl, C., Ferk, F., Bichler, J., Valic, E., Schaffer, A., Schulte-Hermann, R., Fenech, M., Wagner, K.H., Knasmüller, S., 2008. Inhalative exposure to vanadium pentoxide causes DNA damage in workers: results of a multiple end point study. *Environ Health Perspect* 116, 1689-1693.
- Fenech, M., Morley, A.A., 1985. Measurement of micronuclei in lymphocytes. *Mutat Res* 147, 29-36.
- Fontanier, C., Fabri, M.C., Buscail, R., Biscara, L., Koho, K., Reichart, G.J., Cossa, D., Galaup, S., Chabaud, G., Pigot, L., 2012. Deep-sea foraminifera from the Cassidaigne Canyon (NW Mediterranean): assessing the environmental impact of bauxite red mud disposal. *Mar Pollut Bull* 64, 1895-1910.
- Gawu, S.K.Y., Amisshah, E.E., Kuma, J.S., 2012. The proposed alumina industry and how to mitigate against the red mud footprint in Ghana. *Journal of Urban and Environmental Engineering* 6, 48-56.
- Gelencser, A., Kovats, N., Turoczy, B., Rostasi, A., Hoffer, A., Imre, K., Nyiro-Kosa, I., Csakberenyi-Malasics, D., Toth, A., Czitrovsky, A., Nagy, A., Nagy, S., Acs, A., Kovacs, A., Ferincz, A., Hartyani, Z., Posfai, M., 2011. The red mud accident in Ajka (Hungary): characterization and potential health effects of fugitive dust. *Environ Sci Technol* 45, 1608-1615.
- Glen, C.D., Dubrova, Y.E., 2012. Exposure to anticancer drugs can result in transgenerational genomic instability in mice. *Proc Natl Acad Sci U S A* 109, 2984-2988.
- Gundy, S., Farkas, G., Szekely, G., Kasler, M., 2013. No short-term cytogenetic consequences of Hungarian red mud catastrophe. *Mutagenesis* 28, 1-5.
- Heddle, J.A., Fenech, M., Hayashi, M., MacGregor, J.T., 2011. Reflections on the development of micronucleus assays. *Mutagenesis* 26, 3-10.
- IARC, 2006. Vanadium pentoxide, IARC (International Agency for Research on Cancer) Monogr Eval Carcinog Risks Hum, pp. 227-292.
- Klebercz, O., Mayes, W.M., Anton, A.D., Feigl, V., Jarvis, A.P., Gruiz, K., 2012. Ecotoxicity of fluvial sediments downstream of the Ajka red mud spill, Hungary. *J Environ Monit* 14, 2063-2071.

Formatted: German (Austria)

1
2
3 464 Knasmueller, S., Uhl, M., Gottmann, E., Hoelzl, C., Majer, B.J., 2003. The Tradescantia micronucleus
4 465 bioassay, in: Maluszynska, J., Plewa, M. (Eds.), Bioassays in plant cells for improvement of
5 466 ecosystem and human health. Wydawnictwo Uniwersytetu Śląskiego, Katowice, pp. 95-106.
6 467 Knasmueller, S., Gottmann, E., Steinkellner, H., Fomin, A., Pickl, C., Paschke, A., God, R., Kundi, M.,
7 468 1998. Detection of genotoxic effects of heavy metal contaminated soils with plant bioassays.
8 469 Mutat Res 420, 37-48.
9 470 Lehoux, A.P., Lockwood, C.L., Mayes, W.M., Stewart, D.I., Mortimer, R.J.G., Gruiz, K., Burke, I.T., 2013.
10 471 Gypsum addition to soils contaminated by red mud: Implications for aluminium, arsenic,
11 472 molybdenum and vanadium solubility. Environmental Geochemistry and Health 35, 643-656.
12 473 Leme, D.M., Marin-Morales, M.A., 2009. Allium cepa test in environmental monitoring: a review on
13 474 its application. Mutat Res 682, 71-81.
14 475 Lima, M., CARNEIRO, B., COSTA, V., 2009. Caracterização preliminar dos impactos ambientais, danos
15 476 ao ecossistema e riscos a saúde decorrentes do lançamento no rio Murucupi de efluentes do
16 477 processo de beneficiamento de bauxita, Barcarena-Pará.
17 478 2009. Disponível em:<http://iah.iec.pa.gov.br/iah/fulltext/pc/relatorios/barcarena2009.pdf>.
18 479 Ma, T.H., Cabrera, G.L., Chen, R., Gill, B.S., Sandhu, S.S., Vandenberg, A.L., Salamone, M.F., 1994.
19 480 Tradescantia micronucleus bioassay. Mutat Res 310, 221-230.
20 481 Ma, T.H., Davies, B.M., 2009. Effects of extreme low dose radiation in wet condition detected by
21 482 Trad-MCN assay, Genomics in the environmental century. 40th Annual meeting EMS. EMS, St.
22 483 louis Missouri.
23 484 Ma, T.H., Xu, Z., Xu, C., McConnell, H., Rabago, E.V., Arreola, G.A., Zhang, H., 1995. The improved
24 485 Allium/Vicia root tip micronucleus assay for clastogenicity of environmental pollutants. Mutat
25 486 Res 334, 185-195.
26 487 Majer, B.J., Tschерko, D., Paschke, A., Wennrich, R., Kundi, M., Kandeler, E., Knasmueller, S., 2002.
27 488 Effects of heavy metal contamination of soils on micronucleus induction in Tradescantia and
28 489 on microbial enzyme activities: a comparative investigation. Mutat Res 515, 111-124.
29 490 Manerikar, R.S., Apte, A.A., Ghole, V.S., 2008. In vitro and in vivo genotoxicity assessment of Cr(VI)
30 491 using comet assay in earthworm coelomocytes. Environ Toxicol Pharmacol 25, 63-68.
31 492 Marcano, L., Carruyo, I., Fernandez, Y., Montiel, X., Torrealba, Z., 2006. Determination of vanadium
32 493 accumulation in onion root cells (Allium cepa L.) and its correlation with toxicity. Biocell 30,
33 494 259-267.
34 495 Mayes, W., Jarvis, A., Burke, I., Walton, M., Gruiz, K., 2011a. Trace and rare earth element dispersal
35 496 downstream of the Ajka red mud spill, Hungary in: Rűde, R.T., Freund, A., Wolkersdorfer, C.
36 497 (Eds.), Mine Water - Managing the challenges. International Mine Water Association, pp. 29 -
37 498 34
38 499 Mayes, W.M., Jarvis, A.P., Burke, I.T., Walton, M., Feigl, V., Klebercz, O., Gruiz, K., 2011b. Dispersal
39 500 and attenuation of trace contaminants downstream of the Ajka bauxite residue (red mud)
40 501 depository failure, Hungary. Environ Sci Technol 45, 5147-5155.
41 502 Minouflet, M., Ayrault, S., Badot, P.M., Cotelle, S., Ferard, J.F., 2005. Assessment of the genotoxicity
42 503 of 137Cs radiation using Vicia-micronucleus, Tradescantia-micronucleus and Tradescantia-
43 504 stamen-hair mutation bioassays. J Environ Radioact 81, 143-153.
44 505 Misik, M., Ma, T.H., Nersesyan, A., Monarca, S., Kim, J.K., Knasmueller, S., 2011. Micronucleus assays
45 506 with Tradescantia pollen tetrads: an update. Mutagenesis 26, 215-221.
46 507 Misik, M., Micieta, K., Solenska, M., Misikova, K., Pisarcikova, H., Knasmueller, S., 2007. In situ
47 508 biomonitoring of the genotoxic effects of mixed industrial emissions using the Tradescantia
48 509 micronucleus and pollen abortion tests with wild life plants: demonstration of the efficacy of
49 510 emission controls in an eastern European city. Environ Pollut 145, 459-466.
50 511 Peacock, C.L., Sherman, D.M., 2004. Vanadium(V) adsorption onto goethite (alpha-FeOOH) at pH 1.5
51 512 to 12: A surface complexation model based on ab initio molecular geometries and EXAFS
52 513 spectroscopy. Geochimica Et Cosmochimica Acta 68, 1723-1733.

Formatted: German (Austria)

Formatted: German (Austria)

53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 514 Rajeev, Pradhan, J., Das, S.N., Thakur, R.S., 1999. Vanadium sludge - An useful byproduct of alumina
4 515 plant. *Journal of Scientific & Industrial Research* 58, 948-953.
5 516 Renforth, P., Mayes, W.M., Jarvis, A.P., Burke, I.T., Manning, D.A., Gruiz, K., 2012. Contaminant
6 517 mobility and carbon sequestration downstream of the Ajka (Hungary) red mud spill: The
7 518 effects of gypsum dosing. *Sci Total Environ* 421-422, 253-259.
8 519 Rodrigues, G.S., Ma, T.-H., Pimentel, D., Weinstein, L.H., 1997. *Tradescantia* Bioassays as Monitoring
9 520 Systems for Environmental Mutagenesis: A review. *Crit. Rev. Plant Sci.* 16, 325-359.
10 521 Ruyters, S., Mertens, J., Vassilieva, E., Dehandschutter, B., Poffijn, A., Smolders, E., 2011. The red
11 522 mud accident in Ajka (Hungary): plant toxicity and trace metal bioavailability in red mud
12 523 contaminated soil. *Environ Sci Technol* 45, 1616-1622.
13 524 Samal, S., Ray, A.K., Bandopadhyay, A., 2012. Proposal for resources, utilization and processes of red
14 525 mud in India - A review. *International Journal of Mineral Processing*.
15 526 Sarkar, A., Ray, D., Shrivastava, A.N., Sarker, S., 2006. Molecular Biomarkers: their significance and
16 527 application in marine pollution monitoring. *Ecotoxicology* 15, 333-340.
17 528 Shaughnessy, D.T., DeMarini, D.M., 2009. Types and consequence of DNA damage in: Knasmueller, S.,
18 529 DeMarini, D.M., Johnson, I., Gerhauser, C. (Eds.), *Chemoprevention of Cancer and DNA*
19 530 *Damage by Dietary Factors*. Wiley-VCH, Weinheim, pp. 21-34.
20 531 Steinkellner, H., Mun-Sik, K., Helma, C., Ecker, S., Ma, T.H., Horak, O., Kundi, M., Knasmuller, S., 1998.
21 532 Genotoxic effects of heavy metals: comparative investigation with plant bioassays. *Environ Mol*
22 533 *Mutagen* 31, 183-191.
23 534 Taylor, J., Keith, S., Cseh, L., Ingerman, L., Chappell, L., Rhoades, J., Hueber, A., 2012. Toxicological
24 535 Profile for Vanadium. Agency for Toxic Substances and Disease Registry US.
25 536 <http://www.atsdr.cdc.gov/toxprofiles/tp58.pdf>, Atlanta, p. 255.
26 537 Zvereva, E., Toivonen, E., Kozlov, M.V., 2008. Changes in species richness of vascular plants under the
27 538 impact of air pollution: a global perspective. *Global Ecology and Biogeography* 17, 305-319.
28 539
29
30 540
31
32 541
33
34 542
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Formatted: German (Austria)

Table 1 Descriptions of the monitored sites (Kolontar – Devetsen, Hungary) with GPS position for more details see Fig 1.

Sampling site	Description	GPS location	pH of water extract	TOC (%)
Site 1 (S1)	Control site, agricultural area with soil of similar composition and structure as S2 and S3 but located on a hill (20 m above the site of red mud spill)	N47°05.400'	7.18 (0-5cm)	1.36 (0-5cm)
		EO17°28.159'	7.18 (5-15 cm)	1.23 (5-15 cm)
Site 2 (S2)	Agricultural field which was contaminated by red mud, 2.7 km northeast from the reservoir	N47°05.885'	9.21 (0-5 cm)	0.99 (0-5 cm)
		EO 17°27.947'	9.19 (5-15 cm)	1.07 (5-15 cm)
			8.26 (15-25 cm)	1.52 (15-25 cm)
Site 3 (S3)	Agricultural field which was strongly polluted, 200 m from S1 and 4.8 km northeast from the red mud reservoir	N47°05.905'	9.33 (0-5 cm)	0.22 (0-5 cm)
		EO 17° 28.067'	8.32 (5-15 cm)	1.05 (5-15 cm)
			8.24 (15-25 cm)	1.14 (15-25 cm)
Site 4 (S4)	Garden area located at a distance of 4.7 km from the reservoir which was flooded, red mud was removed from the surface by cleanup activities.	N47°05.410'	8.28 (0-5 cm)	
		EO17°28.253'	8.9 (5-15cm)	

Table 2 (A – C). Major elements (2A) and trace elements in soils (2B) and in aqueous leachates (2C) prepared from these samples.^a

Table 2A – Major elements in soil samples collected at sites 1 and 3

Site	SiO ₂	TiO ₂	Al ₂ O ₃	Fe ₂ O ₃	MnO	MgO	CaO	Na ₂ O	K ₂ O	P ₂ O ₅	SO ₃	LOI	Total
S1 (0-5 cm)	82.82	0.58	6.53	2.79	0.089	0.57	0.77	0.67	1.13	0.13	0.01	4.32	100.39
S3 (0-5 cm)	21.76	3.63	14.45	29.81	0.409	1.15	9.90	4.88	0.42	0.23	0.54	11.28	98.46
S3 (5-15 cm)	59.92	0.76	10.82	4.39	0.499	2.33	6.64	1.23	1.83	0.23	0.12	11.29	100.06
S3 (15-25 cm)	60.22	0.76	10.65	4.29	0.494	2.34	6.67	1.21	1.80	0.22	0.11	11.07	99.83

^aall values are indicated as w/w%

Table 2B – Trace elements in soil samples collected at sites 1 and 3.^a

Site	As	Ba	Ce	Co	Cr	Cs	Cu	Ga	La	Mo	Nb	Nd	Ni	Pb
S1 (0-5 cm)	17.8	266.0	44.4	7.3	40.0	1.5	54.1	7.1	24.7	0.9	9.0	21.5	12.5	22.2
S3 (0-5 cm)	201.1	120.4	437.8	54.6	703.3	12.3	112.1	27.0	192.6	9.3	76.9	153.2	283.1	143.0
S3 (5-15 cm)	22.3	414.0	69.7	17.8	76.5	1.7	36.8	12.1	32.3	3.5	13.5	31.3	31.7	23.8
S3 (15-25 cm)	21.7	417.5	70.8	18.5	125.1	3.6	26.0	12.3	34.2	2.4	13.1	31.2	27.7	22.3

Site	Rb	Sb	Sc	Se	Sn	Sr	Th	U	V	W	Y	Zn	Zr
S1 (0-5 cm)	56.4	0.9	7.7	0.5	30.1	232.3	4.8	1.4	57.9	0.8	18	42	180.2
S3 (0-5 cm)	26.8	15.9	94.4	1.1	20.1	357.6	64.6	16.0	898.8	9.1	125.2	123.4	920.5
S3 (5-15 cm)	80.1	1.0	14.4	0.5	12.9	166.1	9.2	3.6	97.4	1.6	32.2	57.8	271.1
S3 (15-25 cm)	76.9	1.0	15.5	1.0	6.9	161.2	8.9	3.3	100.0	1.7	30.5	56.8	273.7

^aall values are indicated as ~~ppm~~mg per kg of dry weight

Table 2C – Trace elements in aqueous leachates prepared from soil samples collected at sites 1 and 3.^a

Site	Al	As	Ba	Ca	Ce	Co	Cr	Cu	Fe	Ga	K	Li	Mg	Mn	Mo
S1 (0-5 cm)	0.010	0.001	0.104	100.178	0.038	0.002	0.002	0.005	0.008	0.013	24.601	0.003	9.574	0.008	0.011
S3 (0-5 cm)	0.951	0.123	0.005	5.253	0.019	0.002	0.068	0.041	0.123	0.02	45.979	0.01	1.703	0.005	0.364
S3 (5-15 cm)	0.000	0.002	0.061	127.421	0.001	0.002	0.0002	0.002	0.517	0.002	8.402	0.003	11.558	11.697	0.052
S3 (15-25 cm)	0.001	0.001	0.044	135.985	0.023	0.002	0.003	0.003	0.073	0.002	26.412	0.005	6.767	0.002	0.004

Site	Na	Ni	P	Pb	S	Se	Si	Sr	Ti	V	Zn	Zr	Hg
S1 (0-5 cm)	2.25	0.002	0.018	0.005	2.681	0.39	3.491	0.278	0.001	0.004	0.017	0.001	0.001
S3 (0-5 cm)	701.321	0.002	0.175	0.005	122.704	0.453	1.862	0.028	0.017	1.169	0.019	0.005	0.001
S3 (5-15 cm)	109.365	0.002	0.101	0.005	14.557	0.05	4.718	0.929	0.000	0.002	0.021	0.001	0.001
S3 (15-25 cm)	19.545	0.002	0.108	0.005	30.032	0.316	2.31	0.478	0.000	0.001	0.001	0.001	0.001

^aall values are indicated as mg/L

ppm

Figure 1




Figure 2A and B




Figure 3A-C






Figure 4





**Institut für
Krebsforschung**

A. o. Univ. Prof. Dr. Siegfried Knasmüller

Abteilung Environmental Toxicology Group

Borschkegasse 8a

A-1090 Wien, Österreich



**MEDIZINISCHE
UNIVERSITÄT
WIEN**

Klinik für Innere Medizin I

A. o. Univ. Prof. Dr. Siegfried Knasmüller

Head Environmental Toxicology Group,

email: siegfried.knasmueller@meduniwien.ac.at

Dear Editor,

On behalf of all authors I declare that there is no conflict of interest and no financial support was received to carry out the study which is describe in the article.

Best personal greetings

Siegfried Knasmueller

Head of Environmental Toxicology Group

20-2-2014 Wien