

Rothamsted Repository Download

A - Papers appearing in refereed journals

Chirinda, N., Loaiza, S., Arenas, L., Ruiz, V., Faverin, C., Alvarez, C., Savian, J. V., Belfon, R., Zuniga, K., Morales, L., Trujillo, C., Arango, M., Rao, I., Arango, J., Peters, M., Barahona, R., Costa, C., Rosenstock, T. S., Richards, M., Martinez-Baron, D. and Cardenas, L. M. 2019.
Adequate vegetative cover decreases nitrous oxide emissions from cattle urine deposited in grazed pastures under rainy season conditions.
Scientific Reports. 9 (908).

The publisher's version can be accessed at:

- <https://dx.doi.org/10.1038/s41598-018-37453-2>

The output can be accessed at: <https://repository.rothamsted.ac.uk/item/84v9x>.

© 29 January 2019, Rothamsted Research. Licensed under the Creative Commons CC BY.

SCIENTIFIC REPORTS



OPEN

Adequate vegetative cover decreases nitrous oxide emissions from cattle urine deposited in grazed pastures under rainy season conditions

Ngonidzashe Chirinda¹, Sandra Loaiza¹, Laura Arenas¹, Verónica Ruiz², Claudia Faverín³, Carolina Alvarez⁴, Jean Víctor Savian⁵, Renaldo Belfon⁶, Karen Zuniga⁷, Luis Alberto Morales-Rincon⁸, Catalina Trujillo¹, Miguel Arango¹⁰, Idupulapati Rao^{1,11}, Jacobo Arango¹, Michael Peters¹, Rolando Barahona⁹, Ciniro Costa Jr.¹², Todd S. Rosenstock¹³, Meryl Richards¹⁴, Deissy Martinez-Baron¹ & Laura Cardenas¹⁵

A decline in pasture productivity is often associated with a reduction in vegetative cover. We hypothesize that nitrogen (N) in urine deposited by grazing cattle on degraded pastures, with low vegetative cover, is highly susceptible to losses. Here, we quantified the magnitude of urine-based nitrous oxide (N₂O) lost from soil under paired degraded (low vegetative cover) and non-degraded (adequate vegetative cover) pastures across five countries of the Latin America and the Caribbean (LAC) region and estimated urine-N emission factors. Soil N₂O emissions from simulated cattle urine patches were quantified with closed static chambers and gas chromatography. At the regional level, rainy season cumulative N₂O emissions (3.31 *versus* 1.91 kg N₂O-N ha⁻¹) and emission factors (0.42 *versus* 0.18%) were higher for low vegetative cover compared to adequate vegetative cover pastures. Findings indicate that under rainy season conditions, adequate vegetative cover through proper pasture management could help reduce urine-induced N₂O emissions from grazed pastures.

The livestock sector accounts for 46% of the agricultural gross domestic product of the Latin America and the Caribbean (LAC) region and grows at 3.7% annually¹. Expanding livestock production is driven by a rapid increase in demand for cattle meat². This increased demand for animal products together with the development of improved forage options to sustain higher levels of cattle productivity increases pressure on grasslands, the dominant cattle production systems of LAC, resulting in overgrazing and degradation of pastures³. According to Kwon³, an estimated 157 million ha (8% of total grazing area) of the grazing area in LAC is degraded. In Brazil

¹International Center for Tropical Agriculture (CIAT), A.A, 6713, Cali, Colombia. ²National Autonomous University of Nicaragua, Managua, Nicaragua. ³National Institute of Agricultural Technology (INTA), Balcarce, Argentina. ⁴National Institute of Agricultural Technology (INTA), Manfredi, Argentina. ⁵Federal University of Rio Grande do Sul (UFRGS), Porto Alegre, Brazil. ⁶The University of the West Indies, St. Augustine, Trinidad and Tobago. ⁷Universidad Nacional de Colombia, sede Palmira, Colombia. ⁸Universidad Nacional de Colombia, Bogotá, Colombia. ⁹Universidad Nacional de Colombia, Medellín, Colombia. ¹⁰The Colombian Corporation for Agricultural Research (AGROSAVIA), Villavicencio, Meta, Colombia. ¹¹Present address: Plant Polymer Research Unit, National Center for Agricultural Utilization Research, Agricultural Research Service, United States Department of Agriculture, 1815 North University Street, Peoria, IL, 61604, USA. ¹²Institute of Agriculture and Forestry Management and Certification (IMAFLOA), Estrada Chico Mendes, Piracicaba, Estrada Chico Mendes, 185, 13426-420, Brazil. ¹³World Agroforestry Center (ICRAF), c/o INERA, Avenue des cliniques, Kinshasa, Democratic Republic of the Congo. ¹⁴Rubenstein School of Environment and Natural Resources, University of Vermont, Burlington, VT, 05405, USA. ¹⁵Rothamsted Research, Sustainable Agriculture Sciences Department, North Wyke, Devon, EX20 2SB, UK. Correspondence and requests for materials should be addressed to N.C. (email: n.chirinda@cgiar.org)

Country	Location	Pasture Condition	Texture	pH	BD (g cm ⁻³)	SOC (%)	SON(%)
Nicaragua	Estelí	AVC	Loam	6.4	1.1	3.0	0.2
		LVC	Clay	7.4	0.6	5.0	0.4
Colombia	Patía	AVC	Clay	6.4	1.5	2.2	0.2
		LVC	Clay	6.3	1.6	2.0	0.2
Colombia	Taluma	AVC	Clay loam	5.8	1.3	1.3	0.1
		LVC	Loam	5.2	1.5	1.3	0.1
Brazil	Rio Grande do Sul	AVC	Clay loam	5.0	1.6	1.4	0.1
		LVC	Clay loam	5.0	1.5	1.3	0.1
Argentina	Balcarce INTA	AVC	Sandy loam	7.5	1.0	3.5	0.3
		LVC	Sandy-clay-loam	8.9	1.1	3.3	0.3
Argentina	Manfredi INTA	AVC	Silt-loam	6.4	1.2	1.8	0.2
		LVC	Silt-loam	6.2	1.2	1.7	0.2
Trinidad and Tobago	St. Augustine	AVC	Sandy-Loam	5.0	*	*	*
		LVC	Sandy-Loam	5.1	*	*	*

Table 1. Soil physical and chemical characteristics of the field sites of study areas. AVC-Adequate vegetative cover; LVC- Low vegetative cover; BD: Bulk density, SOC: Soil organic carbon, SON: Soil organic nitrogen; *Missing data.

half of the 80 million ha of introduced tropical pastures are estimated to be in some state of degradation as they have, among other symptoms, low soil cover⁴.

Cattle excreta deposited on grazed pastures is estimated to represent 16% of global anthropogenic nitrous oxide (N₂O) emissions, a powerful greenhouse gas (GHG)⁵. About 75–95% of cattle ingested N is excreted in either urine or dung, which provides N-rich substrate for nitrification and denitrification^{6,7}. Cattle urine patches can contain very high amounts of soluble N (equivalent to 500–1000 kg N ha⁻¹), more than 2–3 times of the N uptake capacity of pastures⁸. Annually, about 1.5 Tg of total global anthropogenic N₂O emissions (6.7 Tg N₂O-N yr⁻¹) are emitted from excreta produced by grazing cattle^{9,10} through both direct and indirect (from leached and volatilized excreta nitrogen) emissions. About 2% (0.7–6% uncertainty)¹¹ of the nitrogen (N) in deposited urine is lost as N₂O. Lower emission factors (EFs) (<0.7%), reported in other studies have been attributed to differences in climatic conditions, texture, soil moisture, and the N concentration in animal excreta¹².

Pasture degradation may stimulate or constrain N losses. For example low vegetative cover, may reduce N sinks for deposited excreta and thus increase the vulnerability of N to loss through soil microbial processes and leaching. However, the low vegetative cover may also be associated with fewer plant root exudates and thus suppress microbial activity and N₂O emissions¹³. On the other hand, overstocking and overgrazing without time for pasture recovery increases the risk of soil compaction - an indicator of pasture degradation. Soil compaction reduces soil porosity and pore continuity, decreases soil aeration, restricts plant growth and thus, consequently, increases soil N₂O emissions from urine patches^{14,15}. Soil acidification, which could also be an indicator of pasture degradation, has been shown to increase N₂O emissions as acidic conditions generally reduce plant growth and inhibit N₂O reductase enzyme activity which is responsible for transforming N₂O to dinitrogen (N₂)^{16,17}.

Clearly, the effect of pasture degradation on N₂O emissions from urine deposition can influence emission through multiple, often interacting, mechanisms and thus has produced contradictory results in the literature. Previous studies suggest that variations in soil N₂O emissions from deposited urine patches in grazed pastures are driven by differences in several factors including ambient temperature¹⁸, urine volume and urine-N content^{15,19}, soil drainage^{20,21}, and soil moisture^{22,23}. No previous studies have systematically explored the variation in urine-based soil N₂O emissions associated with low vegetative cover in pastures.

Here we tested the hypothesis that N₂O emissions from cattle urine deposited on grazed pastures with adequate vegetative cover are less intense than those from pastures with lower vegetative cover by measuring soil N₂O fluxes from urine patches deposited on different pastures located at seven contrasting sites, spread across five countries in the LAC region during rainy season.

Results

Soil texture at most of the study sites was similar in the low and adequate vegetation cover pastures with the exception of Balcarce (Argentina), Estelí (Nicaragua) and Taluma (Colombia) (Table 1). Soil pH values at the study sites ranged between 5.0 and 8.9, with acidic soils (pH < 6) at Taluma (Colombia), Rio Grande do Sul (Brazil), St. Augustine (Trinidad and Tobago) and neutral to basic soils at other sites. Soil bulk density at the study sites ranged between 0.6 and 1.6 g cm⁻³ and was generally similar between the low and adequate vegetative cover pastures at each study location. The largest differences in bulk density, soil organic carbon and soil organic nitrogen between low and adequate vegetative cover pastures at the Estelí location in Nicaragua (Table 1).

Air temperature and rainfall data for one week before and during the sampling dates are shown in Supplementary Fig. S1. The mean daily temperatures during this period ranged from 19 °C to 24 °C for Estelí (Nicaragua), 23 °C to 31 °C for Patía (Colombia), 27 °C to 29 °C for Taluma (Colombia), 17 °C to 25 °C for Rio Grande do Sul (Brazil), 15 °C to 26 °C for Balcarce (Argentina), 8 °C to 16 °C for Manfredi (Argentina) and 26 °C




Figure 1. Soil N_2O emission from two pasture conditions with cattle urine application at seven field sites in five Latin - America and the Caribbean countries. Error bars represent standard error of the mean. (AVC: Adequate vegetative cover, LVC: Low vegetative cover).

to 30 °C for St. Augustine (Trinidad and Tobago). Rainfall was recorded on less than 19 days during the N_2O monitoring period, with the exception of the Trinidad and Tobago site which received 31 days of rainfall (Fig. S1).

N_2O emission peaks observed in LVC pastures tended to be higher than those in AVC pastures in 5 out of the 7 sites (Fig. 1; Table 2). However, the delayed N_2O peaks observed at Rio Grande do Sul (Brazil) were higher in the AVC (59 ± 11 mg $N_2O-N m^{-2} day^{-1}$) compared to the LVC (45 ± 18 mg $N_2O-N m^{-2} day^{-1}$) pasture. The level of N_2O emissions observed at Balcarce and Manfredi (Argentina) and Taluma (Colombia) sites were lower than 50 mg $N_2O-N m^{-2} day^{-1}$.

Country	Location	Pasture Condition	Nitrogen in applied urine (kg N ha ⁻¹)	Peak N ₂ O emissions (mg N ₂ O-N m ⁻² d ⁻¹)	Cumulative N ₂ O emissions (kg N ₂ O-N ha ⁻¹)
Nicaragua	Estelí	LVC	464	129 (19)	5.82 (0.73) ^a
		AVC		60 (5)	1.85 (0.26) ^b
Colombia	Patía	LVC	789	92 (26)	3.85 (0.71) ^a
		AVC		27 (2)	1.41 (0.51) ^b
Colombia	Taluma	LVC	112	0.2 (0.2)	0.02 (0.01) ^a
		AVC		0.2 (0.1)	0.02 (0.005) ^a
Brazil	Rio Grande do Sul	LVC	619	45 (18)	4.59 (1.23) ^a
		AVC		59 (26)	3.01 (1.88) ^a
Argentina	Balcarce	LVC	1641	29 (10)	1.23 (0.57) ^a
		AVC		22 (2)	0.90 (0.14) ^a
Argentina	Manfredi	LVC	546	4.2 (0.5)	0.21 (0.01) ^a
		AVC		3 (0.5)	0.18 (0.02) ^a
Trinidad & Tobago	St. Augustine	LVC	*	107 (14)	7.49 (1.26) ^a
		AVC		80 (7)	6.00 (0.23) ^a

Table 2. Nitrogen inputs in applied urine, peak N₂O emissions, cumulative N₂O emissions. AVC-Adequate vegetative cover; LVC- Low vegetative cover; number in parenthesis indicates standard error of mean (s.e.m.). At each site values with the same letter for the cumulative N₂O emission are not significantly different ($P < 0.05$).




Figure 2. Emission factor (percent per applied nitrogen) from two pasture types (Adequate vegetative cover and Low vegetative cover) with the application of cattle urine. Error bars represent standard error of the mean.

The N content in applied cattle urine ranged from 112–1,641 kg N ha⁻¹ (Table 2). Over a one-month period, the soil N₂O emission factor of N in applied urine to soil ranged from 0.01 to 1.23%. The highest N₂O emission factor values observed for the LVC pasture (1.23% of applied urine-N) and AVC (0.48% of applied urine-N) pasture were at Estelí (Nicaragua) and Rio Grande do Sul (Brazil), respectively. On the other hand, the lowest N₂O emission factors for LVC (0.02% of applied urine-N) and AVC pasture (0.01% of applied urine-N) were both observed at the Taluma site in Colombia (Fig. 2). At the regional level, mean cumulative N₂O emissions observed in the urine treatments ranged between 0.02 and 7.5 kg N₂O-N ha⁻¹. The highest cumulative N₂O emissions for both treatments (LVC and AVC pasture) were observed at St. Augustine (Trinidad and Tobago) and Estelí (Nicaragua) and the lowest were observed at Taluma (Colombia) (Table 2).

At the regional level, mean N₂O emission factors were significantly, at most 2.5 times, higher ($P < 0.0002$) in LVC ($0.42 \pm 0.19\%$ SEM) than AVC ($0.18 \pm 0.08\%$ SEM) pastures. Also at the regional level mean cumulative N₂O emissions in the LVC (3.31 ± 1.09 kg N₂O-N ha⁻¹ SEM) were higher than those observed in the AVC (1.91 ± 0.78 kg N₂O-N ha⁻¹ SEM) pasture at the 10% level of significance ($P = 0.08$), based on results presented in Table 2. The N₂O emissions at each individual site tended to be higher in LVC pastures than in the AVC pastures, with t-test detecting significant differences at the Nicaragua (Estelí) and Colombia (Patía) sites (Table 2). Rainfall for the measurement period (including data for the period of one week before commencing the monitoring campaigns) explained less of the variation in N₂O emission factors in the LVC (66%) pasture compared to the AVC (88%) pasture (Supplementary Fig. S2). No clear effects of air temperature were observed on N₂O emission factors.

Discussion

The key finding of our study is that, at the regional level of LAC, N_2O emission factors from cattle urine patches in grazed pastures are lower for AVC compared to LVC pastures during rainy season, suggesting the importance of adequate pasture improvement/management in mitigating soil N_2O emissions. We did not have soil moisture data for all the sites to be able to compute the effects of water-filled pore space, which has been shown to be a major driver of N_2O emissions^{24,25}. The strong correlation observed between rainfall and the N_2O emission factor suggests that soil moisture was possibly a key driver of N_2O emissions in the current study. However, the fact that rainfall explained less of the variation in N_2O emissions in the pasture with LVC compared to one with AVC may also imply that other factors, such as vegetative cover, are drivers of N_2O emissions. Yet, interestingly, the steeper slope observed for the LVC (0.0086) compared to the AVC (0.0041) pastures suggests that urine deposited on LVC pastures is more vulnerable to high N_2O losses when exposed to high rainfall. This may explain the significant differences between LVC and AVC pastures that were observed at Estelí (Nicaragua) and Patía (Colombia), where rainfall was high and the observed separation between the LVC and AVC pastures at the Rio Grande do Sul (Brazil) site where rainfall was also high.

High peaks of N_2O emissions observed in LVC pastures compared to AVC pastures were likely due to lower plant N uptake from soil. Moreover, this may in part explain why AVC pastures generally resulted in lower net cumulative N_2O emissions compared to degraded pastures. Despite the significant difference ($P < 0.10$) in net cumulative N_2O emissions between LVC and AVC pastures, at a regional level, site-level comparisons showed that cumulative N_2O emissions from LVC pastures were only significantly higher than those of AVC pastures in two sites, at Estelí (Nicaragua) and Patía (Colombia). The fact that soils within the LVC pastures at the Estelí site were more clayey than those under AVC pastures may have also contributed to the high net cumulative N_2O emission in the former. Previous studies have reported higher N_2O emissions from urine deposited on fine textured soils²⁶. However, at Patía, where both the LVC and AVC pastures were on a clay soil observed differences suggest that despite the obvious influence of soil texture, pasture condition, based on vegetative cover is a driver of N_2O emissions.

Low emissions of N_2O observed at the two study sites in Argentina could have been due to the lower mean air temperatures. At the Balcarce location, mean air temperatures and the amount of N in applied urine were higher than at the Manfredi location. Low temperatures are known to reduce microbial activity and thus the rate of N transformation processes such as nitrification and denitrification in soil and, consequently, N_2O production²⁷. We assume that low temperatures may also be the cause of low net emissions and emission factors at the site in Brazil. It is important to note that the high soil pH (>7) at the Balcarce site, which would be expected to further increase with urine application may have also resulted in the inhibition of nitrification and high ammonia volatilization^{28–30}. In addition, at the Balcarce site, the high urine-N levels may have resulted in microbial stresses with possible impacts on soil N transformation³¹ and thus contributed to the low N_2O emissions and emission factors.

At the Taluma location in Colombia, the absence of a N_2O emission peak following urine application may be because the frequency of measurements at this site was insufficient to capture the expected N_2O emission spike. Several other studies using manual static chambers have reported having missed the N_2O peak, due to the low temporal resolution^{32–34}. This problem can be resolved by increasing the frequency of monitoring using manual static chambers or switching to automated chambers. Alternatively, the N content in applied urine was also the lowest at Taluma, which implies that N_2O fluxes could have been limited by N substrate availability. In addition, the forage grass (*Brachiaria humidicola*) that was used at Taluma had high nitrification inhibition capacity^{35–37}, which could have also contributed to the observed low N_2O emissions.

Absence of significant differences, in N_2O emissions, between LVC and AVC pastures at the St. Augustine (Trinidad and Tobago), Balcarce (Argentina), and Manfredi (Argentina) sites was possibly due to the fact that the spatial variation of vegetative cover of grass (soil cover) in the LVC (50–70%) pasture was high. As a result, local farmers based their classification on animal productivity differences which are influenced by both quantity (biomass) and quality (e.g. digestibility and crude protein content) of the forage on offer to animals. This further suggests that N_2O emission differences between LVC pastures and AVC pastures are driven by differences in soil cover. With high plant density and greater plant vigour, we expect greater uptake of the urine-N by plants which could reduce the amount of N available for microbial transformations in soil such as nitrification and denitrification. It is therefore not surprising that when soil cover was high in the LVC pastures, there was no significant difference in N_2O emissions with AVC pastures. This was however not the case for LVC pasture at the Patía (Colombia) site, which, though having similar soil cover (50–70%) showed significant differences between LVC and AVC pastures. This difference may be due to dissimilar vegetation types at the studied sites which would also affect N uptake and thus N availability for N_2O emissions³⁸.

The IPCC Tier 1 emission factor for urine deposited on grazed forages is 2% with an uncertainty range of 0.7–6%¹¹. During this short-term study, several of the emission factors were below the uncertainty range of the IPCC Tier 1 emission factor. While this may be due to the short gas monitoring period (1 month), several other studies conducted under temperate conditions^{39–42}, reported a similar range of emission factors (0.02–1.63%) as we observed under warm temperate or sub-tropical conditions in Argentina and Brazil (0.02–0.7%). Similarly, the range of emission factors reported from this study under the tropical conditions in Colombia, Nicaragua and Trinidad and Tobago (0.01–1.2%) are in agreement with the range of values that have been reported from studies conducted under tropical conditions^{43,44}.

We conclude that in addition to the known effects of rainfall, temperature and the amount of urine-N, the pasture condition based on vegetative cover also influences N_2O emissions from cattle urine patches. When pasture degradation is associated with a reduction in vegetative cover, N_2O emissions are expected to increase. Therefore, better regional understanding of the state of pasture degradation is vital for a robust understanding of N_2O emissions from cattle urine deposits. More importantly, these findings suggest that improving soil cover/pasture condition through adoption of appropriate grazing and nutrient management practices may contribute towards

mitigating excreta-based soil N₂O emissions from grazed pastures during the rainy season. We expect findings from this regional study to contribute towards reducing uncertainties in future assessments on the importance of improving grassland management to achieve global commitments, such as the Bonn Challenge, 20 × 20 initiative⁴⁵ and the Paris Agreement⁴⁶.

Methods

The experimental plots were located at seven different sites in five countries in LAC spanning diverse climatic conditions and soil types (Supplementary Fig. S3 and Table S1). Rainfall is bimodal in Estelí (Nicaragua) with two seasons from May to June and August to November. In Patía (Colombia), the wet seasons occur during the period from March to May and September to November. Rainfall in St. Augustine (Trinidad and Tobago) is also bimodal with wet seasons occurring from June to August and October to November. At the rest of the study sites rainfall is unimodal with the main rains occurring during the period February to December and September to April at Taluma (Colombia) and the two Argentinean sites, respectively. During the monitoring period rainfall and air temperature data was collected at the nearest weather stations at each of the different study sites.

At each of the study sites, paired experimental plots were set-up on fields with grazed pastures that were classified as having either low vegetative cover (LVC) or adequate vegetative cover (AVC). Pairs of LVC and AVC pastures were not always available at the very same location, but were always less than 1 km apart. We used a qualitative approach including expert knowledge, farmer perceptions and an arbitrary ranking system based on soil cover to define pasture conditions based on vegetative cover using criteria that combined those used by Hollman⁴⁷, Brown⁴⁸ and McCormick and Lodge⁴⁹. Specifically, through visual assessments by forage scientists⁵⁰ we broadly described soil cover as follows: low vegetation cover (<70%) and adequate vegetation cover (>70%) pastures where soil cover is simply the proportion of soil covered by vertical projection of a plant canopy and vegetative biomass (Supplementary Table S2). Aboveground biomass data for the study sites were also obtained from historical records when available. In addition, we also used local farmer assessments to differentiate low and adequate vegetation cover pastures.

On each pasture (low vegetative cover or adequate vegetative cover), experimental plots were organized following a systematic experimental design^{51,52} with five replicates per treatment (Supplementary Fig. S4). The two treatments were urine application and a control-without urine application. The urine was applied to individual independent plots and so there are five replicates for each of the control *versus* urine treatment within each site⁵³. Individual replicate plots (2 m × 2 m) were demarcated within each pasture condition for making measurements of soil properties and N₂O emissions. To simulate grazing, grass in each plot was cut to approximately 5 cm sward height, seven days prior to the beginning of the gas and soil sampling.

Prior to starting the experiment, ten soil subsamples (0–10 cm) were separately collected from LVC and AVC pastures, using augers with 5 cm diameter, and combined to give one composite soil sample for each pasture condition. Collected soil was characterized for texture, pH, total carbon (C) and organic and inorganic N as described by Gee and Bauder⁵⁴, McLean⁵⁵, and Vogel⁵⁶, respectively. A total of 20 cylindrical PVC static chamber bases (10 per treatment) were inserted at the center of each subplot to a depth of 5 cm, five days prior to the start of gas and soil sampling. For each treatment, chamber bases with an internal diameter of 25 cm and a height of 10 cm were distributed in five replicate plots. At each site, cattle urine samples (about 7 L) that were collected from at least 10 local dairy cows were pooled and, immediately, following setting aside a subsample for N analysis, 500 ml of the collected urine was applied to soils to simulate a urination event on soil within each static chamber base at a rate of 1.27 L urine/m². Nitrogen concentration in urine was characterized in each of the study countries using the direct distillation method described by Hoogendoorn *et al.*⁵⁷. Unfortunately, we were unable to quantify urine-N in Trinidad and Tobago as there are currently no laboratories that quantify N in animal urine samples.

Gas measurements were conducted from a total of 20 non-vented PVC static chambers (10 cm height and 25 cm diameter) fitted with two rubber septa (one for gas sampling and another for inserting a thermometer). On each sampling day, PVC chambers were fitted to the chamber bases and sealed with an air-tight rubber belt. Syringes (15 ml) fitted with hypodermic needles were used to collect four gas samples from each of static chambers following chamber closure and at 15, 30 and 45 minutes after chamber closure. Collected gas samples were transferred to pre-evacuated 10 ml headspace glass vials fitted with rubber butyl septa crimp caps. At each site, at least eight gas samples were collected between the months of November to December in 2015 for the localities of Patía (Colombia) and Rio Grande do Sul (Brazil), Estelí (Nicaragua) and St. Augustine (Trinidad and Tobago). For Taluma (Colombia) and Balcarce (Argentina) measurements were conducted between February to March (2016) and at Manfredi, Argentina in the month of May (2016). Sampling months were chosen to coincide with the wet seasons at each of the study sites. The use of non-vented chambers has been reported to cause bias in the flux estimates⁵⁸. Yet, Davidson *et al.*⁵⁹ reported the possibility of artefacts with both vented and non-vented chambers, making it difficult to know which chamber yielded the ‘true’ flux. Since similar non-vented chambers were used at all sites, the chambers did not affect observed differences between AVC and LVC pastures. However, the calculated N₂O emissions factors may differ from those measured in other published studies that used vented chambers.

Gas sampling frequency was as follows: once before the application of urine, 1 hour after urine application, daily for the first three days following urine application, twice a week during the second and third week and once during last week of the experiment. Due to logistical challenges, less frequent measuring campaigns were done at the Taluma site in Colombia. Immediately after the gas sampling campaign, vials were sent to the Greenhouse Gas Laboratory at the International Center for Tropical Agriculture (CIAT) in Colombia, where N₂O concentration was analyzed by gas chromatography (GC-2014 Shimadzu), within a month upon arrival. The daily N₂O fluxes were calculated by regressing N₂O emissions from each chamber on each sampling date against time in order to calculate the hourly flux which was then multiplied by 24 to determine the daily flux. Each calculated flux was

corrected for temperature and barometric pressure according to Ideal Gas Law. Subsequently, cumulative fluxes were calculated from daily N₂O fluxes by interpolation between measurement days⁶⁰.

The N₂O–N emission factor for urine patches was calculated according to Sordi *et al.*⁶¹:

$$EF(\%) = \frac{(N_2O-N \text{ emitted}) - (N_2O-N \text{ control})}{N \text{ applied}} \times 100$$

where EF is the emission factor, N₂O–N_{emitted} and N₂O–N_{Control} are the cumulative N₂O emissions from urine or control patches over the 18 to 24 days monitoring period. N_{applied} represents the amount of N in the applied urine.

Statistical analyses were conducted using the PROC MIXED procedure of SAS⁶². Cumulative N₂O fluxes and emission factors were, correspondingly, log and square-root transformed to achieve normality and obtain homogeneous variances. To determine effects at the region level, cumulative N₂O fluxes were analyzed using a split-plot ANOVA where the main plot was the pasture condition (LVC, AVC) and split-plot was the nitrogen levels (with and without urine application) and the blocking factor was the location and the main plot error term was pasture condition nested within location. In addition, at the regional level the emission factor variable was analyzed using a one-way ANOVA where the treatment effect was the pasture condition and the blocking factor was the location (site). At the individual sites we used the t-test analysis for testing differences in emission factors as influenced by pasture condition.

References

1. Arequipa, P. FAO Regional Office for Latin America and the Caribbean. *Santiago de Chile* (2003).
2. ECLAC-FAO-IICA. The Outlook for Agriculture and Rural Development in the Americas: A Perspective on Latin America and the Caribbean 2015–2016. IICA, San José. (2011).
3. Kwon, H., Nkonya, E., Johnson, T. & Kato, E. Chapter 8: Global Estimate of the Impacts of Grassland Degradation on Livestock Productivity from 2001 to 2011. In E. Nkonya, A. Mirzabaev & J. Von Braun, *Economics of Land Degradation and Improvement – A Global Assessment for Sustainable Development* (pp. 197–213). Washington, DC. (2016).
4. Boddey, R. M. *et al.* Nitrogen cycling in Brachiaria pastures: The key to understanding the process of pasture decline. *Agriculture, Ecosystems and Environment* **103**, 389–403 (2004).
5. Tubiello, F. N. *et al.* Agriculture, forestry and other land use emissions by sources and removals by sinks. *ESS Working Paper* **2**, 4–89 (2014).
6. Eckard, R. J., Grainger, C. & de Klein, C. A. M. Options for the abatement of methane and nitrous oxide from ruminant production: A review. *Livestock Science* **130**, 47–56 (2010).
7. Saggari, S. *et al.* Denitrification and N₂O:N₂ production in temperate grasslands: Processes, measurements, modelling and mitigating negative impacts. *Science of the Total Environment* **465**, 173–95 (2013).
8. Haynes, R. J. & Williams, P. H. Nutrient cycling and soil fertility in the grazed pasture ecosystem. *Advances in Agronomy* **49**, 119–199 (1993).
9. Oenema, O. *et al.* Trends in global nitrous oxide emissions from animal production systems. *Nutrient Cycling in Agroecosystems* **72**, 51–56 (2005).
10. Taghizadeh-Toosi, A. *et al.* Biochar incorporation into pasture soil suppresses in situ nitrous oxide emissions from ruminant urine patches. *Journal of Environmental Quality* **40**, 468–76 (2011).
11. Intergovernmental Panel on Climate Change (IPCC) Guidelines for national greenhouse gas inventories. Greenhouse Gas Inventory Reference Manual, 4. Intergovernmental Panel on Climate Change. Available at, <http://www.ipccnggip.iges.or.jp/public/2006gl/vol4.html> (2006).
12. Rochette, P. *et al.* Soil Nitrous Oxide Emissions Following Deposition of Dairy Cow Excreta in Eastern Canada. *Journal of Environmental Quality* **43**, 829–41 (2014).
13. Henry, S., Texier, S. & Hallet, S. Disentangling the rhizosphere effect on nitrate reducers and denitrifiers: insight into the role of root exudates. *Environmental Microbiology* **10**, 3082–3092 (2008).
14. Hansen, S., Maehlum, J. E. & Bakken, L. R. N₂O and CH₄ fluxes in soil influenced by fertilization and tractor traffic. *Soil Biol. Biochem.* **25**, 621–630 (1993).
15. Van Groenigen, J. W., Kuikman, P. J., De Groot, W. J. M. & Velthof, G. L. Nitrous oxide emission from urine treated soil as influenced by urine composition and soil physical conditions. *Soil Biology and Biochemistry* **37**, 463–473 (2005).
16. Bakken, L. R., Bergaust, L., Liu, B. B. & Frostegård, A. Regulation of denitrification at the cellular level: a clue to the understanding of N₂O emissions from soils. *Philos. Trans. R. Soc. B-Biol. Sci.* **367**, 1226–1234 (2012).
17. Robinson, A., Di, H. J., Cameron, K. C., Podolyan, A. & He, J. Z. The effect of soil pH and dicyandiamide (DCD) on N₂O emissions and ammonia oxidizer abundance in a stimulated grazed pasture soil. *Journal of Soils and Sediments* **14**, 1434–1444 (2014).
18. Selbie, D. *et al.* The effect of urinary nitrogen loading rate and nitrification inhibitor on nitrous oxide emissions from a temperate grassland soil. *Journal of Agricultural Science* **152**, 159–171 (2014).
19. Sordi, A. *et al.* Nitrous oxide emission factors for urine and dung patches in a subtropical Brazilian pastureland. *Agriculture, Ecosystems and Environment* **190**, 94–103 (2014).
20. de Klein, C. A. M. *et al.* Repeated annual use of the nitrification inhibitor dicyandiamide (DCD) does not alter its effectiveness in reducing N₂O emissions from cow urine. *Animal Feed Science and Technology* **166**, 480–491 (2011).
21. Krol, D. J. Improving and disaggregating N₂O emission factors for ruminant excreta on temperate pasture soils. *Science of the Total Environment* **568**, 327–338 (2016).
22. Saggari, S., Bolan, N. S., Bhandral, R., Hedley, C. B. & Luo, J. A review of emissions of methane, ammonia, and nitrous oxide from animal excreta deposition and farm effluent application in grazed pastures. *New Zealand Journal of Agricultural Research* **47**, 513–544 (2004).
23. Cai, Y., Chang, S. X. & Cheng, Y. Greenhouse gas emissions from excreta patches of grazing animals and their mitigation strategies. *Earth-Science Reviews* **171**, 44–57 (2017).
24. Flechard, C. R. *et al.* Effects of climate and management intensity on nitrous oxide emissions in grassland systems across Europe. *Agriculture, Ecosystems & Environment* **121**, 135–152 (2007).
25. Chirinda, N. *et al.* Emissions of nitrous oxide from arable organic and conventional cropping systems on two soil types. *Agriculture, Ecosystems & Environment* **136**, 199–208 (2010).
26. Clough, T. J., Ledgard, S. F., Sprosen, M. S. & Kear, M. J. Fate of N-15 labelled urine on four soil types. *Plant Soil* **199**, 195–203 (1998).
27. Bell, M. J. *et al.* Nitrous oxide emissions from cattle excreta applied to a Scottish grassland: effects of soil and climate conditions and a nitrification inhibitor. *Science of the Total Environment* **508**, 343–353 (2015).
28. Smith, R. V., Doyle, R. M., Burns, L. C. & Stevens, R. J. A model for nitrite accumulation in soils. *Soil Biology and Biochemistry* **29**, 1241–1247 (1997).

29. Selbie, D. R., Buckthought, L. E. & Shepherd, M. A. The challenge of the urine patch for managing nitrogen in grazed pasture systems. *Advances in Agronomy* **129**, 229–292 (2015).
30. Rochette, P. *et al.* NH₃ volatilization, soil concentration and soil pH following subsurface banding of urea at increasing rates. *Canadian Journal of Soil Science* **93**, 261–268 (2013).
31. Petersen, S. O., Simek, M., Stamatiadis, S., & Yamulki, S. Nitrous oxide emissions from grazed grassland: effects of cattle management and soil conditions. Greenhouse gas emissions from agriculture. Mitigation options and strategies, Institute for Energy and Environment. Leipzig, Germany, pp. 75–78 (2004).
32. Wagner-Riddle, C. & Thurtell, G. Nitrous oxide emissions from agricultural fields during winter and spring thaw as affected by management practices. *Nutrient Cycling in Agroecosystems* **52**, 151–163 (1998).
33. Sehy, U., Ruser, R. & Munch, J. Nitrous oxide fluxes from maize fields: Relationship to yield, site-specific fertilization, and soil conditions. *Agriculture, Ecosystems & Environment* **99**, 97–111 (2003).
34. Johnson, J., Archer, D., Weyers, S. & Barbour, N. Do mitigation strategies reduce global warming potential in the northern U.S. corn belt? *Journal of Environmental Quality* **40**, 1551–1559 (2011).
35. Subbarao, G. V. *et al.* Evidence for biological nitrification inhibition in Brachiaria pastures. *Proc. Natl. Acad. Sci. USA* **106**, 17302 (2009).
36. Meena, H. M., Sachdev, M. S., Manhaiah, K. M. & Dotaniya, M. L. Nitrification Inhibition Potential of Brachiaria humidicola. *National Academy Science Letters* **37**, 113–116 (2014).
37. Byrnes, R. C. *et al.* Biological nitrification inhibition by Brachiaria grasses mitigates soil nitrous oxide emissions from bovine urine patches. *Soil Biology & Biochemistry* **107**, 156–163 (2017).
38. Abalos, D., De Deyn, G. B., Kuyper, T. W. & van Groenigen, J. W. Plant species identity surpasses species richness as a key driver of N₂O emissions from grassland. *Global Change Biology* **20**, 265–275 (2014).
39. Yamulki, S., Jarvis, S. C. & Owen, P. Nitrous oxide emissions from excreta applied in a simulated grazing pattern. *Soil Biology and Biochemistry* **30**, 491–500 (1998).
40. Luo, J., Lindsey, S. B. & Ledgard, S. F. Nitrous oxide emissions from animal urine application on a New Zealand pasture. *Biology and Fertility of Soils* **44**, 463–470 (2008).
41. Van der Weerden, T. J. *et al.* Disaggregating nitrous oxide emission factors for ruminant urine and dung deposited onto pastoral soils. *Agriculture Ecosystems and Environment* **141**, 426–436 (2011).
42. Luo, J. *et al.* Nitrous oxide emissions from grazed hill land in New Zealand. *Agriculture, Ecosystems and Environment* **181**, 58–68 (2013).
43. Pelster, D. E. *et al.* Methane and nitrous oxide emissions from cattle excreta on an East African Grassland. *Journal of Environmental Quality* **45**, 1531–1539 (2016).
44. Tully, K. L., Abwanda, S., Thiong'o, M., Mutuo, P. M. & Rosenstock, T. S. Nitrous oxide and methane fluxes from urine and dung deposited on Kenyan pastures. *Journal of Environmental Quality* **46**, 921–929 (2017).
45. World Resources Institute. Initiative 20 × 20. Washington, DC. Available at, <http://www.wri.org/our-work/project/initiative-20x20/about-initiative-20x20> (2015).
46. United Nations Framework Convention on Climate Change. COP 21 Climate Agreement (UNFCCC, Paris) Available at unfccc.int/resource/docs/2015/cop21/eng/l09r01.pdf (2015).
47. Holmann, F. *et al.* Vale la pena recuperar pasturas degradadas. Una evaluación desde la perspectiva de los productores y extensionistas en Honduras. Tegucigalpa (Honduras): CIAT-DICTA-ILRI. **34** (2004).
48. Brown, D. Methods of surveying and measuring vegetation. *Methods of surveying and measuring vegetation. Commonwealth Agricultural Bureaux* **223** (1954).
49. McCormick, L. H., & Lodge, G. M. A field kit for producers to assess pasture health in the paddock. In Proceedings of the 10th Australian agronomy conference, Hobart. Available at <http://www.regional.org.au/au/asa/2001/3/d/mccormick.htm> (verified 20 May 2018).
50. da Trindade, J. K. *et al.* Daily forage intake by cattle on natural grassland: response to forage allowance and sward structure. *Rangeland Ecology and Management* **69**, 59–67 (2016).
51. Cox, D. R. Some recent work on systematic experimental designs. *Journal of the Royal Statistical Society. Series B (Methodological)* **14**, 211–219 (1952).
52. Hurlbert, S. H. Pseudoreplication and the design of ecological field experiments. *Ecological monographs* **54**, 187–211 (1984).
53. Schank, J. C. & Koehnle, T. J. Pseudoreplication is a Pseudoproblem. *Journal Psychology* **123**, 421–433 (2009).
54. Gee, G. W & Bauder, J. W. Particle-size analysis. p. 383–411. In A. Klute (ed.) *Methods of Soil Analysis, Part 1. Physical and Mineralogical Methods*. Agronomy Monograph No. 9 (2ed). American Society of Agronomy/Soil Science Society of America, Madison, WI (1986).
55. McLean, E. O. Soil pH and lime requirement. In *Methods of soil Analysis. Part 2. Chemical and Microbiological properties* 2nd ed. *Am. Soc. Agron., Inc., Madison, Wisconsin* (1983).
56. Vogel, A. W. Compatibility of soil analytical data: determinations of cation exchange capacity, organic carbon, soil reaction, bulk density, and volume percent of water at selected pF values by different methods. *Working Paper and Preprint 7, ISRIC, Wageningen* (1994).
57. Hoogendoorn, C. J., Betteridge, K., Costall, D. A. & Ledgard, S. F. Nitrogen concentration in the urine of cattle, sheep and deer grazing a common ryegrass/cocksfoot/white clover pasture. *New Zealand Journal of Agricultural Research* **53**, 235–243 (2010).
58. Hutchinson, G. L. & Livingston, G. P. Vents and seals in non-steady-state chambers used for measuring gas exchange between soil and the atmosphere. *European Journal of Soil Science* **52**, 675–682 (2001).
59. Davidson, E. A., Savage, K., Verchot, L. V. & Navarro, R. Minimizing artifacts and biases in chamber-based measurements of soil respiration. *Agricultural and Forest Meteorology* **113**, 21–37 (2002).
60. Dobbie, K. E., McTaggart, I. P. & Smith, K. A. Nitrous oxide emissions from intensive agricultural systems: variations between crops and seasons, key driving variables, and mean emission factors. *Journal of Geophysical Research* **104**, 26891–26899 (1999).
61. Sordi, A. *et al.* Nitrous oxide emission factors for urine and dung patches in subtropical Brazilian pastureland. *Agriculture, Ecosystems and Environment* **190**, 91–103 (2014).
62. SAS Institute Inc. SAS/STAT[®] 9.2 User's Guide. Cary, NC: SAS Institute Inc. 6.11. SAS Institute, Cary, NC (2008).

Acknowledgements

This regional study was done in the framework of the Latin America climate change Mitigation Network (LAMNET) implemented as part of the CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS), which is carried out with support from CGIAR Fund Donors and through bilateral funding agreements. For details please visit <https://ccafs.cgiar.org/donors>. Part of this work was also supported BBSRC through the project BBS/E/C/00010320 awarded to Rothamsted Research. The views and opinions expressed in this document are those of the authors and do not necessarily reflect official positions of the sponsoring organizations. We acknowledge the support that the PhD students, who conducted all the experimental work, received from their supervisors namely: Paulo C. F. Carvalho (Brazil); Cimélio Bayer (Brazil); Alejandro O.

Costantini (Argentina); Rodrigo Jimenez, Nubia Rodriguez and the Taluma AGROSAVIA team (Colombia); and Claudio Machado (Argentina). We would also like to thank Jess Evans from the Applied Statistics team at Rothamsted (UK) for the valuable advice on statistical analyses, Dr. Ana Maria Loboguerrero (CCAFS) for supporting the LAMNET initiative and Jhon Freddy Gutierrez (CIAT) and Nelson Vivas (UniCauca) for providing technical and logistical assistance for the work conducted in Patía (Colombia).

Author Contributions

N.C., S.L., L.A., C.T., M.A., I.R., J.A., M.P., R.B., C.C.J., T.S.R., M.R., D.M. designed the experiments and contributed towards data analyses and manuscript writing-up. V.R., C.F., C.A., J.V.S., R.B., K.Z., L.M. conducted the experiments in the different countries and contributed towards data analyses and writing. L.C. contributed towards data analyses and manuscript writing-up.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-018-37453-2>.

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019