



RESEARCH ARTICLE

Open Access

Evaluation of antioxidant capacity with total phenolic content of *Galanthus krasnovii* (Amaryllidaceae)

Galanthus krasnovii (Amaryllidaceae)'nin toplam fenolik içeriği ile antioksidan kapasitenin değerlendirilmesi

Ramazan ERENLER^{a*}, Nusret GENÇ^a, Mahfuz ELMAŞTAŞ^b, Özgür EMİNAĞAOĞLU^c

^a Plant Research Laboratory, Department of Chemistry, Faculty of Arts and Sciences, Tokat Gaziosmanpaşa University, Tokat, Turkey

^b Department of Biochemistry, Faculty of Pharmacy, University of Health Sciences, Istanbul, Turkey

^c Department of Forest Engineering, Faculty of Forestry, Artvin Coruh University, Artvin, Turkey

Article Info

©2018 Ali Nihat Gökyiğit Botanical Garden Application and Research Center of Artvin Coruh University.

*Corresponding author:

e-mail: ramazan.erenler@gop.edu.tr

ORCID: [0000-0002-0505-3190](https://orcid.org/0000-0002-0505-3190)

Article history

Received: February 13, 2019

Received in revised form: March 12, 2019

Accepted: March 13, 2019

Available online: March 15, 2019



This is an Open Access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Keywords:

Galanthus krasnovii, antioxidant, natural product, phenolic contents.

Anahtar kelimeler:

Galanthus krasnovii, antioksidan, doğal ürün, fenolik içerik.

ABSTRACT

Natural products have gained the great interest due to their broad spectrum of biological activities. *Galanthus krasnovii* was dried at shade then extracted with hexane, dichloromethane, and ethyl acetate successively. After removing of solvent by reduced pressure, crude extracts of each solvent were yielded. Antioxidant activity including 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical, 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS) radical cation and reducing power assays were carried out for each extract. Moreover, total phenolic content was determined. Total phenolic content of hexane-, dichloromethane-, and ethyl acetate extracts were found as 60.95 mg GAE/g extract, 71.90 GAE/g extract and 58.90 GAE/g extract respectively. Dichloromethane and ethyl acetate extract revealed the moderate antioxidant activity.

Öz

Doğal ürünler geniş spektrumlu biyolojik aktivite göstermelerinden dolayı yoğun ilgi görmektedirler. *Galanthus krasnovii* gölgede kurutulduktan sonra sırasıyla hekzan, diklorometan ve etil asetat ile ekstrakte edildi. Düşük basınçta çözücü uzaklaştırıldıktan sonra her bir çözücünün ekstraktı elde edildi. Her bir ekstraktın, 1,1-difenil-2-pikralhidrazil(DPPH) radikal, 2,2'-azino-bis(3- etilbenzotiazolin-6-sülfirik asit) (ABTS) radikal katyon ve indirgeme gücü antioksidan çalışmaları gerçekleştirildi. Ayrıca, ekstraktların toplam fenolik içerikleri belirlendi. Hekzan-, diklorometan- ve etil asetat ekstraktlarının toplam fenolik içerikleri sırasıyla 60.95 mg GAE/g ekstrakt, 71.90 GAE/g ekstrakt ve 58.90 GAE/g ekstrakt olarak belirlendi. Diklorometan ve etil asetat ekstraktları orta derece aktivite gösterdi.

Citation:

To cite this article: Erenler R, Genç N, Elmastaş M, Eminağaoğlu Ö (2019). Evaluation of antioxidant capacity with total phenolic content of *Galanthus krasnovii* (Amaryllidaceae). Turk J Biod 2(1): 13-17.

1. INTRODUCTION

Plants play a significant role in drug discovery process. Since they reveal plenty of pharmaceutical properties, isolation of bioactive compounds from plants gains the great interest to find the most effective compounds with activity mechanism (Aksit et al., 2014; Demirtas et al., 2013; Elmastas et al., 2016; Elmastaş et al., 2015; Erenler, Demirtas, et al., 2017; Erenler, Meral, et al., 2017; Erenler, Sen, Aksit, et al., 2016; Erenler, Sen, Yaglioglu, et al., 2016;

Erenler et al., 2018; Guzel et al., 2017; Karan et al., 2017; Karan & Erenler 2017a; Karan & Erenler 2018; Karan & Erenler 2017b; Karan et al., 2018; Topçu et al., 1999; Yaglioglu et al., 2013; Yildiz et al., 2017).

Galanthus L. genus, known as snowdrops, belonging to Amaryllidaceae family consists of 20 species distributed throughout Europe, Asia and Near East (Zubov & Davis 2012). Six species are endemic for Turkey flora (Davis & Özhatay, 2001). Turkey is one of the centers of species

diversity of *Galanthus*. It is the most traded ornamental bulb genus in the world and a great number of bulbs are exported from Turkey each year. This genus is very popular garden plant in Europe and is especially considered as the early messenger of spring. Some of the *Galanthus* species have been included in the red list of threatened species (Convention on International Trade in Endangered Species of Wild Fauna and Flora – CITES) due to the extensive collection, climate change, and habitat destruction.

Previous phytochemical study on *Galanthus* species provided the isolation of bioactive alkaloids such as tazettine, galanthusine, molycorine, galanthamine, gananthine (Berkov et al., 2012; Kintsurashvili & Vachnadze, 2007). *Galanthus woronowii* Losinsk. collected from Trabzon, Turkey included the galanthine, galanthamine, 2-*O*-(3'-hydroxybutanoyl) lycorine, narwedine, *O*-methyleucotamine, sternbergine, lycorine, sanguinine, salsoline (Sarıkaya et al., 2013). *Galanthus* species have been used in homeopathy to treat various illness such as headaches, cardiac failure, mitral disorder (Bokov & Samylina, 2016). Galanthamine, a main constituent of *Galanthus* species, is a licensed drug for the treatment of dementia in Alzheimer's disease. It inhibits the Acetylcholinesterase (AChE) selectively and competitively (Ellis et al., 2009).

Reactive oxygen species are free radicals yielding in human body during the oxidative function. Human body has defense mechanism against oxidative stress such as enzymes and compounds. Natural antioxidants become inadequate under some circumstance such as malnutrition, ultra violet radiation, and smoking. Hence, excess free radical is able to damage to cell membrane leading to degenerative illness and conditions such as Alzheimer's disease, cardiovascular disease, process, diabetes, inflammation and DNA injury leading to carcinogenesis. Antioxidants play an important role for inhibiting or quenching free radicals. Antioxidants have been used in food, cosmetics and medicine to replace synthetic antioxidants forbidden because of their carcinogenicity (Sasaki et al., 2002). There is not any report about the antioxidant activity of *Galanthus krasnovii* A.P.Khokhr.

In this work, it was presented the antioxidant potency of *Galanthus krasnovii* using DPPH*, ABTS**+, and reducing

power assays. In addition, total phenolic content was executed.

2. MATERIAL AND METHOD

2.1. General experimental procedures

UV analysis was carried out with Hitachi U-290 UV–VIS spectrophotometer. Potassium persulphate ($K_2S_2O_8$), trolox, amonyum acetate ($NH_4CO_2CH_3$), butylated hydroxytoluene (BHT), butylated hydroxyanisole (BHA), 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) diammonium salt (ABTS), 1,1-diphenyl-2-picrylhydrazyl (DPPH) were supplied by Sigma (Sigma-Aldrich GmbH, Steinheim, Germany). The solvents with analytical grade, potassium dihydrogen phosphate (KH_2PO_4), potassium hydroxide (KOH), copper (II) chloride ($CuCl_2$), Folin-Ciocalteu reagent, gallic acid were bought from E. Merck (Darmstadt, Germany).

2.2. Plant material

The voucher specimens have been deposited at the Herbarium of Artvin Coruh University (ARTH), Artvin, Turkey. *G. krasnovii* was collected from Turkey: A8 Artvin, Kafkasör, 1100 m, in 05.04.2018 and identified by Prof. Dr. Özgür Eminağaoğlu, specialists of Plant Taxonomy (ARTH 13437).

2.3. Extraction

G. krasnovii (100 g) was extracted with hexane (2 × 100 mL), dichloromethane (2 × 100 mL) and ethyl acetate (2 × 100 mL) sequentially for 2 days. After removing of solvent from each extract solution under reduced pressure, the crude extracts of hexane (1.5 g), dichloromethane (1.2 g) and ethyl acetate (1.3 g) were yielded. The extracts were kept in fridge to analyze the antioxidant activity with total phenolic content (Genç et al., 2019).

2.4. Total phenolic determination

Total phenolic compounds of *G. krasnovii* extracts (hexane, dichloromethane and ethyl acetate) was executed by Folin-Ciocalteu method (Singleton & Slinkard 1977). The extract solution (1.0 mg/1mL) was reacted with Folin ciocalteu (0.1 mL) in distilled water (4.6 mL). After addition of sodium carbonate solution (0.3 mL, 2%), the reaction mixture was incubated for 2 h at room temperature. The absorbance was determined at 765 nm by a spectrophotometer. The standard curve was calculated using gallic acid and the results were presented

as gallic acid equivalents per mg of extract (Erenler et al., 2014). All experiments were carried out in triplicate.

2.5. DPPH free radical scavenging assay

DPPH[•] activity of hexane-, dichloromethane- and ethyl acetate extracts of *G. krasnovii* was carried out by the protocol reported (Blois 1958). DPPH[•] (1.0 mL, 0.26 mM) was reacted with various concentration of each extract solution (3.0 mL) for 15 min at rt. The absorbance was determined at 517 nm. BHA, BHT and Trolox were used as standard controls. IC₅₀ values indicate the concentration of sample that scavenges 50% of DPPH free radical. The DPPH[•] scavenging activity was calculated using the equation:

$$\text{DPPH}^{\bullet} \text{ scavenging effect (\%)} = [(A_1 - A_2) / A_1] \times 100$$

A₁ is the absorbance of the control and A₂ is the absorbance of the sample (Elmastas et al. 2004).

2.6. ABTS radical cation scavenging assay

ABTS radical cation activity was executed by the protocol reported previously (Re et al. 1999). Initially, ABTS^{•+} stock solution was prepared by the treatment of ABTS (2 mM) with potassium persulfate (2.45 mM). Later, it was kept for 6 h in dark at rt. Each extract solution (3.0 mL) was reacted with ABTS^{•+} solution (1.0 mL). The absorbance was determined at 734 nm by a spectrophotometer. The inhibition was calculated for each concentration in comparison to a blank absorbance (Erenler et al. 2015). ABTS^{•+} activity was calculated by the equation: ABTS^{•+} scavenging effect (%) = [(A₁ - A₂) / A₁] × 100 in which, A₁ is ABTS^{•+} initial concentration and A₂ is ABTS^{•+} remaining concentration in the sample. The results were calculated as IC₅₀.

2.7. Cupric ion reducing power assay

Cupric ion reducing power assay was carried out for each extract of *G. krasnovii* (Elmastas et al., 2018). The reduction capacity of each extract can be calculated by the efficient reduction of Cu⁺² to Cu⁺¹. The yellow complex formed after addition of copper (I) chloride. The extract (40-160 µg/mL, 1.0 mL), CuCl₂ (0.01 M, 1.0 mL), neocuproine (1.0 mL, 7.5 × 10⁻³ M), and acetate tampon (1.0 mL, 1.0 M) were added to the reaction flask and incubated for 30 min. The absorbance was recorded at 450 nm (Apak et al., 2004). The result was expressed according to the Trolox equivalent (µmol/g sample).

3. RESULTS AND DISCUSSION

Galanthus species are significant plants including the bioactive compound used in drug. Some species of *Galanthus* genus were reported to reveal the antioxidant activity. Antioxidant activity of various parts (root, leaf, flower and bulb) of *Galanthus elwesii* Hook.f. was executed at different growing stages such as beginning, flowering, after flowering and fruit ripening. The part of the plant and growing stage altered the activity significantly. The leaf at fruit ripening stage of *G. elwesii* displayed the most antioxidant activity (Ay et al., 2018). *G. transcaucasicus* Fomin shoot extract revealed the considerable antioxidant and antimicrobial activity (Karimi et al., 2018). In this study, hexane-, dichloromethane- and ethyl acetate extracts of *Galanthus krasnovii* were investigated for antioxidant activity as well as total phenolic content. Total phenolic content of hexane, dichloromethane and ethyl acetate extracts was found as 60.95 mg GAE/g extract, 71.90 mg GAE/g extract and 58.90 mg GAE/g extract respectively. Hexane extract did not reveal the DPPH free radical activity. However, dichloromethane extract and EtOAc extract exhibited the moderate activity (Figure 1).




Figure 1. Total phenolic content and DPPH free radical activity

In regarding ABTS radical cation activity, dichloromethane extract and EtOAc extract displayed the mild activity with 14.33 (IC₅₀, µg/mL) and 14.98 (IC₅₀, µg/mL) respectively compared to the standard BHA (8.8, IC₅₀, µg/mL). Reducing power activity was executed with respect to trolox equivalent. Dichloromethane extract (1.15 µmol TE / mg extract) displayed the most reducing power effect among the extracts tested. Ethyl acetate and hexane extracts revealed the activity of 0.77 µmol TE / mg extract and 0.75 µmol TE / mg extract, respectively. This result revealed that the compounds in dichloromethane extract had electron donation ability to reduce from Cu⁺² to Cu⁺¹.




Figure 2. ABTS Radical cation assay and reducing power

4. CONCLUSIONS

G. krasnovii has a potency to be used in pharmaceutical and food industry as a natural antioxidant. The cultivation area of *G. krasnovii* should be enlarged. Bioactive compounds should be isolated and alkaloids quantity of *G. krasnovii* should be increased by using various agricultural techniques. Antioxidant effects of isolated compounds should be investigated and activity-structure mechanism should be presented.

REFERENCES

- Aksit H, Çelik SM, Sen Ö, Erenler R, Demirtas I, Telci I, Elmastas M (2014). Complete isolation and characterization of polar portion of *Mentha dumetorum* water extract. *Records of Natural Products* 8:277-280.
- Apak R, Güçlü K, Özyürek M, Karademir SE (2004). Novel total antioxidant capacity index for dietary polyphenols and vitamins C and E, using their cupric ion reducing capability in the presence of neocuproine: CUPRAC method. *Journal of Agricultural and Food Chemistry* 52: 7970-7981.
- Ay EB, Gül M, Açıkgöz MA, Yarılgöz T, Kara ŞM (2018). Assessment of antioxidant activity of giant snowdrop (*Galanthus elwesii* Hook) extracts with their total phenol and flavonoid contents. *Indian Journal of Pharmaceutical Education and Research* 52: 128-132.
- Berkov S, Codina C, Bastida J (2012). The genus *Galanthus*: A source of bioactive compounds. In: *Phytochemicals-A Global Perspective of Their Role in Nutrition and Health*. IntechOpen, 235-254.
- Blois MS (1958). Antioxidant determinations by the use of a stable free radical. *Nature* 181: 1199-1200.
- Bokov D, Samylin I (2016). Identification of Flavonoids in Homeopathic Mother Tinctures of *Galanthus woronowii* Losinsk. And *Galanthus nivalis* L. by Ultra-Performance Liquid Chromatography with Photodiode Arrays and Tandem Quadrupole Mass-Selective Detectors. *Pharmaceutical Chemistry Journal* 50: 458-464.
- Davis AP, Özhatay N (2001). *Galanthus trojanus*: a new species of *Galanthus* (Amaryllidaceae) from north-western Turkey. *Botanical Journal of the Linnean Society* 137: 409-412.
- Demirtas I, Erenler R, Elmastas M, Goktasoglu A (2013). Studies on the antioxidant potential of flavones of *Allium vineale* isolated from its water-soluble fraction. *Food Chemistry* 136: 34-40.
- Ellis J, Nathan P, Villemagne V, Mulligan R, Saunderson T, Young K, Smith C, Welch J, Woodward M, Wesnes K (2009). Galantamine-induced improvements in cognitive function are not related to alterations in $\alpha_4\beta_2$ nicotinic receptors in early Alzheimer's disease as measured in vivo by 2-[¹⁸F] Fluoro-A-85380 PET. *Psychopharmacology* 202: 79-91.
- Elmastas M, Celik SM, Genc N, Aksit H, Erenler R, Gulcin İ (2018). Antioxidant activity of an Anatolian herbal tea—*Origanum minutiflorum*: isolation and characterization of its secondary metabolites. *International Journal of Food Properties* 21: 374-384.
- Elmastas M, Erenler R, Isnac B, Aksit H, Sen O, Genc N, Demirtas I (2016). Isolation and identification of a new neo-clerodane diterpenoid from *Teucrium chamaedrys* L. *Natural Product Research* 30: 299-304.
- Elmastas M, Ozturk L, Gokce I, Erenler R, Aboul-Enein HY (2004). Determination of antioxidant activity of marshmallow flower (*Althaea officinalis* L.). *Analytical Letters* 37: 1859-1869.
- Elmastaş M, Telci İ, Akşit H, Erenler R (2015). Comparison of total phenolic contents and antioxidant capacities in mint genotypes used as spices/Baharat olarak kullanılan nane genotiplerinin toplam fenolik içerikleri ve antioksidan kapasitelerinin karşılaştırılması. *Turkish Journal of Biochemistry* 40: 456-462.
- Erenler R, Demirtas I, Karan T, Altun M, Gul F (2017). Inhibitory Effect of 6, 7-dimethoxy-5-hydroxyflavone on Human Cervix Carcinoma in Vitro. *International Journal of Secondary Metabolite* 4: 512-516.
- Erenler R, Meral B, Sen O, Elmastas M, Aydin A, Eminagaoglu O, Topcu G (2017). Bioassay-guided isolation, identification of compounds from *Origanum rotundifolium* and investigation of their

- antiproliferative and antioxidant activities. *Pharmaceutical Biology* 55: 1646-1653.
- Erenler R, Sen O, Aksit H, Demirtas I, Yaglioglu AS, Elmastas M, Telci İ (2016). Isolation and identification of chemical constituents from *Origanum majorana* and investigation of antiproliferative and antioxidant activities. *Journal of the Science of Food and Agriculture* 96: 822-836.
- Erenler R, Sen O, Yaglioglu AS, Demirtas I (2016). Bioactivity-guided isolation of antiproliferative sesquiterpene lactones from *Centaurea solstitialis* L. ssp. *solstitialis*. *Combinatorial Chemistry & High Throughput Screening* 19: 66-72.
- Erenler R, Telci İ, Elmastaş M, Akşit H, Gül F, Tüfekçi AR, Demirtaş İ, Kayir Ö (2018). Quantification of flavonoids isolated from *Mentha spicata* in selected clones of Turkish mint landraces. *Turkish Journal of Chemistry* 42: 1695-1705.
- Erenler R, Telci I, Ulutas M, Demirtas I, Gul F, Elmastas M, Kayir O (2015). Chemical Constituents, Quantitative Analysis and Antioxidant Activities of *Echinacea purpurea* (L.) Moench and *Echinacea pallida* (Nutt.) Nutt. *Journal of Food Biochemistry* 39: 622-630.
- Erenler R, Yilmaz S, Aksit H, Sen O, Genc N, Elmastas M, Demirtas I (2014). Antioxidant activities of chemical constituents isolated from *Echinops orientalis* Trauv. *Records of Natural Products* 8(1): 32-36.
- Genç N, Yıldız İ, Karan T, Eminağaoğlu Ö, Erenler R (2019). Antioxidant activity and total phenolic contents of *Galanthus woronowii* (Amaryllidaceae). *Turkish Journal of Biodiversity* 2(1): 1-5.
- Guzel A, Aksit H, Elmastas M, Erenler R (2017). Bioassay-guided isolation and identification of antioxidant flavonoids from *Cyclotrichium origanifolium* (Labill.) Manden. and Scheng. *Pharmacognosy Magazine* 13(50): 316-320.
- Karan T, Altuner Z, Erenler R (2017). Growth and Metabolite Production of *Chroococcus minutus* Under Different Temperature and Light Conditions. *Journal of New Results in Science* 6:47-52.
- Karan T, Erenler R (2017a). Effect of Salt and pH Stress of Bioactive Metabolite Production in *Geitlerinema carotinosum*. *International Journal of Secondary Metabolite* 4: 16-19.
- Karan T, Erenler R (2018). Fatty acid constituents and anticancer activity of *Cladophora fracta* (OF Müller ex Vahl) Kützing. *Tropical Journal of Pharmaceutical Research* 17: 1977-1982.
- Karan T, Erenler R (2017b). Screening of norharmine from seven cyanobacteria by high-performance liquid chromatography. *Pharmacognosy Magazine* 13: 723-725.
- Karan T, Yildiz I, Aydin A, Erenler R (2018). Inhibition of various cancer cells proliferation of bornyl acetate and essential oil from *Inula graveolens* (Linnaeus) Desf. *Records of Natural Products* 12: 273-283.
- Karimi E, Mehrabanjoubani P, Homayouni-Tabrizi M, Abdolzadeh A, Soltani M (2018). Phytochemical evaluation, antioxidant properties and antibacterial activity of Iranian medicinal herb *Galanthus transcaucasicus* Fomin. *Journal of Food Measurement and Characterization* 12: 433-440.
- Kintsurashvili L, Vachnadze V (2007). Plants of the Amaryllidaceae family grown and introduced in Georgia: a source of galanthamine. *Pharmaceutical Chemistry Journal* 41: 492-494.
- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C (1999). Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine* 26: 1231-1237.
- Sarikaya BB, Kaya GI, Onur MA, Bastida J, Somer NU (2013). Phytochemical investigation of *Galanthus woronowii*. *Biochemical Systematics and Ecology* 51: 276-279.
- Sasaki YF, Kawaguchi S, Kamaya A, Ohshita M, Kabasawa K, Iwama K, Taniguchi K, Tsuda S (2002). The comet assay with 8 mouse organs: results with 39 currently used food additives. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 519: 103-119.
- Singleton VL, Slinkard K (1977). Total phenol analysis: Automation and comparison with manual methods. *American Journal of Enology and Viticulture* 28: 49-55.
- Topçu G, Erenler R, Çakmak O, Johansson CB, Çelik C, Chai H-B, Pezzuto JM (1999). Diterpenes from the berries of *Juniperus excelsa*. *Phytochemistry* 50: 1195-1199.
- Yaglioglu AS, Akdulum B, Erenler R, Demirtas I, Telci I, Tekin S (2013). Antiproliferative activity of pentadeca-(8E, 13Z) dien-11-yn-2-one and (E)-1, 8-pentadecadiene from *Echinacea pallida* (Nutt.) Nutt. roots. *Medicinal Chemistry Research* 22: 2946-2953.
- Yildiz I, Sen O, Erenler R, Demirtas I, Behcet L (2017). Bioactivity-guided isolation of flavonoids from *Cynanchum acutum* L. subsp. *sibiricum* (willd.) Rech. f. and investigation of their antiproliferative activity. *Natural Product Research* 31: 2629-2633.
- Zubov DA, Davis AP (2012). *Galanthus panjutinii* sp. nov.: A new name for an invalidly published species of *Galanthus* (Amaryllidaceae) from the northern Colchis area of Western Transcaucasia. *Phytotaxa* 50: 55-63.