

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

Instructions:

- (1) Please complete two of these forms, by providing title details and striking out one option from each of the sections 1-3.
- (2) If you select option b of any section you must include a specific time period. The maximum permitted period is 24 months.
- (3) Include the two forms with the copies of your thesis submitted to your supervisor.
- (4) The Library must receive the original of your thesis.
- (5) We strongly recommend that you provide the Library with two copies of your thesis if it has colour plates.

Massey University Library. Thesis Copyright Form

Title of thesis: Development of Methods for the Determination of Bismuth & Thallium in Geological Materials & Their Significance for the Cretaceous - Tertiary Boundary Event

- (1) (a) I give permission for my thesis to be made available to readers in the Massey University Library under conditions determined by the Librarian.
- ~~(b) I do not wish my thesis to be made available to readers without my written consent for \_\_\_\_\_ months.~~
- (2) (a) I agree that my thesis, or a copy, may be sent to another institution under conditions determined by the Librarian.
- ~~(b) I do not wish my thesis, or a copy, to be sent to another institution without my written consent for \_\_\_\_\_ months.~~
- (3) (a) I agree that my thesis may be copied for Library use.
- ~~(b) I do not wish my thesis to be copied for Library use for \_\_\_\_\_ months.~~

Signed *Malcolm [Signature]*

Date 4 November, 1988

The copyright of this thesis belongs to the author. Readers must sign their name in the space below to show that they recognise this. They are asked to add their permanent address.

NAME AND ADDRESS

DATE

\_\_\_\_\_

DEVELOPMENT OF METHODS FOR THE  
DETERMINATION OF BISMUTH AND THALLIUM  
IN GEOLOGICAL MATERIALS  
AND THEIR SIGNIFICANCE FOR  
THE CRETACEOUS-TERTIARY  
BOUNDARY EVENT

A thesis presented in partial fulfilment  
for the degree of

MASTER OF SCIENCE

Department of Chemistry and Biochemistry  
Massey University  
New Zealand

Maki Hoashi

1988

## ACKNOWLEDGEMENTS

I am indebted to the following persons:

My supervisor, Professor R.R. Brooks, for his cheerful encouragement, unfailing assistance, and sense of humor from this project's conception to its maturity;

Dr. C.J. Orth for allowing me to use his INAA data for iridium and for chalcophile elements in the Flaxbourne River K-T boundary samples;

R.B. Gardner for his great patience, support and understanding throughout my studies;

Finally, to J. & T. Hoashi, to whom I dedicate my research and thesis. Without them, I would have accomplished nothing.

## ABSTRACT

Solvent extraction methods have been developed for the determination of thallium and bismuth in rocks. The method for thallium is based on the extraction of the chlorocomplex into methylisobutyl ketone followed by removal of the interfering chloride ion as insoluble silver chloride. The method for bismuth is based on the extraction of a stable metal complex with ammonium pyrrolidine dithiocarbamate into methylisobutyl ketone. Both elements are analysed on a electrothermal graphite furnace atomic absorption spectrometer. Tests on the efficiencies of the methods showed a good precision and accuracy with limits of detection of 9.5 ng/ml (ppb) for thallium and 20 ng/ml (ppb) for bismuth. Recovery studies on synthetic samples showed recoveries of >99.9% for both elements.

A geochemical study was performed on a Cretaceous-Tertiary boundary site located near Flaxbourne River, Ward, New Zealand. Investigations were carried out to determine the enrichment of thallium and bismuth in the stratigraphic column. The results show a good correlation between the iridium and thallium content throughout the rock sequence, but the samples have very low bismuth contents. This is indicative of the different geochemical behavior of bismuth to other chalcophiles. These findings do not contradict the impact theory initially hypothesized by Alvarez et al. (1980), and instead suggest that bismuth's mobility and solubility are far greater than those of thallium and the other chalcophile elements enriched in the Flaxbourne River sequence. This results in the leaching of bismuth from the stratigraphic column by sea water after deposition.

## TABLE OF CONTENTS

Acknowledgments.....	i
Abstract.....	1
Table of Contents.....	ii
List of Figures.....	v
List of Tables.....	vii
List of Appendices.....	viii
I. General Introduction.....	2

## PART ONE

II. Development of Analytical Methods	
II.1. Introduction.....	7
2. List of Reagents.....	8
3. Atomic Absorption Spectrometry	
3.1. Introduction.....	10
2. Flame Spectrometry.....	11
3. Flame-AAS Instrumentation.....	11
4. Graphite Furnace Spectrometry.....	12
5. GF-AAS Instrumentation.....	13
6. Investigation of Parameters for GF-AAS.....	14
6.a. Solubility of Phases and Acid Concentration.....	14
b. Lamp Current.....	15

c. Atomization Temperature.....	15
d. Wavelength.....	15
e. Ramp Times.....	16
f. Background Correction.....	18
g. The Furnace Tube.....	19
h. The Sampling Vial.....	19
i. Digestion of Rock Samples.....	20
II.4. Solvent Extraction of Thallium and Bismuth	
4.1. Introduction.....	22
2. Calibration Curves.....	23
3. The Ammonium Pyrrolidine Dithiocarbamate System	
3.a. Introduction.....	24
b. Experimental.....	27
c. Results and Discussion.....	28
d. Determination of Bismuth in a Standard Rock.....	31
e. Limit of Detection, Precision and Accuracy.....	34
f. Recommended Procedure for the Determination of Bismuth in Rocks.....	34
II.4.4. The Determination of the Thallium Chlorocomplex	
4.a. Introduction.....	36
b. Overcoming Chloride Interference.....	37
c. Determination of Thallium in Reference Rocks.....	45
d. Limit of Detection, Precision and Accuracy.....	46
e. Recommended Procedure for the Determination of Thallium in Rocks.....	47
II.5. Summary of Techniques.....	48

PART TWO

III. Flaxbourne River K-T Boundary: A Case Study

III.1. Introduction.....	50
2. Site Location.....	52
3. Lithology.....	53
4. Biostratigraphy.....	56
5. Geochemistry.....	58
6. Sampling Method.....	60
7. Results.....	60
8. Discussion.....	63
9. Conclusion.....	67
Summary and General Conclusions.....	69
References.....	71
Appendix.....	83



## LIST OF FIGURES

1) Solubility of MIBK in varying HCl concentrations.....	14
2) Determination of optimum atomization temperatures for thallium and bismuth.....	16
3) Temperature profile of sampler program for bismuth.....	18
4) Temperature profile of sampler program for thallium.....	18
5) Calibration curve for bismuth.....	23
6) Calibration curve for thallium.....	23
7) Effect of increasing amounts of APDC on bismuth extraction into MIBK.....	26
8) Extraction of 9 elements in different concentrations of acids with and without addition of APDC.....	28
9) Extraction and $K_d$ values for bismuth with and without APDC.....	30
10) Bismuth and iron extractions in HCl and HNO <sub>3</sub> with APDC.....	32
11) Stability with time of APDC complexes with bismuth and antimony.....	33
12) Extraction and maximum $K_d$ values for Tl(I) and Tl(III) with and without APDC in HCl and HNO <sub>3</sub> .....	37
13) Determination of the optimum amount of AgNO <sub>3</sub> to add to a known amount of thallium for Tl quantification by GF-AAS.....	40
14) Evaporation rates of MIBK and water.....	42
15) Plot of absorbances of thallium and "silver cloud" peaks.....	44

16) Effect of varying atomization temperatures on the  
relative absorbances of thallium and silver.....45

17) Flaxbourne River and nearby K-T boundary site locations..52

18) Lithologic section of the Flaxbourne River K-T  
boundary sequence.....55

19) Results of thallium and bismuth determinations by  
GF-AAS in Flaxbourne River K-T boundary samples with  
iridium, arsenic, antimony, gallium and indium.....61

20) Correlation of thallium versus iridium abundances in  
the Flaxbourne River K-T boundary samples.....62

## LIST OF TABLES

1) Geochemical classification of the elements in relation to the periodic system.....	4
2) Furnace sampler program for bismuth.....	17
3) Furnace sampler program for thallium.....	17
4) Distribution coefficient ( $K_d$ ) for extraction of selected chalcophile ions in 2 M acid with and without APDC.....	29
5) Published values for bismuth concentration in GXR-1.....	31
6) Statistical information for the proposed method of extraction for bismuth.....	34
7) The effect of different halide ions on the absorbance signal of 0.05 ug/ml thallium.....	38
8) Absorption signals for Tl(I) and Tl(III) in the presence and absence of chloride ions.....	39
9) Statistical information for the proposed method of extraction for thallium.....	47
10) Standard K-T boundary event units and their equivalents at Flaxbourne River.....	51
11) GF-AAS determinations of thallium and bismuth for selected intervals in the Flaxbourne River K-T boundary sequence.....	62
12) Enrichment of selected chalcophile elements in coal ash.....	64

## LIST OF APPENDICES

- 1) Concentration of chalcophile elements in the  
Flaxbourne River K-T boundary sequence determined  
by INAA and GF-AAS.....83
- 2) Description of sampling vials considered for  
use with MIBK.....86
- 3) Summary of attempts in the literature to overcome  
the interference of halide ions in GF-AAS  
determinations of thallium.....87
- 4) Summary of procedures and results of colorimetric  
extraction of thallium.....88
- 5) Tl versus Ir Pearson Product Moment Correlation  
calculations for 11 intervals of the Flaxbourne  
River K-T boundary sequence.....89
- 6) Significance of correlation values.....90
- 7) Abundance of selected chalcophiles in 13  
geospheres in ppm ( $\mu\text{g/g}$  or  $\mu\text{g/ml}$ ).....91

## I. GENERAL INTRODUCTION

Very early in the history of modern geology it became convenient to divide regional lithologic sequences into large stratigraphic units that could be distinguished from one another on the basis of fossil content (Raup and Stanley, 1978). There was a natural tendency for early workers to establish system boundaries at major floral and faunal breaks. Many of these boundaries seem especially well-placed, for some of the chosen breaks represent distinct evolutionary changes or extinctions that occurred during short time intervals on a worldwide scale. Explanations of these often dramatic changes in the fossil record have not been totally satisfactory, especially since a global scale and a short, limited time must be considered.

In their benchmark paper, Alvarez et al. (1980) suggested that the extinctions at the terminal Cretaceous were due to catastrophic events caused by the impact of a large (>10 km diameter) asteroid with the earth. This scenario developed from evidence of a significant iridium enrichment in the Cretaceous-Tertiary (K-T) boundary clay in three locations: Gubbio in Italy, Stevns Klint in Denmark, and Woodside Creek in New Zealand. The Alvarez group suggested that the anomalously high iridium concentrations arose from the presence of particulate remains of a large impacting bolide in the boundary sediment. Since then, iridium and other siderophile element anomalies have been found in over 90 K-T boundary sediments worldwide, including several sites in New Zealand (Kyte et al., 1980; Brooks et al., 1986).

Several extinction mechanisms have been suggested as side effects of the initial impact (Alvarez, 1986). These include:

1) Ignition of wildfires and the generation of nitric acid rains by the heat of the meteorite's entry and impact;

2) Meteoritic and terrestrial dust resulting from the impact being thrown into the atmosphere, blocking sunlight and suppressing photosynthesis, resulting in extreme cold and the destruction of food chains;

3) Ejected water vapor remaining in the atmosphere after the impact dust had settled, resulting in a greenhouse effect to follow the extreme cold.

Although some of the elements enriched in the K-T boundary clay are apparently meteoritic (Ir, Ni), others are clearly terrestrial (As, Sb). The latter group are noted to be mainly chalcophile, or sulphur-loving, elements (Strong et al., 1987). Absolute concentrations of all chalcophiles are much higher than would be provided directly from a mantle, crustal or meteoritic source; therefore, an enrichment process must be invoked at the boundary.

Anoxic conditions following the K-T boundary impact would readily cause scavenging of chalcophiles from the water column (Brooks et al., 1984; Kyte et al., 1980). Researchers have noted that chalcophiles can be similarly enriched in coal ash, caused by the low oxidation potential of the medium (Mason and Moore, 1982). Efficient sweep-out of the ejecta would be able to concentrate most of the dissolved chalcophiles into the thin clay boundary layer (Strong et al., 1987).

Thallium and bismuth were chosen as representative and as yet unstudied chalcophile elements regarding the K-T boundary event. All of the chalcophiles in Table 1 have been analysed by instrumental neutron activation analysis with the exceptions of thallium, bismuth, tellurium, cadmium, sulfur and lead for the Flaxbourne River K-T

H		Atmophile: <u>N</u>															<u>He</u>
Li	Be	Lithophile: <u>Na</u>										B	C	<u>N</u>	O	F	<u>Ne</u>
Na	Mg	Chalcophile: <u>Zn</u>										Al	Si	P	<u>S</u>	Cl	<u>A</u>
		Siderophile: <u>Fe</u>															
K	Ca	Sc	Ti	V	Cr	Mn	<u>Fe</u>	<u>Co</u>	<u>Ni</u>	<u>Cu</u>	<u>Zn</u>	<u>Ga</u>	<u>Ge</u>	<u>As</u>	<u>Se</u>	Br	<u>Kr</u>
Rb	Sr	Y	Y	Nb	<u>Mo</u>		<u>Ru</u>	<u>Rh</u>	<u>Pd</u>	<u>Ag</u>	<u>Cd</u>	<u>In</u>	<u>Sn</u>	<u>Sb</u>	<u>Te</u>	I	<u>Xe</u>
Cs	Ba	La-Lu	Hf	Ta	<u>W</u>	<u>Re</u>	<u>Os</u>	<u>Ir</u>	<u>Pt</u>	<u>Au</u>	<u>Hg</u>	<u>Tl</u>	<u>Pb</u>	<u>Bi</u>			
			Th		U												

TABLE 1: Geochemical classification of the elements in relation to the periodic system (Mason and Moore, 1982)

boundary samples (Strong et al., 1987; Orth, 1987, Appendix 1). Difficulties have been noted by previous investigators in the analysis of thallium by graphite furnace atomic absorption spectrometry in the presence of halide ions. Bismuth was chosen from the other five unanalysed chalcophiles to discover if it behaves like the other group VA metals (As, Sb) in the K-T boundary. Analysis of bismuth by hydride generation of bismuthine is possible (Thompson and Thomerson, 1974) but for this study, graphite furnace atomic absorption spectrometry was utilised.

The following project is divided into two main parts:

1) The development of solvent extraction methods to separate thallium and bismuth from rock matrices and their determination by graphite furnace atomic absorption spectrometry;

2) An application of the methods to determine thallium and bismuth levels for samples collected at a K-T boundary site at Flaxbourne River, New Zealand.



It is hoped that this work will provide new, rapid, inexpensive and sensitive methods for the determination of thallium and bismuth in geochemical materials and that a contribution can be made towards clarifying the series of events that destroyed 70% of life forms at the end of the Cretaceous period.

## II. DEVELOPMENT OF ANALYTICAL METHODS