

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

Massey University Library
Thesis Copyright Form

Title of thesis: An Immobilized Cell Bioreactor
For The Malolactic Fermentation
of Wine.

- (1) (a) I give permission for my thesis to be ~~made available~~ to readers in Massey University Library under conditions ~~determined~~ by the Librarian.
- (b) I do not wish my thesis to be made available to readers without my written consent for 24 months.
- (2) (a) I agree that my thesis, or a copy, ~~may be sent to another institution under conditions determined by the Librarian.~~
- (b) I do not wish my thesis, or a copy, to be sent to another institution without my written consent for 24 months.
- (3) (a) ~~I agree that my thesis may be copied for Library use.~~
- (b) I do not wish my thesis to be copied for Library use for ... months.

Signed Deise Jarassé

Date 16/1/92

The copyright of this thesis belongs to the author. Readers must sign their name in the space below to show that they recognise this. They are asked to add their permanent address.

NAME AND ADDRESS

DATE

**AN IMMOBILIZED CELL BIOREACTOR
FOR THE MALOLACTIC FERMENTATION OF WINE**

A THESIS PRESENTED IN PARTIAL FULFILMENT OF
THE REQUIREMENTS FOR THE DEGREE OF
MASTER OF TECHNOLOGY IN BIOTECHNOLOGY AT
MASSEY UNIVERSITY

DENISE E. JANSSEN

1991

ABSTRACT

Malolactic fermentation using immobilized cells of *Leuconostoc oenos* was investigated in order to improve this fermentation at an industrial-scale.

Three strains of bacteria were investigated in some detail, and one was chosen for further work. A satisfactory growth medium for the strain of bacteria used was found to be an apple juice broth. The effect, on both the growth and malic acid bioconversion for *Leuconostoc oenos* strain 1070, of having 6% v/v ethanol in the growth media was tested and found to cause a longer lag phase, and be slightly beneficial, respectively.

Oak chips were decided on as the immobilization media, in preference to bone char, and a synthetic, apple-juice based wine was used to determine operation parameters for a continuous culture bioreactor.

Temperature, pH, ethanol concentration, SO₂, malic acid concentrations, anaerobic conditions and dilution rate were investigated and it was shown that lower malic acid concentrations, and also an interaction between low pH, high temperature and high ethanol concentration affected the malic acid bioconversion adversely. Increasing the dilution rate above 0.35 h⁻¹ caused a 30% drop in the bioconversion rate. The pH level had no effect on bioconversion if the temperature was kept at 21°C or lower. Decreasing the temperature, increasing the ethanol concentration above 10% v/v and increasing SO₂ levels all caused a slight drop in bioconversion rates while strict anaerobic growth and bioconversion conditions caused an increase. The bioconversion rates ranged between 20 and 100 mg malic acid consumed/100ml oak chips/hour.

An industrial prototype bioreactor was built and used at Villa Maria Wineries, Auckland, during the 1991 vintage and successfully processed 200 litres of Chardonnay-style wine in 2 days. The bioconversion rate was between 25 and 30 mg malic acid consumed/100ml oak chips/hour. Informal taste tests showed satisfactory malolactic characteristics in the treated wine.

TABLE OF CONTENTS

ABSTRACT	ii
TABLE OF CONTENTS	iii
LIST OF TABLES	vii
LIST OF FIGURES	viii
CHAPTERS:	
1 INTRODUCTION	1
1.1 ORGANISM	3
1.1.1 Morphology	4
1.1.2 Malolactic Fermentation	4
1.1.2.1 Effect of pH	6
1.1.2.2 Effect of Temperature	6
1.1.2.3 Effect of SO ₂	7
1.1.2.4 Effect of Ethanol	7
1.1.2.5 Effect of Growth Media on Cell Growth	8
1.1.2.6 Inoculation into Wine	8
1.2 IMMOBILIZATION METHODS	
1.2.1 Attachment	9
1.2.2 Entrapment	11
1.2.3 Flocculation	11
1.2.4 Containment	12
1.3 APPLICATION OF CELL IMMOBILIZATION TO MLF	
1.3.1 Methods used for Immobilization	12
1.3.2 Effect of Immobilization on Bioconversion	13
1.4 EXPERIMENTAL OUTLINE	16
2 METHODS AND MATERIALS	
2.1 ORGANISMS	18
2.2 IMMOBILIZATION MEDIA	18
2.3 GROWTH AND BIOCONVERSION MEDIA	

2.3.1	Tomato Juice Broth (TJ Broth)	19
2.3.2	Apple Juice Broth (AJ Broth)	20
2.3.3	Apple Juice Wine (AJ Wine)	20
2.3.4	Yeast Nitrogen Base Wine (YNB Wine)	21
2.4	ANALYTICAL METHODS	
2.4.1	Growth	21
2.4.2	Biomass	21
2.4.3	L-Malic Acid	22
2.4.4	Malolactic Activity	22
2.4.5	Total Reducing Sugar	23
2.4.6	Plate Counts	23
2.4.7	Ethanol	24
2.5	IMMOBILIZED CELL REACTOR	
2.5.1	Equipment	24
2.5.2	Retention Times	26
2.5.3	Sterilization of Reactors	26
2.5.4	Inoculation and Cell Growth	26
2.5.5	Bioconversion Calculations	27
3	PRELIMINARY SCREENING EXPERIMENTS	
3.1	INTRODUCTION	29
3.2	COMPARISON OF DL- AND L- MALIC ACID	
3.2.1	Objective	29
3.2.2	Experimental Conditions	29
3.2.3	Results and Conclusions	30
3.3	SELECTION OF BACTERIAL STRAIN AND GROWTH MEDIUM	
3.3.1	Objectives	33
3.3.2	Experimental Conditions	33
3.3.3	Results and Discussion	34
3.3.4	Conclusions	40
3.4	COMPARISON OF IMMOBILIZATION MEDIA	
3.4.1	Objectives	41

3.4.2	Experimental Conditions	41
3.4.3	Results and Discussion	42
3.4.4	Conclusions	45
3.5	COMPARISON OF REAL WINE AND SYNTHETIC WINE	
3.5.1	Objectives	45
3.5.2	Experimental Conditions	46
3.5.3	Results and Discussion	47
3.5.4	Conclusions	49
3.6	OVERALL CONCLUSIONS	50
4	BIOCONVERSION PARAMETERS	
4.1	ACTIVITY OF FREE CELLS IN BATCH CULTURE USING AJ WINE	
4.1.1	Objectives	51
4.1.2	Experimental Conditions	51
4.1.3	Results and Discussion	51
4.2	COMPARISON OF CHARDONNAY AND AJ WINES, AND EFFECT OF RECYCLING FEED	
4.2.1	Objectives	53
4.2.2	Experimental Conditions	53
4.2.3	Results	53
4.3	COMPARISON OF NON-ANAEROBIC & ANAEROBIC CONDITIONS	
4.3.1	Objectives	56
4.3.2	Experimental Conditions	56
4.3.3	Results	56
4.4	EFFECTS OF TEMPERATURE AND pH	
4.4.1	Objectives	58
4.4.2	Experimental Conditions	58
4.4.3	Results	59
	4.4.3.1 pH	59
	4.4.3.2 Temperature	59
	4.4.3.3 Time	61
4.4.4	Conclusions	61

4.5	EFFECTS OF pH AND ETHANOL	
4.5.1	Objectives	61
4.5.2	Experimental Conditions	62
4.5.3	Results	62
4.6	EFFECT OF MALIC ACID CONCENTRATION AND WINE FLOWRATE:	
	EXPERIMENT 1	
4.6.1	Objectives	64
4.6.2	Experimental Conditions	65
4.6.3	Results	65
4.7	EFFECT OF MALIC ACID CONCENTRATION AND WINE FLOWRATE:	
	EXPERIMENT 2	
4.7.1	Objectives	68
4.7.2	Experimental Conditions	68
4.7.3	Results	68
4.8	OVERALL DISCUSSION	70
4.9	CONCLUSIONS	73
5	INDUSTRIAL PROTOTYPE TRIAL	
5.1	OBJECTIVES	74
5.2	METHODS	74
5.3	RESULTS	77
6	GENERAL DISCUSSION	
6.1	GROWTH	80
6.2	PROTOTYPE TRIAL	81
7	REFERENCES	83
	ACKNOWLEDGEMENTS	89

LIST OF TABLES

Table

- 1 Specific activity of malolactic organisms as given in the literature
- 2 Conditions for initial growth of cells in bioreactors
- 3 Tomato juice/apple juice comparison
- 4 Bioconversion activity at different pH and ethanol concentrations
- 5 Wine feed conditions: Malic/flowrate comparison 1
- 6 Relationship between cell counts and bioconversion rates
- 7 Pilot-scale trial using prototype bioreactor

LIST OF FIGURES

Figure

- 1 Immobilized cell bioreactors
- 2 Growth of *L. oenos* strain 1070 in DL- and L-malic acid broth
- 3 L-malic acid bioconversion by *L. oenos* in DL- and L-malic acid broth
- 4 Growth of *L. oenos* strain 1011 in various media
- 5 Growth of *L. oenos* strain 1041 in various media
- 6 Growth of *L. oenos* strain 1070 in various media
- 7 Comparison of immobilization media
- 8 Transmission electron micrograph of oak chip cross-section
- 9 Scanning electron micrograph of oak chip
- 10 Comparison of real and synthetic wines
- 11 Comparison of Chardonnay and AJ wines, and effect of recycling feed
- 12 Comparison of non-anaerobic and anaerobic conditions
- 13 Effect of temperature and pH on bioconversion
- 14 Effect of pH and ethanol on bioconversion
- 15 Effect of malic acid concentration and flowrate on bioconversion: Experiment 1
- 16 Effect of malic acid concentration and flowrate on bioconversion: Experiment 2
- 17 Prototype bioreactor
- 18 Pilot-scale trial using prototype bioreactor

1.0 INTRODUCTION

Malolactic fermentation (MLF) entails the conversion of L-malic acid to L-lactic acid via decarboxylation using a suitable microorganism. The primary uses of MLF are to lower the acidity in high-acid wine, to give wine enhanced sensory characteristics, especially diacetyl or "buttery" notes, and to improve microbiological stability (Eggenberger, 1988).

Currently, in cool climate wine areas, this fermentation is used for most red wines and some white wines, especially Chardonnays. Normal industrial practice is to use batch fermentation, either initiated naturally by bacteria occurring in the winery or following inoculation with commercially available preparations. Length of fermentation varies greatly with wine type and temperature, and may vary from one week for some red wines to up to one year for a Chardonnay-style wine. Delayed onset of fermentation is seen by winemakers as the primary problem with malolactic fermentation in New Zealand (Pilone, 1988).

Immobilized cells offer several advantages for MLF of wine:

- faster fermentation
- continuous operation
- improved temperature control
- greater tolerance to wines which have high alcohol and sulphur dioxide concentrations and low pH
- better control over the timing and extent of deacidification
- absence of flavour effects caused by bacterial growth in the wine.

Potential disadvantages of the technique include:

- the possibility of microbial contamination of the reactors
- transfer of taints from the reactor to the wine
- loss of activity on prolonged operation
- leakage of cells or immobilization substrate into the wine

(Davis *et al.*, 1985).

For this method to be attractive in commercial practice it must be:

- cheap
- simple
- easily performed in an industrial situation
- not liable to cause oxidation of the wine
- robust
- not susceptible to contamination
- able to impart correct flavour changes to the wine
- must use commercially acceptable supports and organisms

Investigation into any immobilization method must include the effects of major variations in substrate and environmental parameters so that the bioreactor throughput to give effective treatment can be calculated. These parameters include: SO₂, pH, ethanol concentration, temperature and flowrate.

This thesis deals with continuous malolactic fermentation of Chardonnay-style wines using immobilized cells, with the ultimate aim of developing and testing a prototype bioreactor in an industrial situation. Chardonnay-style wine was seen as providing the most difficult conditions for MLF and the style most in need of improved fermentation methodology.