

Michele L. Ybarra, Kelvin Mwaba, Tonya L. Prescott,
Nicolette V. Roman, Bronwyn Rooi & Sheana Bull (2014)
Opportunities for technology-based HIV prevention programming among
high school students in Cape Town, South Africa, *AIDS Care*, 26:12, 1562-1567
DOI: 10.1080/09540121.2014.936814



Opportunities for technology-based HIV prevention programming among high school students in Cape Town, South Africa

Michele L. Ybarra^{a*}, Kelvin Mwaba^b, Tonya L. Prescott^a, Nicolette V. Roman^b, Bronwyn Rooi^b and Sheana Bull^c

^a*Center for Innovative Public Health Research, San Clemente, CA, USA;* ^b*Department of Psychology, University of the Western Cape, Cape Town, South Africa;* ^c*Colorado School of Public Health, University of Colorado Denver, Aurora, CO, USA*

One in three new cases of HIV in South Africa is among adolescents. Given that adolescents are particularly affected, scalable, and cost-effective prevention programs are urgently needed. This study aims to identify opportunities to integrate technology into youth HIV prevention efforts. In 2012, 1107 8th-11th graders completed a paper-and-pencil survey. Respondents were enrolled in one of three public high schools in Langa, a lower income community in Cape Town, South Africa. Eighty-nine percent of respondents have used text messaging (SMS) and 86% have gone online. If an HIV prevention program was offered online, 66% of youth would be somewhat or extremely likely to access it; slightly fewer (55%) felt the same about SMS-based programming. In comparison, 85% said they would be somewhat or extremely likely to access a school-based HIV prevention program. Interest in Internet- (60%) and SMS-based (54%) HIV prevention programming was similar for youth who had a self-appraised risk of HIV compared to youth who appraised their risk to be lower, as it was for youth who were tired of hearing messages about HIV prevention. Technology use is common - even among high school students who live in lower income communities. At the same time, these data reveal that it is not uncommon for youth to be tired of hearing messages about HIV prevention, and many of the typical topics key to HIV prevention have low interest levels among youth. HIV prevention researchers need to be mindful of the extent of existing programming that youth are exposed to. Technology-based programming may be especially amenable to meeting these requirements because of its novelty especially in developing countries, and because interactive functionality can be easily integrated into the program design. Given the preference for school and Internet-based programming, it seems that a hybrid approach is likely feasible and acceptable.

Keywords: HIV prevention; adolescents; technology; needs assessment

*Corresponding author.

Introduction

Despite intensive HIV prevention research (Hargreaves et al., 2007; Rehle et al., 2007) and the development of effective prevention programming (Harrison, Newell, Imrie, & Hoddinott, 2010; Jewkes et al., 2008; Pettifor et al., 2004), adolescent HIV preventive behavior change remains elusive (Hartell, 2005; Shisana et al., 2009; Van der Linde & Human Sciences Research Council, 2013). Of great concern, HIV risk behaviors appear to be increasing among South Africans (Van der Linde & Human Sciences Research Council, 2013).

To promote behavior change, HIV prevention programs not only need to be efficacious but also easily implemented. Data suggest explosive technology growth in South Africa (Central Intelligence Agency, 2013; Smith, 2013). Technology-based interventions lack many of the structural challenges noted in traditional interventions and are less costly to scale up (Cole-Lewis & Kershaw, 2010; Free et al., 2013; Noar, Black, & Pierce, 2009; Perry et al., 2012; Ybarra & Eaton, 2005). Nonetheless, technology is under-utilized to promote HIV preventive behavior in developing countries. To determine whether technology is an appropriate delivery mechanism for adolescent-focused HIV preventive programming in South Africa, a better understanding of both how young people use technology and their interest in engaging with HIV preventive information through technology, is critical.

Methods

The research protocol was reviewed and approved by the University of the Western Cape and the Chesapeake Institutional Review Board. Data were collected between April and August 2012. All participants provided written informed consent.

Location and participants

Respondents attended one of three partner schools in the low-income community, Langa. Because it is the closest black township to Cape Town, Langa has the highest density of people in the region. All students in Grade 8 to Grade 11 who were 16 years of age (the legal age of consent in South Africa) or older were invited to voluntarily participate in the study.

Procedures

Surveys were completed via paper and pencil in the absence of the teachers and school administrators. To assure anonymity, names were not collected on the survey instrument.

Respondents took an average of one hour to complete the survey. The survey was written in English, which is the official language of South Africa and the language of instruction in the high schools. There are multiple living languages in South Africa, however (Intersol Inc., 2010). The survey was first piloted among youth in the target age range to ensure readability and acceptability of sensitive questions.

Measures

Acceptability of information delivery by mode

All youth were asked: “If there was a health education program about HIV/AIDS prevention for teenagers, how likely would you go to it if it was a) at school, b) at a religious organization, c) over e-mail, d) over text (SMS) messages, and e) on the Internet.”

Interest in HIV prevention programming

Topics were chosen to reflect the key components of HIV prevention programs, as well as those that were posited to be salient to youth based upon our previous work with sub-Saharan adolescents (Ybarra, Biringi, Prescott, & Bull, 2012). Youth were asked to rate how much they agreed or disagreed with the statement: “I am tired of hearing about how to prevent HIV/AIDS and other STDs (sexually transmitted diseases).” They also were presented with a list of eight different topics related to HIV/AIDS and were asked whether they would be interested in receiving information about them. Topics included HIV-specific information (e.g., how to use a condom), family planning (e.g., birth control), and relationships (e.g., how to start a relationship).

HIV risk attitudes and behaviors

Youth were asked to appraise their personal risk of getting HIV; and how strongly they agreed or disagreed with the following statement: “I am tired of hearing about how to prevent HIV/AIDS and other STDs (sexually transmitted diseases).”

Vaginal sex was queried: “Have you ever had vaginal sex? We mean when a penis goes into a vagina.” Anal sex was queried: “Have you ever had anal sex? We mean when a penis goes into an anus.”

Data cleaning

All surveys were double entered by project staff to ensure accuracy. Missing data were imputed using the “impute” command in Stata, which estimates missing values using best set regression (StataCorp LP, 2009). Each respondent was required to have valid, non-missing data for at least 70% of the variables included in the analyses. Based on this criterion, 92 respondents were removed and 1015 were retained.

Results

Of the total 1460 students who were enrolled in the three partner schools, 1279 students (88%) were present on the day of the survey. 1107 of the 1191 eligible students completed the survey (response rate = 93%). Demographic and technology use characteristics are shown in Table 1.

Interest in HIV prevention programming

As shown in Table 2, one in three youth (38%) somewhat or strongly agreed that they were tired of hearing about how to prevent HIV/AIDS and other STDs. Few expressed interest in learning: how to “refuse sex” (43%), how to use a condom (42%), and about birth control (32%). Even topics posited to be of interest to adolescents (e.g., how to develop a

relationship) were endorsed by less than half of youth. Males tended to be more fatigued and less interested than females about HIV prevention programming.

Acceptability of information delivery by mode

Of the five access points queried, youth were most likely to indicate that they would be somewhat or very likely to access an HIV prevention program if it were delivered at school (85%, Table 3).

Reaching youth at greater risk of HIV

Sixty percent of youth who appraised their likelihood of getting HIV as above average chance said they were somewhat or very likely to access an HIV prevention program if it were online and 54% if it were via SMS. Similar rates of interest in Internet (60%) and SMS (53%) programming were noted among youth who agreed they were tired of HIV prevention messaging, and for youth who reported ever having vaginal or anal sex (67% and 55%, respectively). As shown in Table 4, among otherwise similar youth, those who were tired of hearing about HIV prevention were about 50% less likely to access a program if it were offered at school or online, with a similar trend noted for email. Youth who had vaginal or anal sex were 34% less likely to access it if it were offered online.

Table 1. Youth characteristics (n = 1015).

Youth characteristics	% (n)
<i>Demographic characteristics</i>	
Female	63.7% (647)
Age (range: 16–24; M:SD)	17.5 (1.2)
<i>Grade</i>	
Grade 8	0.2% (2)
Grade 9	0.1% (1)
Grade 10	52.0% (528)
Grade 11	47.7% (484)
<i>Race</i>	
African	97.7% (992)
Colored ^a	0.6% (6)
Indian	0.1% (1)
White	0.1% (1)
Other	1.5% (15)
<i>Father's education</i>	
No formal education	2.4% (24)
Primary school	6.5% (66)
Secondary school	36.6% (371)
Tertiary institution/university graduate	25.8% (262)
I am not sure	28.8% (292)
<i>Income</i>	
Lower than the average family	29.0% (294)
Similar to the average family	60.8% (617)
Higher than the average family	10.3% (104)
<i>Mother's education</i>	
No formal education	2.6% (26)
Primary school	5.2% (53)
Secondary school	39.8% (404)
Tertiary institution/university graduate	31.1% (316)
I am not sure	21.3% (216)
<i>Importance of religion on respondent's life</i>	
Not at all important	3.0% (30)
Somewhat unimportant	1.4% (14)
Somewhat important	11.8% (120)
Very important	83.8% (851)
<i>Technology use</i>	
Ever used the Internet	85.8% (871)
<i>Ever used SMS (text messaging)</i>	
Do not have a phone	9.9% (100)
Have a phone, do not SMS	1.5% (15)
SMS	88.7% (900)
Self-appraised chance of getting HIV is above average/very strong	12.7 (129)

^aColored refers to youth who are mixed race (White and African).

Discussion

Technology use is common among adolescents attending our three partner schools in the lower income community, Langa, South Africa. These data also reveal that it is not uncommon for youth to be tired of hearing messages about HIV prevention, and many of the typical key HIV prevention topics key have low interest levels among youth. HIV prevention researchers need to be mindful of the extent of existing programming that youth are exposed to. Technology-based programming may be especially amenable to meeting these requirements because of its novelty especially in developing countries, and because interactive functionality can be easily integrated into the program design.

According to youth in this study, the likelihood of accessing an HIV prevention program was greatest if the program were offered at school. Perhaps this is because currently available programming is predominantly delivered in schools, so this is the most familiar option for youth.

Table 2. Interest in specific HIV prevention programming-related topics (*n* = 1015).

Appraisal of HIV prevention programming	All (<i>n</i> = 1015)	Males (<i>n</i> = 368)	Females (<i>n</i> = 647)	<i>p</i> Value
Somewhat/very tired of hearing about how to prevent HIV	37.5% (381)	42.1% (155)	34.9% (226)	0.02
<i>Interested in learning about ...</i>				
How to end a relationship	43.8% (445)	36.7% (135)	47.9% (310)	0.001
How to avoid sex if you don't want to have sex	43.4% (440)	28.8% (106)	51.6% (334)	<0.001
How to use a condom	42.2% (428)	43.5% (160)	41.4% (268)	0.52
How drugs and alcohol affect your decision-making	39.1% (397)	50.3% (185)	32.8% (212)	<0.001
Birth control/family planning	31.5% (320)	27.5% (101)	33.9% (219)	0.04
How to refuse sex from a sugar daddy/mommy	22.7% (230)	22.3% (82)	22.9% (148)	0.83
Where to get an HIV test	21.9% (222)	23.9% (88)	20.7% (134)	0.24
How to develop and maintain a romantic relationship	16.7% (169)	22.0% (81)	13.6% (88)	0.001

p Value based upon a chi-square testing the relative difference in endorsement for males versus females.

Table 3. Likelihood of accessing an HIV prevention program based upon different access points (*n* = 1015).

Likelihood of accessing an HIV prevention program if it were available through...	School	Religious organization	Email	SMS (text messaging)	Internet
Not at all likely	10.1% (102)	17.8% (181)	33.1% (336)	27.5% (279)	20.9% (212)
Somewhat unlikely	4.7% (48)	12.3% (125)	24.6% (250)	17.4% (177)	13.0% (132)
Somewhat likely	38.3% (389)	46.1% (468)	22.2% (225)	30.6% (311)	31.4% (319)
Extremely likely	46.9% (476)	23.7% (241)	20.1% (204)	24.4% (248)	34.7% (352)

This does not mean that youth were opposed to technology-based programs, however: two in three youth said they were at least somewhat likely to access an Internet-based program, and slightly more than one in two youth would access an SMS-based program. Given the preference for school- and Internet-based programming (among the technology-related access points), a hybrid approach may be feasible and acceptable. For example, perhaps an Internet program could be offered as an after-school activity on school grounds. By utilizing the Internet as the program delivery mechanism, one safeguards program fidelity by ensuring all youth have access to the same, accurate sexual health information, while simultaneously allowing youth greater privacy to complete the program when and where they are comfortable. It also takes the onus off the teacher to deliver what can be uncomfortable or embarrassing information to their students, as well as the burden of finding classroom time to teach the material (Kinsman et al., 2001).

Table 4. Relative odds of being somewhat or very likely to access an HIV prevention program^a.

	School (<i>n</i> = 1015)	Religious organization (<i>n</i> = 1015)	Email (<i>n</i> = 1015)	SMS (text messaging; <i>n</i> = 1015)	Internet (<i>n</i> = 1015)
HIV risk attitudes and behaviors	aOR (95% CI)	aOR (95% CI)	aOR (95% CI)	aOR (95% CI)	aOR (95% CI)
Above average chance of getting HIV	0.97 (0.54, 1.73)	1.00 (0.60, 1.66)	0.82 (0.55, 1.23)	0.86 (0.57, 1.32)	1.26 (0.78, 2.04)
Tired of hearing about HIV prevention	0.47 (0.31, 0.72)	0.79 (0.56, 1.11)	<i>0.78 (0.59, 1.03)</i>	0.79 (0.59, 1.06)	0.57 (0.42, 0.80)
Ever had vaginal or anal sex	0.82 (0.51, 1.33)	0.82 (0.56, 1.20)	0.89 (0.66, 1.22)	0.95 (0.69, 1.32)	0.66 (0.45, 0.95)

^aFive different multivariate logistic regression models were estimated, one for each access point (e.g., school). Each model estimates the relative odds of being somewhat or very likely to access an HIV prevention program through the access point in question, given the indicator of HIV risk behavior or attitudes in question. Odds ratios are adjusted for the other two HIV risk attitudes and behaviors, as well as demographic characteristics (i.e., biological sex, age, grade, race, paternal and maternal education, income, i.e., religiosity), technology use (i.e., Internet, SMS), self-reported honesty, and the number of variables imputed. Bold texts denote statistical significance (*p* < 0.05). Italicized text denotes borderline statistical significance (*p* < 0.10).

Limitations

The generalizability of these data to greater South Africa, and youth who do not speak English fluently or are not enrolled in high school is unknown. Furthermore, it is possible that some youth did not answer honestly. Additionally, technology use was lower among the 92 respondents excluded from the analyses. Differences in findings may have been noted if these youth or those who were absent on the day of the survey had been included in the analytical sample.

Conclusions

In South Africa, like many countries, people who live in low-income households are at higher risk of HIV (Government of the Republic of South Africa, 2011). The access to and interest in technology as a delivery mechanism for HIV prevention programming among low income students in the current study suggest this may be an under-utilized opportunity to reach this vulnerable population. Importantly, too, youth with characteristics that may further increase their risk of HIV (e.g., vaginal sex) are generally as likely to be interested in technology based programming as their peers. The current data support a greater exploration of applications of technology in delivering HIV preventive behavior in South Africa.

Funding

This research was funded by a grant from the National Institute of Mental Health at the National Institutes of Health [grant number R03MH094238; PI: Ybarra]. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

- Central Intelligence Agency. (2013). The world factbook: South Africa. Retrieved May 8, 2013, from <https://www.cia.gov/library/publications/the-world-factbook/geos/sf.html>
- Cole-Lewis, H., & Kershaw, T. (2010). Text messaging as a tool for behavior change in disease prevention and management. *Epidemiologic Reviews*, 32, 56–69. doi:10.1093/epirev/mxq004
- Free, C., Phillips, G., Galli, L., Watson, L., Felix, L., Edwards, P., ... Haines, A. (2013). The effectiveness of mobile health technology-based health behaviour change or disease management interventions for health care consumers: A systematic review. *PLoS Medicine*, 10, e1001362. doi:10.1371/journal.pmed.1001362
- Government of the Republic of South Africa. (2011). Getting to success: Improving HIV prevention efforts in South Africa. Pretoria: Government of the Republic of South Africa. Retrieved from http://supportservices.ufs.ac.za/dl/userfiles/Documents/00001/1432_eng.pdf
- Hargreaves, J. R., Bonell, C. P., Morison, L. A., Kim, J. C., Phetla, G., Porter, J. D., ... Pronyk, P. M. (2007). Explaining continued high HIV prevalence in South Africa: Socioeconomic factors, HIV incidence and sexual behaviour change among a rural cohort, 2001–2004. *AIDS*, 21, S39–S48. doi:10.1097/01.aids.0000300534.97601.d6
- Harrison, A., Newell, M., Imrie, J., & Hoddinott, G. (2010). HIV prevention for South African youth: Which interventions work? A systematic review of current evidence. *BMC Public Health*, 10, 102. doi:10.1186/1471-2458-10-102
- Hartell, C. G. (2005). HIV/AIDS in South Africa: A review of sexual behavior among adolescents. *Adolescence*, 40, 171–181.
- Intersol Inc. (2010). Languages of Africa. Retrieved September 11, 2013, from <http://www.intersolinc.com/newsletters/africa.htm>
- Jewkes, R., Nduna, M., Levin, J., Jama, N., Dunkle, K., Puren, A., & Duvvury, N. (2008). Impact of stepping stones on incidence of HIV and HSV-2 and sexual behaviour in rural South Africa: Cluster randomised controlled trial. *British Medical Journal*, 337, a506. doi:10.1136/bmj.a506
- Kinsman, J., Nakiyingi, J., Kamali, A., Carpenter, L., Quigley, M., Pool, R., & Whitworth, J. (2001). Evaluation of a comprehensive school-based AIDS education programme in rural Masaka, Uganda. *Health Education Research*, 16, 85–100. doi:10.1093/her/16.1.85
- Noar, S. M., Black, H. G., & Pierce, L. B. (2009). Efficacy of computer technology-based HIV prevention interventions: A meta-analysis. *AIDS*, 23, 107–115. doi:10.1097/QAD.0b013e32831c5500
- Perry, R. C. W., Kayekjian, K. C., Braun, R. A., Cantu, M., Sheoran, B., & Chung, P. J. (2012). Adolescents' perspectives on the use of a text messaging service for preventive sexual health promotion. *Journal of Adolescent Health*, 51, 220–225. doi:10.1016/j.jadohealth.2011.11.012
- Pettifor, A., Rees, H., Steffenson, A. E., Hlongwa-Madikizela, L., MacPhail, C., Vermaak, K., & Kleinschmidt, I. (2004). Young people's sexual health in South Africa: HIV prevalence and sexual behaviors from a nationally representative household survey. Johannesburg: Reproductive Health Research Unit, University of the Witwatersrand. Retrieved from <http://kaiserfamilyfoundation.files.wordpress.com/2013/01/hiv-and-sexual-behaviour-among-youngsouth-africans-a-national-survey-of-15-24-year-olds.pdf>
- Rehle, T., Shisana, O., Pillay, V., Zuma, K., Puren, A., & Parker, W. (2007). National HIV incidence measures – new insights into the South African epidemic. *South African Medical Journal*, 97, 194–199.
- Shisana, O., Rehle, T., Simbayi, L., Zuma, K., Jooste, S., Simbayi, L., ... the SABSSM III

- Implementation Team. (2009). South African national HIV prevalence, behaviour and communication survey, 2008: A turning tide among teenagers? Cape Town: Human Sciences Research Council Press. Retrieved from <http://www.mrc.ac.za/pressreleases/2009/sanat.pdf>
- Smith, G (2013). The reality of mobile usage and social media growth in South Africa. ImagiNation. Retrieved May 28, 2014, from <http://imagi-nation.co.za/mobile/reality-mobileusage-social-media-growth-south-africa>
- StataCorp LP. (2009). Stata statistical software (release 11) [Computer program]. College Station, TX: Author.
- Van der Linde, I., & Human Sciences Research Council. (2013). Plenary session 3, 20 June 2013: “HIV/AIDS in South Africa: At last the glass is half full”. Retrieved September 11, 2013, from <http://www.hsrc.ac.za/en/media-briefs/hiv-aids-stis-and-tb/plenary-session-3-20-june-2013-hiv-aids-in-south-africa-at-last-the-glass-is-half-full>
- Ybarra, M. L., Biringi, R., Prescott, T., & Bull, S. S. (2012). Usability and navigability of an HIV/AIDS internet intervention for adolescents in a resource limited setting. *Computer Informatics in Nursing*, 30, 587–595. doi:10.1097/NXN.0b013e318266cboe
- Ybarra, M. L., & Eaton, W. W. (2005). Internet-based mental health interventions. *Mental Health Services Research*, 7, 75–87. doi:10.1007/s11020-005-3779-8