

Copyright © 2019 Moller et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

1 **Determinants of Phage Host Range in *Staphylococcus* Species**

2 Abraham G. Moller^{a,c}, Jodi A. Lindsay^b, and Timothy D. Read^{c*}

3 a. Program in Microbiology and Molecular Genetics (MMG), Graduate Division of Biological and Biomedical
4 Sciences (GDBBS), Emory University School of Medicine, Atlanta, Georgia, 30322

5 b. Institute of Infection and Immunity, St George's, University of London, Cranmer Terrace, London, SW17
6 ORE, UK

7 c. Division of Infectious Diseases, Department of Medicine, Emory University School of Medicine, Atlanta,
8 Georgia, 30322

9 *Corresponding author – tread@emory.edu

10 **Abstract**

11 Bacteria in the genus *Staphylococcus* are important targets for phage therapy due to their prevalence
12 as pathogens and increasing antibiotic resistance. Here we review *Staphylococcus* outer surface features and
13 specific phage resistance mechanisms that define host range - the set of strains an individual phage can
14 potentially infect. Phage infection goes through five distinct phases - attachment, uptake, biosynthesis,
15 assembly and lysis. Adsorption inhibition, encompassing outer surface teichoic acid receptor alteration,
16 elimination, or occlusion, limits successful phage attachment and entry. Restriction-modification systems (in
17 particular, type I and IV systems), which target phage DNA inside the cell, serve as the major barriers to
18 biosynthesis as well as transduction and horizontal gene transfer between clonal complexes and species.
19 Resistance to late stages of infection occurs through mechanisms such as assembly interference, in which
20 staphylococcal pathogenicity islands siphon away superinfecting phage proteins to package their own DNA.
21 While genes responsible for teichoic acid biosynthesis, capsule, and restriction-modification are found in most
22 *Staphylococcus* strains, a variety of other host-range determinants (e.g., CRISPRs, abortive infection, and
23 superinfection immunity) are sporadic. Fitness costs of phage resistance through teichoic acid structure
24 alteration could make staphylococcal phage therapies promising, but host range prediction is complex because
25 of the large number of genes involved, many with unknown roles. In addition, little is known about genetic
26 determinants that contribute to host range expansion in the phages themselves. Future research must identify

27 host range determinants, characterize resistance development during infection and treatment, and examine
28 population-wide genetic background effects on resistance selection.

29 **Keywords:** staphylococci, phage resistance, host range, phage therapy, CRISPR

30 Introduction

31 The *Staphylococcus* genus includes commensals and pathogens of humans and animals. *S. aureus*
32 and *S. epidermidis*, in particular, cause diverse infections in humans and have become increasingly antibiotic
33 resistant over the past seventy years. Diseases range from food poisoning to skin and soft tissue infections,
34 pneumonia, osteomyelitis, endocarditis, and septic shock. *S. aureus* is carried by between 20% (persistently)
35 and 60% (intermittently) of the human population (1), primarily on the skin and upper respiratory tract.
36 Methicillin-resistant *S. aureus* (MRSA) emerged in the mid-1960s (2) and has reduced the options for
37 treatment with beta-lactam antibiotics. The combination of high carriage rates, diverse pathologies, prevalent
38 antimicrobial resistance, and lack of a licensed vaccine (3) makes staphylococcal species important targets for
39 new therapies.

40 Bacteriophage (phages) are natural killers of *Staphylococcus* bacteria lysing bacterial cells through
41 expression of holins, which permeabilize the membrane and release endolysins (4, 5) that degrade the
42 peptidoglycan of the cell wall (6). Phage therapy is a promising alternative to antibiotics for treating infections
43 because of the large number of diverse phages with low toxicity to humans and non-target species (7, 8).

44 Phage therapy has a long history, reaching back before the antibiotic era to shortly after the discovery
45 of phages themselves by Frederick Twort and Felix d'Herelle in the 1910s (9–11). While overshadowed by the
46 subsequent discovery of antibiotics and generally abandoned in the West for many years, phage therapy
47 persisted as a bacterial treatment in eastern Europe and the nations that composed the former Soviet Union
48 (9, 10). There, phage cocktails were developed for sepsis, osteomyelitis, and burn wounds, among other
49 staphylococcal diseases, with complete recovery reported in some cases (12). Polish and Soviet studies
50 showed that phage lysates effectively treated staphylococcal skin and lung infections (13). More recently, the
51 emergence of multi-drug resistance in bacterial pathogens has renewed interest in phage therapy and phage
52 biology (8, 14). Safety studies on the staphylococcal phage lysate (SPL) as well as phage cocktails containing
53 *S. aureus*-specific phages indicated that they had no adverse effects when administered intranasally,

54 intravenously, orally, topically, or subcutaneously (14). Phages have also been recently approved by the FDA
55 as a treatment to clear another Gram-positive species (*Listeria monocytogenes*) present in food (15) and
56 approved as personalized treatment for burn wound infections (16).

57 All known staphylococcal phages are members of the order *Caudovirales* with linear dsDNA virion
58 genomes. Staphylococcal phages are divided into three families with distinctive morphologies – the long,
59 noncontractile-tailed *Siphoviridae*, the contractile-tailed *Myoviridae*, and the short, noncontractile-tailed
60 *Podoviridae* (17, 18). *Siphoviridae* genomes are 39-43 kb in size, while those of the *Myoviridae* are 120-140 kb
61 and *Podoviridae* are 16-18 kb (17). Currently reported *Siphoviridae* are typically temperate phages that encode
62 lysogeny functions within a genomic module, while reported *Myoviridae* and *Podoviridae* are virulent. The
63 virulent phages are the strongest potential candidates for phage therapy, given that they are not known to
64 lysogenize and thus obligately kill their targets. Lytic staphylococcal phages have surprisingly broad host
65 ranges (19–22), anti-biofilm activity (19, 23), and varying effectiveness against infection (24–26). The
66 *Siphoviridae* are agents of horizontal gene transfer (HGT) through transduction (27) into recipient strains (17)
67 and activation of staphylococcal pathogenicity islands (SaPIs) (28). The *Siphoviridae* have been subdivided
68 into “integrase types” based on the sequence of the integrase gene necessary for lysogenic insertion into the
69 chromosome (17, 29). Certain integrase type phages introduce specific virulence factors (17). Integrase type 3
70 (Sa3int) phages encode the immune evasion cluster (IEC), which includes the staphylokinase (*sak*),
71 staphylococcal complement inhibitor (*scn*), chemotaxis inhibitory protein (*chp*), and enterotoxin S (*sea*). In
72 addition, Sa2int phage often encode Panton-Valentine leukocidin (*lukFS-PV*), while Sa1int phages often
73 encode exfoliative toxin A (*eta*). Temperate staphylococcal phages can also disrupt chromosomal virulence
74 factors (17). Sa3int and Sa6int phages, for example, integrate into sites in the beta-hemolysin (*hlyb*) or lipase
75 (*geh*) genes, respectively (30, 31).

76 No single phage can kill every *Staphylococcus* strain. Instead, each phage has a particular host range,
77 defined as the set of strains permissive for its infection. Host range can be limited by active host resistance
78 mechanisms such as CRISPR or restriction-modification that actively suppress phage infection or by passive
79 mechanisms such as loss of receptors for phage adsorption. It is unclear whether these host range limiting
80 factors have arisen through specific adaptation against phage infection or are byproducts of selection against
81 other stresses. There are, however, specific phage counteracting mechanisms to host resistance that serve to

82 broaden phage host range. Phage host range has great importance to phage therapy because it defines the
83 potential scope of treatable strains, thus informing selection of phages for rational, personalized cocktail
84 development.

85 Mechanisms of resistance to phage have been reviewed previously across bacteria generally (32, 33)
86 and in lactic acid bacteria (34), but this is the first article to focus on the particular features of *Staphylococcus*
87 (**Figure 1**). By far, the majority of the literature has focused on two species: *S. epidermidis*, and especially, *S.*
88 *aureus*. However, we include studies on other species (e.g. *S. simulans*) where appropriate. We then reflect on
89 possible consequences of resistance on phage host range and potential phage therapy for staphylococcal
90 infections, given that phage resistance elements determine host range and thus provide one criterion for phage
91 efficacy in therapy. We also consider the evolutionary trade-offs of phage resistance in a therapeutic context
92 due to the potential effects of phage resistance on either virulence or antibiotic resistance.

93 Host resistance can occur at different points in the phage life cycle (**Figure 1**) (32, 33). There are no
94 reports in *Staphylococcus* of mechanisms that limit host range at the uptake and host lysis phases. We
95 therefore concentrate on the attachment, biosynthesis, and assembly phases.

96 **Attachment**

97 **Wall teichoic acid is the primary staphylococcal phage receptor**

98 Attachment of phages to the outside of the *Staphylococcus* cell (**Figure 2A**) is the first stage of infection
99 (**Figure 1**). *Staphylococcus* may be resistant to phage adsorption if the receptor molecule is not present, not
100 recognized by the phage, or blocked. Mutations that alter components of the outer surface can have the effect
101 of inhibiting adsorption and thus conferring resistance. Through genetic and biochemical studies on a small
102 range of staphylococcal phages, the polyribitol phosphate (poly-RboP) polymer of wall teichoic acid (WTA) or
103 N-acetylglucosamine (GlcNAc) modifications at the 4 positions of ribitol phosphate monomers in WTA appear
104 to be the primary targets (35–41).

105 In an early *S. aureus* phage resistance study published in 1969, N-methyl-N'-nitro-N-nitrosoguanidine-
106 mutagenized strain H (Multi Locus Sequence Type 30; ST30) (42) phage-resistant mutants were selected by
107 plating on agar plates containing lawns of 52A (siphovirus) (40). Mutants also found resistant to phage K

108 (myovirus) were deficient in N-acetylglucosamine, cell wall phosphorus, and ester-linked D-alanine in their
109 envelopes, presumably due to a loss of wall teichoic acid production. Further biochemical characterization
110 showed that the mutants lacked UDP-GlcNAc:polyribitol phosphate transferase activity and WTA.
111 Counterintuitively, they did show the relevant biochemical activity for the last known step in WTA biosynthesis
112 (phosphoribitol transferase – TarL, **Figure 2B**) (38). This surprising result suggested the double resistant
113 mutants produced ribitol phosphate but either failed to properly polymerize WTA or attach it to the cell wall.
114 These mutants had pleiotropic phenotypic differences from their parent strain (41), including a longer
115 generation time than its parent; cell growth in clumps; irregular, rough, gray colonies; and increased levels of
116 wall-bound autolysin. A later study characterizing spontaneous *S. aureus* strain A170 (ST45) mutants resistant
117 to siphovirus M^{Sa} found similar phenotypic defects (43) and biochemical assays also showed that resistance
118 was likely due to the lack of GlcNAc-modified WTA.

119 Peschel and colleagues identified genes responsible for phage adsorption in a series of elegant
120 molecular genetic studies in the RN4220 (ST8) (44) background (35, 36, 45). Deletion of undecaprenyl-
121 phosphate N-acetylglucosaminyl 1-phosphate transferase (*tagO*), the first gene involved in WTA biosynthesis,
122 conferred resistance and reduced adsorption to tested *Myoviridae* (Φ812 and ΦK), while a transposon
123 insertion mutant in the *tarM* gene had resistance and reduced adsorption to *Siphoviridae* (ΦSa2mw, Φ47, Φ13,
124 and Φ77). Complementation of wild-type alleles rescued these phenotypes (35). TarM is a glycosyltransferase
125 responsible for attaching α-O-GlcNAc to the 4 position of the ribitol phosphate WTA monomer (46, 47). The
126 *tarM* mutant was previously shown to lack GlcNAc-modified WTA in its envelope (46). TarS, the
127 glycosyltransferase responsible for attaching β-O-GlcNAc to the 4 position of the ribitol phosphate WTA
128 monomer (48), was specifically required for podovirus adsorption (45). Deletion of *tarS* conferred resistance
129 and reduced adsorption to tested *Podoviridae* (Φ44AHJD, Φ66, and ΦP68) (45), but only deletion of both *tarS*
130 and *tarM* conferred reduced adsorption to tested *Siphoviridae* (Φ11) in the same RN4220 background used in
131 prior studies (49, 50). On the other hand, even *tarS*⁺, *tarM*⁺ strains were resistant to *Podoviridae*, suggesting
132 WTA decorated with α-O-GlcNAc by TarM impeded podovirus adsorption (45). Taken together, these findings
133 suggested, for the small number of representatives that were tested, elimination of WTA confers resistance to
134 all classes of phage, elimination of GlcNAc modifications confers resistance to the *Siphoviridae* and
135 *Podoviridae*, and elimination of β-O-GlcNAc modification confers resistance specifically to the *Podoviridae*.

136 Given the conservation of wall teichoic acid biosynthesis genes amongst *S. aureus* genomes (51) and the
137 cross-species activity of staphylococcal phages such as phage K (52), these conclusions could be expected to
138 hold in staphylococci beyond *S. aureus*.

139 Recent studies have suggested that as the number of strains and phages expands we may find a larger
140 number of genes influencing host range through attachment. Azam et al. conducted a long-term evolution
141 experiment in which they selected *S. aureus* SA003 (ST352) mutants resistant to myovirus Φ SA012 (53).
142 Resistant mutants gained missense mutations in five genes (*tagO*, RNase adapter protein *rapZ*, putative
143 membrane protein *yozyB*, guanylate kinase *gmk*, and alpha subunit of DNA-dependent RNA polymerase *rpoA*),
144 a nonsense mutation in one gene (UDP-N-acetylglucosamine 1-carboxyvinyltransferase *murA2*), and a 1,779
145 bp deletion that included the C-terminal region of the teichoic acid glycosyltransferase *tarS*, a non-coding
146 region, and the N-terminal region of the iron-sulfur repair protein *scd*. Complementation of mutations in genes
147 *scd*, *tagO*, *rapZ*, and *murA2* restored Φ SA012 sensitivity and adsorption, while only complementation of
148 mutations in *tarS* restored sensitivity and adsorption of another myovirus, Φ SA039. The results suggested that
149 while Φ SA012 recognized the WTA backbone, Φ SA039 was unusual in recognizing β -O-GlcNAc-modified
150 WTA, hinting that there may be more variability in receptor targets within phage groups than the limited number
151 of earlier studies suggested.

152 The carriage of a prophage in certain *S. aureus* CC5 and CC398 strains that encodes alternative WTA
153 glycosyltransferase *tarP* (54) adds further complications. TarP attaches GlcNAc to the 3 position of ribitol
154 phosphate rather than the 4 position, thus conferring *Siphoviridae* (Φ 11, Φ 52a, Φ 80) sensitivity but
155 *Podoviridae* (Φ 44, Φ 66, and Φ P68) resistance. It is interesting in the light of host range evolution that a gene
156 carried on a prophage can change the properties of the *S. aureus* surface and thus affect the host ranges of
157 other phages.

158 Although the majority of staphylococcal phage tested bind WTA and GlcNAc receptors, there is one
159 known exception. Siphovirus Φ 187 binds WTA glycosylated with N-acetyl-D-galactosamine (GalNAc), the
160 unusual WTA synthesized by *S. aureus* ST395 (55). The α -O-GalNAc transferase *tagN*, the nucleotide sugar
161 epimerase *tagV*, and the short GroP WTA polymerase *tagF* genes are required specifically for synthesis of
162 ST395 WTA. Homologs of these genes were found in genomes of multiple Coagulase-Negative *Staphylococci*
163 (CoNS) strains, such as *S. pseudointermedius* ED99, *S. epidermidis* M23864:W1, and *S. lugdunensis*

164 N920143. Complementation of a *S. aureus* PS187 *tagN* C-terminal glycosyltransferase deletion with the wild-
165 type *tagN* gene or that from *S. carnosus* (*tagN*-Sc) successfully restored the wild-type phenotype, suggesting
166 *tagN* homologs in other CoNS genomes had similar functions to that in *S. aureus* PS187 (ST395).
167 Complementation of the *tagN* C-terminal deletion with either PS187 or *S. carnosus tagN* also restored wild-
168 type Φ 187 sensitivity. This difference in WTA structure was shown to prohibit transduction between ST395 and
169 other *S. aureus* lineages (56). Staphylococcal pathogenicity island (SaPI) particles prepared in a ST1, 5, 8, 22,
170 25, or 30 strain with phages Φ 11 or Φ 80 α failed to transduce any ST395 strains. SaPI particles prepared in a
171 ST395 strain, on the other hand, transduced other ST395 strains as well as CoNS species and *Listeria*
172 *monocytogenes*. These findings suggest the unique ST395 WTA restricts phage host range to strains of the
173 same sequence type or Gram-positives with a related WTA structure, such as *Listeria monocytogenes*.

174 There has been one study showing that staphylococcal phages (siphovirus Φ SLT) can bind lipoteichoic
175 acid (LTA), the lipid-anchored, polyglycerol phosphate (GroP) TA polymer (57) (**Figure 2A**). However,
176 subsequent elimination of LTA biosynthesis through *ltaS* deletion had no effect on phage adsorption or
177 sensitivity (35) and therefore the potential significance of LTA as an alternative receptor is currently unknown.

178 **The effects of surface proteins and extracellular polysaccharides on attachment**

179 Although proteins serve as receptors for many Gram-positive phages (for example, the YueB receptor
180 for *Bacillus subtilis* phage SPP1 (58)), there is no evidence to suggest *S. aureus* proteins serve as its phage
181 receptors. Phage interaction protein (Pip) homologs exist throughout the Gram-positives, serving as protein
182 receptors to which phage irreversibly bind (59). There are Pip surface protein homologs anchored to the
183 staphylococcal cell wall through the action of the sortase enzyme in *Staphylococcus* (60, 61). However, neither
184 deletion of the Pip homologs in RN4220 (ST8) (49) nor sortase A in Newman (ST254) (62, 63) affected
185 sensitivity to phage Φ 11 and phages Φ NM1, Φ NM2, and Φ NM4, respectively.

186 Some classes of proteins or extracellular polysaccharides have been shown to block phage adsorption
187 in the staphylococci through occlusion of the WTA receptors. Overproduction of surface protein A in *S. aureus*
188 was shown to reduce phage adsorption through this mechanism (64), but work on surface protein occlusion
189 remains limited. Capsule types 1 and 2 - strains M (ST1254) (42) and Smith diffuse (ST707) (42), respectively -
190 were shown to occlude adsorption (65), but the most common capsule types, 5 and 8, showed inconclusive

191 results (66, 67). Differences in capsule thickness between strains may account for these variable results. Type
192 1 and 2 strains are mucoid and heavily encapsulated, while type 5 and 8 are non-mucoid despite
193 encapsulation (68). The CoNS species *Staphylococcus simulans* also showed capsule-dependent inhibition of
194 phage adsorption (69).

195 The exopolysaccharides (EPS) of staphylococcal biofilms have not been shown to occlude adsorption.
196 Surface proteins, such as biofilm-associated protein (Bap), exopolysaccharides (polysaccharide intercellular
197 adhesin - PIA - composed of poly-N-acetylglucosamine – PNAG – and synthesized by the products of the
198 *icaADBC* operon), and extracellular DNA (eDNA) compose staphylococcal biofilms, which can form by PIA-
199 dependent or protein (Bap)-dependent mechanisms (70, 71). Other surface proteins more common than Bap
200 can also mediate biofilm formation, such as FnbA/FnbB (72, 73) and SasG (74) in *S. aureus* and Aap in *S.*
201 *epidermidis* (70). Both *S. aureus* (19, 75) and *S. epidermidis* (52, 76, 77) biofilms are susceptible to phage
202 predation. Phage resistance in staphylococcal biofilms may instead be associated with altered biofilm diffusion
203 or metabolism, the latter of which resembles stationary phase growth. Studies on *S. epidermidis* suggested
204 phage susceptibility was similar in biofilms and stationary phase cultures (52). Phages may in fact promote
205 bacterial persistence in *S. aureus* biofilms by releasing nutrients from lysed cells for remaining live ones to
206 utilize (78).

207 **Biosynthesis**

208 **Superinfection immunity**

209 Staphylococcal temperate phages encode homologs of the *cl* repressor (17, 18). In *E. coli*, this protein
210 represses expression of the lytic cycle in newly infecting phages with the same *cl* protein-binding sites, thus
211 stopping new infections through a mechanism called superinfection immunity. Molecular and evolutionary
212 studies on the *E. coli* phage lambda model suggest many superinfection immunity groups (in which member
213 temperate phages confer immunity to each other upon integration) coexist in nature (79), with *cl* repressor –
214 operator coevolution driving the emergence of new immunity groups (80). Superinfection immunity as a
215 determining factor in phage host range in staphylococcal species appears not to have been studied yet, but
216 since prophages are common (most sequenced *S. aureus* genomes contain 1-4 prophages) (18, 81), it may be
217 a significant barrier to phage infection.

218 Restriction-modification (R-M) systems

219 Bacteria can resist phage infection by degrading injected phage DNA before it has the chance to
220 replicate and enter the lytic or lysogenic cycle (**Figure 1**). Restriction-modification (R-M) is a prominent phage
221 infection barrier in the *Staphylococcus* genus. R-M systems are modular operons containing combinations of
222 host specificity determinant (*hsd*) genes encoding three types of functions: restriction endonuclease activity
223 (*hsdR*) responsible for destroying unmodified DNA, DNA adenosine or cytosine methyltransferase activity
224 (*hsdM*) responsible for modifying host DNA so that it is not cleaved by restriction endonucleases, and
225 specificity DNA-binding proteins (*hsdS*) responsible for recognizing sequence motifs targeted for cleavage or
226 modification (82).

227 There are four known types of R-M systems in bacteria, all of which have been found in the
228 staphylococci (83). In type I systems, the restriction enzyme cleaves unmodified DNA adjacent to its binding
229 site, sometimes separated by as much as 1000 bp from the binding site, while the modification enzyme
230 methylates host DNA at the target site specified by the specificity protein. A complex containing all three types
231 of subunits restricts unmodified exogenous DNA, while HsdSHsdM complexes only modify DNA. In type II
232 systems, the restriction enzyme (HsdR₂) cleaves unmodified DNA at its binding site, while the modification
233 enzyme (HsdM) modifies DNA at this site. In type III systems, the restriction enzyme cleaves unmodified DNA
234 roughly 24-28 bp downstream from its asymmetric target site, while the modification enzyme methylates a
235 single strand of host DNA at the target site. The modification subunit (Mod) modifies one strand of DNA either
236 by itself (Mod₂) or in complex with the restriction subunit (e.g., Mod₂Res₁ or Mod₂Res₂), while the restriction
237 subunit (Res) cleaves unmodified DNA only in complex with modification subunits (Mod₂Res₁ or Mod₂Res₂). In
238 type IV systems, the restriction enzyme only cleaves modified, methylated DNA. Type IV systems do not
239 include a modification enzyme. These systems have been well studied in *S. aureus* (and in *S. epidermidis*, to a
240 more limited extent) due to their role in restricting natural horizontal gene transfer and genetic manipulation of
241 the organism (83–86).

242 Type I R-M systems are the most abundant class of R-M systems reported in *S. aureus*, followed by
243 type IV and then type II systems (83). Type III systems appear to be rare, with only two described in the genus
244 (83). Analyses of the restriction enzyme genomic database REBASE in 2014 showed that all completed *S.*

245 *aureus* genomes encode a type I R-M system and that most *S. aureus* genomes annotated with R-M genes
246 encode a type I system (83, 87). The most common type I R-M locus found in *S. aureus* is Sau1 (88).
247 Expressing a functional Sau1 *hsdR* gene in restriction-deficient *S. aureus* strain RN4220 greatly reduced
248 electroporation, conjugation, and transduction frequencies (88). *S. aureus* genomes generally encode two
249 Sau1 *hsdS* genes that specify two distinct DNA motif targets for restriction or modification (89). The Sau1
250 HsdS subunit determines target specificity through its two target recognition domains (TRDs), which each bind
251 to one part of the target sequence (90). TRDs are the least conserved portions of the HsdS amino acid
252 sequences (88), and vary in carriage between strains with lineage and/or clonal complex-specific variant
253 associations, as microarray hybridization studies indicate (88, 89). The Sau1 system prevented transfer of
254 plasmid DNA from one clonal complex (CC5) to another (CC8) with a different target recognition site (89),
255 showing that restriction defines barriers between clonal complexes. Sau1 also affected susceptibility of two
256 CC8 strains (NCTC8325-4 and RN4220 *phsdR*) but not the *hsdR*-deficient RN4220 to phage Φ 75 (siphovirus)
257 propagated in a CC51 strain (879R4RF), suggesting Sau1 can control phage host range (88). Sau1 variation is
258 a powerful marker of lineage/clonal complex (88, 91) and likely drives the independent evolution of clonal
259 complexes. Sau1 would therefore be predicted to be a major host range limitation to phages grown in a strain
260 of a different clonal complex. Since the target sites of nearly all *S. aureus* Sau1 R-Ms from each of the different
261 clonal complexes have now been identified (90), it should be possible to bioinformatically predict the Sau1-
262 defined clonal complex host range of any sequenced bacteriophage.

263 Type IV R-M system SauUSI is estimated to be found in 90% of *S. aureus* strains (83, 92) and, in
264 combination with Sau1, presents an effective restriction barrier for resisting phage infection (93). SauUSI
265 specifically restricts DNA methylated or hydroxymethylated at the C5 position of cytosine (92). The preferred
266 binding site for SauUSI is Sm5CNGS, where S represents either cytosine or guanine (92). Type II R-M
267 systems have been estimated to be in ~33% of strains and display a range of target sites (83, 94–96). The
268 most common type II R-M system found in *S. aureus* is called Sau3A (94). The Sau3A restriction enzyme
269 cleaves 5' to the guanine in unmodified 5'-GATC-3' sequences. The Sau3A modification enzyme, on the other
270 hand, methylates the restriction site at the C5 position of cytosine (97). Some type II systems, such as Sau42I,
271 are encoded by phages. Sau42I is an example of a type IIS R-M system, which binds asymmetric DNA
272 sequences and cleaves outside the recognition site, unlike most type II systems (82). Unlike type I and type IV,

273 type II systems are often carried on mobile genetic elements which are capable of frequent transfer between
274 strains and are not conserved amongst all members of the same clonal complex, so they present a more
275 strain-specific and variable limit to host range (87). Certain *S. aureus* type II R-M systems (e.g., Sau96I) serve
276 to negate the Type IV SauUSI system because they methylate cytosines and guanines in sequences SauUSI
277 targets for cleavage. This is an interesting example of how R-M systems acquired by HGT can have
278 unpredictable interactions with existing systems.

279 If unmodified phages can survive restriction enzyme degradation upon cell entry, the phage DNA
280 molecules acquire protective DNA methylation as they replicate. While survival of restriction can happen
281 stochastically at high multiplicities of infection, phages have also been shown to have evolved or acquired
282 adaptations for restriction evasion. Anti-restriction mechanisms include restriction site alteration, restriction site
283 occlusion, indirect subversion of restriction-modification activity, and direct inhibition of restriction-modification
284 systems (98). Restriction site alteration can include both incorporation of alternative bases, such as 5-
285 hydroxymethyluracil (5hmU) and 5-hydroxymethylcytosine (5hmC), and loss of restriction sites through
286 selection. A clear example of the latter in the staphylococci is the elimination of GATC sites in the 140 kb
287 phage K genome, enabling its avoidance of Sau3A restriction (99). Another example is the evolution of
288 particular antimicrobial resistance-carrying conjugative plasmids which have lost specific Sau1 R-M sites
289 allowing their transfer between common MRSA lineages (88). Restriction site occlusion refers to DNA-binding
290 proteins preventing restriction enzymes from binding and digesting DNA (98, 100, 101). R-M subversion either
291 occurs through stimulation of host modification enzymes or destruction of restriction cofactors (e.g., SAM) (98,
292 102, 103). R-M inhibition occurs most often in type I systems (but also in some type II systems) through the
293 binding of specific anti-restriction proteins, such as ArdA, ArdB, and Ocr (98, 104, 105). There is no literature
294 specifically characterizing anti-restriction in *Staphylococcus*, but an *E. coli* *ardA* homolog has been identified in
295 the staphylococcal Tn916 and Tn5801 transposons (106).

296 **Clustered regularly interspaced short palindromic repeat (CRISPR) systems**

297 CRISPRs confer immunity to phage infection through the cleavage of extrinsic DNA in a sequence-
298 specific manner. Unlike R-M systems, which target specific DNA sequence motifs, CRISPRs adaptively
299 incorporate target sequences from phages they have destroyed to increase the efficiency of protection. After

300 integrating short segments of foreign DNA as spacers of CRISPR arrays, CRISPR-associated (Cas) nucleases
301 process the transcribed CRISPR array RNA into CRISPR RNAs (crRNAs) used to target new incursions of
302 identical foreign DNA elements for destruction (107, 108). Surveys of *S. aureus* and *S. epidermidis* genomes
303 indicate CRISPRs are not common in these species (109, 110). These surveys looked for the presence of *cas6*
304 and *cas9* genes, which are nucleases required for Type I/III and Type II CRISPR-mediated resistance,
305 respectively. Cas6 is an endoribonuclease found in Type I and III CRISPR systems that cleaves pre-crRNA
306 transcripts within the 3' end of the repeat region to produce mature guide crRNAs (111, 112), while Cas9 is an
307 endonuclease found in Type II CRISPR systems that cleaves DNA in a crRNA-guided manner (112, 113). Only
308 12 of 300 published *S. epidermidis* genomes searched encoded the Cas6 nuclease, 18 of 130 *S. epidermidis*
309 isolates from Denmark (Copenhagen University Hospital) tested positive for *cas6* via PCR, and 14 of nearly
310 5000 published *S. aureus* genomes encoded CRISPR/Cas systems (109). Another study specifically
311 examining *S. aureus* found that 2 of 32 *S. aureus* strains encoded CRISPR/Cas systems (110). These
312 CRISPRs were similar to those found in two *S. lugdunensis* strains, suggesting they were recombined with *S.*
313 *lugdunensis* or derived from a common ancestor (110). CRISPR/Cas systems have also occasionally been
314 reported in strains of other species (*S. capitis*, *S. schleiferi*, *S. intermedius*, *S. argenteus*, and *S. microti*) (109).
315 Only a single *S. aureus* strain has been reported to encode Cas9, which is found in an SCCmec-like region
316 (114). Nonetheless, CRISPR systems have been shown to be important in resisting introduction of foreign
317 DNA in *S. epidermidis* RP62a (115, 116). Anti-CRISPR mechanisms, such as proteins that prevent CRISPR-
318 Cas systems from binding DNA target sites, are being discovered in many phages (117–119), although not yet
319 in those specific for staphylococci. Currently discovered anti-CRISPR mechanisms have been shown to target
320 both type I and type II CRISPR systems (117–120).

321 **Assembly**

322 Assembly interference is the parasitization of superinfecting phage by chromosomal phage-like
323 elements and has been demonstrated experimentally in *S. aureus* pathogenicity island (SaPI)-helper phage
324 interactions. SaPIs encode important virulence factors, such as toxic shock syndrome toxin (TSST), but are
325 only mobilized by superinfecting helper siphoviruses (28, 121). The *Dut* dUTPase protein expressed by helper
326 phages derepresses the *Stl* SaPI repressor, activating the SaPI lytic cycle (28). The derepressed SaPIs then
327 take advantage of the superinfection to proliferate at the expense of the helper phage. SaPIs interfere with

328 helper phage assembly through several mechanisms (122) - remodeling phage capsid proteins to fit the small
329 SaPI genome (123–127), encoding phage packaging interference (Ppi) proteins that prevent helper phage
330 DNA packaging into new SaPI particles (123), and disrupting phage late gene activation (128). All known
331 SaPIs encode phage packaging interference (Ppi) proteins, which divert phage DNA packaging toward SaPIs
332 by inhibiting helper phage terminase small subunits (TerS_P) but not corresponding SaPI subunits (TerS_S) (123).
333 Ppi proteins are divided into two classes based on sequence that differ in helper phage specificity – Class I
334 interferes with $\Phi 80\alpha$ and $\Phi 11$, while Class II interferes with $\Phi 12$ (123). The PtiM-modulated PtiA and the PtiB
335 SaPI2 proteins inhibit expression of the LtrC-activated phage 80 late gene operon (packaging and lysis genes),
336 thus interfering with later steps of the helper phage life cycle (128). The SaPI particles then go on to infect new
337 *S. aureus* hosts, integrating their DNA into the chromosome instead of killing the cell. Helper phages and
338 SaPIs are thought to gain and lose resistance to each other in a 'Red Queen' scenario, given the observed
339 rapid co-evolution of their respective *dut* and *stl* genes (129). SaPIs are found throughout *Staphylococcus*
340 species and beyond; therefore, they may be a common strain-specific modifier of siphovirus infection potential.

341 **Other phage host range limiting factors**

342 Several uncommon or less well-understood mechanisms may contribute to phage host range limitation
343 in *Staphylococcus*. One abortive infection (Abi) system, the eukaryotic-like serine/threonine kinase Stk2, has
344 been characterized in *S. aureus* and *S. epidermidis* (130). In this case, siphovirus infection results in self-
345 induced killing of the host cell, preventing the amplification and spread of phages in the population. Stk2 was
346 found to be activated by a phage protein of unknown function and caused cell death by phosphorylating host
347 proteins involved in diverse core cellular functions. Only *S. epidermidis* RP62A and a few *S. aureus* strains
348 encode Stk2, however, suggesting limited genus-wide importance. The recent long-term evolution study on *S.*
349 *aureus* strain SA003 uncovered two genes involved in post-adsorption resistance to myovirus $\Phi SA012$ (53).
350 Missense mutations in guanylate kinase and the alpha subunit of DNA-dependent RNA polymerase conferred
351 resistance but not corresponding decreases in adsorption rate, suggesting some post-adsorption role in
352 resisting infection. More phage resistance systems likely remain undiscovered. A genome-wide association
353 study of 207 clinical MRSA strains and 12 phage preparations identified 167 gene families putatively
354 associated with phage-bacterial interactions (131). While these families included restriction-modification genes,

355 transcriptional regulators, and genes of prophage and SaPI origin, most were accessory gene families of
356 unknown function.

357 **Phage host range in *Staphylococcus* is determined by a hierarchical** 358 **combination of host factors**

359 In summary, we have described how host range of a *Staphylococcus* phage is determined by a
360 combination of both host and phage-encoded genes, as well as the epigenetic DNA methylation patterns
361 conferred on its DNA from the last strain it infected. Bacterial encoded factors can be conceived as affecting
362 host range at different levels within the species (**Figure 3**). At the highest level, most phages' target for
363 receptor binding (WTA) is highly conserved across *Staphylococcus* species. Strains with unusual WTAs, such
364 as *S. aureus* ST395 and CoNS strains with poly-GroP WTA (55, 56), would be expected to be genetically
365 isolated within the genus. Type I and IV R-M HsdS allotypes and capsule type are conserved between most
366 strains of the same CC but differ between isolates of different CC groups and thus contribute to defining host
367 range in a large subset of *S. aureus* strains. At the level of individual strains, inserted prophages and SaPIs,
368 Stk2, type II systems acquired by HGT, and other as yet unknown functions may all serve to limit host range.
369 We know even less about phage-encoded systems that counteract host resistance. The finding that lytic
370 phages (*Myoviridae* and *Podoviridae*) tend to have broader host ranges than *Siphoviridae* when challenged
371 against the same set of *Staphylococcus* strains suggests the former encode an array of uncharacterized genes
372 that work against host defenses.

373 **Future directions**

374 Although much progress has been made in the past five decades toward understanding the
375 mechanisms that define staphylococcal phage host range, numerous important questions remain. We need to
376 know more about species other than *S. aureus* and *S. epidermidis*, and even within these species, we need to
377 make sure that rarer and non-methicillin resistant strains are included in studies (132). We also need to ensure
378 that our collections reflect the true diversity of phages that infect *Staphylococcus* species. Even within the two
379 main species only a relatively small number of phages have been tested. This will lead us to consider the

380 questions of phage ecology when understanding what types of phages are found in different environments and
381 with what abundance.

382 Discovering novel phage resistance mechanisms would aid the effort to understand determinants of
383 host range. Many phage resistance mechanisms have been identified and characterized in other Gram-
384 positives and other bacteria generally but not in the staphylococci. Superinfection exclusion (Sie) and abortive
385 infection (Abi) systems, for example, are well-characterized in the lactococci (133–135). In addition, a recent
386 publication describes some 26 new anti-phage defense systems identified in bacteria (136), not including the
387 recently discovered bacteriophage exclusion (BREX) and defense island system associated with restriction-
388 modification (DISARM) phage defenses (137–139). Six of the ten verified, newly discovered anti-phage
389 defense systems (Thoeris, Hachiman, Gabija, Septu, Lamassu, and Kiwa) have orthologs in staphylococcal
390 genomes (136).

391 Understanding phage host range to the point that we can make accurate predictions based on the host
392 genome will be important for developing phage therapies against *Staphylococcus* strains. Ideally, cocktail
393 formulations for therapy consist of phages with broad, non-overlapping host ranges against the target species
394 (or clonal complex) to be treated. As there are many more genome sequences available than strains that can
395 be tested for sensitivity in the laboratory (e.g > 40,000 for *S. aureus*) (140), with a predictive model we could
396 run *in silico* tests on genome sequences to model the efficacy of the cocktail. With the potential for genome
397 sequencing to be used in the future as a primary clinical diagnostic, we could modify the cocktail to contain
398 phages that specifically target the bacterium causing the infection.

399 Knowledge of phage host range will also lead us to understand the fitness costs of resistance and its
400 potential trade-offs with virulence and antibiotic resistance of *Staphylococcus*. Strains with null mutations in
401 biosynthetic genes are rare, given WTA's roles in cell division, autolysis, virulence, and antibiotic resistance
402 (36, 37). Although knocking out the genes involved in the first two steps of WTA biosynthesis has no fitness
403 cost in *S. aureus* (at least in laboratory conditions) (141, 142), WTA has many critical physiological roles,
404 especially in environments subject to phage therapy. Staphylococcal WTA is required for nasal colonization
405 (141, 143), cell division (41, 43), regulating autolysis (144, 145), lysozyme resistance through cell wall
406 crosslinking (132, 146), resistance to cationic antimicrobial peptides and fatty acids (147, 148), and biofilm
407 formation (149). WTA-altered or negative phage-resistant mutants would in turn become less virulent (43) and

408 even antibiotic sensitive – highly unfit in the natural habitat colonizing mammalian hosts or in an infection site
409 subject to treatment. Given that methicillin resistance requires WTA (50), phage/beta-lactam combination
410 therapies could be particularly promising. Mutants resistant to either phage or beta-lactams would be sensitive
411 to the other treatment, assuming the infecting strain is sensitive to the phage treatment. Nonetheless, as we
412 note for host range, strains containing minor but fitness-neutral resistance mechanisms, such as R-M systems
413 – rather than costly mutations – may be the most recalcitrant to phage therapy. Staphylococcal phage
414 therapies must then overcome both immediate, emerging mutational resistance and intrinsic resistance
415 mechanisms (e.g., R-M systems) specific to strains or clonal complexes. These resistance limitations,
416 however, could be overcome by selecting phage host range mutants that escaped host resistance
417 mechanisms, thus isolating more useful phages that would form more effective phage cocktails (150, 151).

418 Phage-resistant mutants isolated so far, such as those described in the adsorption studies, were
419 typically selected in rich, aerated laboratory medium. The consequences for fitness of the same mutations
420 occurring during *in vivo* infection might be more severe. In addition, both the relevance of various resistance
421 mechanisms *in vivo* and the effect of strain genetic background on resistance selection - especially on a
422 species-wide scale – have been left unexamined in most previous work. One study in mammalian hosts
423 showed that environment altered phage transfer frequency and selection (152), leading to spread of prophage
424 and selection of phage resistance by superimmunity. In laboratory media, phage transfer frequency was lower
425 and spread of prophage was less pronounced (152). It will be important to know both how quickly and in which
426 loci mutations emerge as well as the more general distribution of resistance gene families.

427 Finally, it is interesting to consider what phage host range studies reveal about the hosts themselves.
428 Staphylococci seem to be unusual among Gram-positives in requiring conserved WTA receptors for
429 attachment and having no reported role for protein receptors. Differences in the outer surface of
430 *Staphylococcus* and/or a feature of the phage ecology within the genus requiring highly conserved receptors
431 may account for this fact. Another interesting question is why CRISPRs play a much-reduced role for
432 intercepting extrinsic phage DNA than R-M systems in this genus compared to other bacteria. It could be that
433 CRISPR systems have a finite capacity for carrying fragments of mobile genetic elements, while R-M systems
434 can attack a wider range of incoming DNA, relevant to rapidly evolving populations. Future studies that probe

435 these questions may reveal some of the differential evolutionary forces that shape the genomes of pathogenic
436 bacteria.

437 **Conclusions**

438 Staphylococcal phage resistance mechanisms have been identified at three stages of infection
439 (attachment, biosynthesis, and assembly) and regulate host range in a hierarchical manner depending on
440 mechanism conservation. Nonetheless, staphylococcal phage-bacterial interactions certainly present many
441 open questions that must be addressed to accurately develop and evaluate possible phage therapies. We
442 need further studies to objectively identify the contribution of individual phage resistance mechanisms to host
443 range. Such work would provide the information needed not only to formulate phage cocktails effective against
444 a wide variety of strains but also to overcome remaining obstacles to cocktail development (e.g., highly
445 effective R-M or Abi systems). Future studies relevant to phage therapy should also characterize phage
446 resistance development during infection and therapy as well as the effects of resistance on mutant fitness.
447 Taken together, this future work will inform the rational design of phage cocktails to treat staphylococcal
448 infections alone or in combination with antibiotics.

449 **Acknowledgments**

450 We thank Michelle Su and Robert Petit for critically reading the manuscript and providing helpful comments.
451 AGM was supported by the National Science Foundation (NSF) Graduate Research Fellowship Program
452 (GRFP). JAL was supported by the Medical Research Council (grant MR/P028322/1). TDR was supported by
453 the National Institutes of Health (NIH) grant R21 AI121860.

454 **References**

- 455 1. Kluytmans J, Belkum A van, Verbrugh H. 1997. Nasal carriage of *Staphylococcus aureus*: epidemiology,
456 underlying mechanisms, and associated risks. *Clin Microbiol Rev* 10:505–520.
- 457 2. Kong EF, Johnson JK, Jabra-Rizk MA. 2016. Community-Associated Methicillin-Resistant
458 *Staphylococcus aureus*: An Enemy amidst Us. *PLOS Pathogens* 12:e1005837.

- 459 3. Bagnoli F, Bertholet S, Grandi G. 2012. Inferring Reasons for the Failure of *Staphylococcus aureus*
460 Vaccines in Clinical Trials. *Front Cell Infect Microbiol* 2.
- 461 4. Wang I-N, Smith DL, Young R. 2000. Holins: The Protein Clocks of Bacteriophage Infections. *Annu Rev*
462 *Microbiol* 54:799–825.
- 463 5. Young R, Bläsi U. 1995. Holins: form and function in bacteriophage lysis. *FEMS Microbiology Reviews*
464 17:195–205.
- 465 6. Loessner MJ. 2005. Bacteriophage endolysins — current state of research and applications. *Current*
466 *Opinion in Microbiology* 8:480–487.
- 467 7. Pirisi A. 2000. Phage therapy—advantages over antibiotics? *The Lancet* 356:1418.
- 468 8. Nobrega FL, Costa AR, Kluskens LD, Azeredo J. 2015. Revisiting phage therapy: new applications for
469 old resources. *Trends in Microbiology* 23:185–191.
- 470 9. Stone R. 2002. Stalin's Forgotten Cure. *Science* 298:728–731.
- 471 10. Bradbury J. 2004. "My enemy's enemy is my friend." *The Lancet* 363:624–625.
- 472 11. Kutter E, De Vos D, Gvasalia G, Alavidze Z, Gogokhia L, Kuhl S, Abedon ST. 2010. Phage Therapy in
473 Clinical Practice: Treatment of Human Infections. *Current Pharmaceutical Biotechnology* 11:69–86.
- 474 12. Kutateladze M, Adamia R. 2010. Bacteriophages as potential new therapeutics to replace or supplement
475 antibiotics. *Trends in Biotechnology* 28:591–595.
- 476 13. Sulakvelidze A, Alavidze Z, Morris JG. 2001. Bacteriophage Therapy. *Antimicrob Agents Chemother*
477 45:649–659.
- 478 14. Lu TK, Koeris MS. 2011. The next generation of bacteriophage therapy. *Current Opinion in Microbiology*
479 14:524–531.
- 480 15. Ly-Chatain MH. 2014. The factors affecting effectiveness of treatment in phages therapy. *Front Microbiol*
481 5.

- 482 16. Jault P, Leclerc T, Jennes S, Pirnay JP, Que Y-A, Resch G, Rousseau AF, Ravat F, Carsin H, Le Floch
483 R, Schaal JV, Soler C, Fevre C, Arnaud I, Bretaudeau L, Gabard J. 2018. Efficacy and tolerability of a
484 cocktail of bacteriophages to treat burn wounds infected by *Pseudomonas aeruginosa* (PhagoBurn): a
485 randomised, controlled, double-blind phase 1/2 trial. *The Lancet Infectious Diseases*.
- 486 17. Xia G, Wolz C. 2014. Phages of *Staphylococcus aureus* and their impact on host evolution. *Infection,
487 Genetics and Evolution* 21:593–601.
- 488 18. Deghorain M, Van Melder L. 2012. The Staphylococci Phages Family: An Overview. *Viruses* 4:3316–
489 3335.
- 490 19. Alves DR, Gaudion A, Bean JE, Esteban PP, Arnot TC, Harper DR, Kot W, Hansen LH, Enright MC,
491 Jenkins ATA. 2014. Combined Use of Bacteriophage K and a Novel Bacteriophage To Reduce
492 *Staphylococcus aureus* Biofilm Formation. *Appl Environ Microbiol* 80:6694–6703.
- 493 20. Hsieh S-E, Lo H-H, Chen S-T, Lee M-C, Tseng Y-H. 2011. Wide Host Range and Strong Lytic Activity of
494 *Staphylococcus aureus* Lytic Phage Stau2. *Appl Environ Microbiol* 77:756–761.
- 495 21. Synnott AJ, Kuang Y, Kurimoto M, Yamamichi K, Iwano H, Tanji Y. 2009. Isolation from Sewage Influent
496 and Characterization of Novel *Staphylococcus aureus* Bacteriophages with Wide Host Ranges and
497 Potent Lytic Capabilities. *Appl Environ Microbiol* 75:4483–4490.
- 498 22. O'Flaherty S, Ross RP, Meaney W, Fitzgerald GF, Elbreki MF, Coffey A. 2005. Potential of the
499 Polyvalent Anti-*Staphylococcus* Bacteriophage K for Control of Antibiotic-Resistant *Staphylococci* from
500 Hospitals. *Appl Environ Microbiol* 71:1836–1842.
- 501 23. Gutiérrez D, Briers Y, Rodríguez-Rubio L, Martínez B, Rodríguez A, Lavigne R, García P. 2015. Role of
502 the Pre-neck Appendage Protein (Dpo7) from Phage vB_SepiS-phiPLA7 as an Anti-biofilm Agent in
503 *Staphylococcal* Species. *Front Microbiol* 6.
- 504 24. Matsuzaki S, Yasuda M, Nishikawa H, Kuroda M, Ujihara T, Shuin T, Shen Y, Jin Z, Fujimoto S,
505 Nasimuzzaman MD, Wakiguchi H, Sugihara S, Sugiura T, Koda S, Muraoka A, Imai S. 2003.

- 506 Experimental Protection of Mice against Lethal *Staphylococcus aureus* Infection by Novel Bacteriophage
507 ϕ MR11. *J Infect Dis* 187:613–624.
- 508 25. Wills QF, Kerrigan C, Soothill JS. 2005. Experimental Bacteriophage Protection against *Staphylococcus*
509 *aureus* Abscesses in a Rabbit Model. *Antimicrobial Agents and Chemotherapy* 49:1220–1221.
- 510 26. Verstappen KM, Tulinski P, Duim B, Fluit AC, Carney J, Nes A van, Wagenaar JA. 2016. The
511 Effectiveness of Bacteriophages against Methicillin-Resistant *Staphylococcus aureus* ST398 Nasal
512 Colonization in Pigs. *PLOS ONE* 11:e0160242.
- 513 27. Chen J, Quiles-Puchalt N, Chiang YN, Bacigalupe R, Fillol-Salom A, Chee MSJ, Fitzgerald JR, Penadés
514 JR. 2018. Genome hypermobility by lateral transduction. *Science* 362:207–212.
- 515 28. Novick RP, Christie GE, Penadés JR. 2010. The phage-related chromosomal islands of Gram-positive
516 bacteria. *Nat Rev Microbiol* 8:541–551.
- 517 29. McCarthy AJ, Witney AA, Lindsay JA. 2012. *Staphylococcus aureus* Temperate Bacteriophage:
518 Carriage and Horizontal Gene Transfer is Lineage Associated. *Front Cell Infect Microbiol* 2.
- 519 30. Carroll JD, Cafferkey MT, Coleman DC. 1993. Serotype F double- and triple-converting phage
520 insertionally inactivate the *Staphylococcus aureus* β -toxin determinant by a common molecular
521 mechanism. *FEMS Microbiology Letters* 106:147–155.
- 522 31. Iandolo JJ, Worrell V, Groicher KH, Qian Y, Tian R, Kenton S, Dorman A, Ji H, Lin S, Loh P, Qi S, Zhu
523 H, Roe BA. 2002. Comparative analysis of the genomes of the temperate bacteriophages ϕ 11, ϕ 12 and
524 ϕ 13 of *Staphylococcus aureus* 8325. *Gene* 289:109–118.
- 525 32. Labrie SJ, Samson JE, Moineau S. 2010. Bacteriophage resistance mechanisms. *Nat Rev Microbiol*
526 8:317–327.
- 527 33. Hyman P, Abedon ST. 2010. Bacteriophage Host Range and Bacterial Resistance. *Advances in Applied*
528 *Microbiology* 70:217–248.

- 529 34. Allison GE, Klaenhammer TR. 1998. Phage Resistance Mechanisms in Lactic Acid Bacteria.
530 International Dairy Journal 8:207–226.
- 531 35. Xia G, Corrigan RM, Winstel V, Goerke C, Gründling A, Peschel A. 2011. Wall Teichoic Acid-Dependent
532 Adsorption of Staphylococcal Siphovirus and Myovirus. J Bacteriol 193:4006–4009.
- 533 36. Xia G, Kohler T, Peschel A. 2010. The wall teichoic acid and lipoteichoic acid polymers of
534 Staphylococcus aureus. International Journal of Medical Microbiology 300:148–154.
- 535 37. Brown S, Santa Maria JP, Walker S. 2013. Wall Teichoic Acids of Gram-Positive Bacteria. Annu Rev
536 Microbiol 67.
- 537 38. Shaw DRD, Mirelman D, Chatterjee AN, Park JT. 1970. Ribitol Teichoic Acid Synthesis in
538 Bacteriophage-resistant Mutants of Staphylococcus aureus H. J Biol Chem 245:5101–5106.
- 539 39. Shaw DRD, Chatterjee AN. 1971. O-Acetyl Groups as a Component of the Bacteriophage Receptor on
540 Staphylococcus aureus Cell Walls. J Bacteriol 108:584–585.
- 541 40. Chatterjee AN. 1969. Use of Bacteriophage-resistant Mutants to Study the Nature of the Bacteriophage
542 Receptor Site of Staphylococcus aureus. J Bacteriol 98:519–527.
- 543 41. Chatterjee AN, Mirelman D, Singer HJ, Park JT. 1969. Properties of a Novel Pleiotropic Bacteriophage-
544 Resistant Mutant of Staphylococcus aureus H1. J Bacteriol 100:846–853.
- 545 42. NCTC 3000 Project.
- 546 43. Capparelli R, Nocerino N, Lanzetta R, Silipo A, Amoresano A, Giangrande C, Becker K, Blaiotta G,
547 Evidente A, Cimmino A, Iannaccone M, Parlato M, Medaglia C, Roperto S, Roperto F, Ramunno L,
548 Iannelli D. 2010. Bacteriophage-Resistant Staphylococcus aureus Mutant Confers Broad Immunity
549 against Staphylococcal Infection in Mice. PLOS ONE 5:e11720.
- 550 44. Nair D, Memmi G, Hernandez D, Bard J, Beaume M, Gill S, Francois P, Cheung AL. 2011. Whole-
551 Genome Sequencing of Staphylococcus aureus Strain RN4220, a Key Laboratory Strain Used in

- 552 Virulence Research, Identifies Mutations That Affect Not Only Virulence Factors but Also the Fitness of
553 the Strain. *Journal of Bacteriology* 193:2332–2335.
- 554 45. Li X, Gerlach D, Du X, Larsen J, Stegger M, Kühner P, Peschel A, Xia G, Winstel V. 2015. An accessory
555 wall teichoic acid glycosyltransferase protects *Staphylococcus aureus* from the lytic activity of
556 Podoviridae. *Sci Rep* 5.
- 557 46. Xia G, Maier L, Sanchez-Carballo P, Li M, Otto M, Holst O, Peschel A. 2010. Glycosylation of wall
558 teichoic acid in *Staphylococcus aureus* by TarM. *J Biol Chem* 285:13405–13415.
- 559 47. Sobhanifar S, Worrall LJ, Gruninger RJ, Wasney GA, Blaukopf M, Baumann L, Lameignere E,
560 Solomonson M, Brown ED, Withers SG, Strynadka NCJ. 2015. Structure and mechanism of
561 *Staphylococcus aureus* TarM, the wall teichoic acid α -glycosyltransferase. *Proc Natl Acad Sci USA*
562 112:E576-585.
- 563 48. Sobhanifar S, Worrall LJ, King DT, Wasney GA, Baumann L, Gale RT, Nosella M, Brown ED, Withers
564 SG, Strynadka NCJ. 2016. Structure and Mechanism of *Staphylococcus aureus* TarS, the Wall Teichoic
565 Acid β -glycosyltransferase Involved in Methicillin Resistance. *PLOS Pathogens* 12:e1006067.
- 566 49. Li X, Koç C, Kühner P, Stierhof Y-D, Krismer B, Enright MC, Penadés JR, Wolz C, Stehle T, Cambillau
567 C, Peschel A, Xia G. 2016. An essential role for the baseplate protein Gp45 in phage adsorption to
568 *Staphylococcus aureus*. *Sci Rep* 6.
- 569 50. Brown S, Xia G, Luhachack LG, Campbell J, Meredith TC, Chen C, Winstel V, Gekeler C, Irazoqui JE,
570 Peschel A, Walker S. 2012. Methicillin resistance in *Staphylococcus aureus* requires glycosylated wall
571 teichoic acids. *PNAS* 109:18909–18914.
- 572 51. Qian Z, Yin Y, Zhang Y, Lu L, Li Y, Jiang Y. 2006. Genomic characterization of ribitol teichoic acid
573 synthesis in *Staphylococcus aureus*: genes, genomic organization and gene duplication. *BMC Genomics*
574 7:74.

- 575 52. Cerca N, Oliveira R, Azeredo J. Susceptibility of *Staphylococcus epidermidis* planktonic cells and
576 biofilms to the lytic action of staphylococcus bacteriophage K. *Letters in Applied Microbiology* 45:313–
577 317.
- 578 53. Azam AH, Hoshiga F, Takeuchi I, Miyanaga K, Tanji Y. 2018. Analysis of phage resistance in
579 *Staphylococcus aureus* SA003 reveals different binding mechanisms for the closely related Twort-like
580 phages ϕ SA012 and ϕ SA039. *Appl Microbiol Biotechnol* 1–15.
- 581 54. Gerlach D, Guo Y, Castro CD, Kim S-H, Schlatterer K, Xu F-F, Pereira C, Seeberger PH, Ali S, Codée J,
582 Sirisarn W, Schulte B, Wolz C, Larsen J, Molinaro A, Lee BL, Xia G, Stehle T, Peschel A. 2018.
583 Methicillin-resistant *Staphylococcus aureus* alters cell wall glycosylation to evade immunity. *Nature* 1.
- 584 55. Winstel V, Sanchez-Carballo P, Holst O, Xia G, Peschel A. 2014. Biosynthesis of the Unique Wall
585 Teichoic Acid of *Staphylococcus aureus* Lineage ST395. *mBio* 5:e00869-14.
- 586 56. Winstel V, Liang C, Sanchez-Carballo P, Steglich M, Munar M, Bröker BM, Penadés JR, Nübel U, Holst
587 O, Dandekar T, Peschel A, Xia G. 2013. Wall teichoic acid structure governs horizontal gene transfer
588 between major bacterial pathogens. *Nature Communications* 4:2345.
- 589 57. Kaneko J, Narita-Yamada S, Wakabayashi Y, Kamio Y. 2009. Identification of ORF636 in Phage ϕ SLT
590 Carrying Panton-Valentine Leukocidin Genes, Acting as an Adhesion Protein for a
591 Poly(Glycerophosphate) Chain of Lipoteichoic Acid on the Cell Surface of *Staphylococcus aureus*. *J*
592 *Bacteriol* 191:4674–4680.
- 593 58. Baptista C, Santos MA, São-José C. 2008. Phage SPP1 reversible adsorption to *Bacillus subtilis* cell
594 wall teichoic acids accelerates virus recognition of membrane receptor YueB. *J Bacteriol* 190:4989–
595 4996.
- 596 59. Duerkop BA, Huo W, Bhardwaj P, Palmer KL, Hooper LV. 2016. Molecular Basis for Lytic Bacteriophage
597 Resistance in Enterococci. *mBio* 7:e01304-16.
- 598 60. Mazmanian SK, Liu G, Ton-That H, Schneewind O. 1999. *Staphylococcus aureus* Sortase, an Enzyme
599 that Anchors Surface Proteins to the Cell Wall. *Science* 285:760–763.

- 600 61. Ton-That H, Liu G, Mazmanian SK, Faull KF, Schneewind O. 1999. Purification and characterization of
601 sortase, the transpeptidase that cleaves surface proteins of *Staphylococcus aureus* at the LPXTG motif.
602 PNAS 96:12424–12429.
- 603 62. Baba T, Bae T, Schneewind O, Takeuchi F, Hiramatsu K. 2008. Genome Sequence of *Staphylococcus*
604 *aureus* Strain Newman and Comparative Analysis of Staphylococcal Genomes: Polymorphism and
605 Evolution of Two Major Pathogenicity Islands. *Journal of Bacteriology* 190:300–310.
- 606 63. Bae T, Baba T, Hiramatsu K, Schneewind O. 2006. Prophages of *Staphylococcus aureus* Newman and
607 their contribution to virulence. *Molecular Microbiology* 62:1035–1047.
- 608 64. Nordström K, Forsgren A. 1974. Effect of Protein A on Adsorption of Bacteriophages to *Staphylococcus*
609 *aureus*. *J Virol* 14:198–202.
- 610 65. Wilkinson BJ, Holmes KM. 1979. *Staphylococcus aureus* cell surface: capsule as a barrier to
611 bacteriophage adsorption. *Infect Immun* 23:549–552.
- 612 66. Sutra L, Audurier A, Poutrel B. 1988. Relationship between capsular types 5 and 8 and phage types in
613 *Staphylococcus aureus* isolates from cow, goat and ewe milk. *FEMS Microbiol Lett* 55:83–85.
- 614 67. Sompolinsky D, Samra Z, Karakawa WW, Vann WF, Schneerson R, Malik Z. 1985. Encapsulation and
615 capsular types in isolates of *Staphylococcus aureus* from different sources and relationship to phage
616 types. *J Clin Microbiol* 22:828–834.
- 617 68. O’Riordan K, Lee JC. 2004. *Staphylococcus aureus* Capsular Polysaccharides. *Clin Microbiol Rev*
618 17:218–234.
- 619 69. Ohshima Y, Schumacher-Perdreau F, Peters G, Pulverer G. 1988. The role of capsule as a barrier to
620 bacteriophage adsorption in an encapsulated *Staphylococcus simulans* strain. *Med Microbiol Immunol*
621 177:229–233.
- 622 70. O’Gara JP. 2007. *ica* and beyond: biofilm mechanisms and regulation in *Staphylococcus epidermidis*
623 and *Staphylococcus aureus*. *FEMS Microbiol Lett* 270:179–188.

- 624 71. Götz F. 2002. Staphylococcus and biofilms. *Molecular Microbiology* 43:1367–1378.
- 625 72. McCourt J, O'Halloran DP, McCarthy H, O'Gara JP, Geoghegan JA. 2014. Fibronectin-binding proteins
626 are required for biofilm formation by community-associated methicillin-resistant *Staphylococcus aureus*
627 strain LAC. *FEMS Microbiol Lett* 353:157–164.
- 628 73. O'Neill E, Pozzi C, Houston P, Humphreys H, Robinson DA, Loughman A, Foster TJ, O'Gara JP. 2008.
629 A Novel *Staphylococcus aureus* Biofilm Phenotype Mediated by the Fibronectin-Binding Proteins,
630 FnBPA and FnBPB. *Journal of Bacteriology* 190:3835–3850.
- 631 74. Geoghegan JA, Corrigan RM, Gruszka DT, Speziale P, O'Gara JP, Potts JR, Foster TJ. 2010. Role of
632 Surface Protein SasG in Biofilm Formation by *Staphylococcus aureus*. *Journal of Bacteriology*
633 192:5663–5673.
- 634 75. Kelly D, McAuliffe O, Ross RP, Coffey A. Prevention of *Staphylococcus aureus* biofilm formation and
635 reduction in established biofilm density using a combination of phage K and modified derivatives. *Letters*
636 *in Applied Microbiology* 54:286–291.
- 637 76. Curtin JJ, Donlan RM. 2006. Using Bacteriophages To Reduce Formation of Catheter-Associated
638 Biofilms by *Staphylococcus epidermidis*. *Antimicrob Agents Chemother* 50:1268–1275.
- 639 77. Gutiérrez D, Martínez B, Rodríguez A, García P. 2012. Genomic characterization of two *Staphylococcus*
640 *epidermidis* bacteriophages with anti-biofilm potential. *BMC Genomics* 13:228.
- 641 78. Resch A, Fehrenbacher B, Eisele K, Schaller M, Götz F. Phage release from biofilm and planktonic
642 *Staphylococcus aureus* cells. *FEMS Microbiology Letters* 252:89–96.
- 643 79. Kameyama L, Fernandez L, Calderon J, Ortiz-Rojas A, Patterson TA. 1999. Characterization of Wild
644 Lambdoid Bacteriophages: Detection of a Wide Distribution of Phage Immunity Groups and Identification
645 of a Nus-Dependent, Nonlambdoid Phage Group. *Virology* 263:100–111.
- 646 80. Berngruber TW, Weissing FJ, Gandon S. 2010. Inhibition of Superinfection and the Evolution of Viral
647 Latency. *J Virol* 84:10200–10208.

- 648 81. Goerke C, Pantucek R, Holtfreter S, Schulte B, Zink M, Grumann D, Bröker BM, Doskar J, Wolz C.
649 2009. Diversity of Prophages in Dominant *Staphylococcus aureus* Clonal Lineages. *Journal of*
650 *Bacteriology* 191:3462–3468.
- 651 82. Wilson GG, Murray NE. 1991. Restriction and Modification Systems. *Annual Review of Genetics*
652 25:585–627.
- 653 83. Sadykov MR. 2014. Restriction–Modification Systems as a Barrier for Genetic Manipulation of
654 *Staphylococcus aureus*, p. 9–23. *In* Bose, JL (ed.), *The Genetic Manipulation of Staphylococci*. Springer
655 New York, New York, NY.
- 656 84. Monk IR, Shah IM, Xu M, Tan M-W, Foster TJ. 2012. Transforming the Untransformable: Application of
657 Direct Transformation To Manipulate Genetically *Staphylococcus aureus* and *Staphylococcus*
658 *epidermidis*. *mBio* 3:e00277-11.
- 659 85. Monk IR, Foster TJ. 2012. Genetic manipulation of *Staphylococci* – Breaking through the barrier. *Front*
660 *Cell Infect Microbiol* 2.
- 661 86. Costa SK, Donegan NP, Corvaglia A-R, François P, Cheung AL. 2017. Bypassing the Restriction
662 System To Improve Transformation of *Staphylococcus epidermidis*. *J Bacteriol* 199.
- 663 87. Roberts RJ, Vincze T, Posfai J, Macelis D. 2015. REBASE—a database for DNA restriction and
664 modification: enzymes, genes and genomes. *Nucleic Acids Res* 43:D298–D299.
- 665 88. Waldron DE, Lindsay JA. 2006. *Sau*1: a Novel Lineage-Specific Type I Restriction-Modification System
666 That Blocks Horizontal Gene Transfer into *Staphylococcus aureus* and between *S. aureus* Isolates of
667 Different Lineages. *J Bacteriol* 188:5578–5585.
- 668 89. Roberts GA, Houston PJ, White JH, Chen K, Stephanou AS, Cooper LP, Dryden DTF, Lindsay JA.
669 2013. Impact of target site distribution for Type I restriction enzymes on the evolution of methicillin-
670 resistant *Staphylococcus aureus* (MRSA) populations. *Nucleic Acids Res* 41:7472–7484.

- 671 90. Cooper LP, Roberts GA, White JH, Luyten YA, Bower EKM, Morgan RD, Roberts RJ, Lindsay JA,
672 Dryden DTF. 2017. DNA target recognition domains in the Type I restriction and modification systems of
673 *Staphylococcus aureus*. *Nucleic Acids Res* 45:3395–3406.
- 674 91. Stegger M, Lindsay JA, Moodley A, Skov R, Broens EM, Guardabassi L. 2011. Rapid PCR Detection of
675 *Staphylococcus aureus* Clonal Complex 398 by Targeting the Restriction-Modification System Carrying
676 *sau1-hsdS1*. *Journal of Clinical Microbiology* 49:732–734.
- 677 92. Xu S, Corvaglia AR, Chan S-H, Zheng Y, Linder P. 2011. A type IV modification-dependent restriction
678 enzyme *SauUSI* from *Staphylococcus aureus* subsp. *aureus* USA300. *Nucleic Acids Res* 39:5597–5610.
- 679 93. Veiga H, Pinho MG. 2009. Inactivation of the *SauI* Type I Restriction-Modification System Is Not
680 Sufficient To Generate *Staphylococcus aureus* Strains Capable of Efficiently Accepting Foreign DNA.
681 *Appl Environ Microbiol* 75:3034–3038.
- 682 94. Seeber S, Kessler C, Götz F. 1990. Cloning, expression and characterization of the *Sau3AI* restriction
683 and modification genes in *Staphylococcus carnosus* TM300. *Gene* 94:37–43.
- 684 95. Sussenbach JS, Steenbergh PH, Rost JA, van Leeuwen WJ, van Embden JD. 1978. A second site-
685 specific restriction endonuclease from *Staphylococcus aureus*. *Nucleic Acids Res* 5:1153–1163.
- 686 96. Dempsey RM. 2005. *Sau42I*, a *BcglI*-like restriction-modification system encoded by the *Staphylococcus*
687 *aureus* quadruple-converting phage 42. *Microbiology* 151:1301–1311.
- 688 97. Lebenka Ai, Rachkus I. 1989. [DNA-methylase *Sau 3A*: isolation and various properties]. *Biokhimiia*
689 54:1009–1014.
- 690 98. Tock MR, Dryden DT. 2005. The biology of restriction and anti-restriction. *Current Opinion in*
691 *Microbiology* 8:466–472.
- 692 99. O'Flaherty S, Coffey A, Edwards R, Meaney W, Fitzgerald GF, Ross RP. 2004. Genome of
693 *Staphylococcal Phage K*: a New Lineage of *Myoviridae* Infecting Gram-Positive Bacteria with a Low
694 G+C Content. *Journal of Bacteriology* 186:2862–2871.

- 695 100. Iida S, Streiff MB, Bickle TA, Arber W. 1987. Two DNA antirestriction systems of bacteriophage P1,
696 darA, and darB: characterization of darA- phages. *Virology* 157:156–166.
- 697 101. Kruger DH, Barcak GJ, Reuter M, Smith HO. 1988. EcoRII can be activated to cleave refractory DNA
698 recognition sites. *Nucleic Acids Res* 16:3997–4008.
- 699 102. Zabeau M, Friedman S, Van Montagu M, Schell J. 1980. The ral gene of phage λ . *Molec Gen Genet*
700 179:63–73.
- 701 103. Studier FW, Movva NR. 1976. SAMase gene of bacteriophage T3 is responsible for overcoming host
702 restriction. *Journal of Virology* 19:136–145.
- 703 104. Zavilgelsky GB, Kotova VY, Rastorguev SM. 2008. Comparative analysis of anti-restriction activities of
704 ArdA (Collb-P9) and Ocr (T7) proteins. *Biochemistry Moscow* 73:906.
- 705 105. McMahon SA, Roberts GA, Johnson KA, Cooper LP, Liu H, White JH, Carter LG, Sanghvi B, Oke M,
706 Walkinshaw MD, Blakely GW, Naismith JH, Dryden DTF. 2009. Extensive DNA mimicry by the ArdA
707 anti-restriction protein and its role in the spread of antibiotic resistance. *Nucleic Acids Res* 37:4887–
708 4897.
- 709 106. Sansevere EA, Robinson DA. 2017. Staphylococci on ICE: Overlooked agents of horizontal gene
710 transfer. *Mobile Genetic Elements* 7:1–10.
- 711 107. Barrangou R, Marraffini LA. 2014. CRISPR-Cas Systems: Prokaryotes Upgrade to Adaptive Immunity.
712 *Molecular Cell* 54:234–244.
- 713 108. Barrangou R, Fremaux C, Deveau H, Richards M, Boyaval P, Moineau S, Romero DA, Horvath P. 2007.
714 CRISPR Provides Acquired Resistance Against Viruses in Prokaryotes. *Science* 315:1709–1712.
- 715 109. Li Q, Xie X, Yin K, Tang Y, Zhou X, Chen Y, Xia J, Hu Y, Ingmer H, Li Y, Jiao X. 2016. Characterization
716 of CRISPR-Cas system in clinical *Staphylococcus epidermidis* strains revealed its potential association
717 with bacterial infection sites. *Microbiological Research* 193:103–110.

- 718 110. Yang S, Liu J, Shao F, Wang P, Duan G, Yang H. 2015. Analysis of the features of 45 identified
719 CRISPR loci in 32 *Staphylococcus aureus*. *Biochemical and Biophysical Research Communications*
720 464:894–900.
- 721 111. Carte J, Wang R, Li H, Terns RM, Terns MP. 2008. Cas6 is an endoribonuclease that generates guide
722 RNAs for invader defense in prokaryotes. *Genes Dev* 22:3489–3496.
- 723 112. Jiang F, Doudna JA. 2015. The structural biology of CRISPR-Cas systems. *Current Opinion in Structural*
724 *Biology* 30:100–111.
- 725 113. Gasiunas G, Barrangou R, Horvath P, Siksnys V. 2012. Cas9–crRNA ribonucleoprotein complex
726 mediates specific DNA cleavage for adaptive immunity in bacteria. *PNAS* 109:E2579–E2586.
- 727 114. Kinnevey PM, Shore AC, Brennan GI, Sullivan DJ, Ehricht R, Monecke S, Slickers P, Coleman DC.
728 2013. Emergence of Sequence Type 779 Methicillin-Resistant *Staphylococcus aureus* Harboring a
729 Novel Pseudo Staphylococcal Cassette Chromosome mec (SCCmec)-SCC-SCCCRISPR Composite
730 Element in Irish Hospitals. *Antimicrob Agents Chemother* 57:524–531.
- 731 115. Gill SR, Fouts DE, Archer GL, Mongodin EF, DeBoy RT, Ravel J, Paulsen IT, Kolonay JF, Brinkac L,
732 Beanan M, Dodson RJ, Daugherty SC, Madupu R, Angiuoli SV, Durkin AS, Haft DH, Vamathevan J,
733 Khouri H, Utterback T, Lee C, Dimitrov G, Jiang L, Qin H, Weidman J, Tran K, Kang K, Hance IR,
734 Nelson KE, Fraser CM. 2005. Insights on Evolution of Virulence and Resistance from the Complete
735 Genome Analysis of an Early Methicillin-Resistant *Staphylococcus aureus* Strain and a Biofilm-
736 Producing Methicillin-Resistant *Staphylococcus epidermidis* Strain. *J Bacteriol* 187:2426–2438.
- 737 116. Marraffini LA, Sontheimer EJ. 2008. CRISPR Interference Limits Horizontal Gene Transfer in
738 *Staphylococci* by Targeting DNA. *Science* 322:1843–1845.
- 739 117. Bondy-Denomy J, Garcia B, Strum S, Du M, Rollins MF, Hidalgo-Reyes Y, Wiedenheft B, Maxwell KL,
740 Davidson AR. 2015. Multiple mechanisms for CRISPR–Cas inhibition by anti-CRISPR proteins. *Nature*
741 526:136–139.

- 742 118. Pawluk A, Staals RHJ, Taylor C, Watson BNJ, Saha S, Fineran PC, Maxwell KL, Davidson AR. 2016.
743 Inactivation of CRISPR-Cas systems by anti-CRISPR proteins in diverse bacterial species. *Nature*
744 *Microbiology* 1:16085.
- 745 119. Maxwell KL. 2016. Phages Fight Back: Inactivation of the CRISPR-Cas Bacterial Immune System by
746 Anti-CRISPR Proteins. *PLOS Pathogens* 12:e1005282.
- 747 120. Rauch BJ, Silvis MR, Hultquist JF, Waters CS, McGregor MJ, Krogan NJ, Bondy-Denomy J. 2017.
748 Inhibition of CRISPR-Cas9 with Bacteriophage Proteins. *Cell* 168:150-158.e10.
- 749 121. Lindsay JA, Ruzin A, Ross HF, Kurepina N, Novick RP. 1998. The gene for toxic shock toxin is carried
750 by a family of mobile pathogenicity islands in *Staphylococcus aureus*. *Molecular Microbiology* 29:527–
751 543.
- 752 122. Seed KD. 2015. Battling Phages: How Bacteria Defend against Viral Attack. *PLOS Pathogens*
753 11:e1004847.
- 754 123. Ram G, Chen J, Kumar K, Ross HF, Ubeda C, Damle PK, Lane KD, Penadés JR, Christie GE, Novick
755 RP. 2012. Staphylococcal pathogenicity island interference with helper phage reproduction is a
756 paradigm of molecular parasitism. *PNAS* 109:16300–16305.
- 757 124. Ruzin A, Lindsay J, Novick RP. 2001. Molecular genetics of SaPI1 – a mobile pathogenicity island in
758 *Staphylococcus aureus*. *Molecular Microbiology* 41:365–377.
- 759 125. Damle PK, Wall EA, Spilman MS, Dearborn AD, Ram G, Novick RP, Dokland T, Christie GE. 2012. The
760 roles of SaPI1 proteins gp7 (CpmA) and gp6 (CpmB) in capsid size determination and helper phage
761 interference. *Virology* 432:277–282.
- 762 126. Ubeda Carles, Maiques Elisa, Barry Peter, Matthews Avery, Tormo María Ángeles, Lasa Íñigo, Novick
763 Richard P., Penadés José R. 2007. SaPI mutations affecting replication and transfer and enabling
764 autonomous replication in the absence of helper phage. *Molecular Microbiology* 67:493–503.

- 765 127. Poliakov A, Chang JR, Spilman MS, Damle PK, Christie GE, Mobley JA, Dokland T. 2008. Capsid size
766 determination by *Staphylococcus aureus* pathogenicity island SaPI1 involves specific incorporation of
767 SaPI1 proteins into procapsids. *J Mol Biol* 380:465–475.
- 768 128. Ram G, Chen J, Ross HF, Novick RP. 2014. Precisely modulated pathogenicity island interference with
769 late phage gene transcription. *Proc Natl Acad Sci U S A* 111:14536–14541.
- 770 129. Frígols B, Quiles-Puchalt N, Mir-Sanchis I, Donderis J, Elena SF, Buckling A, Novick RP, Marina A,
771 Penadés JR. 2015. Virus Satellites Drive Viral Evolution and Ecology. *PLoS Genet* 11.
- 772 130. Depardieu F, Didier J-P, Bernheim A, Sherlock A, Molina H, Duclos B, Bikard D. 2016. A Eukaryotic-like
773 Serine/Threonine Kinase Protects *Staphylococci* against Phages. *Cell Host & Microbe* 20:471–481.
- 774 131. Zschach H, Larsen MV, Hasman H, Westh H, Nielsen M, Międzybrodzki R, Jończyk-Matysiak E, Weber-
775 Dąbrowska B, Górski A. 2018. Use of a Regression Model to Study Host-Genomic Determinants of
776 Phage Susceptibility in MRSA. *Antibiotics (Basel)* 7.
- 777 132. Atilano ML, Pereira PM, Yates J, Reed P, Veiga H, Pinho MG, Filipe SR. 2010. Teichoic acids are
778 temporal and spatial regulators of peptidoglycan cross-linking in *Staphylococcus aureus*. *PNAS*
779 107:18991–18996.
- 780 133. Mahony J, McGrath S, Fitzgerald GF, van Sinderen D. 2008. Identification and Characterization of
781 Lactococcal-Prophage-Carried Superinfection Exclusion Genes. *Appl Environ Microbiol* 74:6206–6215.
- 782 134. Sun X, Göhler A, Heller KJ, Neve H. 2006. The *Itp* gene of temperate *Streptococcus thermophilus* phage
783 TP-J34 confers superinfection exclusion to *Streptococcus thermophilus* and *Lactococcus lactis*. *Virology*
784 350:146–157.
- 785 135. Chopin M-C, Chopin A, Bidnenko E. 2005. Phage abortive infection in lactococci: variations on a theme.
786 *Current Opinion in Microbiology* 8:473–479.
- 787 136. Doron S, Melamed S, Ofir G, Leavitt A, Lopatina A, Keren M, Amitai G, Sorek R. 2018. Systematic
788 discovery of antiphage defense systems in the microbial pangenome. *Science* eaar4120.

- 789 137. Goldfarb T, Sberro H, Weinstock E, Cohen O, Doron S, Charpak-Amikam Y, Afik S, Ofir G, Sorek R.
790 2015. BREX is a novel phage resistance system widespread in microbial genomes. *The EMBO Journal*
791 34:169–183.
- 792 138. Barrangou R, van der Oost J. 2015. Bacteriophage exclusion, a new defense system. *EMBO J* 34:134–
793 135.
- 794 139. Ofir G, Melamed S, Sberro H, Mukamel Z, Silverman S, Yaakov G, Doron S, Sorek R. 2018. DISARM is
795 a widespread bacterial defence system with broad anti-phage activities. *Nat Microbiol* 3:90–98.
- 796 140. Petit III RA, Read TD. 2018. *Staphylococcus aureus* viewed from the perspective of 40,000+ genomes.
797 *PeerJ* 6:e5261.
- 798 141. Weidenmaier C, Kokai-Kun JF, Kristian SA, Chanturiya T, Kalbacher H, Gross M, Nicholson G,
799 Neumeister B, Mond JJ, Peschel A. 2004. Role of teichoic acids in *Staphylococcus aureus* nasal
800 colonization, a major risk factor in nosocomial infections. *Nature Medicine* 10:243–245.
- 801 142. D'Elia MA, Henderson JA, Beveridge TJ, Heinrichs DE, Brown ED. 2009. The N-Acetylmannosamine
802 Transferase Catalyzes the First Committed Step of Teichoic Acid Assembly in *Bacillus subtilis* and
803 *Staphylococcus aureus*. *J Bacteriol* 191:4030–4034.
- 804 143. Winstel V, Kühner P, Salomon F, Larsen J, Skov R, Hoffmann W, Peschel A, Weidenmaier C. 2015.
805 Wall Teichoic Acid Glycosylation Governs *Staphylococcus aureus* Nasal Colonization. *mBio* 6:e00632-
806 15.
- 807 144. Schlag M, Biswas R, Krismer B, Kohler T, Zoll S, Yu W, Schwarz H, Peschel A, Götz F. 2010. Role of
808 staphylococcal wall teichoic acid in targeting the major autolysin Atl. *Molecular Microbiology* 75:864–
809 873.
- 810 145. Biswas R, Martinez RE, Göhring N, Schlag M, Josten M, Xia G, Hegler F, Gekeler C, Gleske A-K, Götz
811 F, Sahl H-G, Kappler A, Peschel A. 2012. Proton-binding capacity of *Staphylococcus aureus* wall
812 teichoic acid and its role in controlling autolysin activity. *PLoS ONE* 7:e41415.

- 813 146. Bera A, Biswas R, Herbert S, Kulauzovic E, Weidenmaier C, Peschel A, Götz F. 2007. Influence of Wall
814 Teichoic Acid on Lysozyme Resistance in *Staphylococcus aureus*. *J Bacteriol* 189:280–283.
- 815 147. Kohler T, Weidenmaier C, Peschel A. 2009. Wall Teichoic Acid Protects *Staphylococcus aureus* against
816 Antimicrobial Fatty Acids from Human Skin. *J Bacteriol* 191:4482–4484.
- 817 148. Peschel A, Otto M, Jack RW, Kalbacher H, Jung G, Götz F. 1999. Inactivation of the *dlt* Operon in
818 *Staphylococcus aureus* Confers Sensitivity to Defensins, Protegrins, and Other Antimicrobial Peptides. *J*
819 *Biol Chem* 274:8405–8410.
- 820 149. Holland LM, Conlon B, O’Gara JP. 2011. Mutation of *tagO* reveals an essential role for wall teichoic
821 acids in *Staphylococcus epidermidis* biofilm development. *Microbiology* 157:408–418.
- 822 150. Yu L, Wang S, Guo Z, Liu H, Sun D, Yan G, Hu D, Du C, Feng X, Han W, Gu J, Sun C, Lei L. 2018. A
823 guard-killer phage cocktail effectively lyses the host and inhibits the development of phage-resistant
824 strains of *Escherichia coli*. *Appl Microbiol Biotechnol* 102:971–983.
- 825 151. Gu J, Liu X, Li Y, Han W, Lei L, Yang Y, Zhao H, Gao Y, Song J, Lu R, Sun C, Feng X. 2012. A Method
826 for Generation Phage Cocktail with Great Therapeutic Potential. *PLOS ONE* 7:e31698.
- 827 152. McCarthy AJ, Loeffler A, Witney AA, Gould KA, Lloyd DH, Lindsay JA. 2014. Extensive Horizontal Gene
828 Transfer during *Staphylococcus aureus* Co-colonization In Vivo. *Genome Biol Evol* 6:2697–2708.
- 829
- 830
- 831
- 832
- 833
- 834

835

836

837

838


839

840

841 **Figures**

842 Figure 1: Stages of phage infection and corresponding examples of resistance mechanisms at each stage.

843 Examples not yet identified in the staphylococci are listed in red.



844

845

846

847

848

849

850


851 Figure 2: A - The structure of the staphylococcal cell envelope. Lipoteichoic acid is shown in orange (glycerol

852 phosphate), a surface protein in black, wall teichoic acid in orange (glycerol phosphate) and yellow (ribitol

853 phosphate), capsule in blue, and cell wall carbohydrates in green (N-acetylglucosamine – GlcNAc) and purple

854 (N-acetylmuramic acid – MurNAc). Staphylococcal phages bind WTA and/or its ribitol phosphate modifications

855 (i.e., GlcNAc). B – Outline of the wall teichoic acid (WTA) biosynthesis pathway with proteins corresponding to
 856 each step listed in the blue arrows. Abbreviations are defined as follows - C_{55} -P, undecaprenyl phosphate;
 857 GlcNAc, N-acetylglucosamine; UDP-GlcNAc, uridine-5-diphosphate-N-acetylglucosamine; ManNAc, N-
 858 acetylmannosamine; UDP-ManNAc, uridine-5-diphosphate-N-acetylmannosamine; Gro-P, glycerol phosphate;
 859 CDP-Gro, cytidyl diphosphate-glycerol; Rbo-P, ribitol phosphate; CDP-Rbo, cytidyl diphosphate-ribitol; ABC,
 860 ATP-binding cassette; and LCP, LytR-CpsA-Psr.



861


862

863

864

865 Figure 3: Phage host range for an individual strain is the combination of multiple factors that have different
 866 levels of conservation within the species. This is illustrated by a hypothetical phylogenetic tree. Mechanisms
 867 can be present throughout strains (1, most conserved – red), present in many strains but with considerable
 868 allelic variation (2, conserved but polymorphic – shades of green), or present in a few strains, possibly with
 869 allelic variation (3a-3c, less conserved with potential polymorphism – blue, purple, and yellow). Branches

870 where mechanisms evolved by mutation or homologous recombination in the case of 1 and 2 or were acquired
 871 by HGT, in the case of 3a-3c, are annotated with colored stars. The table on the right summarizes the
 872 mechanisms (1-3c) present in each strain (A-J) using shaded boxes with corresponding colors. Strain J has a
 873 mutation that results in the null phenotype for the red mechanism. Host range is the result of the combination
 874 of mechanisms present, so strains A-C as well as F, H, and I would be predicted to have identical host ranges,
 875 but phage-specific factors could also introduce variability.



876