

Northumbria Research Link

Citation: Egidi, Eleonora, Wood, Jennifer, Fox, Edward, Liu, Wuxing and Franks, Ashley (2016) Draft Genome Sequence of *Rhodococcus erythropolis* NSX2, an Actinobacterium Isolated from a Cadmium-Contaminated Environment. *Genome Announcements*, 4 (5). e01147-16. ISSN 2169-8287

Published by: American Society for Microbiology

URL: <http://dx.doi.org/10.1128/genomeA.01147-16>
<<http://dx.doi.org/10.1128/genomeA.01147-16>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/38752/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

www.northumbria.ac.uk/nrl



Draft Genome Sequence of *Rhodococcus erythropolis* NSX2, an Actinobacterium Isolated from a Cadmium-Contaminated Environment

Eleonora Egidi,^a Jennifer L. Wood,^a Edward M. Fox,^b Wuxing Liu,^c Ashley E. Franks^a

Department of Physiology, Anatomy and Microbiology, La Trobe University, Bundoora, Victoria, Australia^a; CSIRO Agriculture and Food, Werribee, Victoria, Australia^b; Key Laboratory of Soil Environment and Pollution Remediation, Institute of Soil Science, Chinese Academy of Sciences, Nanjing, China^c

***Rhodococcus erythropolis* NSX2 is a rhizobacterium isolated from a heavy metal–contaminated environment. The 6.2-Mb annotated genome sequence shows that this strain harbors genes associated with heavy-metal resistance and xenobiotics degradation.**

Received 21 August 2016 Accepted 31 August 2016 Published 20 October 2016

Citation Egidi E, Wood JL, Fox EM, Liu W, Franks AE. 2016. Draft genome sequence of *Rhodococcus erythropolis* NSX2, an actinobacterium isolated from a cadmium-contaminated environment. *Genome Announc* 4(5):e01147-16. doi:10.1128/genomeA.01147-16.

Copyright © 2016 Egidi et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ashley E. Franks, a.franks@latrobe.edu.au.

Actinobacteria represent a dominant fraction of the metabolically active rhizosphere population in several heavy-metal (HM) accumulators (1). Members of this phylum are able to produce both metal-mobilizing and metal-immobilizing compounds (2), a crucial requirement to increase the HM extractability (3). This suggests an important role of *Actinobacteria* in metal uptake and translocation during phytoextraction (4). Nevertheless, our knowledge on the mechanisms of HM solubilization and resistance in the phylum is still limited. In this paper, we report the draft genome sequence of *Rhodococcus erythropolis* NSX2, a Gram-positive actinobacterium isolated from the rhizosphere of the heavy-metal hyperaccumulator *Sedum X Graptosedum* grown in a cadmium-contaminated environment (5). NSX2 was shown to be resistant to a MIC of 600 mg·L⁻¹ for Cd (5). The draft genome sequence will shed new light on the mechanism underlying the heavy-metal resistance, plant-growth promotion, and degradation abilities of *R. erythropolis* NSX2.

The strain was grown in cell culture, and the total genomic DNA was extracted from purified *R. erythropolis* NSX2 and converted to sequencing libraries using the Nextera XT DNA library preparation kit (Illumina). Libraries were normalized and pooled before sequencing on an Illumina MiSeq with 2 × 300-bp paired-end reads. For each isolate, the A5-miseq pipeline (6) was used to perform read trimming and correction, contig assembly, crude scaffolding, misassembly correction, and final scaffolding. The total length of the assembled genome was 6,279,737 bp, with a G+C content of 62.4% and median coverage of 66×. The final number of contigs was 37 with an *N*₅₀ value of 608,915.

A total of 5,963 candidate protein-coding genes were identified by automated annotation of the *R. erythropolis* NSX2 draft genome sequence using RAST (7). Comparative genome analysis revealed that *R. erythropolis* PR4 234621.6 (GenBank accession no. GCA_000010105.1) is NSX2's closest neighbor (score = 532). The NSX2 genome retains several traits associated with the uptake, efflux, reduction, and oxidation of metal ions. These include genes coding for the cobalt-zinc-cadmium resistance proteins CzcA and CzcD, the cation efflux system protein CusA, an arsenate reductase, an arsenic efflux pump protein, the copper resistance protein

CopC, and a transcriptional regulator for the mercury resistance protein MerR.

Interestingly, the NSX2 genome also harbors genes associated with the metabolism of aromatic compounds. In particular, we identified predicted genes encoding for proteins involved in the catabolism of protocatechuate and catechol (e.g., beta-ketoadipyl CoA thiolase, protocatechuate 3,4-dioxygenase beta chain, and beta-ketoadipate enol-lactone hydrolase), as well as proteins linked to the biphenyl degradation process (e.g., biphenyl-2,3-diol 1,2-dioxygenase, and 2-hydroxy-6-oxo-6-phenylhexa-2,4-dienoate hydrolase). The presence of these genetic markers suggests the ability of *R. erythropolis* NSX2 to metabolize many plant-derived compounds, as well as potentially degrade environmental pollutants (8).

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number MDCH00000000. The version described in this paper is the first version, MDCH01000000.

FUNDING INFORMATION

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

REFERENCES

- Gremion F, Chatzinotas A, Harms H. 2003. Comparative 16S rDNA and 16S rRNA sequence analysis indicates that *Actinobacteria* might be a dominant part of the metabolically active bacteria in heavy metal-contaminated bulk and rhizosphere soil. *Environ Microbiol* 5:896–907. <http://dx.doi.org/10.1046/j.1462-2920.2003.00484.x>.
- Kuffner M, De Maria S, Puschenreiter M, Fallmann K, Wieshammer G, Gorfer M, Strauss J, Rivelli AR, Sessitsch A. 2010. Culturable bacteria from Zn- and Cd-accumulating *Salix caprea* with differential effects on plant growth and heavy metal availability. *J Appl Microbiol* 108:1471–1484. <http://dx.doi.org/10.1111/j.1365-2672.2010.04670.x>.
- Wood JL, Liu W, Tang C, Franks AE. 2016. Microorganisms in heavy metal bioremediation: strategies for applying microbial-community engineering to remediate soils. *AIMS Bioengineering* 3:211–229. <http://dx.doi.org/10.3934/bioeng.2016.2.211>.
- Ma Y, Rajkumar M, Zhang C, Freitas H. 2016. Beneficial role of bacterial endophytes in heavy metal phytoremediation. *J Environ Manage* 174:14–25. <http://dx.doi.org/10.1016/j.jenvman.2016.02.047>.
- Liu W, Wang Q, Wang B, Hou J, Luo Y, Tang C, Franks AE. 2015. Plant

- growth-promoting rhizobacteria enhance the growth and Cd uptake of *Sedum plumbizincicola* in a Cd-contaminated soil. *J Soils Sediments* 15: 1191–1199. <http://dx.doi.org/10.1007/s11368-015-1067-9>.
6. Coil D, Jospin G, Darling AE. 2015. A5-miseq: an updated pipeline to assemble microbial genomes from Illumina MiSeq data. *Bioinformatics* 31:587–589. <http://dx.doi.org/10.1093/bioinformatics/btu661>.
 7. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9: <http://dx.doi.org/10.1186/1471-2164-9-75>.
 8. Okoh AI. 2006. Biodegradation alternative in the cleanup of petroleum hydrocarbon pollutants. *Biotechnol Mol Biol Rev* 1:38–50.