



Immunization with recombinant FliD confers protection against *Helicobacter pylori* infection in mice

Amir Ghasemi^{a,b,*}, Nazanin Mohammad^a, Josef Mautner^c, Mehrnaz Taghipour Karsabet^a, Abolfazl Ardjmand^d, Rezvan Moniri^{a,e}

^a Department of Microbiology and Immunology, Faculty of Medicine, Kashan University of Medical Sciences, Kashan, Iran

^b Department of Infectious Disease and Immunology, College of Veterinary Medicine, University of Florida, FL, USA

^c Technische Universität München & Helmholtz Zentrum München, Munich, Germany

^d Physiology Research Center, Kashan University of Medical Sciences, Kashan, Iran

^e Anatomical Sciences Research Center, Kashan University of Medical Sciences, Kashan, Iran

ARTICLE INFO

Keywords:

Helicobacter pylori
Immunization
Recombinant
Adjuvant
Protection

ABSTRACT

Nearly half of the world's population is infected with *Helicobacter pylori*. Clinical manifestations of this infection range from gastritis and peptic ulcers to gastric adenocarcinoma and lymphoma. Due to the emerging of antibiotic resistant strains and poor patient compliance of the antibiotic therapy, there is increasing interest in the development of a protective vaccine against *H. pylori* infection. The bacterial protein FliD forms a capping structure on the end of each flagellum which is critical to prevent depolymerization and structural degradation. In this study, the potential of FliD as a prospective *H. pylori* subunit vaccine was assessed. For this purpose, immunogenicity and protective efficacy of recombinant FliD (rFliD) from *H. pylori* was evaluated in C57BL/6 mice. Purified rFliD was formulated with different adjuvants and administered via subcutaneous or oral route. Subcutaneous immunization with rFliD elicited predominantly mixed Th1 and Th17 immune responses, with high titers of specific IgG₁ and IgG_{2a}. Splenocytes of immunized mice exhibited strong antigen-specific memory responses, resulting in the secretion of high amounts of IFN- γ and IL-17, and low levels of IL-4. Immunization with rFliD caused a significant reduction in *H. pylori* bacterial load relative to naïve control mice ($p < 0.001$), demonstrating a robust protective effect. Taken together, these results suggest that subcutaneous vaccination with rFliD formulated with CpG or Addavax could be considered as a potential candidate for the development of a subunit vaccine against *H. pylori* infection.

1. Introduction

H. pylori is a spiral-shaped, extracellular Gram-negative, and microaerophilic bacterium that has colonized the stomach of approximately 50% of the worldwide human population (Correa and Piazuelo, 2011). In 1994, *H. pylori* was classified as a class I carcinogen by the World Health Organization (Anderl and Gerhard, 2014). Infection is strongly associated with the development of several gastrointestinal diseases, including duodenal and gastric ulcers, gastric adenocarcinoma and gastric mucosa-associated lymphoid tissue lymphoma (MALT) (Suzuki et al., 2009). Antibiotic therapy, although effective, is associated with several disadvantages including low patient compliance due to the drugs' side effects, treatment failure as a result of the emergence of drug-resistant strains, high costs, and the failure to prevent reinfection (Boyanova et al., 2002; Frenck and Clemens, 2003; Gisbert, 2005; Wheeldon et al., 2005). Therefore, alternative approaches to combat *H.*

pylori infection are currently being pursued, including vaccination. Several *H. pylori* proteins have been identified as immunogenic in preclinical models, including Urease B (UreB) (Yang et al., 2005), Vacuolating toxin A (VacA) (Ghiara et al., 1997), *H. pylori* adhesion A (HpaA) (Sutton et al., 2007), neutrophil-activating protein A (NapA) (Satin et al., 2000), outer membrane protein (Omp) (Czinn and Blanchard, 2011), cytotoxin-associated antigen (CagA) (Ghiara et al., 1997), heat-shock proteins (Hsp) (Chionh et al., 2014a; Ferrero et al., 1995), OipA (Chen et al., 2012) catalase (Radcliff et al., 1997) and chimeric genes (Zhang et al., 2015; Li et al., 2012; Lv et al., 2014). Each of these antigens has the ability to reduce the bacterial load in animal models, but the elicited protection provided by them is less than ideal. Current strategies to enhance vaccine efficacy include the identification of most suitable immune targets and the combination of immunodominant antigens into multivalent formulations. Khalifeh Gholi et al. (Khalifeh Gholi et al., 2013) demonstrated that FliD, the flagellar

* Corresponding author at: Department of Microbiology and Immunology, Faculty of Medicine, Kashan University of Medical Sciences, P.O. Box: 8715988141, Kashan, Iran.
E-mail address: amghasemi@ufl.edu (A. Ghasemi).

Table 1
Vaccine formulations and route of application.

Route	Vaccine	Antigen dose	Adjuvant dose	Total volume	Number of mice
Subcutaneous	rFliD + CpG	30 µg	20 µg	100 µl	18
	rFliD + Addavax	30 µg	50 µl	100 µl	18
	rFliD + PBS	30 µg	0	100 µl	18
	CpG	0	20 µg	100 µl	18
	Addavax	0	50 µl	100 µl	18
	PBS	0	0	100 µl	18
Oral	rFliD + CTB	30 µg	5 µg	100 µl	18
	CTB	0	5 µg	100 µl	18

hook-associated protein 2, reacts with approximately 97 percent of sera obtained from patients infected with *H. pylori*, but not with sera from uninfected individuals, and reacts very weakly with sera from patients whose infections had been eradicated, suggesting that this is a common immune target in the infected human host (Khalifeh Gholi et al., 2013). FliD plays a crucial role in flagella assembly. Flagellin is important for bacterial motility and is essential for colonization and persistence of *H. pylori* in the stomach niche (Eaton et al., 1996).

The objective of the current study was to investigate the protective efficacy and underlying mechanisms of recombinant FliD (rFliD) antigen. Owing to inherent safety and the low risk of adverse reactions (Ghasemi et al., 2015), protein subunit immunization is an attractive vaccination approach. However, immune responses elicited by the administration of pure antigens are usually low. Therefore, we investigated whether formulations of FliD with different adjuvants, CpG, Cholera toxin subunit B (CTB), or Addavax, enhance protection of mice against *H. pylori* infection.

2. Materials and methods

2.1. Bacterial strains

E. coli strains BL21 and TOP10 were used for expression of FliD. Bacteria were routinely grown at 37 °C in LB broth or agar. The *H. pylori* strain SS1 was grown on *Brucella* agar supplemented with 5% sheep blood, 5 µg/mL trimethoprim, 161.5 µg/mL polymyxin B (Sigma), 10 µg/mL vancomycin (sigma) and 2.5 µg/mL amphotericin B (Sigma), in an anaerobic jar with microaerophilic gas generating kit (Merck, Germany) for 3 days at 37 °C. For infection experiments, bacteria were subcultured in brain heart infusion broth (BHI, Merck) containing 10 µg/mL vancomycin and supplemented with 10% Fetal Bovine Serum (FBS) (Sigma), and grown in microaerophilic conditions for 72 h at 37 °C.

2.2. Cloning, expression, and purification of recombinant FliD

Cloning and expression of FliD from *H. pylori* in BL21 (DE3) and its purification has been described previously (Khalifeh Gholi et al., 2013). Briefly, the *fliD* gene was amplified from genomic DNA of *H. pylori* by PCR (forward primer: 5'-ATGGCAATAGGTTTCATTAAGCT-3', reverse primer: 5'-ATTCTTTTAGCCGCCGCTT-3'). The DNA fragment was directly inserted into pTZ57R (InsTAclone PCR Cloning Kit, Thermo-Fisher, USA) and then subcloned into the pET28a+ vector (Novagen, USA) to add a 6xHis tag. To express FliD, 1 mM isopropyl-β-D-thiogalactopyranoside (IPTG) was used. Upon induction, rFliD was expressed in soluble form and purified under native conditions. Following SDS-PAGE separation, purity and identity of the recombinant protein was evaluated by Coomassie blue staining and Western blotting (Ghasemi et al., 2014a; Ghasemi et al., 2013). For immunodetection, size-separated proteins were transferred to a PVDF membrane and rFliD detected using anti-6xHis peroxidase (Roche, Germany) (1/40,000). Finally, the membrane was developed using 3, 3'-Diaminobenzidine tetrahydrochloride (Sigma). To preclude effects caused by residual

endotoxin present in the protein preparation, only batches of purified protein with an endotoxin content of less than 0.05 endotoxin units per mg of protein (evaluated by Limulus amoebocyte lysate analysis kit, Lonza, Basel, Switzerland) were used in further studies. The Bradford method was used to determine the concentration of the recombinant protein (Stoscheck, 1990).

2.3. Mice

Six- to eight-week-old female C57BL/6 mice were purchased from the Pasteur Institute of Iran and maintained under specific pathogen-free conditions. Mice were handled under optimal conditions of temperature, hygiene, humidity and light (cycles of 12 h dark/light). All experimental animal procedures were approved by the ethical committee of the Kashan University of Medical Science.

2.4. Immunization and *H. pylori* inoculation of mice

One hundred and forty-four mice were randomly divided into eight groups (n = 18 each). Five groups were immunized subcutaneously (s.c.) three times at 2-week intervals with either 30 µg rFliD formulated with 20 µg/mouse CpG (CpG, ODN1826 5'-TCCATGACGTTCTGAC GTT- 3', synthesized by TAG Copenhagen, Denmark), 30 µg rFliD formulated with (1:1) Addavax (Invivogen, USA), or PBS, or 20 µg/mouse CpG, Addavax, and PBS alone. The two remaining groups were orally immunized with rFliD and 20 µg/mouse CTB (Sigma, Germany) or CTB alone (Table 1). To enable immunization in a small volume, the recombinant antigen was lyophilized and then reconstituted in a volume of 100 µl PBS containing the respective adjuvant.

Two weeks after the final immunization, eight mice from each group were challenged orally thrice in 2-day intervals with 5×10^8 CFU mouse-adapted *H. pylori* strain SS1 in 100 µl brain heart infusion broth. Five mice were bled to obtain sera at days 0, 15, 30, 45, and 75 after the first immunization. The remaining five mice were sacrificed to evaluate immune responses including cytokine production, sIgA secretion, and IgG immune responses on the day of the challenge. For measuring sIgA secretion, gastric fluid was collected as described previously (Chionh et al., 2014b).

2.5. Humoral and mucosal immune responses

To measure rFliD-specific serum IgG₁, IgG_{2a} and gastric fluid sIgA titers in immunized mice, an enzyme-linked immunosorbent assay (ELISA) was used. 96-well polystyrene plates (Greiner Bio-One, Frickenhausen, Germany) were coated with the purified rFliD (1 µg/mL). After overnight incubation, TBST buffer (Tris-buffered saline, pH 7.4, and containing 0.05% Tween 20) were used to wash the plates three times followed by blocking with 300 µl 10% FBS in PBS for 2 h at 37 °C. After that, plates were incubated with serial dilutions of mouse sera or gastric and intestinal fluid for 2 h at room temperature and then washed. HRP-conjugated goat-anti-mouse IgG₁, IgG_{2a} or sIgA antibodies (BD Pharmingen, USA) were added to the wells for another 90 min at 37 °C. After the last washing step, specific reactivity was

measured by the addition of 50 μ l/well of the enzyme substrate TMB (Pishtaz Teb, Tehran, Iran). The reaction was stopped by adding 30 μ l 20% H₂SO₄ to each well. Optical density at 495 nm was measured using an ELISA plate reader (Bio-Tek Instruments, Winooski, USA).

2.6. Cellular immune responses

Assessment of cellular immune responses has been described elsewhere (Ghasemi et al., 2015; Ghasemi et al., 2014b). Briefly, fifteen days after the last immunization, the mice were euthanized and spleens removed under sterile conditions. The spleens were minced and homogenized and then mononuclear cells (MNCs) were isolated by Ficoll–Paque (GE Healthcare, Uppsala, Sweden) density centrifugation. 2×10^6 splenocytes were plated in one well of a 24-well plate in 2 ml of complete medium (RPMI-1640 supplemented with 2 mM L-glutamine, 100 U/ml penicillin, 100 μ g/ml streptomycin and 10% heat inactivated FBS) and 10 μ g/ml rFliD were added to each well and the plates incubated at 37 °C in 5% CO₂ for 48 h. Control wells received PBS instead of the antigen. After 48 h, supernatants were collected and stored at –70 °C for cytokine assay. Concentrations of interferon-gamma (IFN- γ), interleukin-17 (IL-17), and –4 (IL-4) were determined by ELISA according to the manufacturer's instructions (R&D Systems, USA).

2.7. Protection experiments

H. pylori CFU were quantified six weeks after the last immunization. For this purpose, the copy number of *H. pylori* 16s gene within mouse gastric tissues was measured by a quantitative PCR as described previously (Roussel et al., 2007). A recombinant plasmid carrying the *H. pylori* 16S gene was used as a standard to measure copy numbers of the 16S gene in DNA extracted from half of a mouse stomach and the obtained Colony-Forming Units (CFU) calculated as CFU per stomach.

2.8. Statistical analysis

Statistical analysis was carried out using SPSS computer software, version 16.0. All data were represented as the mean \pm standard deviation (SD). Differences among data of the five groups were analyzed by one factor analysis of variance (ANOVA) and Tukey's post hoc test in SPSS. P-values < 0.05 were considered as statistically significant.

3. Results

3.1. Expression and purification of the recombinant FliD protein

To obtain sufficient amounts of rFliD for protection experiments and immunological analyses, the *H. pylori* *fliD* gene was amplified by PCR and fused to a 6 x histidine tag in the pET28 bacterial expression vector. *E. coli* BL21 cells were transformed with the resulting plasmid, and protein expression was induced with IPTG. Upon purification with Ni-NTA agarose, rFliD expression product was verified via SDS-PAGE and Western blot (Ghasemi et al., 2013). Consistent with molecular weight prediction, recombinant rFliD had a size of approximately 84 kDa (Fig. 1). From one liter of liquid culture, approximately 35 mg of rFliD protein was obtained.

3.2. Subcutaneous vaccination with rFliD formulated with adjuvants provides protection against *H. pylori* infection

To investigate whether immunization with rFliD provided protection against subsequent infection with *H. pylori*, groups of mice were immunized with rFliD formulated with different adjuvants, rFliD alone, or PBS. Two weeks after the last immunization, mice were challenged thrice at one day intervals with 5×10^8 *H. pylori* SS1. Bacterial DNA load was assessed by qPCR four weeks post infection.

The copy numbers of *H. pylori* DNA in groups immunized with rFliD

formulated with adjuvants were significantly lower than in mice immunized with rFliD in PBS or control mice. The highest degree of protection among animals vaccinated with rFliD plus adjuvants was observed in mice vaccinated with rFliD plus CpG, albeit without reaching statistical significance (Fig. 2). Compared to the negative control group, even mice vaccinated with rFliD in PBS showed a significant reduction in CFU, implicating a high immunogenicity of this antigen.

3.3. Vaccination with rFliD and adjuvants induces vigorous antibody responses

Specific antibody titers produced in mice immunized with rFliD formulated with different adjuvants were evaluated by ELISA in sera obtained at different time points after the first immunization. Vaccination with rFliD induced strong immunoglobulin G (IgG) responses, with IgG₁ titers usually exceeding those of the IgG_{2a} subtype (Fig. 3). The IgG₁/IgG_{2a} ratio was 1–2 and 1–3.44 in mice immunized subcutaneously and orally, respectively. These results suggested that CpG and Addavax induced a Th1-biased immune response.

Although different immune responses were observed for CpG and CTB, the total amount of IgG detected was similar for both adjuvants. Responses induced in mice orally immunized with rFliD formulated with CTB were markedly lower than in mice immunized subcutaneously with rFliD plus CpG or Addavax. However, the IgG responses of mice immunized with rFliD formulated with CTB were significantly increased 75 days after the initial immunization. Mice vaccinated with rFliD formulated with either CpG or Addavax demonstrated increased IgG₁ and IgG_{2a} titers commencing at the second week post-immunization, peaking at the sixth week and remaining elevated until the eleventh week after the first immunization. Compared to all adjuvant formulations, rFliD alone induced only low titers of both IgG₁ and IgG_{2a} antibodies in immunized mice (Fig. 3). To assess the mucosal immune responses elicited by immunization, gastric fluid IgA production was determined for each vaccine. As shown in Fig. 3, the three groups immunized with rFliD plus adjuvants showed significantly elevated gastric mucosal IgA titers compared to the control group which had only received PBS (P < 0.001). The group immunized with rFliD and PBS exhibited no difference when compared to the PBS control group.

Taken together, these findings suggest that the combination of rFliD with CpG, Addavax, or CTB induced robust antibody responses.

3.4. Vaccination with rFliD and adjuvants elicits a mixed Th1 and Th17 type of immune response

To further characterize the induced immune responses, splenocytes from mice immunized with rFliD formulated with different adjuvants, or rFliD alone, or PBS were isolated 45 days after the first immunization. Following incubation with rFliD, cytokine production was assessed. Secretion of IFN- γ , IL-4, and IL-17 from splenocytes of immunized mice after re-stimulation was assessed two weeks after the final immunization boost. As shown in Fig. 4, the concentrations of all tested cytokines were significantly increased in mice vaccinated with rFliD formulated with adjuvants compared to the PBS control group, yet the amount of IL-4 was comparable low. Splenocytes of mice immunized with rFliD without adjuvants secreted significantly higher amounts of IFN- γ and IL-17 than splenocytes from PBS control mice. These results indicate that rFliD alone is capable of eliciting cellular immune responses mostly of Th1 and Th17 type. Moreover, these immune responses were markedly enhanced by the use of adjuvants.

It is important to note that there were no discernable differences in the patterns of cytokines secreted by splenocytes from mice immunized with rFliD formulated with the three different adjuvants (Fig. 4).




Fig. 1. Expression and purification of recombinant FliD. (a) Analysis of rFliD protein expression by SDS-PAGE. Protein expression in bacteria transformed with the *fliD* gene was induced by the addition of IPTG to the media. Four hours after induction, the bacteria were harvested, the bacterial lysates run over Nickel-NTA columns and bound proteins eluted with Imidazole. Aliquots of the different fractions were size-separated by SDS-PAGE and the resulting gel stained with Coomassie-Blue. Lane 1: molecular weight marker, lane 2: bacterial lysate after IPTG induction, lane 3: flow through of Nickel-NTA column, lanes 4 and 5: column eluate with buffer containing 20 mM Imidazole, lanes 6 and 7: eluate with buffer containing 40 mM Imidazole, Lane 8: eluate with buffer containing 1 M Imidazole. (b) Western blot analysis of rFliD using a monoclonal antibody directed against the 6xHis-tag attached to rFliD. Lane 1: Molecular weight marker, lane 2: purified rFliD, lane 3: lysate of untransformed bacteria. Expected size of rFliD: 84 kDa.




Fig. 2. *H. pylori* colonization of stomach tissue in immunized mice. C57BL/6 mice were immunized on day 0, 15 and 30 with three doses of 30 μ g rFliD with CpG, Addavax, CTB or PBS. The control groups (8 mice per each group) only received adjuvants or PBS. Two weeks after final vaccination (on day 45), mice were challenged orally with *H. pylori*. Four weeks post challenge, levels of gastric *H. pylori* colonization were determined by real-time quantitative PCR. Significance (ANOVA) was evaluated with reference to the PBS control. ns: not significant, ***: $p < 0.001$, *: $p < 0.05$.


4. Discussion

The development of an effective vaccine against *H. pylori* has been confounded by several aspects including choice of the antigen, route of vaccination, and selection of adjuvants to boost immunogenicity (Li et al., 2015; Brown et al., 2014). Motility enables *H. pylori* to reach the gastric epithelium, attach via adhesion factors, and establish infection in the epithelium (Sheu et al., 2010). Polar flagella is considered an important virulence factor for *H. pylori*, as studies done with a non-motile *fliD* mutant demonstrated that FliD, and consequently a functional flagellum, is required for the colonization of mice (Haiko and Westerlund-Wikstrom, 2013). Moreover, it has been shown that FliD promotes biofilm formation, thereby contributing toward environmental resilience (Rathawongjirakul et al., 2016). The association


between the presence of antibodies against FliD and gastric cancer is well documented. Based on these findings, it has been suggested that FliD can serve as a biomarker in screening for GC patients (Li et al., 2016). We hypothesized that highly expressed proteins that are critical for *H. pylori* infection might constitute optimal candidates for subunit vaccines. In this regard, rFliD was administered orally or subcutaneously and formulated with Addavax, CpG, or CTB adjuvants. Addavax is a squalene-based oil-in-water emulsion capable of eliciting both cellular and humoral immune responses (Ren et al., 2017; Calabro et al., 2013). MF59, which has similar composition as Addavax, has been approved for clinical use in the US with a seasonal flu vaccine (Tsai, 2011). By contrast, CpG oligodeoxynucleotides are toll-like receptor 9 (TLR9) agonists known to stimulate innate immune defenses and antigen-specific T cell responses (Bode et al., 2011). CTB is a potent adjuvant which can stimulate innate and antigen-specific responses upon oral or intranasal administration (Stratmann, 2015).

We looked at the protective capability of rFliD by challenging vaccinated mice with live *H. pylori* SS1. Our results demonstrate that mice immunized with rFliD in combination with any of the tested adjuvants conferred significant protective immunity to mice as compared to the control group. Immunization with non-adjuvanted rFliD resulted in partial protection against *H. pylori* but significant in comparison to the naïve control. Taken together, our results suggest that FliD is capable of eliciting strong and protective immune responses in mice when combined with adjuvants. Our results also demonstrated that the strongest immune responses were elicited by subcutaneous rather than oral immunization. Since both CpG and Addavax have been approved for human application, they may form the basis of future subunit vaccines against *H. pylori* (Mohan et al., 2013; Mbow et al., 2010).

To investigate how immunity against *H. pylori* infection was established, cellular immune responses were assessed by measuring cytokine production of splenocytes from immunized mice after re-stimulation *in vitro*. These cells showed robust IFN- γ secretion, which is indicative of a Th1 type of response. Furthermore, high quantities of IL-17 were produced by splenocytes of immunized mice upon re-stimulation. These findings are consistent with earlier studies by other investigators who reported mixed Th1 and Th17 responses against a set of distinct *H. pylori* proteins (Lv et al., 2014; Chionh et al., 2014b; Sjkovist Ottsjo



a



b

Fig. 3. Analysis of the rFliD-specific antibody response in immunized animals. (a) Kinetics of the IgG₁ and IgG_{2a} responses after immunization with rFliD plus different adjuvants. Animals (5 mice per each group) were bled retroorbitally on the indicated days and specific IgG₁ and IgG_{2a} antibody titers against rFliD were evaluated by ELISA. Titer values represent the mean ± SD of sera from three analyses of five animals each. (b) Gastric scrapings were collected 45 days post first immunization and rFliD-specific IgA titers analyzed by ELISA. Ψ: Comparison of antigen-specific anti-rFliD IgG₁ and IgG_{2a} in mice immunized with rFliD formulated with and without adjuvant. †: Comparison of anti-rFliD IgA in mice immunized with rFliD versus PBS. ns: not significant, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$.




Fig. 4. Cytokine production by splenocytes from immunized mice after restimulation with rFliD. Spleen cells of mice from all five groups (5 mice per each group) were stimulated *in vitro* with 10 µg/mL rFliD for 48 h. Cytokine levels in culture supernatant were measured by sandwich ELISA. The data are the mean ± SD of five individual mice from each group with two repeats. Significance was assessed between mice immunized with rFliD +/- adjuvants and mice treated with PBS. ***: $p < 0.001$.

et al., 2015; Liu et al., 2011; Chen et al., 2014). In addition, increased amounts of IL-4 were produced by splenocytes of rFliD-immunized versus naïve mice. However, the amount of IL-4 produced by the cells was comparable low. Therefore, it remains unknown whether the vaccines also elicited a significant Th2 type of immune response. Adjuvants are able to modulate strength and type of the T cell response against a

given antigen. For example, adjuvants like CTB can skew immune response from Th1 to Th2 whereas MF59[®] is a strong stimulator of both Th1 and Th2 immune responses (Ott et al., 1995; Berin and Shreffler, 2008). As determined by cytokine secretion assays, all adjuvants employed in this study induced very similar types of cellular immune responses against rFliD that encompassed different immune effector functions, which may explain their protective effect. Although the specific functions of the secreted cytokines in the control of *H. pylori* infection still need to be elucidated in more detail, Th1 and Th17 cells have been implicated in inducing local inflammation and in promoting protective immune responses (Larussa et al., 2015). Moreover, Th17 responses are involved in the recruitment of neutrophils, the release of antimicrobial peptides and the induction of IL-17-driven Th1 immunity (Ray et al., 2014).

To further characterize the Th1 and Th2 profile of the elicited immune responses, production of antigen-specific IgG₁ and IgG_{2a} antibodies was assessed. Although rFliD-induced humoral immune responses were predominantly of IgG₁ isotype, significant levels of IgG_{2a} titers were also detected. Specific antibodies against rFliD of both isotypes were also produced in mice immunized with rFliD alone, but titers were significantly lower in comparison to mice vaccinated with rFliD and adjuvants. The elicited antibody responses in animals vaccinated with rFliD and adjuvants were more pronounced when applied subcutaneously than through the oral route. To assess the mucosal

humoral immune response elicited by the different vaccines, gastric IgA production was evaluated. Immunizations with rFlid and Addavax, CpG, or CTB induced significant levels of antigen-specific gastric IgA compared to control mice. These findings suggest that vaccination with rFlid and adjuvant, regardless of subcutaneous or oral route of delivery, efficiently induced gastric mucosal immune responses.

Although the immune mechanisms that provide protective immunity against *H. pylori* infection are still poorly defined, cellular immune responses seem to be critical and sufficient to provide protection even in the absence of B cells (Ernst and Pappo, 2001). Which type of cellular immune response is most protective is still matter of debate. While Th1 type of responses were considered most effective in previous studies, Mohammadi et al. (Mohammadi et al., 1997) recently showed that Th1 cells increased the severity of gastritis and Th2 cells were required for reducing bacterial load. Likewise, Th17 type of responses were shown to favor bacterial growth (Shi et al., 2010), while other studies emphasized the importance of Th-17 cells and IL-17 in reducing bacterial load. For example, DeLyria et al. (DeLyria et al., 2009) showed that mice vaccinated against *H. pylori* showed robust Th-17 responses and that protection was dependent on neutrophils. Moreover, the secretion of IL-17 in mice vaccinated against *H. pylori* was shown to be critical for the reduction of the bacterial load (Velin et al., 2009). In the present study, the ratio of IgG1/IgG2a and the cytokine profile of restimulated splenocytes from vaccinated mice are indicative of a mixed Th1 and Th17 type of response. How precisely protective immunity is established by the vaccine-induced immune responses is currently not known and requires analysis of the local recall response, e.g. the production of cytokines, especially IFN- γ and IL-17, and antimicrobial peptides in the gastric tissue after challenge in immunized mice. Additional experiments are also needed to assess whether vaccine efficacy can be further increased by incorporating additional *H. pylori* antigens.

Acknowledgments

We would like to thank you Homayun Pourbabae and Mohammad Khalifeh Gholi for technical supports. The present study was supported by a grant from Kashan University of Medical Sciences (Grant no. 93178).

References

- Anderl, F., Gerhard, M., 2014. Helicobacter pylori vaccination: is there a path to protection. *World J. Gastroenterol.* 34, 11939–11949.
- Berin, M.C., Shreffler, W.G., 2008. T(H)2 adjuvants: implications for food allergy. *J. Allerg Clin. Immunol.* 121, 1311–1320 (quiz 1321–1312).
- Bode, C., Zhao, G., Steinhagen, F., Kinjo, T., Klinman, D.M., 2011. CpG DNA as a vaccine adjuvant. *Expert Rev. Vaccines* 10, 499–511.
- Boyanova, L., Mentis, A., Gubina, M., Rozynek, E., Gosciniaik, G., Kalenic, S., Göral, V., Kupcinskas, L., Kantareken, B., Aydin, A., 2002. The status of antimicrobial resistance of Helicobacter pylori in eastern Europe. *Clin. Microbiol. Infect.* 8, 388–396.
- Brown, L.J., Rosatte, R.C., Fehner-Gardiner, C., Ellison, J.A., Jackson, F.R., Bachmann, P., Taylor, J.S., Franka, R., Donovan, D., 2014. Oral vaccination and protection of striped skunks (*Mephitis mephitis*) against rabies using ONRAB(R). *Vaccine* 32, 3675–3679.
- Calabro, S., Tritto, E., Pezzotti, A., Taccone, M., Muzzi, A., Bertholet, S., De Gregorio, E., O'Hagan, D.T., Baudner, B., Seubert, A., 2013. The adjuvant effect of MF59 is due to the oil-in-water emulsion formulation, none of the individual components induce a comparable adjuvant effect. *Vaccine* 31, 3363–3369.
- Chen, J., Lin, M., Li, N., Lin, L., She, F., 2012. Therapeutic vaccination with Salmonella-delivered codon-optimized outer inflammatory protein DNA vaccine enhances protection in Helicobacter pylori infected mice. *Vaccine* 30, 5310–5315.
- Chionh, Y.T., Arulmuruganar, A., Venditti, E., Ng, G.Z., Han, J.-X., Entwisle, C., Ang, C.-S., Colaco, C.A., McNulty, S., Sutton, P., 2014a. Heat shock protein complex vaccination induces protection against Helicobacter pylori without exogenous adjuvant. *Vaccine* 32, 2350–2358.
- Chionh, Y.T., Arulmuruganar, A., Venditti, E., Ng, G.Z., Han, J.X., Entwisle, C., Ang, C.S., Colaco, C.A., McNulty, S., Sutton, P., 2014b. Heat shock protein complex vaccination induces protection against Helicobacter pylori without exogenous adjuvant. *Vaccine* 32, 2350–2358.
- Chen, J., Li, N., She, F., 2014. Helicobacter pylori outer inflammatory protein DNA vaccine-loaded bacterial ghost enhances immune protective efficacy in C57BL/6 mice. *Vaccine* 32, 6054–6060.
- Correa, P., Piazuelo, M.B., 2011. Helicobacter pylori infection and gastric adenocarcinoma. *US Gastroenterol. Hepatol. Rev.* 7, 59.
- Czinn, S.J., Blanchard, T., 2011. Vaccinating against Helicobacter pylori infection. *Nat. Rev. Gastroenterol. Hepatol.* 8, 133–140.
- DeLyria, E.S., Redline, R.W., Blanchard, T.G., 2009. Vaccination of mice against H pylori induces a strong Th-17 response and immunity that is neutrophil dependent. *Gastroenterology* 136, 247–256.
- Eaton, K.A., Suerbaum, S., Josenhans, C., Krakowka, S., 1996. Colonization of gnotobiotic piglets by Helicobacter pylori deficient in two flagellin genes. *Infect. Immun.* 64, 2445–2448.
- Ernst, P.B., Pappo, J., 2001. T-cell-mediated mucosal immunity in the absence of antibody: lessons from Helicobacter pylori infection. *Acta Odontol. Scand.* 59, 216–221.
- Ferrero, R.L., Thiberge, J.-M., Kansau, I., Wuscher, N., Huerre, M., Labigne, A., 1995. The GroES homolog of Helicobacter pylori confers protective immunity against mucosal infection in mice. *Proc. Natl. Acad. Sci.* 92, 6499–6503.
- Frenc, R.W., Clemens, J., 2003. Helicobacter in the developing world. *Microbes Infect.* 5, 705–713.
- Ghasemi, A., Salari, M.H., Pourmand, M.R., Zarnani, A.H., Ahmadi, H., Shirazi, M.H., Jeddi-Tehrani, M., 2013. Optimization and efficient purification in production of *Brucella melitensis* recombinant HSP a and TF proteins with low endotoxin contents. *Jundishapur J. Microbiol.* 6, e6875.
- Ghasemi, A., Jeddi-Tehrani, M., Mautner, J., Salari, M.H., Zarnani, A.H., 2014a. Immunization of mice with a novel recombinant molecular chaperon confers protection against *Brucella melitensis* infection. *Vaccine* 32, 6659–6666.
- Ghasemi, A., Zarnani, A.H., Ghoojani, A., Rezaei, S., Salari, M.H., Jeddi-Tehrani, M., 2014b. Identification of a new immunogenic candidate conferring protection against *Brucella melitensis* infection in mice. *Mol. Immunol.* 62, 142–149.
- Ghasemi, A., Jeddi-Tehrani, M., Mautner, J., Salari, M.H., Zarnani, A.H., 2015. Simultaneous immunization of mice with Omp31 and TF provides protection against *Brucella melitensis* infection. *Vaccine* 33, 5532–5538.
- Ghiara, P., Rossi, M., Marchetti, M., Di Tommaso, A., Vindigni, C., Ciampolini, F., Covacci, A., Telford, J.L., De Magistris, M.T., Pizzi, M., 1997. Therapeutic intra-gastric vaccination against Helicobacter pylori in mice eradicates an otherwise chronic infection and confers protection against reinfection. *Infect. Immun.* 65, 4996–5002.
- Gisbert, J.P., 2005. The recurrence of Helicobacter pylori infection: incidence and variables influencing it. A critical review. *Am. J. Gastroenterol.* 100, 2083–2099.
- Haiko, J., Westerlund-Wikstrom, B., 2013. The role of the bacterial flagellum in adhesion and virulence. *Biology* 2, 1242–1267.
- Khalifeh Gholi, M., Kalali, B., Formichella, L., Gottner, G., Shamsipour, F., Zarnani, A.H., Hosseini, M., Busch, D.H., Shirazi, M.H., Gerhard, M., 2013. Helicobacter pylori Flid protein is a highly sensitive and specific marker for serologic diagnosis of H. pylori infection. *Int. J. Med. Microbiol.: IJMM* 303, 618–623.
- Larussa, T., Leone, I., Suraci, E., Imeneo, M., Luzzo, F., 2015. Helicobacter pylori and T helper cells: mechanisms of immune escape and tolerance. *J. Immunol. Res.* 2015, 981328.
- Li, H.B., Zhang, J.Y., He, Y.F., Chen, L., Li, B., Liu, K.Y., Yang, W.C., Zhao, Z., Zou, Q.M., Wu, C., 2012. Systemic immunization with an epitope-based vaccine elicits a Th1-biased response and provides protection against Helicobacter pylori in mice. *Vaccine* 31, 120–126.
- Li, H., Zhang, J., He, Y., Li, B., Chen, L., Huang, W., Zou, Q., Wu, C., 2015. Intranasal immunization with an epitope-based vaccine results in earlier protection but not better protective efficacy, against Helicobacter pylori compared to subcutaneous immunization. *Immunol. Res.* 62, 368–376.
- Li, H., Zhang, B., Hu, X., Dong, Y., Fan, Q., Guo, F., Ren, X., Zhou, H., Tian, W., Zhao, Y., 2016. Serum Helicobacter pylori Flid antibody and the risk of gastric cancer. *Oncotarget* 7, 22397–22408.
- Liu, K.Y., Shi, Y., Luo, P., Yu, S., Chen, L., Zhao, Z., Mao, X.H., Guo, G., Wu, C., Zou, Q.M., 2011. Therapeutic efficacy of oral immunization with attenuated Salmonella typhimurium expressing Helicobacter pylori CagA, VacA and UreB fusion proteins in mice model. *Vaccine* 29, 6679–6685.
- Lv, X., Yang, J., Song, H., Li, T., Guo, L., Xing, Y., Xi, T., 2014. Therapeutic efficacy of the multi-epitope vaccine CTB-UE against Helicobacter pylori infection in a Mongolian gerbil model and its microRNA-155-associated immuno-protective mechanism. *Vaccine* 32, 5343–5352.
- Mbow, M.L., De Gregorio, E., Valiante, N.M., Rappuoli, R., 2010. New adjuvants for human vaccines. *Curr. Opin. Immunol.* 22, 411–416.
- Mohammadi, M., Nedrud, J., Redline, R., Lycke, N., Czinn, S.J., 1997. Murine CD4 T-cell response to Helicobacter infection: TH1 cells enhance gastritis and TH2 cells reduce bacterial load. *Gastroenterology* 113, 1848–1857.
- Mohan, T., Verma, P., Rao, D.N., 2013. Novel adjuvants & delivery vehicles for vaccines development: a road ahead. *Indian J. Med. Res.* 138, 779–795.
- Ott, G., Barchfeld, G.L., Chernoff, D., Radhakrishnan, R., van Hoogevest, P., Van Nest, G., 1995. MF59. Design and evaluation of a safe and potent adjuvant for human vaccines. *Pharm. Biotechnol.* 6, 277–296.
- Radcliff, F.J., Hazell, S.L., Kolesnikow, T., Doidge, C., Lee, A., 1997. Catalase, a novel antigen for Helicobacter pylori vaccination. *Infect. Immun.* 65, 4668–4674.
- Rathawangjirakul, P., Thongkerd, V., Chaicumpa, W., 2016. The impacts of a flid mutation on the biofilm formation of Helicobacter pylori. *Asia. Pac. J. Trop. Biomed.* 6, 1008–1014.
- Ray, S., De Salvo, C., Pizarro, T.T., 2014. Central role of IL-17/Th17 immune responses and the gut microbiota in the pathogenesis of intestinal fibrosis. *Curr. Opin. Gastroenterol.* 30, 531–538.
- Ren, S.T., Zhang, X.M., Sun, P.F., Sun, L.J., Guo, X., Tian, T., Zhang, J., Guo, Q.Y., Li, X., Guo, L.J., Che, J., Wang, B., Zhang, H., 2017. Intranasal immunization using mannan as a novel adjuvant for an inactivated influenza vaccine and its adjuvant effect compared with MF59. *PLoS One* 12, e0169501.

- Roussel, Y., Harris, A., Lee, M.H., Wilks, M., 2007. Novel methods of quantitative real-time PCR data analysis in a murine *Helicobacter pylori* vaccine model. *Vaccine* 25, 2919–2929.
- Satin, B., Del Giudice, G., Della Bianca, V., Dusi, S., Laudanna, C., Tonello, F., Kelleher, D., Rappuoli, R., Montecucco, C., Rossi, F., 2000. The neutrophil-activating protein (HP-NAP) of *Helicobacter pylori* is a protective antigen and a major virulence factor. *J. Exp. Med.* 191, 1467–1476.
- Sheu, B.S., Yang, H.B., Yeh, Y.C., Wu, J.J., 2010. *Helicobacter pylori* colonization of the human gastric epithelium: a bug's first step is a novel target for us. *J. Gastroenterol. Hepatol.* 25, 26–32.
- Shi, Y., Liu, X.F., Zhuang, Y., Zhang, J.Y., Liu, T., Yin, Z., Wu, C., Mao, X.H., Jia, K.R., Wang, F.J., Guo, H., Flavell, R.A., Zhao, Z., Liu, K.Y., Xiao, B., Guo, Y., Zhang, W.J., Zhou, W.Y., Guo, G., Zou, Q.M., 2010. *Helicobacter pylori*-induced Th17 responses modulate Th1 cell responses, benefit bacterial growth, and contribute to pathology in mice. *J. Immunol.* 184, 5121–5129.
- Sjokvist Ottsjo, L., Flach, C.F., Nilsson, S., Malefyt Rde, W., Walduck, A.K., Raghavan, S., 2015. Defining the roles of IFN-gamma and IL-17A in inflammation and protection against *Helicobacter pylori* infection. *PLoS One* 10, e0131444.
- Stoscheck, C.M., 1990. Quantitation of protein. *Methods Enzymol.* 182, 50–68.
- Stratmann, T., 2015. Cholera toxin subunit B as Adjuvant—an accelerator in protective immunity and a break in autoimmunity. *Vaccines* 3, 579–596.
- Sutton, P., Doidge, C., Pinczower, G., Wilson, J., Harbour, S., Swierczak, A., Lee, A., 2007. Effectiveness of vaccination with recombinant HpaA from *Helicobacter pylori* is influenced by host genetic background. *FEMS Immunol. Med. Microbiol.* 50, 213–219.
- Suzuki, H., Saito, Y., Hibi, T., 2009. *Helicobacter pylori* and Gastric Mucosa-associated Lymphoid Tissue (MALT) Lymphoma: Updated Review of Clinical Outcomes and the Molecular Pathogenesis. *Gut and liver* 3, 81–87.
- Tsai, T.F., 2011. MF59 adjuvanted seasonal and pandemic influenza vaccines, *Yakugaku zasshi*. *J. Pharm. Soc. Japan* 131, 1733–1741.
- Velin, D., Favre, L., Bernasconi, E., Bachmann, D., Pythoud, C., Saiji, E., Bouzourene, H., Michetti, P., 2009. Interleukin-17 is a critical mediator of vaccine-induced reduction of *Helicobacter* infection in the mouse model. *Gastroenterology* 136, 2237–2246 (e2231).
- Wheeldon, T.U., Hoang, T., Phung, D., Björkman, A., Granström, M., Sörberg, M., 2005. Long-term follow-up of *Helicobacter pylori* eradication therapy in Vietnam: reinfection and clinical outcome. *Aliment. Pharmacol. Ther.* 21, 1047–1053.
- Yang, X., Liu, W., Yang, W., Zhong, D., Liu, Y., Zhang, J., Jiang, J., Li, S., 2005. Oral immunization of mice with vaccine of attenuated *Salmonella typhimurium* expressing *Helicobacter pylori* urease B subunit. *Biomed. Environ. Sci.* 18, 411.
- Zhang, X., Zhang, J., Yang, F., Wu, W., Sun, H., Xie, Q., Si, W., Zou, Q., Yang, Z., 2015. Immunization with heat shock protein a and gamma-glutamyl transpeptidase induces reduction on the *Helicobacter pylori* colonization in mice. *PLoS One* 10, e0130391.