



Article (refereed) - postprint

Horton, Alice A.; Vijver, Martina G.; Lahive, Elma; Spurgeon, David J.; Svendsen, Claus; Heutink, Roel; van Bodegom, Peter M.; Baas, Jan. 2018. **Acute toxicity of organic pesticides to *Daphnia magna* is unchanged by co-exposure to polystyrene microplastics.** *Ecotoxicology and Environmental Safety*, 166. 26-34. <https://doi.org/10.1016/j.ecoenv.2018.09.052>

© 2018 Elsevier Inc.

This manuscript version is made available under the CC-BY-NC-ND 4.0  <https://creativecommons.org/licenses/by-nc-nd/4.0/>

This version available <http://nora.nerc.ac.uk/id/eprint/521060/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

NOTICE: this is the author's version of a work that was accepted for publication in *Ecotoxicology and Environmental Safety*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Ecotoxicology and Environmental Safety*, 166. 26-34. <https://doi.org/10.1016/j.ecoenv.2018.09.052>

www.elsevier.com/

Contact CEH NORA team
at noraceh@ceh.ac.uk

1 **Acute toxicity of organic pesticides to *Daphnia magna* is unchanged by co-exposure to**
2 **polystyrene microplastics**

3

4 Alice A. Horton ^{ab*}, Martina G. Vijver ^b, Elma Lahive ^a, David J. Spurgeon ^a,

5 Claus Svendsen ^a, Roel Heutink ^b, Peter M. van Bodegom ^b, Jan Baas ^{ab}

6

7 ^aCentre for Ecology and Hydrology, Maclean Building, Benson Lane, Wallingford,

8 Oxfordshire, OX10 8BB. UK.

9 ^bInstitute of Environmental Sciences, University of Leiden, P.O. Box 9518, 2300 RA Leiden,

10 The Netherlands.

11

12 *Corresponding author. Email address: alihort@ceh.ac.uk (A. A. Horton)

13

14

15 **Keywords**

16 Ecotoxicology, microbeads, log Kow, insecticide, dimethoate, deltamethrin.

17

18

19

20

21 **Abstract**

22 *Daphnia magna* were exposed to two pesticides in the presence or absence of
23 microplastics (300 000 particles ml⁻¹ 1 µm polystyrene spheres) and to microplastics alone.
24 The pesticides were dimethoate, an organophosphate insecticide with a low log Kow, and
25 deltamethrin, a pyrethroid insecticide with a high log Kow. *Daphnia* were exposed to a
26 nominal concentration range of 0.15, 0.31, 0.63, 1.25, 2.5, 5 mg l⁻¹ dimethoate and 0.016,
27 0.08, 0.4, 2, 5 and 10 µg l⁻¹ deltamethrin. Exposure to polystyrene microplastics alone showed
28 no effects on *Daphnia magna* survival and mobility over a 72 hour exposure. In the
29 dimethoate exposures, mobility and survival were both affected from a concentration of 1.25
30 mg l⁻¹, with effects were seen on mobility from 28 hours and survival from 48 hours, with
31 greater effects seen with increasing concentration and exposure time. In deltamethrin
32 exposures, survival was affected from a concentration of 0.4 µg l⁻¹ and mobility from a
33 concentration of 0.08 µg l⁻¹. Effects of deltamethrin on mobility were seen from 5 hours and
34 on survival from 28 hours, with greater effects on survival and mobility seen with increasing
35 concentration and exposure time. Contrary to expectations, pesticide toxicity to *Daphnia*
36 *magna* was not affected by the presence of microplastics, regardless of chemical binding
37 affinity (log Kow). This therefore suggests that polystyrene microplastics are unlikely to act
38 as a significant sink, nor as a vector for increased uptake of pesticides by aquatic organisms.

39

40

41 **Capsule**

42 Polystyrene microplastics are unlikely to act as vector for increased uptake of pesticides by
43 aquatic organisms

44

45 **Introduction**

46 Microplastics are a pollutant of increasing environmental concern based on their
47 ubiquitous and persistent nature. It is widely recognised that microplastics will form
48 biological and chemical associations within the environment. For example microplastics may
49 become associated with algae or bacteria (biofilms) (Hoellein et al., 2016; McCormick et al.,
50 2014) or may sorb organic chemicals due to their hydrophobic nature (Bakir et al., 2012;
51 Koelmans et al., 2016; Mato et al., 2001). The potential for association of hydrophobic
52 organic chemicals (HOCs) with microplastics has been recognised and has prompted studies
53 on whether this association will affect the bioavailability of HOCs, and thus their toxicity to
54 organisms. Studies have shown that microplastics can make HOCs either more bioavailable,
55 by acting as a vector for uptake following ingestion (Avio et al., 2015; Chen et al., 2017;
56 Rochman et al., 2013b), or less bioavailable due to strong irreversible binding of HOCs to
57 microplastics, removing HOCs from solution and remaining bound even if ingested
58 (Beckingham and Ghosh, 2016). It has even been suggested that microplastics may lead to
59 the removal of HOCs from body tissues following the ingestion of clean plastics by a
60 previously contaminated organism (Koelmans et al., 2013). The majority of studies on
61 microplastics and chemical associations to date have focussed on the marine environment.
62 However, concentrations of HOCs and microplastics in continental terrestrial and freshwater
63 environments are expected to be higher than marine environments due to their proximity to
64 the sources combined with limited dispersal and dilution, thus highlighting the importance of
65 studying terrestrial and freshwater systems (Dris et al., 2015; Horton et al., 2017).

66 The capacity for a chemical to bind to microplastics is, among other factors,
67 determined by its hydrophobicity, usually expressed as the log Kow value. Kow represents
68 the partition coefficient between octanol and water (Brooke, 2014). A chemical with a high
69 log Kow will have a lower water solubility than less hydrophobic substances (with a lower

70 log Kow), meaning that it will preferentially bind to organic particulate matter within the
71 system rather than remaining within solution (Lee et al., 2014; Mackay et al., 1980). It is
72 therefore expected that a chemical with a high log Kow (high hydrophobicity) will also have
73 a higher affinity for binding to microplastics in an aqueous system than a chemical with a
74 lower log Kow (higher hydrophilicity) (Wang et al., 2018). Such interactions can potentially
75 remove the chemical from solution and concentrate it on the surface of the plastic, thereby
76 changing bioavailability (Gouin et al., 2011; Lee et al., 2014; Velzeboer et al., 2014). The
77 aim of this study was therefore to investigate how the presence of microplastics would affect
78 the toxicity of high and low log Kow organic pesticides to a relevant freshwater organism, the
79 cladoceran *Daphnia magna*. Pesticides were chosen as their toxicity is well-documented. The
80 starting hypothesis was that the presence of microplastics within an aquatic solution would
81 reduce the toxicity of a pesticide with a high log Kow, due to its high binding capacity to the
82 microplastics making it less bioavailable (Beckingham and Ghosh, 2016; Koelmans et al.,
83 2013), whereas the toxicity of a low log Kow pesticide would be less affected by the presence
84 of microplastics.

85

86 **Materials and methods**

87 *The test chemicals*

88 We chose two pesticides to represent chemicals with high and low log Kow, both with
89 known toxicity to *Daphnia magna*. Dimethoate and deltamethrin were chosen both for their
90 differing chemical properties (specifically log Kow) and because they are environmentally
91 relevant, being representative of two widely used classes of insecticides. Both pesticides
92 target receptors associated with nervous system function to cause neurotoxicity. Dimethoate
93 is an organophosphate insecticide with a low log Kow (0.704) (Pesticide Properties Database,

94 2017b). It is relatively soluble in water (between 23.5-39.8 g l⁻¹ at 25°C) (Pesticide Properties
95 Database, 2017b; Sigma-Aldrich, 2017). It was first registered for use in 1962 and is still
96 widely applied to agricultural land worldwide (Van Scoy et al., 2016). Deltamethrin is a
97 pyrethroid insecticide also widely used in agriculture (Ren et al. 2009) and aquaculture (Ernst
98 et al. 2014). Deltamethrin is very poorly soluble in water, with a solubility between 0.2-2 µg
99 l⁻¹ at 25°C (Mestres and Mestres, 1992; Pesticide Properties Database, 2017a). Due to this
100 hydrophobic nature, with a log Kow reported between 4.6 (Kaneko, 2010) and 6.2 (PubChem
101 Compound Database, 2017), deltamethrin entering a water body would be expected to adsorb
102 readily to particulate matter such as microplastics, in addition to sediment and organic matter
103 (Lee et al., 2014; Lee et al., 2002).

104

105 *The test organism*

106 *Daphnia magna* is commonly used for ecotoxicological testing and as such, toxicity
107 data are readily available for *D. magna* for both deltamethrin and dimethoate toxicity
108 (Andersen et al., 2006; Toumi et al., 2013), as well as information on microplastic uptake and
109 toxicity (Besseling et al., 2014; Jemec et al., 2016; Rehse et al., 2016). This makes them an
110 ideal species for investigating how toxicity may be influenced by the interaction of these
111 pesticides with microplastics.

112 *D. magna* were taken from the Leiden University culture which has been continuously
113 maintained for over six years in the laboratory. According to the OECD guideline 202, *D.*
114 *magna* were cultured in glass containers with Artificial ElendtM4 medium at a density of 1
115 individual/10 ml of ElendtM4 medium (OECD, 2004). The culture medium was refreshed
116 twice a week. The test organisms were fed *ad libitum* with *Raphidocelis subcapitata* algae
117 and maintained inside a temperature controlled chamber (20 ± 1 °C) under a 16:8 light-dark

118 cycle. Throughout the duration of culturing, sensitivity of the test species was checked every
119 six months using the standardized toxicity test conducted with $K_2Cr_2O_7$ as a reference
120 compound (OECD, 2004).

121

122 *Preparation of the microplastic beads*

123 Microplastics as fluorescent polystyrene beads were purchased from Phosphorex
124 (USA) with a nominal size of 1 μm , as a solution containing DI water, an anti-microbial
125 agent (sodium azide) and a surfactant (Tween 20). The size of particles was confirmed by
126 TEM as being $1.2 \pm 0.2 \mu m$ (mean \pm SD) (Fig S1). Previous experimental studies have shown
127 that microplastics within the size range 20 nm – 5 μm are commonly ingested by *D. magna*,
128 as they represent a similar size range as their common algal food sources (Besseling et al.,
129 2014; Ogonowski et al., 2016; Rehse et al., 2016; Rist et al., 2017; Rosenkranz et al., 2009).
130 Both sodium azide and Tween 20 may act as toxicants and so the beads were washed in order
131 to remove these from the solution used for microplastic spiking. For washing, the supplied
132 stock of beads (1 ml) was diluted to approximately 12 ml with Milli-Q water, vortexed to mix
133 and then centrifuged at 5180 g (5000 rpm) (Beckman Coulter Avanti J-E centrifuge, USA)
134 for 5 minutes. The supernatant was then carefully pipetted leaving approximately 1 ml of
135 solution containing the particles at the bottom. These cleaning steps of dilution and
136 centrifuging were then repeated twice more to ensure maximum removal of the sodium azide
137 and Tween20. Following the final cleaning step the solution was diluted with Milli-Q water
138 to give a total stock solution volume of 10 ml. The number of beads per ml of this new bead
139 stock was measured using a flow cytometer (BD Accuri C6, BD Biosciences, USA). This
140 bead stock was used for spiking the test medium to a nominal concentration of 300 000
141 particles ml^{-1} . This concentration is roughly equivalent to the number of algal cells that

142 daphnids would be exposed to in an excess food situation (*i.e.* under culture conditions) and
143 equates to approximately $0.29 \mu\text{g ml}^{-1}$ ($287.7 \mu\text{g l}^{-1}$, calculations in SI).

144

145 *Preparation of the test solutions*

146 A dimethoate (PESTANAL[®], analytical standard, Sigma Aldrich Ltd, UK) stock
147 solution of 1 g l^{-1} was prepared directly in Elendt artificial freshwater. In order to produce the
148 required concentrations, the appropriate amount of stock solution was made up to 250 ml
149 with Elendt artificial freshwater. Based on toxicity values of dimethoate to *D. magna*, with 48
150 h LC₅₀ ranging from $0.86\text{-}2 \text{ mg l}^{-1}$ (Beusen and Neven, 1989; Syberg et al., 2008), exposure
151 concentrations were made in the range $0.156, 0.313, 0.625, 1.25, 2.5, 5 \text{ mg l}^{-1}$ ($0.68, 1.36,$
152 $2.73, 5.45, 10.9, 21.8 \mu\text{M}$).

153 To spike the test medium with deltamethrin it was necessary to dissolve it in a solvent
154 carrier due to its low solubility in water. Deltamethrin (certified reference material, Sigma-
155 Aldrich Ltd, UK) was dissolved in acetone to prepare a stock solution of $10\,000 \mu\text{g l}^{-1}$. A
156 serial dilution of this stock, was made by further dilution in acetone to create a deltamethrin
157 concentration series for spiking into artificial freshwater. A volume of $375 \mu\text{l}$ of the relevant
158 stock was added to 250 ml Elendt artificial freshwater (giving an acetone concentration of
159 0.15% within the exposure solution) in order to give the required exposure concentration
160 range: $0.016, 0.08, 0.4, 2, 5$ and $10 \mu\text{g l}^{-1}$ ($0.03, 0.16, 0.79, 3.96, 9.9, 19.79 \text{ nM}$). These
161 exposure concentrations were based on literature toxicity data for *D. magna* with 48 h LC₅₀s
162 ranging from $0.038\text{-}0.45 \mu\text{g l}^{-1}$ (Ren et al., 2009; Xiu et al., 1989) and 24 h LC₅₀s ranging
163 from $0.113\text{-}9.4 \mu\text{g l}^{-1}$ (Toumi et al., 2013; Xiu et al., 1989).

164 For both pesticides, treatments were prepared with and without microplastics. For the
165 microplastic treatments, the polystyrene bead stock solution was added to the exposure

166 solutions after the artificial freshwater had been spiked with the chemicals. The appropriate
167 volume of stock solution (as determined using the flow cytometer) was added to a volume of
168 250 ml of spiked solution to give a nominal concentration of 300 000 particles ml⁻¹. Four
169 replicates of 40 ml exposure solution held in 50 ml glass jars were prepared for each
170 treatment. With an average particle size of 1.2 µm ± 0.2 µm, the average surface area of the
171 microplastics within 40 ml was calculated as approx. 38-74 cm² dependent on variation in
172 particle size (surface area calculations are in SI). This concentration of particles provides a
173 comparable surface area to that of the glass vessel (40 ml water was calculated to cover
174 approx. 63 cm² of the internal surface area). Thus introduction of microplastics at this
175 concentration effectively doubles the surface area available for chemical binding. Each jar
176 was allowed to equilibrate for 24 hours before introduction of the organisms (Lee et al.,
177 2002).

178 Control treatments consisted of artificial freshwater only (further referred to as
179 'control'), artificial freshwater with microplastics only (equal to microplastic concentrations
180 in pesticide exposures: 300 000 particles ml⁻¹, further referred to as 'microplastic control'),
181 artificial freshwater with acetone (0.1 %, further referred to as 'acetone control'), and
182 artificial freshwater with both microplastics (300 000 particles ml⁻¹) and acetone (0.1%)
183 (further referred to as 'microplastic and acetone control'). These solutions were made and
184 distributed to glass jars 24 hours prior to introduction of daphnids as per pesticide treatments.

185

186 *Acute Toxicity Tests*

187 Following the equilibration period, five neonates (< 24 hours old) were added to each
188 jar. Errors were made in some vessels with 4 neonates added to a vessel (4 vessels overall) or
189 6 neonates added to a vessel (3 vessels overall). This was taken into account during the data

190 analysis. Jars were completely randomised throughout the exposure to avoid systematic bias.
191 *Daphnia* were observed at 5, 8, 21, 28, 48 and 72 hours. To enable resuspension of any
192 settled particles, each test jar was gently mixed at each observation point by drawing approx.
193 1-2 ml of exposure media in and out of a glass pipette three times. Aqueous pH was measured
194 in one jar from each concentration at the beginning and the end of the test. The organisms
195 were not fed for the duration of the experiment. Mortality was recorded as per OECD
196 protocol 202 (OECD, 2004). Impaired mobility was also recorded at each time point. This
197 was defined as an individual that was alive, as seen by the clear movement of limbs, but was
198 not able to swim effectively *i.e.* swimming erratically or not swimming effectively in a
199 forward direction, and additionally showing no response to gentle agitation with a glass
200 pipette tip. Sub-lethal behavioural effects are commonly seen in organisms when testing
201 pesticides with a neurotoxic mode of action (Desneux et al., 2007; Haynes, 1988; Sørensen et
202 al., 1995).

203

204 *Chemical analysis*

205 Water samples for chemical analysis were taken (1 ml dimethoate, 2 ml deltamethrin)
206 at 0, 24 and 72 hours after preparation of the solutions for deltamethrin treatments and 0 and
207 72 hours for dimethoate treatments. Fewer dimethoate measurements were taken than for
208 deltamethrin, as dimethoate was expected to be less complex in terms of chemistry, with
209 concentrations not expected to change over time (Eichelberger and Lichtenberg, 1971; Roast
210 et al., 1999). Samples were spun in 1ml glass tubes (2 tubes per sample) in a centrifuge at
211 approx. centrifugation 6000 G (8000 rpm) for 5 minutes (Eppendorf 24-place Fixed-angle
212 rotor, FA-45-24-11-HS) to remove microplastics and samples were subsequently stored in a
213 fridge at 5°C in the dark prior to analysis. Three replicate samples were taken from a medium

214 and a high nominal concentration for each chemical (0.625 and 5 mg l⁻¹ dimethoate, 0.4 and
215 10 µg l⁻¹ deltamethrin) at each of the above specified time points. Chemical analysis was
216 carried out by Wageningen Environmental Research (Alterra), and full details of chemical
217 sampling and analytical procedures are available in the Supplementary Information (SI).

218

219 *Data analysis*

220 To determine differences between treatments with and without microplastics at
221 different time points for each chemical, survival frequency data for each chemical were
222 analysed using a Chi-squared (χ^2) test (Microsoft Excel), where treatments without
223 microplastics were the ‘expected’ and those with microplastics were the ‘observed’. Mobility
224 frequency data were analysed using Fisher’s exact test (R statistical software) due to a
225 number of zero values (no daphnids swimming normally) which would not be accurately
226 represented using the χ^2 . Both tests accounted for any odd numbers where too few or too
227 many neonates had been added initially. Effects on survival and mobility with respect to
228 chemical concentrations and time were evaluated using ANOVA for each endpoint and each
229 chemical, with time points and concentrations considered as factors (R statistical software). A
230 *post-hoc* Tukey HSD test was carried out to determine pairwise differences with time and
231 concentration (R statistical software). Chemical data were analysed using ANOVA with time
232 considered as a factor. A *post-hoc* Tukey HSD test was carried out to determine pairwise
233 differences with time and nominal concentration (R statistical software).

234 Further analyses of the survival data over time were carried out using a process-based
235 survival model. The model assumes that the toxicant must be first taken up in the organism
236 before it can exert an effect. The kinetics are described with a one-compartment model and
237 the effects is described with the ‘stochastic death’ model. The model is extensively described

238 in Jager et al. (2006) and Kooijman and Bedaux (1996). This model is accepted by the OECD
239 (OECD, 2006), where an additional elaborate (mathematical) description can be found with
240 examples of the use of the model. The model links exposure concentrations to a survival
241 probability using three parameters for the whole time-course of the exposure (the No Effect
242 Concentration (NEC): a threshold for toxic effects, the killing rate (k_r): a measure for the
243 toxic potency of the compound, and the elimination rate (k_e) as a kinetic parameter).

244 Parameter values for dimethoate were calculated using the known (measured)
245 chemical exposure concentrations and the survival data. The parameter values were
246 subsequently compared to independent values obtained from literature for verification. For
247 deltamethrin, the uncertainties related to the actual exposure concentrations prompted a
248 ‘reverse modelling’ approach. Literature toxicity values for deltamethrin to *D. magna* (Xiu et
249 al., 1989) were used to derive the model parameters, which were subsequently used to fit the
250 model output to the survival data, allowing back-calculation of actual exposure
251 concentrations (further details on this approach are available in the SI). The benefits of
252 including the modelling are threefold: 1) to validate the results of the traditional statistical
253 analysis, 2) to calculate the actual concentrations of pesticides that the *Daphnia* are exposed
254 to and 3) to determine toxicity effects over time, allowing for extrapolation of toxicity
255 estimates beyond the timeframe of the experiments. Together, these benefits allowed us to
256 better understand the dynamics of toxicity within the experiment.

257

258 **Results**

259 *Daphnia survival*

260 *Daphnia* survival in the controls without microplastics or chemicals, and in the
261 acetone controls, was 100%. This high control survival validates the criteria of the toxicity

262 test according to OECD guidelines for *Daphnia magna* acute toxicity testing (OECD, 2004).
263 Microplastics alone did not affect survival over the 72 hour test period with only one
264 mortality in the microplastic control treatment (5%) after the 72 hour exposure period and
265 100% survival in the microplastics and acetone control treatments. Without the use of a
266 microscope, microplastics were clearly visible within the guts of daphnids as a white mass.

267 There was a significant effect of pesticide exposure concentration on survival ($p <$
268 0.01 for both pesticides, ANOVA). There were also a significant effect of exposure time on
269 survival ($p < 0.01$ for both pesticides, ANOVA) and a significant interaction between
270 concentration and time also occurred ($p < 0.01$ for both pesticides, ANOVA). Over the 72 h
271 exposure, significant effects were seen on survival at exposure concentrations above 1.25 mg
272 l^{-1} for dimethoate ($p < 0.01$, ANOVA + Tukey HSD) and above $0.4 \text{ } \mu\text{g l}^{-1}$ for deltamethrin (p
273 < 0.05 , ANOVA + Tukey HSD). When considering time, significant effects on survival were
274 seen from 48 hours in dimethoate treatments above 2.5 mg l^{-1} ($p < 0.01$, ANOVA + Tukey
275 HSD, Table 1a) and from 28 hours in deltamethrin treatments above $2 \text{ } \mu\text{g l}^{-1}$ ($p < 0.01$,
276 ANOVA + Tukey HSD, Table 2a). For both pesticides there was no significant difference in
277 the survival of organisms based on the presence or absence of microplastics ($p > 0.05$ at
278 every time point, χ^2) To give a visual representation of this similarity, the survival and
279 mobility probability was calculated and the deviance between treatments with and without
280 microplastics depicted (Figs. 1a and 2a). Deviance was calculated as the difference in
281 survival (or mobility) probabilities for treatments without MPs (– MP) vs. those with MPs (+
282 MP) at given concentrations.

283
284
285

Table 1. Survival probabilities (Table 1a) and probabilities of normal mobility (Table 1b) for *D. magna* exposed to dimethoate at each concentration and time point, calculated by dividing the remaining surviving neonates by the original 20 to give a probability between 0-1.

Table 1a)		Time (h)						
Dimethoate exposure concentration (mg l ⁻¹)		0	5	8	21	28	48	72
0	Without MP	1	1	1	1	1	1	1
0	With MP	1	1	1	1	1	1	1
0.156	Without MP	1	1	1	1	1	1	1
0.156	With MP	1	1	1	1	1	0.9	0.8
0.313	Without MP	1	1	1	1	1	1	1
0.313	With MP	1	1	1	1	1	0.6	0.5
0.625	Without MP	1	1	1	1	1	1	1
0.625	With MP	1	1	1	1	1	0.9	0.6
1.25	Without MP	1	1	1	1	1	1	0.6
1.25	With MP	1	1	1	1	1	0.9	0.7
2.5	Without MP	1	1	1	0.8	0.8	0.4	0
2.5	With MP	1	1	1	1	1	0.6	0
5	Without MP	1	1	1	0.7	0.7	0.2	0.1
5	With MP	1	1	1	0.9	0.8	0.2	0

Table 1b)		Time (h)						
Dimethoate exposure concentration (mg l ⁻¹)		0	5	8	21	28	48	72
0	Without MP	1	1	1	1	1	1	1
0	With MP	1	1	1	1	1	1	1
0.156	Without MP	1	1	1	1	1	1	1
0.156	With MP	1	1	1	1	1	1	0.8
0.313	Without MP	1	1	1	1	1	1	1
0.313	With MP	1	1	1	1	1	0.8	0.4
0.625	Without MP	1	1	1	1	1	1	1
0.625	With MP	1	1	1	1	1	0.9	0.5
1.25	Without MP	1	1	1	1	1	0.9	0.4
1.25	With MP	1	1	1	1	1	0.7	0.6
2.5	Without MP	1	1	1	1	0.5	0	0
2.5	With MP	1	1	1	0.9	0.7	0.2	0
5	Without MP	1	1	1	0.7	0.3	0	0
5	With MP	1	1	1	0.4	0.4	0	0

Table 2. Survival probabilities (Table 2a) and probabilities of normal mobility (Table 2b) for *D. magna* exposed to deltamethrin at each concentration and time point, calculated by dividing the remaining surviving neonates by the original 20 to give a probability between 0-1.

Table 2a)		Time (h)						
Deltamethrin exposure concentration (µg l ⁻¹)		0	5	8	21	28	48	72
0	Without MP	1	1	1	1	1	1	1
0	With MP	1	1	1	1	1	0.9	1
0.016	Without MP	1	1	1	1	1	1	1
0.016	With MP	1	1	1	1	1	1	1
0.08	Without MP	1	1	1	1	1	1	0
0.08	With MP	1	1	1	1	1	0.9	0
0.4	Without MP	1	1	1	0.9	1	0.9	0
0.4	With MP	1	1	1	1	0.9	0.7	0
2	Without MP	1	1	1	0.9	0.7	0.5	0
2	With MP	1	1	1	1	0.7	0.6	0
5	Without MP	1	1	1	0.7	0.7	0.2	0
5	With MP	1	1	1	0.8	0.8	0.5	0
10	Without MP	1	1	1	1	0.7	0.3	0
10	With MP	1	1	1	1	0.6	0.2	0

Table 2b)		Time (h)						
Deltamethrin exposure concentration (µg l ⁻¹)		0	5	8	21	28	48	72
0	Without MP	1	1	1	1	1	1	1
0	With MP	1	1	1	1	1	0.9	1
0.016	Without MP	1	1	1	1	1	1	1
0.016	With MP	1	1	1	1	1	1	1
0.08	Without MP	1	1	1	0.9	1	0.7	0
0.08	With MP	1	1	1	1	1	0.6	0
0.4	Without MP	1	1	1	0.7	0.4	0.2	0
0.4	With MP	1	1	1	0.7	0.3	0	0
2	Without MP	1	0.9	0.8	0.1	0.1	0	0
2	With MP	1	1	0.8	0.1	0	0	0
5	Without MP	1	0.9	0.7	0.1	0.1	0	0
5	With MP	1	1	0.6	0.1	0.2	0.1	0
10	Without MP	1	0.4	0.6	0.1	0	0	0
10	With MP	1	0.4	0.6	0.2	0	0	0

286 *Daphnid mobility*

287 There were also concentration-dependent effects on daphnid mobility. There was a
288 significant effect of pesticide exposure concentration on mobility ($p < 0.01$ for both
289 pesticides, ANOVA). There were also a significant effect of exposure time on mobility for
290 both chemicals ($p < 0.01$ for both pesticides, ANOVA) and a significant interaction between
291 concentration and time also occurred for both chemicals (ANOVA, $p < 0.01$ for both
292 chemicals). Over the 72 h exposure, significant mobility impairment was observed in
293 *Daphnia* exposed to dimethoate at concentrations of 1.25 mg l^{-1} and above ($p < 0.01$,
294 ANOVA + Tukey HSD). Similarly *Daphnia* exposed to $0.08 \text{ } \mu\text{g l}^{-1}$ deltamethrin and above
295 suffered significant mobility impairment ($p < 0.05$, ANOVA + Tukey HSD). When
296 considering time, significant effects on mobility were seen from 21 hours for dimethoate at 5
297 mg l^{-1} ($p < 0.01$, ANOVA + Tukey HSD, Table 1b) and from 5 hours for deltamethrin at 10
298 $\mu\text{g l}^{-1}$ ($p < 0.01$, ANOVA + Tukey HSD, Table 2b). The presence of microplastics resulted in
299 no significant difference in the number of daphnids suffering impaired mobility for either
300 chemical at any time point ($p > 0.05$, Fisher's exact test). As for survival, plots for deviance
301 were created to give a visual representation of this similarity using deviance in probability of
302 normal mobility of treatments with vs. without microplastics (Figs 1b and 2b). Effects on
303 mobility were seen at earlier time points than effects on survival, as would be expected given
304 that sublethal behavioural effects are a precursor to mortality.

305

306 *Chemical concentrations*

307 The pH remained consistent throughout the test with a mean pH of $7.81 (\pm 0.17 \text{ SD})$
308 across treatments at 0 hrs and $7.9 (\pm 0.05 \text{ SD})$ at 72 hours.

309 All measured dimethoate concentrations were lower than the nominal concentrations,
310 ranging from (average) 59-63% of nominal values, although this difference was not
311 significant ($p > 0.05$, t-test, Table S1). Measured concentrations of dimethoate did not vary
312 significantly over time ($p > 0.05$, ANOVA) and there was no effect of microplastics on the
313 measured concentrations of dimethoate ($p > 0.05$, ANOVA) (Figs. 3a and 3b). There was no
314 significant effect of microplastics on concentration over time (interaction $p > 0.05$, ANOVA).

315 There was a significant difference between nominal and measured deltamethrin
316 concentrations ($p < 0.01$, t-test), with average measured concentrations ranging from 3.7-
317 20.5% of the nominal concentrations (Table S2). Due to an apparent difference in trend
318 between the low and high nominal concentrations measured (Figs. 4a and 4b), these were
319 analysed separately to tease apart concentration-dependent effects. At the low nominal
320 concentration ($0.4 \mu\text{g l}^{-1}$), there was no effect of microplastics or time on the measured
321 concentrations (both $p > 0.05$, ANOVA), nor an interaction of time and microplastics ($p >$
322 0.05 , ANOVA). At the highest nominal concentration ($10 \mu\text{g l}^{-1}$), both microplastics and time
323 significantly influenced the measured concentrations, with concentrations lower when
324 microplastics were present (both microplastics and time $p < 0.01$, ANOVA), and with an
325 initial significant decrease in concentration up to 24 hours (0-24 h, $p < 0.01$, ANOVA +
326 Tukey HSD, 24-72 h, $p > 0.05$, ANOVA + Tukey HSD). There was no significant effect of
327 microplastics on concentration over time (interaction $p > 0.05$, ANOVA).

328

329 *Model analysis*

330 Fitting of separate stochastic death models for both dimethoate and deltamethrin gave
331 an estimation of toxicity over time at the experimental exposure concentrations and provided
332 a consistent fit with the survival data (Figs. S2 and S3). For dimethoate, the model-derived

333 LC₅₀ was 0.5 mg l⁻¹ (the full range of model-derived LC_x values for dimethoate available in
 334 Table S6). For deltamethrin, the model-derived LC₅₀ was 0.023 µg l⁻¹ (the full range of
 335 model-derived LC_x values for deltamethrin are available in Table S7). For both pesticides, the
 336 model shows no difference in pesticide exposure, or survival, with or without microplastics.
 337 For deltamethrin, using the reverse modelling approach, the survival data were used to
 338 determine the actual exposure concentrations as an indirect and complementary assessment of
 339 the measured concentrations (Table 3).

340

341 Table 3. Nominal concentration range of deltamethrin compared to modelled exposure
 342 concentrations and measured concentrations.

343

Nominal Concentration (µg l ⁻¹)	Nominal Concentration (nM)	Modelled Concentration (µg l ⁻¹)	Modelled Concentration (nM)	Measured Concentration (µg l ⁻¹)
0.016	0.03	0.012	0.024	-
0.08	0.16	0.03	0.06	-
0.4	0.79	0.04	0.079	0.05
2	3.96	0.08	0.16	-
5	9.9	0.08	0.16	-
10	19.79	0.09	0.18	0.40

344

345 The reverse modelling to predict actual exposure concentrations indicated that the
 346 concentrations in the three highest test treatments are more or less equal. This is likely
 347 governed by the solubility limit, which would therefore be around 0.08-0.09 µg l⁻¹ (close to
 348 the reported value of 0.2 µg l⁻¹ (Mestres and Mestres, 1992; Pesticide Properties Database,
 349 2017a). The reported 48 h LC₅₀ taken from literature that informed the parameters used for
 350 this model estimation was at the lower end of the scale: 0.038 µg l⁻¹ (Xiu et al., 1989),
 351 compared to 0.32-0.63 µg l⁻¹ reported by (Toumi et al., 2013), although is comparable to that
 352 reported in other studies (0.05-0.6 µg l⁻¹ reported by (Day and Maguire, 1990). With higher
 353 input values the calculated exposure concentrations may have been higher.

354

355 **Discussion**

356 *Biological effects*

357 Although microplastics are commonly implicated in causing physiological damage to
358 organisms, leading to reduced fitness and mortality (Lee et al., 2013; Rehse et al., 2016;
359 Wright et al., 2013), no microplastic-specific effects on mobility or survival were seen in this
360 acute test, despite the high concentration of microplastics used and visual confirmation of
361 ingestion. This result is in accordance with a number of other studies where high
362 concentrations of microplastics were shown to cause no observable detrimental effects
363 (Hämer et al., 2014; Kaposi et al., 2014; Weber et al., 2018). Although other acute studies
364 have measured subtle effects of exposure to microplastics that may have occurred, for
365 example immune responses, gut blockage, reduced assimilation efficiency or reduced scope
366 for growth (Blarer and Burkhardt-Holm, 2016; Cole et al., 2015; Jeong et al., 2016; Lo and
367 Chan, 2018), these were beyond the scope of this study which was not planned to determine
368 the effects of microplastics alone, but to determine whether the presence of microplastics
369 influenced the toxic effects of pesticides.

370 Contrary to the hypothesis that microplastics would lead to a reduction in toxic effect
371 of the high log Kow pesticide deltamethrin, the results showed no alteration in the acute
372 toxicity of either deltamethrin or dimethoate to *D. magna*, regardless of the chemical binding
373 capacity (log Kow) (Figs. 1 and 2, Tables 1 and 2). Mortality and mobility impairment
374 increased with concentration and time for both pesticides, as expected, however the
375 concentrations at which detrimental effects occurred were not influenced by the presence of
376 microplastics. This is also highlighted by the results of the stochastic death modelling.

377

378 *Linking biological effects to chemical exposure*

379 The measured concentrations for deltamethrin were significantly lower than expected
380 across all treatments, on average between 3.7-20.5 % the nominal concentration, depending
381 on the time the sample was taken and the presence of microplastics (Fig. 3). Measured
382 concentrations were highly variable, especially at the lower measured concentrations when
383 microplastics were present (Fig. 4a). Additional replicate samples would have helped to
384 reduce this variability and may have helped to clarify whether the lack of significance was
385 simply due to high variability. However, regardless of the significant differences found in
386 measured deltamethrin concentrations between treatments with and without microplastics at
387 higher concentrations (Fig. 4b), no differences in toxicity were observed. This highlights that
388 the chemical dynamics within the system were complex and that while some binding of
389 pesticides to microplastics may have occurred, this did not reduce the bioavailability of the
390 two pesticides enough to lower the resulting observed toxicity. As predicted, there was no
391 significant difference in water concentration with or without microplastics for dimethoate,
392 supporting the lack of difference in the survival and mobility data, and no significant change
393 in concentrations over time (Fig. 3). This difference between deltamethrin and dimethoate
394 highlights that hydrophobicity of chemicals can influence binding and removal from solution,
395 influencing different chemicals in different ways, however toxicity is more complex to
396 predict.

397 Due to the high hydrophobicity of deltamethrin, it is likely that this pesticide bound
398 strongly to both the glass vessel and the microplastic particles (where present) (Lee et al.,
399 2002; Sethi et al., 2014; Wheelock et al., 2005). To overcome this we introduced a 24 h
400 equilibrium period following the suggestion made by Lee et al. (2002). Nonetheless it turned
401 out extremely difficult to make accurate quantifications of the deltamethrin concentrations in
402 water, as deltamethrin is also likely bind to organic matter including the *Daphnia* and any
403 associated organic detritus or excreta. This means that, despite the 24 h equilibration phase,

404 the equilibrium likely shifted when the *Daphnia* were introduced to the solution, highlighted
405 by the significant reduction in concentration within the aqueous solution within the first 24
406 hours. This is a highly dynamic system and the equilibrium is likely to continue to shift over
407 time leading the chemical to be associated with different substrates at different times. This
408 highlights the complexity of working with deltamethrin, with binding, availability and ease of
409 chemical extraction dependent on substrates available and methods used.

410 Due to the discrepancy between measured and nominal concentrations for
411 deltamethrin, we were not able to directly relate toxicity to nominal or measured chemical
412 concentrations. It was for these reasons that we carried out the reverse modelling approach to
413 determine the likely exposure concentrations the *Daphnia* were exposed to (Table 3) and thus
414 enable us to determine the toxicity of deltamethrin (SI). The model showed that, probably as
415 a result of the limit of solubility of the hydrophobic insecticide, the top three concentrations
416 of deltamethrin (nominal concentrations 2, 5, and 10 $\mu\text{g l}^{-1}$) were in fact likely to have been
417 almost identical at 0.08-0.09 $\mu\text{g l}^{-1}$ (Table 3). This was reflected in the survival and mobility
418 matrices showing survival and mobility to be comparable across the top three concentrations
419 (comparing top three concentrations across survival and mobility, all $p > 0.05$ ANOVA +
420 Tukey HSD, Table 2). This highest calculated exposure concentration was below the
421 expected lower limit of solubility (0.2 $\mu\text{g l}^{-1}$ at 25°C). This could be due to the combined
422 effects of a lower temperature than stated for maximum solubility (experiments were run at
423 20°C \pm 1°C) and additional dissolved constituents in the Elendt artificial freshwater, both of
424 which may have led to a decreased capacity for dissolution.

425 Although the highest concentrations of deltamethrin used in this study were above
426 solubility, the actual value for solubility is uncertain, reported between 0.2-2 $\mu\text{g l}^{-1}$ (Mestres
427 and Mestres, 1992). EC₅₀ values for deltamethrin for effects on mortality and immobilisation
428 in *D. magna* reported in the literature are highly variable, ranging from 0.11 to 9.4 $\mu\text{g l}^{-1}$ at 24

429 h and 0.03 to 0.63 $\mu\text{g l}^{-1}$ at 48 h (Toumi et al., 2013; Xiu et al., 1989). The highest of these
430 values, particularly for the 24 h exposure time, hence are well above stated solubility. In this
431 study, the modelled 96 h LC_{50} of 0.023 $\mu\text{g l}^{-1}$ is in the same order of magnitude as the
432 literature value of 0.01 $\mu\text{g l}^{-1}$ calculated by Xiu et al. (1989), although it should be noted that
433 their calculation was based on nominal concentrations. Many studies focus solely on nominal
434 concentrations, not taking into account solubility or binding issues, while studies that do seek
435 to determine concentrations find measured concentrations to be vastly reduced from nominal
436 values (Lee et al., 2002; Toumi et al., 2013; Wheelock et al., 2005).

437 The modelling allowed us to compare the toxicity observed in this study to literature
438 data (SI) and enabled us to develop a better understanding of the biological effects seen under
439 given chemical and microplastics exposures. For dimethoate, measured concentrations were
440 much closer to stated nominal concentrations, and were consistent over time. Model
441 estimations for toxicity of dimethoate in this study based on the measured chemical data
442 showed exposures to be comparable with or without microplastics, with our LC_{50} results
443 shown to be comparable to literature values (SI).

444

445 *Binding of pesticides to microplastics*

446 Different polymers have different affinities for chemical binding and therefore may
447 have differing propensities for altering the toxicity of associated chemicals. For example, it
448 has been reported that polyethylene and polypropylene will have greater affinities for
449 chemical sorption than polyvinyl chloride (PVC) or polyethylene terephthalate (PET)
450 (Rochman et al., 2013a). Polystyrene has been suggested as having a lower affinity for
451 hydrophobic chemical sorption than polyethylene, but higher than PVC (Wang and Wang,
452 2018). It is nonetheless recognised that polystyrene will associate with hydrophobic organic

453 chemicals within the environment (Liu et al., 2015; Rochman et al., 2013c). The
454 concentration of polystyrene particles used in this experiment ($300\,000\text{ particles ml}^{-1}$) is far
455 above the concentrations that will likely be found within the freshwater environment (see
456 Horton et al. (2017) for an overview of freshwater microplastic studies), although this
457 exposure level is within the range of other experimental studies using microplastics (Lu et al.,
458 2016; Ogonowski et al., 2016; Rehse et al., 2016; Setälä et al., 2014). This study was
459 therefore intended to give a representation of the possible effects of interactions between
460 microplastics, pesticides and freshwater organisms in a scenario where microplastics were
461 highly abundant.

462 The presence of microplastics would have provided an increased surface area
463 available for chemical binding (in this instance the surface area of the microplastics was
464 calculated to be approximately equivalent to that of the vessel, effectively doubling the
465 surface area). Therefore a lower concentration of deltamethrin would have been expected in
466 the water when microplastics were present. The chemical measurement results confirm this
467 effect, as at the highest exposure concentration of deltamethrin (nominal concentration of $10\text{ }\mu\text{g/l}$),
468 water concentrations were significantly lower when microplastics were present (Fig.
469 4b). This implies that deltamethrin was binding to microplastics (inferred by a reduced
470 concentration in water when compared to an equivalent nominal concentration without
471 microplastics). However, it is important to note that despite the difference with and without
472 microplastics at the highest concentration of deltamethrin (nominal concentration $10\text{ }\mu\text{g l}^{-1}$),
473 the reduced concentration in the presence of microplastics was not observed at the lower
474 concentration measured (nominal concentration $0.4\text{ }\mu\text{g l}^{-1}$) (Fig. 4a). In the higher nominal
475 exposure levels ($10\text{ }\mu\text{g l}^{-1}$), the decline in measured concentration continues after the 24 h
476 equilibration period highlighting the complex chemical dynamics within the solution, with
477 the introduction of daphnia likely to alter the equilibrium. Questions remain surrounding the

478 dynamics and kinetics of chemical behaviour and toxicity in relation to the presence of
479 microplastics. However, as there were no significant effects on survival and mobility between
480 microplastic and non-microplastic treatments in this study, these complex dynamics do not
481 appear to affect the overall bioavailability, and as a result, acute toxicity of the chemicals.

482

483 *Outlook*

484 If effects are to be seen with respect to chemicals in association with microplastics,
485 especially their facilitation of chemical uptake and toxicity, it is most likely that these would
486 be seen under controlled laboratory conditions where uncontaminated organisms are exposed
487 to contaminated plastics (of a size that enables ingestion), as opposed to in the environment
488 where organisms will already have been exposed to a variety of different chemicals
489 (Koelmans et al., 2016). This study was designed to enable optimum chemical binding and
490 ingestion of microplastics by *D. magna*. Given the high concentration of microplastics in this
491 study and, thus, the high surface area available for binding, an alteration in the bioavailability
492 and toxicity of hydrophobic deltamethrin (high log K_{ow}) would have been expected, whereas
493 dimethoate (low log K_{ow}) would be expected to be consistently bioavailable and toxic
494 regardless of the presence of microplastics (Cole et al., 2011; Teuten et al., 2009). In contrast,
495 our results show that there was no effect of microplastics on the response of daphnids to
496 either of the two pesticides, despite the very different chemical characteristics. The vector
497 effects, or so-called ‘Trojan Horse’ effects, as ascribed to microplastics (Rochman et al.,
498 2014; Rochman et al., 2013c) were not observed. It is therefore unlikely that microplastics
499 will exert short-term effects on pesticide toxicity under real field conditions where sediment
500 and organic matter would compete with microplastics for binding of chemicals. Additionally,
501 in areas highly polluted with pesticides or other organic chemicals, the presence of

502 microplastics is unlikely to alter the availability of these pollutants (Tanaka et al., 2018). In
503 terms of chemical toxicity associated with microplastics, it is feasible that plasticisers will
504 pose a greater chemical risk to organisms than sorbed hydrophobic chemicals (Devriese et al.,
505 2017; Lohmann, 2017). Although polymer, particle and chemical-specific, these data are a
506 valuable contribution to the wider understanding of microplastic and chemical associations,
507 and the complexities underlying these mechanisms.

508

509 **Acknowledgements**

510 We would like to thank Laura Buijse and Steven Crum at Wageningen Alterra for conducting
511 the chemical analysis, plus Els Baalbergen and Marinda van Pomeran for additional
512 assistance in the lab. Many thanks to Kerstin Jurkschat, University of Oxford, for
513 characterising the polystyrene particles by TEM. This work was funded by the UK Natural
514 Environment Research Council through National Capability funding of the Centre for
515 Ecology and Hydrology Pollution and Environmental Risk Program.

516

517 **References**

- 518 Andersen, T. H., et al., 2006. Acute and chronic effects of pulse exposure of *Daphnia magna* to
519 dimethoate and pirimicarb. *Environmental Toxicology and Chemistry*. 25, 1187-1195.
- 520 Avio, C. G., et al., 2015. Pollutants bioavailability and toxicological risk from microplastics to marine
521 mussels. *Environmental Pollution*. 198, 211-22.
- 522 Bakir, A., et al., 2012. Competitive sorption of persistent organic pollutants onto microplastics in the
523 marine environment. *Marine Pollution Bulletin*. 64, 2782-9.
- 524 Beckingham, B., Ghosh, U., 2016. Differential bioavailability of polychlorinated biphenyls associated
525 with environmental particles: Microplastic in comparison to wood, coal and biochar.
526 *Environmental Pollution*. 220, 150-158.
- 527 Besseling, E., et al., 2014. Nanoplastic affects growth of *S. obliquus* and reproduction of *D. magna*.
528 *Environmental Science & Technology*. 48, 12336-43.
- 529 Beusen, J.-M., Neven, B., 1989. Toxicity of Dimethoate to *Daphnia magna* and freshwater fish.
530 *Bulletin of Environmental Contamination and Toxicology*. 42, 126-133.
- 531 Blarer, P., Burkhardt-Holm, P., 2016. Microplastics affect assimilation efficiency in the freshwater
532 amphipod *Gammarus fossarum*. *Environmental Science and Pollution Research*
533 *International*. 23, 23522-23532.
- 534 Brooke, N., Physicochemical Properties. In: N. Bradley, et al., Eds.), *Essentials of Environmental*
535 *Public Health Science: A Handbook for Field Professionals*. OUP Oxford, 2014, pp. 9-19.
- 536 Chen, Q., et al., 2017. Pollutants in Plastics within the North Pacific Subtropical Gyre. *Environmental*
537 *Science & Technology*.
- 538 Cole, M., et al., 2015. The impact of polystyrene microplastics on feeding, function and fecundity in
539 the marine copepod *Calanus helgolandicus*. *Environmental Science & Technology*. 49, 1130-
540 7.
- 541 Cole, M., et al., 2011. Microplastics as contaminants in the marine environment: a review. *Marine*
542 *Pollution Bulletin*. 62, 2588-97.
- 543 Day, K. E., Maguire, R. J., 1990. Acute toxicity of isomers of the pyrethroid insecticide deltamethrin
544 and its major degradation products to *Daphnia magna*. *Environmental Toxicology and*
545 *Chemistry*. 9, 1297-1300.
- 546 Desneux, N., et al., 2007. The sublethal effects of pesticides on beneficial arthropods. *Annual Review*
547 *of Entomology*. 52, 81-106.
- 548 Devriese, L. I., et al., 2017. Bioaccumulation of PCBs from microplastics in Norway lobster (*Nephrops*
549 *norvegicus*): An experimental study. *Chemosphere*. 186, 10-16.
- 550 Dris, R., et al., 2015. Beyond the ocean: contamination of freshwater ecosystems with (micro-)
551 plastic particles. *Environmental Chemistry*. 12, 539-550.
- 552 Eichelberger, J. W., Lichtenberg, J. J., 1971. Persistence of pesticides in river water. *Environmental*
553 *Science & Technology*. 5, 541-544.
- 554 Gouin, T., et al., 2011. A Thermodynamic Approach for Assessing the Environmental Exposure of
555 Chemicals Absorbed to Microplastic. *Environmental Science & Technology*. 45, 1466-1472.
- 556 Hämer, J., et al., 2014. Fate of microplastics in the marine isopod *Idotea emarginata*. *Environmental*
557 *Science & Technology*. 48, 13451-8.
- 558 Haynes, K. F., 1988. Sublethal Effects of Neurotoxic Insecticides on Insect Behavior. *Annual Review of*
559 *Entomology*. 33, 149-168.
- 560 Hoellein, T., et al., Consider a source: Microplastic in rivers is abundant, mobile, and selects for
561 unique bacterial assemblages. *Ocean Sciences Meeting New Orleans, Louisiana, USA, 2016*.
- 562 Horton, A. A., et al., 2017. Microplastics in freshwater and terrestrial environments: Evaluating the
563 current understanding to identify the knowledge gaps and future research priorities. *Science*
564 *of the Total Environment*. 586, 127-141.
- 565 Jager, T., et al., 2006. Making sense of ecotoxicological test results: towards application of process-
566 based models. *Ecotoxicology*. 15, 305-314.

567 Jemec, A., et al., 2016. Uptake and effects of microplastic textile fibers on freshwater crustacean
568 *Daphnia magna*. Environmental Pollution. 219, 201-209.

569 Jeong, C. B., et al., 2016. Microplastic Size-Dependent Toxicity, Oxidative Stress Induction, and p-JNK
570 and p-p38 Activation in the Monogonont Rotifer (*Brachionus koreanus*). Environmental
571 Science & Technology. 50, 8849-57.

572 Kaneko, H., Pyrethroid Chemistry and Metabolism. In: R. Krieger, (Ed.), Hayes' Handbook of Pesticide
573 Toxicology. Elsevier Science, 2010, pp. 1635-1663.

574 Kaposi, K. L., et al., 2014. Ingestion of microplastic has limited impact on a marine larva.
575 Environmental Science & Technology. 48, 1638-45.

576 Koelmans, A. A., et al., 2016. Microplastic as a Vector for Chemicals in the Aquatic Environment:
577 Critical Review and Model-Supported Reinterpretation of Empirical Studies. Environmental
578 Science & Technology. 50, 3315-26.

579 Koelmans, A. A., et al., 2013. Plastic as a carrier of POPs to aquatic organisms: a model analysis.
580 Environmental Science & Technology. 47, 7812-20.

581 Kooijman, S., Bedaux, J., 1996. Analysis of toxicity tests on *Daphnia* survival and reproduction. Water
582 Research. 30, 1711-1723.

583 Lee, H., et al., 2014. Sorption capacity of plastic debris for hydrophobic organic chemicals. Science of
584 the Total Environment. 470-471, 1545-52.

585 Lee, K. W., et al., 2013. Size-dependent effects of micro polystyrene particles in the marine copepod
586 *Tigriopus japonicus*. Environmental Science & Technology. 47, 11278-83.

587 Lee, S., et al., 2002. Recovery of synthetic pyrethroids in water samples during storage and
588 extraction. Journal of Agricultural and Food Chemistry. 50, 7194-7198.

589 Liu, L., et al., 2015. Sorption of polycyclic aromatic hydrocarbons to polystyrene nanoplastic.
590 Environmental Toxicology and Chemistry. 35, 1650-1655.

591 Lo, H. K. A., Chan, K. Y. K., 2018. Negative effects of microplastic exposure on growth and
592 development of *Crepidula onyx*. Environmental Pollution. 233, 588-595.

593 Lohmann, R., 2017. Microplastics are not important for the cycling and bioaccumulation of organic
594 pollutants in the oceans-but should microplastics be considered POPs themselves? Integr
595 Environ Assess Manag. 13, 460-465.

596 Lu, Y., et al., 2016. Uptake and Accumulation of Polystyrene Microplastics in Zebrafish (*Danio rerio*)
597 and Toxic Effects in Liver. Environmental Science & Technology. 50, 4054-4060.

598 Mackay, D., et al., 1980. Relationships between aqueous solubility and octanol-water partition
599 coefficients. Chemosphere. 9, 701-711.

600 Mato, Y., et al., 2001. Plastic resin pellets as a transport medium for toxic chemicals in the marine
601 environment. Environmental Science & Technology. 35, 318-324.

602 McCormick, A., et al., 2014. Microplastic is an abundant and distinct microbial habitat in an urban
603 river. Environmental Science & Technology. 48, 11863-71.

604 Mestres, R., Mestres, G., Deltamethrin: Uses and Environmental Safety. In: G. W. Ware, (Ed.),
605 Reviews of Environmental Contamination and Toxicology: Continuation of Residue Reviews.
606 Springer New York, New York, NY, 1992, pp. 1-18.

607 OECD, 2004. Test No. 202: *Daphnia* sp. Acute Immobilisation Test. OECD Publishing.

608 OECD, 2006. Current Approaches in the Statistical Analysis of Ecotoxicity Data: A Guidance to
609 Application (ENV/JM/MONO(2006)18). OECD Publishing, Paris.

610 Ogonowski, M., et al., 2016. The Effects of Natural and Anthropogenic Microparticles on Individual
611 Fitness in *Daphnia magna*. PLOS ONE. 11, e0155063.

612 Pesticide Properties Database, Deltamethrin. University of Hertfordshire, UK, 2017a.

613 Pesticide Properties Database, Dimethoate. University of Hertfordshire, UK, 2017b.

614 PubChem Compound Database, Deltamethrin. National Center for Biotechnology Information, USA,
615 2017.

616 Rehse, S., et al., 2016. Short-term exposure with high concentrations of pristine microplastic
617 particles leads to immobilisation of *Daphnia magna*. Chemosphere. 153, 91-9.

618 Ren, Z., et al., 2009. Behavioral responses of *Daphnia magna* to stresses of chemicals with different
619 toxic characteristics. Bulletin of Environmental Contamination and Toxicology. 82, 310-6.
620 Rist, S., et al., 2017. Ingestion of micro- and nanoplastics in *Daphnia magna* – Quantification of body
621 burdens and assessment of feeding rates and reproduction. Environmental Pollution. 228,
622 398-407.
623 Roast, S. D., et al., 1999. Toxicity of the organophosphate pesticides chlorpyrifos and dimethoate to
624 *Neomysis integer* (Crustacea: Mysidacea). Water Research. 33, 319-326.
625 Rochman, C. M., et al., 2013a. Long-term field measurement of sorption of organic contaminants to
626 five types of plastic pellets: implications for plastic marine debris. Environmental Science and
627 Technology. 47, 1646-54.
628 Rochman, C. M., et al., 2013b. Ingested plastic transfers hazardous chemicals to fish and induces
629 hepatic stress. Scientific Reports. 3, 3263.
630 Rochman, C. M., et al., 2014. Early warning signs of endocrine disruption in adult fish from the
631 ingestion of polyethylene with and without sorbed chemical pollutants from the marine
632 environment. Science of the Total Environment. 493, 656-61.
633 Rochman, C. M., et al., 2013c. Polystyrene plastic: a source and sink for polycyclic aromatic
634 hydrocarbons in the marine environment. Environmental Science and Technology. 47,
635 13976-84.
636 Rosenkranz, P., et al., 2009. A comparison of nanoparticle and fine particle uptake by *Daphnia*
637 *magna*. Environmental Toxicology and Chemistry. 28, 2142-2149.
638 Setälä, O., et al., 2014. Ingestion and transfer of microplastics in the planktonic food web.
639 Environmental Pollution. 185, 77-83.
640 Sethi, P. K., et al., 2014. Measurement of plasma protein and lipoprotein binding of pyrethroids.
641 Journal of Pharmacological and Toxicological Methods. 70, 106-111.
642 Sigma-Aldrich, Dimethoate Safety Data Sheet. Sigma-Aldrich Co. LLC, 2017.
643 Sørensen, F. F., et al., 1995. The effects of sublethal dimethoate exposure on the locomotor behavior
644 of the collembolan *Folsomia candida* (Isotomidae). Environmental Toxicology and Chemistry.
645 14, 1587-1590.
646 Syberg, K., et al., 2008. Mixture toxicity of three toxicants with similar and dissimilar modes of action
647 to *Daphnia magna*. Ecotoxicology and Environmental Safety. 69, 428-36.
648 Tanaka, K., et al., 2018. Transfer of Hazardous Chemicals from Ingested Plastics to Higher-Trophic-
649 Level Organisms. The handbook of environmental chemistry. Springer, Berlin, Heidelberg.
650 Teuten, E. L., et al., 2009. Transport and release of chemicals from plastics to the environment and
651 to wildlife. Philosophical Transactions of the Royal Society B: Biological Sciences. 364, 2027-
652 45.
653 Toumi, H., et al., 2013. Effects of deltamethrin (pyrethroid insecticide) on growth, reproduction,
654 embryonic development and sex differentiation in two strains of *Daphnia magna* (Crustacea,
655 Cladocera). Science of the Total Environment. 458-460, 47-53.
656 Van Scoy, A., et al., Environmental Fate and Toxicology of Dimethoate. In: W. P. de Voegt, (Ed.),
657 Reviews of Environmental Contamination and Toxicology Volume 237. Springer International
658 Publishing, Cham, 2016, pp. 53-70.
659 Velzeboer, I., et al., 2014. Strong sorption of PCBs to nanoplastics, microplastics, carbon nanotubes,
660 and fullerenes. Environmental Science & Technology. 48, 4869-76.
661 Wang, F., et al., 2018. Interaction of toxic chemicals with microplastics: a critical review. Water
662 Research.
663 Wang, W., Wang, J., 2018. Comparative evaluation of sorption kinetics and isotherms of pyrene onto
664 microplastics. Chemosphere. 193, 567-573.
665 Weber, A., et al., 2018. PET microplastics do not negatively affect the survival, development,
666 metabolism and feeding activity of the freshwater invertebrate *Gammarus pulex*.
667 Environmental Pollution. 234, 181-189.

668 Wheelock, C. E., et al., 2005. Influence of container adsorption upon observed pyrethroid toxicity to
669 *Ceriodaphnia dubia* and *Hyalella azteca*. *Aquatic Toxicology*. 74, 47-52.
670 Wright, S. L., et al., 2013. Microplastic ingestion decreases energy reserves in marine worms. *Current*
671 *Biology*. 23, 1031-1033.
672 Xiu, R., et al., 1989. Toxicity of the new pyrethroid insecticide, deltamethrin, to *Daphnia magna*.
673 *Hydrobiologia*. 188, 411-413.

674

675 **Figure captions**

676 Fig. 1. Data for dimethoate showing 1a) a comparison of survival probabilities (the deviance
677 in survival probability based on a ratio of survival probability without microplastics and with
678 microplastics) and 1b) a comparison of normal mobility probabilities (calculated as for 1a).
679 Deviations from 0 indicate the extent of the difference when microplastics were present. The
680 closer to 0, the more similar the data. Full survival and mobility probability values for
681 dimethoate are presented in Tables 1a and 1b respectively.

682

683 Fig. 2. Data for deltamethrin showing 2a) a comparison of survival probabilities (the
684 deviance in survival probability based on a ratio of survival probability without microplastics
685 and with microplastics) 2b) a comparison of normal mobility probabilities (calculated as for
686 2a). Deviations from 0 indicate the extent of the difference when microplastics were present.
687 The closer to 0, the more similar the data. Full survival and mobility probability values for
688 deltamethrin are presented in Tables 2a and 2b respectively.

689


690 Fig. 3. Average measured concentrations based on three replicate samples of dimethoate (\pm
691 SD) at different time points taken from treatments with nominal concentrations (a) 0.625 mg
692 l^{-1} and (b) 5 mg l^{-1} , with or without microplastics, at each time point. '- MP' = no
693 microplastics, '+ MP' = with microplastics.

694

695 Fig. 4. Average measured concentrations based on three replicate samples of deltamethrin (\pm
696 SD) at different time points taken from treatments with nominal concentrations (a) 0.4 $\mu\text{g l}^{-1}$
697 and (b) 10 $\mu\text{g l}^{-1}$ b, with or without microplastics, at each time point. '- MP' = no
698 microplastics, '+ MP' = with microplastics.

Figure 1

1a



1b





Figure 2

2a



2b




Figure 3






Figure 4



Acute toxicity of organic pesticides to *Daphnia magna* is unchanged by co-exposure to polystyrene microplastics

Alice A. Horton ^{ab*}, Martina G. Vijver ^b, Elma Lahive ^a, David J. Spurgeon ^a, Claus Svendsen ^a, Roel Heutink ^b, Peter M. van Bodegom ^b, Jan Baas ^{ab}

^a Centre for Ecology and Hydrology, Maclean Building, Benson Lane, Wallingford, Oxfordshire, OX10 8BB. UK.

^b Institute of Environmental Sciences, University of Leiden, P.O. Box 9518, 2300 RA Leiden, The Netherlands.

SUPPLEMENTARY INFORMATION

S1. Area and mass calculations

S1.1. Surface area calculations

Particles were calculated using TEM as being $1.2 \mu\text{m} \pm 0.2 \mu\text{m}$ in diameter (fig S3). Surface area was therefore calculated for particles of $1 \mu\text{m}$ and $1.4 \mu\text{m}$ to account for variation, using the equation:

$$A = 4\pi r^2 \text{ (equation 1)}$$

Calculated surface area ranged from $3.14 \mu\text{m}^2$ for a $1 \mu\text{m}$ particle and $6.15 \mu\text{m}^2$ for a $1.4 \mu\text{m}$ particle (median $1.2 \mu\text{m} \pm 0.2 \mu\text{m}$). Given a concentration of $300\,000 \text{ particles ml}^{-1}$, the number in 40 ml solution was approximately $12\,000\,000$. This therefore gave a total particle surface area per vessel of between 37.7 cm^2 and 73.9 cm^2 .

The surface area of the inside of the vessel was calculated to be approximately 62.8 cm^2 based on a depth of 3.8 cm and a diameter of 4.2 cm when filled with 40 ml water.




Fig. S1. TEM image of polystyrene particles used in the exposures.

S1.2. Particle mass calculations

Particle mass was calculated by taking the known particle density: 1.06 g cm⁻³, and the mean particle radius: 0.6 μm (0.00006 cm). The volume of an individual sphere was calculated using the equation:

$$V = \frac{4}{3} \pi r^3 \text{ (equation 2)}$$

This gave a particle volume of 9.05 x 10⁻¹³ cm³. Volume was then multiplied by density to give the mass of one particle: 9.59 x 10⁻¹³ g (9.59 x 10⁻⁷ μg). This could then be multiplied by 300 000 to give the mass of particles per ml: 2.88 x 10⁻⁷ g ml⁻¹ (0.29 μg ml⁻¹) and then by 1000 to give the mass of particles per l: 0.00029 g l⁻¹ (287.7 μg l⁻¹).

S2. Chemical analysis methods

For the dimethoate treatments, 1 ml samples were taken from three replicate vessels of two different nominal concentrations (5 mg l⁻¹ and 0.625 mg l⁻¹) at 0 and 72 hours. Following removal, the microplastic samples were immediately spun in 1ml glass tubes (2 tubes per sample) in a centrifuge at approx. 6000 G (8000 rpm) for 5 minutes (Eppendorf 24-place Fixed-angle rotor, FA-45-24-11-HS) From the centrifuged microplastic samples, 800 μl was carefully pipetted into a glass vial to avoid resuspending the particles and 400 μl methanol added. The non-microplastic samples were not centrifuged and 500 μl methanol was added to the 1 ml sample. Vials were tightly sealed with a cap (phenolic cap with aluminium liner) and were then shaken well to mix.

For the deltamethrin treatments, 2 ml samples were taken from three replicate vessels of two different nominal concentrations (10 μg l⁻¹ and 0.04 μg l⁻¹) at 0, 24 and 72 hours (based on times of daphnia exposure) and the microplastic treatments centrifuged as before. The 1.6 ml (800 μl per tube) supernatant carefully pipetted off to avoid resuspending the particles. This

was transferred to a glass vial and 1.6 ml hexane added. The non-microplastic samples were not centrifuged and 2 ml hexane was added to the 2 ml sample. The microplastic and non-microplastics samples were then treated the same by shaking the sample with the hexane vigorously for 1 minute in a glass vial tightly sealed with aluminium foil and parafilm and then pipetting 1.2 ml of the hexane fraction into a 2ml brown glass vial (Sigma Aldrich). Vials were tightly sealed with a cap (phenolic cap with aluminium liner, Sigma Aldrich).

All chemical samples were analysed at Wageningen Environmental Research (Alterra). The analytical method was developed at the laboratory of the Environmental Risk Assessment team.

Dimethoate samples were diluted 100 times with acetonitrile-ultrapure water by using a Dilutor Hamilton 600 series. The diluted samples were analysed using an Agilent LC-MS×MS suite (Agilent 6460 Triple Quad LC/MS) equipped with autosampler (Agilent G1329B), pump (Agilent G1311B (Quat. pump)), an ESI (+Agilent Jet Stream) source and a column thermostat (Agilent G1316A). The separation was performed in reverse phase LC (Column: Agilent Zorbax Eclipse XDB C18; 4.6 mm x 150 mm, 5 micron) under gradient elution of Eluents C (Milli-Q water (Advantage A10) + 0.1 % v/v formic acid) and Eluent D (Acetonitrile + 0.1 % formic acid). The initial composition of the mobile phase (40%:60%, C:D) was first held for 2 mins, then changed in 1 min to 20%:80% (C:D) (between 2 and 3 minutes run time), held for 3 minutes (between 3 and 6 minutes run time), changed back to the initial composition over 1 minute (between 6 and 7 minutes) and held there 1 more minute (between 7 and 8 minutes). The flow rate and column temperature were fixed at 0.7 mL.min⁻¹ and 35°C, respectively. Dimethoate retention time was ca. 2.5 minutes and was detected by monitoring the 230 m/z – 198.9 m/z transition (quantifier), qualified with additional peaks at m/z = 171 and 125. Injected samples were quantified by peak area using the calibration curve constructed from calibration standards included in the same sample sequence.

Deltamethrin was measured in the hexane extract by using an Agilent 6890 gas chromatograph equipped with an electron capture detector (ECD). Three microliters of the extract was injected via split injection and analysed in a wall-coated open tubular (WCOT) fused silica column (Varian CP Sil5) using He gas as the mobile phase. The oven temperature was programmed so that the initial temperature of 50°C was held for 7 minutes after which, the temperature was ramped at a rate of 50°C min⁻¹ to a final temperature of 300°C minutes and held for 15:30 minutes. Retention time for deltamethrin was approximately 25.3 minutes. Injected samples were quantified by peak area using the calibration curve constructed from calibration standards included in the same sample sequence.

Table S1. Nominal and average measured concentrations (three replicate samples) for dimethoate treatments

Nominal concentration (mg l ⁻¹)	Microplastic treatment	Time point	Average measured concentration (mg l ⁻¹)	Standard deviation
0.625	NO	0	0.383	0.011
0.625	NO	72	0.378	0.007
0.625	YES	0	0.376	0.002
0.625	YES	72	0.369	0.014
5	NO	0	3.112	0.021
5	NO	72	3.149	0.027
5	YES	0	3.134	0.049
5	YES	72	3.067	0.051

Table S2. Nominal and average measured concentrations (three replicate samples) for deltamethrin treatments

Nominal concentration ($\mu\text{g l}^{-1}$)	Microplastic treatment	Time point	Average measured concentration ($\mu\text{g l}^{-1}$)	Standard deviation
0.4	NO	0	0.082	0.054
0.4	NO	24	0.076	0.044
0.4	NO	72	0.050	0.015
0.4	YES	0	0.050	0.006
0.4	YES	24	0.029	0.004
0.4	YES	72	0.072	0.016
10	NO	0	1.657	0.234
10	NO	24	1.077	0.161
10	NO	72	0.544	0.089
10	YES	0	0.892	0.322
10	YES	24	0.475	0.035
10	YES	72	0.375	0.021

S3. DEB modelling methods

S3.1. Modelling approach

The Stochastic Death model was used to model the data. This model is extensively described in the original paper by Kooijman and Bedaux (1996) and is accepted by the OECD (OECD, 2006). In addition see Jager et al. (2011) for an extensive review on the different survival models.

The model needs three parameters to describe the whole time course of toxic effects:

- 1) No Effect Concentration (NEC): a toxicological threshold for effects
- 2) Killing rate (k_r): a measure for the toxicity of the compound
- 3) Elimination rate (k_e): a kinetic parameter determining the kinetics of the compound

There is an additional parameter (the blank killing rate (BKR)) to take control mortality into account. The NEC is the most important parameter as this reflects the inherent sensitivity of the species for a toxicant. Usually this parameter is also the parameter value with the smallest confidence interval.

Parameter values can be estimated from the raw data of a survival experiment (e.g. Hesketh et al. (2016)), given multiple points in time, as the approach is basically a TK-TD approach. The model can also be used, if the parameter values are known, to back-estimate the exposure concentrations if the survival probabilities are taken from the experiments.

S3.1.1. Dimethoate

Actual concentrations were measured for two nominal concentrations (5 and 0.625 mg/L nominal) at the start of the exposure and at the end of the exposure (24 hrs and 96 hrs after preparing the exposure solutions). Concentrations were stable over the measurement period


and there is a constant fraction of the nominal concentrations for the two measured concentrations (0.625 and 5 mg l⁻¹), this fraction equals 61% of the nominal concentrations both for treatments with and without microplastics. The exposure concentrations calculated based on measured values therefore gave a range of 0, 0.08, 0.15, 0.3, 0.6, 1.2, 2.4 mg l⁻¹. There appears to be no effect of the microplastics on the actual concentrations. This was the starting point for the parameter estimates. The results of the parameter estimates are summarised in Table S3 (all expressed in μ moles).

Table S3. Estimated parameter values for dimethoate with and without microplastics. Where present, numbers in brackets represent 95% confidence intervals.

Experiment	BKR (hr⁻¹)	NEC (mg l⁻¹)	NEC (μM)	k_r (mg l⁻¹ hr⁻¹)	k_r (μM hr⁻¹)	k_e (hr⁻¹)
Dimethoate without microplastics	1.7E-04	0.147 (0.101)	0.64 (0.44)	0.0053 (0.0039)	0.023 (0.017)	0.011 (0.009)
Dimethoate with microplastics	2.7E-03	0.105 (0.039)	0.46 (0.17)	0.023*	0.1*	0.004 (0.001)

* fixed in model

a)



b)




Figure S2. Model fit to dimethoate survival data (+ symbols). Each line represents a different concentration, although for visual clarity, some concentrations have been removed. Fig. S2a shows the model fit to the data without microplastics, fig. S2b shows the model fit to the data with microplastics

The estimated parameter values are identical with and without microplastics (as could be expected as there are no differences in the survival matrices (see the results section of the main text). In addition, the value found for the No Effect Concentration in this research is in perfect agreement with an earlier estimate of 0.63 μM (Baas et al., 2016). LC_x values were calculated (table S6) and compared to literature values (section S3.1.).

S3.1.2. Deltamethrin

As there was a large discrepancy between nominal and actual exposure concentrations for deltamethrin, the nominal chemical exposure concentrations cannot be used to inform the parameters of the model and obtain a reliable estimate of deltamethrin toxicity. We therefore needed to carry out reverse modelling based on known toxicity data, to allow us to estimate actual exposure concentrations and toxicity within our experiment. An independent estimate of the parameter values can be carried out if we have at least three LC_{50} values at different points in time that can be taken from the available literature. In the US-EPA ECOTOX database (US EPA, 2017) we can find 24, 48 and 96 hr LC_{50} values for *Daphnia magna* exposed to deltamethrin (most of the reported data contain only one point in time and are therefore of no use for a TK-TD approach). There is a significant range in the 48 hr LC_{50} values in different publications (Toumi et al., 2013; Xiu et al., 1989), but the numbers presented here (Table S4) are in line with the general picture that emerges from the database. With these values a NEC, killing rate and elimination rate could be derived (Table S5). From these parameters, a model was fit using survival over time (including 96 h, beyond the scope of the test) and thus extrapolating to a realistic exposure concentration range (table 1). LC_x values were calculated (table S7) and compared to literature values as validation of the concentration measurements (section S3.2.).

Table S4. Toxicity data for daphnia exposed to deltamethrin over a 96-hour time period (Xiu et al., 1989)


hr	LC ₅₀ (ug l ⁻¹)
24	0.13
48	0.038
96	0.01

Table S5. Estimated parameter values for deltamethrin.

Experiment	BKR (hr ⁻¹)	NEC (ug l ⁻¹)	NEC (nM)	k _r (ug l ⁻¹ hr ⁻¹)	k _r (nM hr ⁻¹)	k _e (hr ⁻¹)
Deltamethrin	1.7E-04	0.004	0.008	0.56	1.1	0.32

For the purposes of comparison to, and extrapolation from, other studies, for deltamethrin we can only focus on the data without microplastics. As the survival data shows no significant difference whether microplastics are present or not it is therefore reasonable to assume these are the same and therefore only one set of parameter values are presented (Table S5).

a)



b)




Fig. S3. Model fit to deltamethrin survival data (+ symbols). Each line represents a different concentration although for visual clarity, some concentrations have been removed. Fig. S3a shows the model fit to the data without microplastics, fig. S3b shows the model fit to the data with microplastics.

S4. Model-based LC₅₀ values

S4.1. Dimethoate

The 48 h LC₅₀ for dimethoate based on measured values was 1.22 mg l⁻¹ which very closely resembles the 48 h LC₅₀ value of 1.1 mg l⁻¹ reported by Andersen et al. (2006). Beusen and Neven (1989) reported LC₅₀ values of 1.7 and 2 mg l⁻¹ for open and closed experimental systems respectively, values which are also very similar to our 48 h LC₅₀. Although all reported literature values are based on nominal concentrations, the limited difference between nominal and actual concentrations means these can be accurately compared.

Table S6. Modelled LC_x values for dimethoate at different time points based on calculated exposure concentrations.

LC _x (mg l ⁻¹)	Time (hr)			
	24	48	72	96
1	0.8	0.41	0.3	0.25
5	1.05	0.5	0.34	0.28
10	1.31	0.57	0.39	0.3
50	3.48	1.22	0.71	0.5
90	9.08	2.77	1.47	0.99

S4.2. Deltamethrin

The 48 h LC₅₀ value of 0.046 µg l⁻¹ as calculated by the model is comparable to the 48 h LC₅₀ value of 0.12 µg l⁻¹ reported on the deltamethrin safety data sheet (Sigma-Aldrich, 2017). The result is also within a similar range to that reported by Toumi et al. (2013) who calculated 48 h LC₅₀ values of 0.32 µg l⁻¹ and 0.63 µg l⁻¹ based on measured concentrations, with variation dependent on the strain of *D. magna*. The modelled value for 96 h LC₅₀ is 0.023 µg l⁻¹, which is in the same order of magnitude as the literature value of 0.01 µg l⁻¹ calculated by Xiu et al. (1989). However these values should be treated with caution as these

concentrations are approaching/exceeding the solubility limit of deltamethrin, and are often based on nominal concentrations.

Table S7. Modelled LC_x values for deltamethrin at different time points based on calculated exposure concentrations.

LC _x (µg l ⁻¹)	Time (hr)			
	24	48	72	96
1	0.024	0.015	0.012	0.011
5	0.032	0.018	0.014	0.012
10	0.040	0.021	0.016	0.013
50	0.118	0.046	0.029	0.023
90	0.321	0.109	0.064	0.046

Although 48 and 96 hour LC_{50s} for deltamethrin can be broadly compared to those of other studies, there is huge variability within the literature which suggests that determining LC_{50s} for deltamethrin is complicated, as solubility and LC₅₀ can both be influenced by factors such as temperature, pH and vessel material.

References

- Andersen, T. H., et al., 2006. Acute and chronic effects of pulse exposure of *Daphnia magna* to dimethoate and pirimicarb. *Environmental Toxicology and Chemistry*. 25, 1187-1195.
- Baas, J., et al., 2016. Comparison and evaluation of pesticide monitoring programs using a process-based mixture model. *Environmental Toxicology and Chemistry*. 35, 3113-3123.
- Beusen, J.-M., Neven, B., 1989. Toxicity of Dimethoate to *Daphnia magna* and freshwater fish. *Bulletin of Environmental Contamination and Toxicology*. 42, 126-133.
- Hesketh, H., et al., 2016. Extending standard testing period in honeybees to predict lifespan impacts of pesticides and heavy metals using dynamic energy budget modelling. *Scientific Reports*. 6, 37655.
- Jager, T., et al., 2011. General Unified Threshold Model of Survival - a Toxicokinetic-Toxicodynamic Framework for Ecotoxicology. *Environmental Science & Technology*. 45, 2529-2540.
- Kooijman, S., Bedaux, J., 1996. Analysis of toxicity tests on *Daphnia* survival and reproduction. *Water Research*. 30, 1711-1723.
- OECD, 2006. Current Approaches in the Statistical Analysis of Ecotoxicity Data: A Guidance to Application (ENV/JM/MONO(2006)18). OECD Publishing, Paris.
- Sigma-Aldrich, Dimethoate Safety Data Sheet. Sigma-Aldrich Co. LLC, 2017.
- Toumi, H., et al., 2013. Effects of deltamethrin (pyrethroid insecticide) on growth, reproduction, embryonic development and sex differentiation in two strains of *Daphnia magna* (Crustacea, Cladocera). *Science of the Total Environment*. 458-460, 47-53.
- US EPA, ECOTOXicology Knowledgebase System. Version 4.0. 2017.
- Xiu, R., et al., 1989. Toxicity of the new pyrethroid insecticide, deltamethrin, to *Daphnia magna*. *Hydrobiologia*. 188, 411-413.