

Multiple timescale constraints for high-flux magma chamber assembly prior to the Late Bronze Age eruption of Santorini (Greece)

T. Flaherty¹, T.H. Druitt¹, H. Tuffen², M.D. Higgins³, F. Costa⁴, A. Cadoux⁵

¹ Laboratoire Magmas et Volcans, Université Clermont Auvergne-CNRS-IRD, OPGC, F-63100, Clermont-Ferrand, France

² Lancaster Environment Centre, Lancaster University, Lancaster, UK

³ Sciences de la Terre, Université du Québec à Chicoutimi, 555 Boulevard de l'Université, Chicoutimi, Québec, G7H
2B1, Canada

⁴ Earth Observatory of Singapore, Nanyang Technological University, Singapore 639798, Singapore

⁵ GEOPS, Univ. Paris-Sud, CNRS, Université Paris-Saclay, 91405 Orsay, France

Corresponding author: T.H. Druitt (tim.druitt@uca.fr)

ABSTRACT

The rhyodacitic magma discharged during the 30-80 km³ DRE (dense rock equivalent) Late Bronze Age (LBA; also called 'Minoan') eruption of Santorini caldera is known from previous studies to have had a complex history of polybaric ascent and storage prior to eruption. We refine the timescales of these processes by modelling Mg-Fe diffusion profiles in orthopyroxene and clinopyroxene crystals. The data are integrated with previously published information on the LBA eruption (phase equilibria studies, melt inclusion volatile barometry, Mg-in-plagioclase diffusion chronometry), as well as new plagioclase crystal size distributions and the established pre-LBA history of the volcano, to reconstruct the events that led up to the assembly and discharge of the LBA magma chamber. Orthopyroxene, clinopyroxene and plagioclase crystals in the rhyodacite have compositionally distinct rims, overgrowing relict, probably source-derived, more magnesian (or calcic) cores, and record one or more crystallization (plag>>opx>cpx) events during the few centuries to years prior to eruption. The crystallization event(s) can be explained by the rapid transfer of rhyodacitic melt from a dioritic/gabbroic region of the subcaldera pluton (mostly in the 8-12 km depth range), followed by injection, cooling and mixing in a large contiguous melt lens at 4-6 km depth (the pre-eruptive magma chamber). Since crystals from all eruptive phases yield similar timescales, the melt transfer event(s), the last of which took place less than two years before the eruption, must have involved most of the magma subsequently erupted. The data are consistent with a model in which prolonged generation, storage and segregation of silicic melts was followed by gravitational instability in the subcaldera pluton, causing the rapid interconnection and amalgamation of melt-rich domains. The melts then drained to the top of the pluton, at fluxes of up to 0.1-1 km³.y⁻¹, where steep vertical gradients of density and rheology probably caused them to inject laterally, forming a short-lived holding chamber prior to eruption. This interpretation is consistent with growing evidence that some large silicic magma chambers are transient features on geological timescales. A similar process preceded at least one earlier caldera-forming eruption on Santorini, suggesting that it may be a general feature of this rift-hosted magmatic system.

Keywords: diffusion chronometry, crystal size distribution, Santorini, Minoan eruption, magma chamber, magma ascent

INTRODUCTION

Caldera-forming eruptions of silicic magma, such as the magnitude 7, Late Bronze Age (LBA; also called 'Minoan') eruption of Santorini, are low-frequency, high-impact events with effects at global scales (Mason et al. 2004; Self 2006; Oppenheimer 2013). The processes leading to reactivation of long-dormant caldera volcanoes are of societal interest, in particular for interpreting geophysical signals in times of unrest. Recent studies have shown that such systems can build up to eruption on geologically short timescales, raising the question of how this occurs (Morgan et al. 2006; Wark et al. 2007; Wilson and Charlier 2009; Gualda et al. 2012; Druitt et al. 2012; Matthews et al. 2012; Allan et al. 2013; 2017; Fabbro et al. 2013; 2017; Chamberlain et al. 2014; Pamukcu et al. 2015; Barker et al. 2016).

Silicic magmas are believed to be generated in bodies of crystal mush, where they can be stored in the crust for long periods of time (Bacon and Druitt 1988; Brophy 1991; Hildreth and Wilson 2007; Bachmann and Bergantz 2004; 2008; Deering et al. 2010; Bégué et al. 2014; Bergantz et al. 2015; Bachmann and Huber 2016; Cooper 2017; Rubin et al. 2017). Whereas large volumes of melt lose heat rapidly by convection, locked-in mushes cool slowly by conduction (Koyaguchi and Kaneko 1999), providing a long-lived source of melt that can be periodically tapped. Creation of large chambers of melt-dominated silicic magma requires (1) production and incubation of the melt within mush bodies, (2) extraction of that melt, and (3) its accumulation in magma chambers (Annen et al. 2006; Brown 2013, and references therein). Low rates of melt accumulation lead to pluton construction through freezing of successive batches, whereas the runaway addition of mass and heat associated with high accumulation rates generates magma chambers (Annen 2009). The critical accumulation rate for magma chamber formation probably lies between 10^{-3} and $10^{-1} \text{ km}^3 \cdot \text{y}^{-1}$ for a Santorini-sized system (Schopa and Annen 2013; Gelman et al. 2013; Karakas et al. 2017). The flow rate necessary to transport the melt through a dyke up to the magma chamber without freezing provides an additional constraint ($\sim 10^{-2} \text{ km}^3 \cdot \text{y}^{-1}$; Menand et al. 2015). Theoretical models of silicic melt extraction from crystal mushes by intergranular porous flow can explain magma chamber formation on timescales of 10^4 - 10^6 y (Bachmann and Bergantz 2004; Solano et al. 2012). However, silicic magma chambers in some cases are assembled much faster than predicted theoretically, at rates as high as 0.05 - $1 \text{ km}^3 \cdot \text{y}^{-1}$ and on timescales as low as 10 - 10^3 y (Druitt et al. 2012; Allan et al. 2013; 2017; Barker et al. 2016; Cooper et al. 2017; Fabbro et al. 2017). A possible explanation is that melt-rich domains present in the mush body interconnect and amalgamate very rapidly to form a magma chamber (Christopher et al. 2015; Cashman et al. 2017; Sparks and Cashman 2017). Some questions arise from these concepts. How common is rapid magma chamber assembly in silicic systems? What is the mechanism driving high-flux melt transfer? How does the timescale of magma chamber development scale with the size of the system? Following chamber assembly, what finally triggers the eruption?

The LBA eruption of Santorini has played a role in the development of these ideas. Cottrell et al. (1999) proposed, based on a combination of phase equilibria experiments and melt inclusion H_2O barometry, that

the LBA silicic magma underwent polybaric ascent from a deep storage zone to a shallow magma chamber about 300 years prior to eruption. Their model was extended by Druitt et al. (2012), who used Mg-in plagioclase diffusion modelling to reduce the timescale from 300 years to a few decades. While Cottrell et al. (1999) proposed that the entire erupted magma volume was transferred in this time, Druitt et al. (2012) envisaged the late-stage replenishment of a pre-existing chamber by several km³ of silicic magma. In both interpretations, however, magma chamber assembly took place on timescales that were geologically very short. Other data have subsequently been published on the LBA magma: micro-Sr isotopic analyses of plagioclase (Martin et al. 2010), phase equilibria experiments on Santorini silicic magmas (Cadoux et al. 2014) and the basaltic/andesitic magmas parental to them (Andújar et al. 2015; 2016), H₂O+CO₂ melt inclusion barometry (Druitt et al. 2016), and the chemistry and petrology of minor juvenile and cognate components in the LBA eruption products (Druitt 2014).

In the present paper we present new data on the compositions and zoning patterns of orthopyroxene and clinopyroxene crystals and, in particular, on timescales of pre-eruptive crystal residence estimated by modelling of Mg-Fe diffusion gradients in these minerals (Morgan et al. 2004; Saunders et al. 2012; Allan et al. 2013; Costa et al. 2013; Chamberlain et al. 2014; Kilgour et al. 2014; Petrone et al. 2016; Singer et al. 2016; Krimer and Costa 2017; Fabbro et al. 2017). We also derive updated timescale information from plagioclase crystals. We then integrate the new, and previously published, information into a model for the architecture and dynamics of the Santorini plumbing system in the period leading up to the LBA eruption, and place constraints on the physical mechanisms at play.

THE LATE BRONZE AGE ERUPTION: CONTEXT AND PRODUCTS

Santorini is the southernmost volcanic centre of the South Aegean Volcanic Arc in Greece. It lies in a 100-km-long, 45-km-wide SW-NE rift zone (Nomikou et al. 2016; Hooft et al. 2017; Fig. 1a) on 25-km-thick continental crust with seismically distinct upper (0-15 km) and lower (15-25 km) layers (Konstaninou 2010).

The earliest preserved volcanic products on Santorini date from 650 ka. Since 350 ka the volcano has produced twelve Plinian eruptions (Druitt et al. 1999), at least four of which caused edifice collapse resulting in the present-day 10 x 7 km, multistage caldera. The Plinian eruptions were separated by inter-Plinian periods of 10-30 ky durations and characterised by intracaldera effusive volcanism and lower intensity (sub-Plinian or less) explosive activity. Following over half a million years of activity, Santorini must be underlain by a large pluton, probably zoned from granodiorite through diorite to gabbro with increasing depth. Mantle-derived basalt fractionates to andesite beneath Santorini near the upper-lower crustal boundary (~400 MPa), and andesite fractionates to silicic magma in the upper crust (200-400 MPa) (Andújar et al. 2015; 2016). Santorini magmas are calcalkaline to mildly tholeiitic in nature (Huijsmans 1985; Huijsmans and Barton 1989; Druitt et al. 1999).

The Plinian eruption prior to the LBA eruption (the $>10 \text{ km}^3$ DRE Cape Riva eruption) has been dated at $21.8 \pm 0.4 \text{ ka}$ and formed a caldera located in the northern basin of the present-day caldera (Athanasiss et al. 2016, and references therein; Fig. 1c). A sequence of events then occurred within the crust during the subsequent 17.2 ky inter-Plinian period that culminated in the LBA eruption (Fig. 1b). There is evidence for three eruptive events during this period. (1) Immediately following the Cape Riva eruption, an andesitic edifice at least 2.5 km^3 in volume was built within the Cape Riva caldera. Blocks of this former edifice occur as black glassy lithic blocks in the LBA ejecta (Druitt 2014); one block has been dated at $20.2 \pm 1.0 \text{ ka}$ (Karátson et al. 2018). The andesite was a distinctive high-Ba magma unrelated to either the Cape Riva or LBA silicic magmas (Druitt 2014). (2) At 12-13 ka, a small eruption of LBA-like silicic magma took place, forming a crypto-ash layer in marine and terrestrial sediments (St. Seymour et al. 2004; Satow et al. 2015). (3) A number of small basaltic eruptions of unknown age also took place (Vespa et al. 2006; Vaggelli et al. 2009).

The LBA eruption occurred in the late 17th century BCE, and discharged 30-80 km^3 DRE of magmatic and accidental materials (Pyle 1990; Sigurdsson et al. 2006; Johnston et al. 2014). The eruption has been reconstructed in detail (Bond and Sparks 1976, Heiken and McCoy 1984; Sigurdsson et al. 1990; Pyle 1990; Sparks and Wilson 1990; Cioni et al. 2000; Druitt 2014). It comprised five phases, here numbered 0 to 4: (0) a precursory Vulcanian phase, (1) a Plinian phase, (2) a phase of syn-Plinian base surges, (3) a phase of phreatomagmatic activity with low-temperature pyroclastic flows, and (4) a phase of hot, fluidized pyroclastic flows, which accounted for most of the magma discharged (Fig. 1d). Phases 0 and 1 took place from a subaerial vent (star, Fig. 1c). The vent then migrated during phase 2 into the flooded Cape Riva caldera, such that phase 3 was strongly phreatomagmatic. Phase 4 appears to have been fed from multiple subaerial vents of poorly constrained locations. Caldera collapse probably took place during phase 4, deepening and enlarging the extant Cape Riva caldera.

Since the LBA eruption, repeated effusion of dacitic magma ($4.85 \pm 0.7 \text{ km}^3$ in total) has built up the largely submerged Kameni intracaldera edifice on the floor of the present-day caldera, with its subaerial summits forming the islands of Palaea and Nea Kameni (Pyle and Elliot 2006; Nomikou et al. 2014). The first recorded eruption of Kameni took place in 197 BCE, and the most recent in 1950. The mean eruption rate over this time has been $1.3 \times 10^{-3} \text{ km}^3 \cdot \text{y}^{-1}$ (Nomikou et al. 2014), similar to that estimated for a previous inter-Plinian period between 70 and 55 ka ($\sim 1 \times 10^{-3} \text{ km}^3 \cdot \text{y}^{-1}$; Druitt et al. 1999). The Kameni vents lie along a NE-SW lineament called the Kameni Line, which lies parallel to the regional fault trend and which has played a central role in Santorini volcanism (Fig. 1c). Several Plinian eruptions, including the LBA, began at a vent on or near the Kameni Line. Seismicity was strongly concentrated along the Kameni Line during a period of bradyseismic unrest in 2011-2012 (Newman et al. 2012).

SUMMARY OF PREVIOUSLY PUBLISHED DATA ON THE LBA RHYODACITE

Petrology and geochemistry

Analyses of twenty-three visually homogeneous pumice clasts of the rhyodacite presented by Druitt (2014) reveal a restricted range of whole-rock compositions (70-71 wt% SiO₂, recalculated volatile-free; FeO/MgO of 3 to 5), irrespective of eruptive phase. The crystalline phases are plagioclase, orthopyroxene, clinopyroxene (weight proportions plag : opx : cpx of 0.84 : 0.10 : 0.06; Cadoux et al. 2014), magnetite and ilmenite set in 73-74 wt% SiO₂ rhyolitic glass. Apatite and pyrrhotite occur mostly as inclusions in pyroxenes and magnetite. Trace quantities of zircon are present (Schmitt et al. 2016). Crystal contents based on K mass balance in nine pumice-glass pairs range from <5 to 20 wt%, the lowest values occurring in phase 1 (<5-12 wt%), and the highest in phase 4 (12-20 wt%) (Online Resource 1). New measurements of plagioclase contents in fifteen thin sections by image analysis (3-16 vol%; Online Resource 2) are consistent with this range.

Plagioclases (≤3 mm in length) were described by Druitt et al. (2012) based on BSE images of 300 crystals (Fig. 2a). Most have compositions of An_{40±3} (with rare thin zones as sodic as An₃₀) and are weakly normally zoned to euhedral edges of An₃₇₋₄₀. Superimposed on the overall normal zoning are multiple (typically 3-6), 50-100 μm wide, sawtooth-zoned packets, each consisting of normally zoned plagioclase terminated by a subtle resorption surface and a small (<few molar %) increase in An content. About half of the crystals have An₃₀₋₈₈ composition cores that are variably euhedral to corroded (Fig. 2a), and exhibit complex combinations of normal, reverse, sawtooth, oscillatory or patchy zoning, with multiple resorption surfaces, indicative of long and complex histories prior to overgrowth by the An_{40±3} rims. The majority of the cores have compositions of An_{50±7}, sometimes with inner cores of An₅₈₋₈₈ (Fig. 2b). Very rarely (<<1%) the cores are more sodic (An₃₀₋₃₆) than the rims. An additional group of plagioclase crystals occurs in small quantities, and uniquely in pumices from eruptive phase 4. Their distribution is very heterogeneous; they are present in some phase-4 pumices but not in others. They have the same compositions as the calcic cores (An_{50±7}), but lack the An_{40±3} rims (Fig. 2a). Druitt et al. (2012) referred to them as '*rimless An_{50±7} crystals*'. These rimless crystals have thin selvages of a glass that is slightly less evolved than the normal interstitial glass (Online Resource 1).

Martin et al. (2010) measured the ⁸⁷Sr/⁸⁶Sr ratios of plagioclase crystals in phase-1 pumices to be 0.7049-0.7051, slightly less Sr-radiogenic than the interstitial glass (0.7051-7054). Seven of their eight isotopic analyses were from the An_{40±3} plagioclase, and one was from a An_{50±7} core; there was no difference in ⁸⁷Sr/⁸⁶Sr between the two compositions. Di Salvo et al. (2013) also isotopically analysed some rims, calcic cores and highly calcic inner cores, and found that they all had similar range of ⁸⁷Sr/⁸⁶Sr. Martin et al. (2010) interpreted the relatively high ⁸⁷Sr/⁸⁶Sr of the interstitial glass to post-crystallization contamination of the rhyodacitic magma by a more Sr-radiogenic melt.

Melt inclusions (MIs) in $An_{40\pm 3}$ plagioclase contain 69-75 wt% SiO_2 (Online Resource 1). Secondary ion mass spectrometry (SIMS) analysis of >60 isolated, crack-free MIs hosted in plagioclase, opx and cpx revealed volatile concentrations of 4.0-6.9 wt% H_2O , 20-200 ppm CO_2 , 40-100 ppm S, and 3000-4500 ppm Cl (Cadoux et al. 2014; Druitt et al. 2016). MIs within plagioclase $An_{50\pm 7}$ cores are more H_2O -rich (4.2-6.9 wt%) than those within $An_{40\pm 3}$ rims (4.0-5.0 wt%) (Fig. 2c). Cottrell et al. (1999) also measured ≤ 6.5 wt% H_2O in seven MIs within $An_{50\pm 7}$ cores. If corrected for ~ 10 wt% post-entrapment crystallization (in the cooler environment of rim overgrowth; Online Resource 1), then the highest H_2O content of the core-hosted MIs *at the moment of entrapment* fall to 6.1 wt%. CO_2 , S and Cl concentrations of MIs in the cores and rims are similar. Although $An_{50\pm 7}$ cores with H_2O -rich MIs occur in pumices from eruptive phase 1, 2 and 4, they are particularly abundant in phase 1 (Fig. 2d).

Druitt et al. (2016) used MI H_2O+CO_2 contents and the solubility models of Newman and Lowenstern (2002) and Papale et al. (2006) to estimate entrapment pressures, which were converted to entrapment depths using a mean upper crust density of $2640 \text{ kg}\cdot\text{m}^{-3}$ (Konstantinou, 2010) assuming volatile saturation (Cadoux et al. 2014; Druitt et al. 2016). MI entrapment in the $An_{40\pm 3}$ plagioclase occurred at 100-160 MPa (4-6 km depth). However, MI entrapment within $An_{50\pm 7}$ plagioclase cores extended to higher inferred pressures (240 MPa for 6.1 wt% H_2O and 100 ppm CO_2 , corresponding to a depth of 9 km). The proposed explanation is that the cores were derived from depth and transferred to shallower levels where rim overgrowth occurred, in accordance with the earlier model of Cottrell et al. (1999).

Pre-eruptive temperatures and oxygen fugacities (fO_2) based on touching Fe-Ti oxide pairs in contact with the glass were estimated to be $885 \pm 7 \text{ }^\circ\text{C}$ and $10^{-6.7 \pm 0.2} \text{ Pa}$ by Cottrell et al. (1999) using the algorithm of Andersen and Lindsley (1988). Cadoux et al. (2014) and Druitt (2014) estimated $853 \pm 7 \text{ }^\circ\text{C}$ and $10^{-7.8 \pm 0.2} \text{ Pa}$ using the same algorithm, and $866 \pm 9 \text{ }^\circ\text{C}$ and $10^{-7.86 \pm 0.07} \text{ Pa}$ using that of Ghiorso and Evans (2008). There is no systematic variation of temperature with eruptive phase.

Pre-eruptive magma chamber

The LBA rhyodacite was erupted from a single interconnected magma reservoir (henceforth termed the *magma chamber*), because similar whole rock and glass compositions, phenocryst edge compositions, phenocryst MI compositions, and Fe-Ti oxide temperatures are found for pumices from all eruptive phases (Cottrell et al. 1999; Cadoux et al. 2014; Druitt 2014). Cottrell et al. (1999) used phase equilibria, MI H_2O contents, and the absence of stable amphibole to infer pre-eruptive storage of the magma at 50 MPa, $885 \text{ }^\circ\text{C}$, and H_2O -saturated conditions. These values were revised by Cadoux et al. (2014), who experimentally reproduced the phenocryst-glass assemblage at higher pressure (≥ 200 MPa) and lower temperature ($860 \text{ }^\circ\text{C}$), with melt saturated with respect to $H_2O + Cl + CO_2$, but slightly undersaturated with respect to pure H_2O ($X_{H_2O} = H_2O/[H_2O+CO_2]$ of 0.8-0.9). Cadoux et al. (2014) also proposed that the melt co-existed with a H_2O+Cl -rich hydrosaline liquid. The MI $H_2O + CO_2$ data of Druitt et al. (2016) yielded pre-eruptive storage pressures (100-160 MPa; 4-6 km) intermediate between the estimates of the previous authors. Combining

the different constraints, the best-fit pre-eruptive conditions for the LBA rhyodacite are 850-880 °C, $fO_2 \sim$ NNO, 100-160 MPa (4-6 km depth), melt saturated with 4.0-5.0 wt% H₂O, <200 ppm CO₂, <150 ppm S and 3000-5000 ppm Cl, a free H₂O-rich vapour phase, and possibly a H₂O+Cl-rich hydrosaline liquid. The magma chamber was less than 2 km thick (4-6 km depth).

Source region of the rhyodacite

Cottrell et al. (1999) proposed a rhyodacite source reservoir at 210-240 MPa and 825 ±25 °C. However, phase equilibrium experiments on a typical 59 wt% SiO₂ Santorini andesite (Andújar et al. 2016) generated 68-70 wt% SiO₂ melt, similar to LBA rhyodacite, at 900-950 °C and under relatively oxidising conditions (NNO+1.5). Andújar et al. noted that, while most LBA pumices are calcalkaline rhyodacite with Fe/Mg ratios of about 3, some are more tholeiitic with Fe/Mg ratios of up to 5. The main, calcalkaline rhyodacite was produced experimentally at 200 MPa and 6 wt% H₂O in an assemblage containing plagioclase (An_{50±5}), cpx (Mg# 70-75) and opx (Mg# 70-75). More tholeiitic rhyodacitic melts were produced at higher pressure (400 MPa). The LBA rhyodacite was thus generated at 900-950 °C over the pressure range 200-400 MPa, but mostly at ~200 MPa (Andújar et al. 2016). The high H₂O contents and entrapment pressures recorded by plagioclase core-hosted MIs (up to 240 MPa, or 9 km) agree with the experimental constraints. The best estimate is that the LBA rhyodacitic melts were extracted mostly from a 900-950 °C, dioritic/gabbroic region of the subcaldera pluton at 8-12 km depth (200-300 MPa), possibly with lesser quantities of more tholeiitic silicic melts extracted from deeper levels (~400 MPa).

Amphibole may have been present at 200 MPa in the source region, but only at temperatures of <940 °C (Fig. 3a, c of Andújar et al. 2016). Any source-derived amphiboles must have been resorbed prior to eruption because amphibole was not stable in the magma chamber (Cottrell et al. 1999; Cadoux et al. 2014). The presence of amphibole in the source region is consistent with the low Y and Dy/Yb of the LBA rhyodacite (Klaver et al. 2016)

Minor andesitic components

Two minor (<1 %) andesitic components were co-erupted with the rhyodacite (Sparks et al. 1977; Druitt et al. 1999; Cottrell et al. 1999; Druitt 2014) (Fig. 1d):

- A suite of microphenocryst-rich andesitic pumices and dioritic plutonic nodules (55-100 vol% crystals) that were erupted at the start of phase 0, and throughout phase 1, but as only trace amounts in phases 2 to 4 (Fig. 1d). These have the same high-Ba composition as the post-Cape Riva intracaldera edifice, and are interpreted as samples of a variably crystalline residual pocket of the same magma that was pushed out by the ascending LBA rhyodacite (Druitt 2014).

- Enclaves of andesitic magma that were mingled into the rhyodacite within a few months of the eruption, quenched, then released during eruptive fragmentation (Cottrell et al. 1999). They were erupted mainly during phase 1. These enclaves are also enriched in Ba (Druitt 2014).

METHODS AND ASSUMPTIONS

Orthopyroxene (opx) and clinopyroxene (cpx) crystals were extracted from the 250-400 and 400-700 μm size fractions of ten gently crushed pumice lumps (2-4 per eruptive phase) and mounted in epoxy resin. Crystals with adhering grey, microlite-rich groundmass derived from disaggregated andesitic enclaves were excluded. Pyroxene zoning textures were studied using greyscale back-scattered electron (BSE) images calibrated for Mg number ($\text{Mg\#} = 100 \times \text{Mg}/[\text{Mg} + \text{Fe}^{\text{Total}}]$ atomic). Only crystals judged by visual inspection to be symmetrically mounted and polished through their centres were imaged. A total of 65-120 opx crystals (eruptive phases 1-4) and 12 cpx crystals (eruptive phases 1, 2 and 4) were examined. All the opx and cpx crystals had their long (c) axes parallel to the surface of the grain mount. Compositional images were made of selected opx crystals. Full details of sample preparation and analysis are given in Online Resource 3.

Diffusive re-equilibration timescales were determined for opx and cpx using binary element diffusion modelling of Mg-Fe profiles across inter-zonal boundaries (Morgan et al. 2004; 2006; Allan et al. 2013; 2017; Chamberlain et al. 2014; Barker et al. 2016) (Online Resource 4). We implemented an analytical solution to the one-dimensional diffusion equation, assuming an initial step function in Mg#, then fitted the measured profiles by a least squares method, with maximum and minimum uncertainties and a correlation index, R^2 (Online Resource 3). Diffusion modelling was carried out perpendicular to crystal length (c axis) in both pyroxene species. In opx it was carried out parallel to the a axis, sections through the c - a plane being recognized by the presence of {201} crystal terminations. Orientations of the cpx crystals were determined using the SHAPE software (Dowty 1987) by matching measured interfacial angles with those of modelled cpx sections. The diffusion coefficients of Ganguly and Tazzoli (1994) and Dohmen et al. (2016) were used for opx, and that of Müller et al. (2013) was used for cpx, with corrections for diffusion anisotropy of opx and cpx explained in Online Resource 3, respectively. Uncertainties on the diffusion timescales were calculated using standard error propagation (Online Resource 3).

The timescale derived by modelling of Mg-Fe diffusion across a particular zone boundary is the time elapsed between zone boundary formation and eruption quench, at the assumed temperature of residence. The modelling was carried out at 855 ± 15 °C and $f\text{O}_2 = 10^{-7.8 \pm 0.2}$ Pa ($\sim\text{NNO}$), representative of the pre-eruptive conditions (Cadoux et al. 2014; Druitt 2014).

Crystal residence timescales obtained by 1D diffusion modelling are maximum estimates (Costa et al. 2008; Shea et al. 2015; Krimer and Costa 2017). However, modelling along short profiles perpendicular to the long (c) axis in prismatic crystals of high aspect ratio (up to 4 in cpx, up to 8 in opx) is pseudo-1D, reducing any overestimation due to this effect. By assuming an initial step function, the approach also

ignores the presence of any growth zoning, also resulting in an overestimation of residence time (Costa et al. 2008; Costa and Morgan 2010; Chamberlain et al. 2014; Till et al. 2015). However, any growth-generated zoning is less important perpendicular to the *c* axis than parallel to it.

NEW INSIGHTS FROM PYROXENES

Pyroxene zoning patterns

In the following descriptions, compositionally distinct inner and outer zones of a crystal are referred to as the *core* and *rim*. The outer limit of the crystal in contact with glass is referred to as the crystal (or rim) *edge*.

Orthopyroxene crystals (≤ 3 mm in length) occur with length/width ratios of up to 8 (Fig. 3). Compositions span the range $Wo_{2-3}En_{52-70}Fs_{28-45}$ (Mg# 53-71), with Al_2O_3 contents of 0.4-0.7 wt% (rarely up to 1.4 wt%). Most crystals have restricted compositions (Mg# 57-61), and diffuse, short-wavelength (5-15 μm) oscillatory zoning, particularly evident parallel to the *c* axis. Some contain sharply defined, 2-30 μm -thick growth zones of slightly more Fe-rich compositions (Mg# 53-55; e.g., bright zone in Fig. 3c), although the euhedral edge composition is always in the range Mg# 57-61. Many crystals that appear homogeneous at low BSE resolutions contain complex, low-magnitude oscillatory zoning visible at high BSE resolutions. Of five opx crystals imaged compositionally, four exhibit sector zoning of Al. (Fig. 4a). Al, being trivalent, diffuses two orders of magnitude slower than Mg and Fe in opx (Nakagawa et al. 2005), and so may preserve the original small-scale growth zoning that is partially re-equilibrated in more rapidly diffusing Mg and Fe. While Al and Mg# are correlated at large scales they are de-correlated at shorter scales, Al exhibiting sharper fluctuations than Mg# (Fig. 4b). This suggests that growth-generated compositional gradients are narrow, justifying a step-function initial condition for Mg-Fe in the diffusion modelling.

One-fifth of the opx crystals have magnesian cores with Mg# 63-71 (Fig. 3e), commonly containing multiple resorption surfaces, melt inclusions, and patchy zoning. The melt inclusions are surrounded by streaky compositional zoning elongated parallel to the *c* axis, formed by diffusional Mg-Fe exchange between the melt inclusions and host opx (e.g., Tomiya and Takahashi 2005; Allan et al. 2013; Barker et al. 2016).

Clinopyroxene crystals (≤ 2 mm in length) have aspect ratios of up to 4 (Fig. 5). Compositions span the range $Wo_{39-44}En_{36-44}Fs_{14-22}$ (Mg# 62-78), but most crystals have Mg# of 62-70, with edges of 66-70 (Fig. 5a). Both normal and reverse zoning are observed in these crystals, but 2-30 μm -scale oscillatory zoning is most common. Al_2O_3 contents are typically 0.7-1.0 wt%, but can reach 2.0-2.5 wt% in more Fe-rich zones. Four of the twelve crystals examined in detail contain magnesian cores with Mg# 72-78 and Al_2O_3 contents of 1.4-1.7 wt% (Figs. 5c and e). Some cores are riddled with melt inclusions, and have a wispy zoning due to partial diffusive Mg-Fe overprinting by the Mg# ~ 67 composition around melt inclusions and along fractures and cleavage planes (Fig. 5c).

Diffusion timescales of pyroxene residence

Diffusion timescales were measured across (1) the inner limit (core-rim boundary) of the crystal rim, and (2) zones boundaries *within* the rims. The aim was to estimate the residence time of the crystal rim of each pyroxene species since its formation. The results are given in full in Online Resource 4, and summarized in Table 1.

A total of 85 zone boundaries in 16 opx crystals were modelled, but application of quality-control criteria (Online Resource 3) retained only 32 of the profiles for timescale calculation, all models fitting the actual profiles with $R^2 > 0.95$. The Mg-Fe gradients ranged in width from 0.4 to 48 μm . Three examples are shown in Fig. 3. Diffusion profiles with $R^2 > 0.99$ give visually good fits (Figs. 3b and d). Those with R^2 of 0.98-0.99 give less good fits, commonly due to the presence of micron-scale fluctuations of Mg# in the crystal (e.g., Fig. 3f). These fluctuations may show that the entire profile was generated by growth, not diffusion. However such fluctuations are also observed in experimental studies of diffusion due to the presence of crystal microfractures not visible on BSE images (J. Ganguly, written communication). Profiles with $R^2 < 0.98$, on the other hand, probably have significant components of growth zoning. We place emphasis on those profiles with $R^2 > 0.99$.

Diffusion timescales across the core-rim boundaries in two opx crystals give 138-459 y using the Dohmen et al. (2016) diffusion coefficient (Fig. 6a). Timescales from zone boundaries within the crystal rims range from <1 to 420 y, largely consistent with the core-rim boundaries, with a single outlier at 852 y (Fig. 6a). However, all except one of the timescales with the highest quality fit ($R^2 > 0.99$) are <55 y (Figs. 3b and d). The timescales from within the rims themselves suggests that rim growth continued until eruption-driven quench took place. The single longer timescale (852 y) comes from near the centre of a core-free crystal (Fig. 6), and may record an earlier phase of rim growth. There are no systematic differences in the timescale ranges in opx crystals from the different eruptive phases.

Orthopyroxene diffusion timescales calculated using the Ganguly and Tazzoli (1994) diffusion coefficient are five times shorter than those using Dohmen et al. (2016). The core-rim boundaries give timescales of 42-110 y, and zone boundaries within the rims give <1 -36 y (with the best fits being <10 y), with the single outlier at 157 y.

We have no firm basis for favouring one opx diffusion coefficient over the other (as discussed in Online Resource 3); however, by placing emphasis on the Dohmen et al. (2016) results, we are inferring *maximum* timescales. Moreover the opx timescales derived using Dohmen et al. (2016) are similar to those obtained from cpx (see below). We therefore conclude that the main phase of opx crystallization began *no more than* 450 years prior to eruption (possibly with an earlier phase at ~ 850 y; Fig. 6a) and was still ongoing within the last few years prior to the eruption.

A total of 26 profiles in 7 cpx crystals were modelled perpendicular to the *c* axis assuming initial step functions in Mg# and distinguishing between profiles across the wispy diffusion fronts defining the core-rim boundary (Figs. 5c and e), and profiles across zone boundaries within the rims (Figs. 5a, c and e; Fig. 6b).

The wispy core-rim boundaries of three cpx crystals give 543-1700 y, but are probably significantly overestimated because the boundaries are not planar and hence not well represented by 1D modelling (Figs. 5c and e). This is unlike the core-rim boundaries in the opx crystals, which are approximately planar (Fig. 3e). Zone boundaries within the cpx rims, all of which are planar, give timescales of 2 to 500 y, with the best-quality fits ($R^2 > 0.99$) giving 2 to 120 years (Fig. 6b). Fuzzier planar boundaries deeper in the rims give 1173-2031 y.

The diffusion timescales for cpx are longer than those of opx. However, if the wispy core-rim boundary timescales are excluded, then the timescales for cpx range from <1 to 500 years prior to eruption (possibly extending to 1000-2000 years; Fig. 6b), which is consistent with the opx timescales using the diffusion coefficient of Dohmen et al. (2016). We accept this interpretation, while recognising that some inner zones of the cpx rims may pre-date the rims on the opx crystals.

Two endmember situations can be envisaged for crystal rim growth. In one, growth occurred very rapidly and the crystal then resided at high temperature until eruption (case $t_{\text{growth}} \ll t_{\text{residence}}$; Fig. 7a). Alternatively, the rims grew continuously from initiation until eruption (case $t_{\text{growth}} = t_{\text{residence}}$; Fig. 7a). The diffusion timescales for both opx and cpx generally increase from small values near crystal edges to much larger values in the interior (Fig. 7b and c). Although this may in part be an artefact due to the higher probability of more oblique intersection of zone boundaries at depth than near the crystal surface, it suggests that the scenario $t_{\text{growth}} = t_{\text{residence}}$ is more realistic for LBA magma. Visual extrapolation of the timescales back to the core-rim boundary (or to the centre of the phenocryst, if core-free) suggests the onset of rim growth as early as >1000-2000 y to as late as <100 y, with most lying in the range 600-100 y. Converting this to growth rate gives 10^{-11} - 10^{-10} mm.s⁻¹ for both pyroxene species.

ADDITIONAL TIMESCALE INFORMATION FROM PLAGIOCLASE

Three additional timescale constraints are provided by studies involving plagioclase. First, Druitt et al. (2012) placed constraints on the pre-eruptive residence timescale of An_{40±3} plagioclase by characterising Mg contents across core-rim boundaries using SIMS, reconstructing the initial growth profiles, and then modelling subsequent Mg diffusion. The An_{40±3} rims were proposed to have overgrown the An_{50±7} cores 0.3-12 y prior to eruption, with the rimless An_{50±7} crystals residing in the rhyodacite for ≤0.3 y prior to eruption. However, application of a new, better-constrained plagioclase Mg diffusion coefficient (Van Orman et al. 2014) raises the timescale of An_{40±3} plagioclase residence from 0.3-12 y to 1.5-60 y, and that of the rimless An_{50±7} crystals from <0.3 y to <1.5 y (Online Resource 2).

Second, crystal size distributions (CSD) can provide time information if assumptions are made concerning crystal growth rate (see reviews by Higgins 2006 and Armienti 2008). New plagioclase CSDs (Online Resource 2), when combined with a crystal growth rate derived from the Mg diffusion data, give a characteristic plagioclase growth timescale of <40-800 y, which is similar to the residence timescales

derived from the pyroxenes. This approach is circular in that we require a growth rate based on diffusion modelling to obtain a timescale from the CSD data. However it shows that the CSDs and diffusion profiling return mutually consistent timescales.

Third, some MIs in plagioclase $An_{50\pm 7}$ cores contain higher H_2O (≤ 6.9 wt%) than those in the $An_{40\pm 3}$ rims (4.0-5.0 wt%) (Druitt et al. 2016). We have used the H_2O diffusion model of Qin et al. (1992) to estimate that a core-hosted MI with 6.9 wt% H_2O requires <2 years to diffusively re-equilibrate through its host plagioclase with ambient melt containing 4.5 wt% H_2O (Online Resource 2). This shows that cores containing H_2O -rich MIs must have arrived in the magma chamber <2 years preceding the eruption to have avoided diffusive re-equilibration.

DISCUSSION

We now bring together previously published data on the LBA rhyodacite with the new timescale constraints to interpret the processes that took place during build-up to the LBA eruption. The term '*phenocryst*' is used for a crystal that grew from its co-erupted host melt (allowing for small compositional variations), '*antecryst*' is used for a crystal grown from a melt different from the host, but related to the same magmatic system (e.g., Hildreth and Wilson 2007), and '*xenocryst*' is used for a crystal derived from an earlier magmatic system.

Origins of the crystal populations

Phenocryst crystallization

The main plagioclase ($An_{40\pm 3}$), opx (Mg# 57-61) and cpx (Mg# 63-68) components (constituting the rims on core-bearing crystals, and core-free crystals of rim composition) are interpreted as phenocrystic, for the following reasons. (1) Values of K_D ($= [Fe^{tot}/Mg]_{mineral} / [Fe^{tot}/Mg]_{melt}$; where Fe^{tot} is total iron) for opx and cpx derived from the phase equilibria data of Cadoux et al. (2014) on LBA pumice are 0.29 ± 0.07 and 0.22 ± 0.05 , respectively. The opx therefore crystallized from melt with Mg# of 24-31, and the cpx from melt with Mg# of 26-32, which span the compositional range between bulk rhyodacite pumice (Mg# 26-32) and its interstitial glass (Mg# of 18-24). Assuming equilibrium between the rims of opx and cpx crystals, the two-pyroxene thermometer of Putirka (2008; eq. 36) yields a temperature of ~ 850 °C, which is consistent with the Fe-Ti oxide estimates. (2) Plagioclase $An_{40\pm 3}$ compositions are also consistent with crystallization from this range of melts, based on their Sr contents (Druitt et al. 2012) and on phase equilibria (Cadoux et al. 2014). (3) Melt inclusions in all three minerals have compositions ranging from bulk pumice to interstitial glass (Online Resource 1). Although the plagioclase crystals have slightly different Sr isotopic ratios than the host glass, this is attributed to post-crystallization contamination of the melt by a more radiogenic component (Martin et al. 2010; see below).

The phenocrysts are interpreted as recording one or more events of crystallization of the rhyodacitic magma in the proportions $\text{plag} \gg \text{opx} > \text{cpx}$. All three phenocryst phases exhibit oscillatory zoning indicative either of varying P-T-X conditions, or of kinetic effects, during growth (Streck 2008). Multiple sawtooth zoning in $\text{An}_{40 \pm 3}$ plagioclase phenocrysts may record repeated cycles of dissolution-precipitation due to fluctuations of temperature, $P_{\text{H}_2\text{O}}$, and/or melt composition, probably during slow growth (Streck 2008). On the other hand, the presence of Al sector zoning in a small proportion of opx phenocrysts is indicative of fast growth, at least initially. Sector zoning of Al has been generated experimentally in opx by Schwandt and McKay (2006), who showed that it is characteristic of rapid crystal growth with differential entrapment of highly charged cations such as Al^{3+} .

Antecrystic cores of plagioclase, opx and cpx

The compositional and textural diversity of the plagioclase, opx and cpx cores - patchy zoning, multiple dissolution-reaction-precipitation textures, and (in cpx) wispy diffusional overprinting (Fig. 5) - are suggestive of protracted open-system histories prior to incorporation into the rhyodacite. The corroded forms of many cores record a perturbation of P-T-X conditions prior to rim overgrowth.

Possible origins for the cores include (1) xenocrysts derived from older volcanic formations, (2) in-mingled antecrysts from the co-erupted andesitic components (which indeed contain crystals that overlap compositionally with the cores; see Table 4 in Druitt 2014), or (3) antecrysts derived from the source region of the rhyodacite. In some of the plagioclase crystals, the $\text{An}_{50 \pm 7}$ cores and An_{58-88} inner cores have the same ranges of Sr isotopic compositions as the $\text{An}_{40 \pm 3}$ rims (Martin et al. 2010; Di Salvo et al. 2013), which would tend to rule out origin 1. It would also rule out origin 2, since plagioclase in the co-erupted high-Ba andesitic components is more radiogenic ($^{87}\text{Sr}/^{86}\text{Sr}$ of 0.7051-0.7055) than that in the rhyodacite (0.7049-0.7051) (Martin et al. 2010). The plagioclase cores would thus have had to have resided long enough in the rhyodacite to come to isotopic equilibrium, which for a 100 μm -diameter An_{50} crystal would take 10^3 y at 860 °C (Davidson et al. 2001).

We tentatively interpret many of the plagioclase, opx and cpx cores as antecrysts derived from the source region of the rhyodacite. Several lines of evidence are consistent with this. (1) The compositions of the cores (plag $\text{An}_{50 \pm 7}$; cpx Mg# 70-74; opx Mg# 65-71) are very similar to those of the crystal assemblage in equilibrium with LBA-like, 69 wt% SiO_2 melt (plag $\text{An}_{50 \pm 5}$; cpx Mg# 70-75; opx Mg# 70-75) at the postulated source conditions in the experiments of Andújar et al. (2016). (2) Compositions of MIs within the plagioclase $\text{An}_{50 \pm 7}$ cores are consistent with equilibrium with 69 wt% SiO_2 melt (Online Resource 1). (3) Rimless $\text{An}_{50 \pm 7}$ plagioclases have glass selvages of this same composition (Online Resource 1). (4) Sr contents of $\text{An}_{50 \pm 7}$ plagioclase cores and rimless $\text{An}_{50 \pm 7}$ plagioclase imply equilibrium with melts of 63-69 wt% SiO_2 (see Fig. 3d of Druitt et al. 2012).

The rimless An_{50±7} plagioclase

The rimless An_{50±7} plagioclase crystals were introduced (immersed in 69 wt% SiO₂ melt) into the magma within 1.5 years of the eruption, probably immediately prior to, or during, eruptive phase 4. Their heterogeneous distribution implies that intermingling was still ongoing at the time of eruption. We do not have H₂O analyses of MIs in these crystals, so their depth of origin is unconstrained. These crystals lack An_{40±3} rims, implying insufficient time for rim growth prior to eruption quench. They may have been entrained into the magma chamber from underlying crystal mushes during phase 4 caldera collapse.

Rapid magma transfer and chamber assembly

The collective dataset shows that: (1) the LBA rhyodacite was derived mostly from a source region of gabbroic/dioritic composition at 8-12 km depth in the subcaldera pluton, and was transferred to a well-mixed magma chamber at 4-6 km, from where it was erupted; (2) almost all discharged plagioclase, opx and cpx phenocrysts had crystallized within 1000-2000 y of the eruption, mostly within 600-100 y, and that they had grown continuously from initiation until eruption quench; (3) growth of phenocrysts with antecrystic cores containing H₂O-rich melt inclusions must have occurred ≤2 years prior to eruption.

Why did most of the crystals form so late relative to the 17,000-year preceding inter-Plinian period? One possibility is that the LBA magma chamber grew incrementally in the upper crust over many thousands of years, and that crystals were progressively removed by sedimentation. In this case only the most recently formed phenocrysts would have been retained by the magma and erupted. In order to test this, we have modelled in Online Resource 5 the sedimentation rate of plagioclase and pyroxene crystals in the rhyodacite, taking reasonable values for the magma properties (viscosity of 3×10^4 Pa.s and density of 2300 kg.m⁻³) and chamber thickness (~1 km), and accounting for convection. The results show that the characteristic timescale of crystal extraction by sedimentation, even for the largest crystals, is at least an order of magnitude longer than the residence timescales recorded by most of the crystals themselves. We therefore conclude that the data are not consistent with phenocryst growth in a slowly cooling and progressively crystallising magma chamber over many thousands of years, but instead imply the existence of a crystallization event (or several events) within a few centuries to years of the eruption.

We consider three processes that could have caused such a crystallization event: (1) CO₂ flushing from underplated mafic magma (Caricchi et al. 2018), (2) polybaric ascent and cooling (Cottrell et al. 1999), and (3) injection of a hotter, antecryst-bearing melt into a cooler magma chamber (Druitt et al. 2012). A fourth possibility, overpressure release during eruption, is not viable because the timescales are much longer than those of the eruption (probably months to weeks for phase P0, days to hours for phases 1-4; Druitt 2014).

Inspection of Figure 3 of Cadoux et al. (2014) shows that (at 200 MPa) 13 % crystallization could be produced by a drop of H₂O content from 6.3 to 5.2 wt%, which could be achieved by adding about 400 ppm CO₂ (process 1; estimated using the algorithm VolatileCalc of Newman and Lowenstern 2002). However, MIs in the rhyodacite phenocrysts contain much lower CO₂ (all <200 ppm and most <100 ppm; Druitt et al.

2016), so while we cannot firmly exclude some CO₂ flushing from underplated basalt, it was probably not the main process driving crystallization. Some combination of processes 2 and 3 must have been mainly responsible.

In order to test this, we have plotted on a pressure-temperature diagram (Fig. 8) the conditions of the source region (green box) and pre-eruptive magma chamber (orange box). We have also superimposed the liquidus of the LBA rhyodacite and contours of crystallinity from the experiments of Cadoux et al. (2014) at the slightly H₂O-undersaturated conditions appropriate for the LBA magma chamber. Ascent of the rhyodacitic melt from the source region to the magma chamber would have involved 40-140 MPa of decompression and 40-100 °C of cooling, resulting in growth of 5-15 wt% of phenocrysts of An₃₅₋₄₀ plag >> opx > cpx (Cadoux et al. 2014), in agreement with observations. Plagioclase-melt temperatures given by MIs in the An_{40±3} phenocrysts and phenocrystic rims span a range of 30-60 °C (Table 3 in Druitt et al. 2016), broadly consistent with this amount of cooling. Had melt ascent been slow (dashed green arrow, Fig. 8), much crystal growth may have occurred during decompression prior to injection into the magma chamber, whereas had it been rapid (solid green arrow, Fig. 8), most crystal growth would have occurred subsequently by cooling in the chamber.

Our observations favour mainly rapid melt transfer. First, the antecrystic plagioclase An_{50±7} cores containing H₂O-rich melt inclusions were transferred from depth into the magma chamber (and overgrown by the An_{40±3} rims) less than two years before the eruption onset. Hence at least that late magma batch must have ascended rapidly. Second, many pyroxene phenocrysts appear to have grown continuously from their initiation until the eruption (Fig. 7b and c). Had growth been driven partly by decompression (dashed green arrow), then ascent of the magma from depth to the magma chamber would have had to have taken many centuries to decades, which is inconsistent with the first line of evidence and also seems unlikely given current models of silicic magma transfer through the crust (Petford et al. 1993; Menand et al. 2015). Third, some opx crystals exhibit Al sector zoning, possibly consistent with rapid initial crystallization followed by prolonged residence in the magma chamber accompanied by further growth. The data therefore favour an interpretation involving rapid magma ascent followed by prolonged phenocryst growth in the cooler environment of the upper crust. Sawtooth zoning in plagioclase phenocrysts may record repeated precipitation-dissolution cycles in the magma chamber due to convective turnover and heating by the arrival of successive batches of hotter silicic melt.

We note that if phenocryst growth took place mainly by cooling over a 30-60 °C window, then the residence timescales calculated at the final, pre-eruptive temperature of 855 °C may be overestimated. For example, a 30 °C higher mean residence temperature (i.e. 885 °C), would half the modelled diffusion timescales.

The magma chamber was assembled over ~10³ years, and mostly over a few centuries to decades prior to the eruption. While Cottrell et al (1999) assumed that this involved most of the erupted magma, Druitt et al. (2012) suggested more conservatively that it involved just the last silicic melt batch to enter the

chamber. However, our pyroxene timescale data now lead us to prefer the interpretation of Cottrell et al. (1999), since most of the opx and cpx phenocrysts form a single timescale population (Fig. 6).

Reconstruction of the events

Figure 9 summarises the different events inferred to have occurred over the 17 ky-long inter-Plinian period prior to the LBA eruption. Following the 22 ka Cape Riva eruption, repeated crustal intrusion of hydrous basaltic to andesitic melts generated LBA-like silicic melts through a combination of fractional crystallization, crustal melting and pluton defrosting. Explosive eruption of small volumes of this silicic magma at 12-13 ka shows that these LBA-like melts were already present at depth by this time (Satow et al. 2015). By several hundred years prior to the LBA eruption a region of variably mushy gabbro/diorite extended from an unknown upper limit to at least 12 km beneath the volcano. The region consisted of plagioclase, cpx, opx and calcalkaline silicic melts with 6-7 wt% H₂O. Amphibole was probably present in cooler (<940 °C) parts of the body. Smaller quantities of more tholeiitic silicic melts may have been present at greater depths. Thermal waxing and waning during prolonged incubation of crystals and melt may have imparted disequilibrium textures on the crystals that were subsequently erupted as phenocryst cores.

Following possible initial upwellings of LBA-like silicic melt, runaway magma chamber assembly began a few centuries prior to the eruption. Rupturing of the mush structure by ascending melts caused entrainment of plagioclase (An_{50±7}), cpx (Mg# 70-74) and opx (Mg# 65-71) crystals (antecrystic cores). Decompression of volatile-saturated melts with 6-7 wt% dissolved H₂O from 240 MPa to 100-160 MPa caused exsolution of 10-15 vol% of vapour during ascent (Druitt et al. 2016). Upon reaching 4-6 km depth, bubbly melts spread out and accumulated in a single large lens (the magma chamber).

Growth of plagioclase, opx and cpx in the ascending melts was driven partly by volatile-saturated decompression during ascent, but mostly by several tens of °C cooling in the new upper crustal environment. Any amphibole inherited from the source was resorbed. Silicic melts arriving in the magma chamber were progressively homogenized by convection (Druitt et al. 2012, supplementary discussion). Arrival of silicic melts (carrying source-derived plagioclase antecrysts with H₂O-rich MIs) were still ongoing within two years of the eruption, and a small volume of andesite was introduced in the final months to days before eruption (Cottrell et al. 1999). At some late stage, after the growth of most crystals, the rhyodacite mixed with a radiogenic, aphyric melt (Martin et al. 2010).

Finally, conditions were attained within the magma chamber for the propagation of a dyke to the surface, and the eruption began. During ascent, the dyke intersected a variably crystallized shallow intrusion of high-Ba, amphibole-bearing diorite, pushing out some of its contents (see Fig. 12 in Druitt 2014). It is possible that the radiogenic, aphyric melt recognized by Martin et al. (2010) resided within this intrusion, which was indeed more radiogenic than the rhyodacite. Entrainment of rimless An_{50±7} plagioclase and 69 wt % SiO₂ melt may have taken place from crystal mushes of the magma chamber floor as the caldera collapsed during phase 4. Assuming a 45 km² area comparable to that of the caldera, a large

proportion of the <2-km-thick magma chamber must have been emptied during the eruption to account for the 30-80 km³ DRE volume discharged.

Dynamics of melt extraction and ponding

Our evidence for the LBA eruption agrees with an emerging consensus that large silicic magma chambers can be assembled on geologically short timescales through high-flux melt transfer events (Allan et al. 2013; 2017; Wotzlaw et al. 2014; Barker et al. 2016; Cooper et al. 2017; Fabbro et al. 2017). It is also consistent with the mush instability model, in which dispersed melt-rich domains rapidly amalgamate to form one or more large lenses of eruptible magma (Christopher et al. 2015; Cashman et al. 2017; Sparks and Cashman 2017). The time-averaged flux of melts into the LBA magma chamber during assembly is estimated at 0.1-1 km³.y⁻¹ (50 km³ in 500-50 y). These estimates are higher than that of Druitt et al. (2012) (>0.05 km³.y⁻¹), who envisaged that only a few km³ of the magma ascended on these timescales, not all of it. Rapid transfer of such large volumes of viscous, silicic magma would require three conditions.

1. The melt was already segregated into macroscopic melt-rich pockets and lenses in the source region, perhaps due to porous flow instabilities (Solano et al. 2012) or tectonic stresses (Katz et al. 2006; Weinberg et al. 2015), as observed in some fossilized intrusions (Brown 2013; Holness 2018 and references therein); another possibility is that the melt was stored as a sill complex deep within the subcaldera pluton (Jaxybulatov et al. 2014).
2. Compaction-driven melt flow took place through a network of high-permeability conduits or dykes (Petford et al. 1993; Menand et al. 2015), since intergranular flow of silicic melt would be far too slow (e.g., Solano et al. 2012).
3. Unless compensated by basaltic intrusion at depth, rapid extraction of large volumes of silicic melt from the source region was accompanied by (or driven by) subsidence, creating space for the inflating magma chamber higher up (Weibe and Collins 1998; Cruden 1998; Grocott et al. 2009).

Instability of the source region would have initiated a complex chain of magmatic processes consistent with our observations: mixing of compositionally and thermally diverse melt batches, thermal cycling of entrained crystals, polybaric melt ascent, and decompression- and/or cooling-driven crystal growth as the heterogeneous melt mixture drained upwards. The appeal of this mechanism is that it might have occurred on timescales governed by the rheology of the crystal mush, possibly as short as a few years (Sparks and Cashman 2017). More research is required to understand what caused the LBA source region to unlock and compact. It may have occurred spontaneously once (following melt production over many thousands of years) the melt fraction exceeded some threshold. Alternatively, some combination of basaltic intrusion (Cottrell et al. 1999), seismic shaking (Davis et al. 2007; Gottsman et al. 2009) and faulting (Allan et al. 2017) due to a regional rifting event may have been responsible.

It is also unclear why the ascending melts injected to form a short-lived holding chamber at 4-6 km depth, rather than vent directly at the surface. One possibility is that the top of the subcaldera pluton was a

level of steep vertical gradients in density and rheology between underlying ductile mushes and overlying brittle subsolidus rock. Sill formation occurs where dykes encounter layers of low density or high rigidity, causing magma to inject laterally (Menand 2011). Epicentres of earthquakes beneath the caldera during unrest in 2011-12 were limited to the upper few km of the crust (Newman et al. 2012), suggesting that the brittle-ductile transition indeed lies a few km beneath the caldera. The LBA magma chamber might be visualized as a 'tear' near the top of the foundering subcaldera pluton, into which the ascending melts injected. Other possibilities are that the ascending melts intruded by hydrofracturing along a level of crustal weakness (Castro et al. 2016), or that they injected along a low-angle shear zone, as observed in some exhumed intrusive complexes in the Aegean region (Rabillard et al. 2015). We envisage that the bubbly magma remained in this transient holding chamber until sufficient overpressure had developed to drive a dyke to the surface through the cool, brittle uppermost crust and initiate eruption (Degruyter et al. 2015). Uplift and rifting of the surface above the magma chamber may have caused reactivation of caldera faults like the Kameni line, along which the eruption began (Fig. 1c).

The eruption 'trigger'

The evidence for high-flux transfer of large volumes of bubbly silicic melt into the upper crust over the centuries and decades prior to the LBA eruption implies that the eruption itself was almost certainly a consequence of that process. The final input of silicic magma occurred less than two years before the eruption onset; the bubbly magma may have risen to the top of the chamber and been discharged first, explaining the presence of H₂O-rich plagioclase-hosted MIs early in the eruption. It was conceivably this final melt batch that tipped the magma overpressure over the threshold for dyke propagation, triggering the eruption. However, in another sense the eruption had been 'triggered' as much as several centuries previously by the onset of instability deep within the subcaldera pluton. The LBA eruption was probably not the result of a single trigger process, but a complex chain of events.

The presence of quenched andesitic enclaves in the LBA magma led Sparks et al. (1977) to propose that replenishment by andesitic magma played a fundamental role in causing the eruption by pressurizing the chamber. Indeed, the concept of 'mafic' triggering of silicic eruptions is widespread (Pallister et al. 1992; Murphy et al. 1998; Snyder 2000; Leonard et al. 2002; Ruprecht and Bachmann 2010). Cottrell et al. (1999) argued that the andesite was introduced into the magma chamber within a few months to days of the eruption. However our evidence suggests that the volume of silicic melt flooding into the magma chamber was probably much larger than that of the associated andesitic magma. While deep injection of andesite (along with other factors) may have initially played a role in driving the pluton into a state of instability, once silicic melt transfer was underway the andesite may have simply been carried passively from depth by its silicic host, or entrained from nearby intrusions, and have contributed little to pressurizing the chamber.

High-flux melt transfer events at Santorini

Study of the dacitic products of the Cape Riva eruption has revealed a similar story to that proposed for the LBA eruption (Fabbro et al. 2013; 2017). The Cape Riva eruption was preceded by prolonged leakage of a chemically different dacite, showing that the Cape Riva magma chamber did not exist for more than 2800 ± 1400 y, making it a short-lived feature on geological timescales. Furthermore, phenocryst rim timescales for Cape Riva dacite (1 to 25 y for plagioclase using the Van Ormen et al. 2014 diffusion coefficient; 4 to ~ 1000 y for opx using the Dohmen et al. 2016 diffusion coefficient) are broadly similar to those for the LBA magma, and record phenocryst residence over the centuries to years prior to eruption. Drawing on MI $\text{H}_2\text{O}+\text{CO}_2$ barometry and phase equilibria data, Fabbro et al. (2017) postulated that the Cape Riva dacitic magma ascended in a runaway process from a deep source reservoir to a short-lived, shallow holding chamber, from which it subsequently erupted.

It therefore seems likely that rapid magma chamber assembly prior to Plinian eruptions is a general process at Santorini volcano. This behaviour resembles that of Taupo caldera, where studies also imply rapid magma chamber assembly prior to large eruptions (Allan et al. 2013; 2017; Barker et al. 2016; Cooper et al. 2017). Both Santorini and Taupo lie in active continental rift environments (Nomikou et al. 2016; Beaven et al. 2016), suggesting possible roles of seismicity and crustal extension in triggering high-flux melt transfer events at the two systems.

While the normal ascent rate of silicic magma into the upper crust below Santorini through slow compaction of crystal mushes may be sufficient to form small, shallow magma pockets, major destabilization events may be required to build the large magma chambers that feed Plinian eruptions and form calderas. Indeed, if magma chamber formation is driven and accommodated by deep instabilities, then Santorini caldera might be the surface expression of foundering extending down to the middle crust, both above and below the upper crustal chamber (Fig. 10). The estimated flux of melt to the inflating magma chamber prior to the LBA eruption ($0.1\text{-}1 \text{ km}^3 \cdot \text{y}^{-1}$) is two to three orders of magnitude higher than eruption rates during inter-Plinian periods ($0.001 \text{ km}^3 \cdot \text{y}^{-1}$). Repeated destabilisation events in the pluton beneath Santorini may provide a way of reconciling a continuous supply of basaltic magma and heat to the base of the crust with the intermittency of large silicic eruptions.

CONCLUSIONS

Integration of mineral zoning patterns, modelling of multi-mineral diffusion profiles, analysis of crystal size distributions, mineral thermometry, laboratory phase equilibria experiments, and mixed-volatile melt inclusion barometry have enabled us to unravel the history of the rhyodacitic magma of the $30\text{-}80 \text{ km}^3$ DRE Late Bronze Age (LBA) caldera-forming eruption, and to refine and reconcile the different models that have been proposed previously for this system (Cottrell et al. 1999; Martin et al. 2010; Druitt et al. 2012).

Diffusion profile modelling provides timescales for high-temperature residence of three main phenocryst

phases (plagioclase, opx, and cpx) that are consistent to within an order of magnitude, and records one or more events of phenocryst crystallization that took place over the several centuries to years prior to the eruption.

The LBA magma had a two-step history, in accordance with recent conceptual models (Fig. 10; Annen et al. 2006; Brown 2013, and references therein). (1) Prolonged production over at least 8-9 ky (from 12-13 ka to eruption) of silicic melt, mostly in a dioritic/gabbroic region at 8-12 km depth in the subcaldera pluton. (2) Rapid transfer of large volumes of silicic melt to the top of the pluton, where it accumulated to form a magma chamber (4-6 km depth). The crystallization event records the ascent of the vapour-saturated silicic melt and its subsequent cooling in the magma chamber. Magma chamber assembly may have been driven by sudden instability and compaction of the source region (Christopher et al. 2015; Sparks and Cashman 2017), with extraction of melt and associated fluids over a geologically short period of time. The magma chamber may have been a transient holding level near the top of the subcaldera pluton, at the boundary between underlying ductile mushes and overlying brittle, subsolidus crust.

The location of Santorini in a seismically active continental rift zone raises the possibility that magma chamber assembly was triggered by an episode of tectonic rifting and seismicity, possibly accompanied by basaltic injection into the deep crust. The high rates of upper crustal melt accumulation inferred prior to the LBA eruption ($0.1-1 \text{ km}^3 \cdot \text{y}^{-1}$) are two to three orders of magnitude higher than the time-averaged eruptive rates during inter-Plinian periods ($0.001 \text{ km}^3 \cdot \text{y}^{-1}$), and are consistent with accumulation rates necessary to build shallow magma chambers in thermal models (Annen 2009; Gelman et al. 2013; Menand et al. 2015).

Rapid magma chamber assembly also took place prior to the 22 ka Cape Riva eruption (Fabbro et al. 2013; 2017), and may be a general process for caldera-forming eruptions at Santorini. Repeated deep destabilisation events provide a way of reconciling a continuous supply of basaltic magma and heat to the base of the crust with the intermittency of caldera-forming eruptions. It is consistent with the emerging view that large melt-dominated magma chambers are transient features on geologic timescales (Glazner et al. 2004; Druitt et al. 2012; Cooper and Kent 2014; Barboni and Schoene 2014). Had modern geophysical techniques been available a few centuries or decades prior to the LBA eruption, they might have detected a thick column of crystal mush (e.g., Chu et al. 2010; Huang et al. 2015) or stacked sills (Jaxybulatov et al. 2014) beneath Santorini, but probably not a large shallow chamber of melt.

ACKNOWLEDGEMENTS

We thank Jean-Luc Devidal and Jean-Marc Hénot for their expertise in electron microprobe analysis and electron microscopy, respectively, Gareth Fabbro for allowing us to modify his pyroxene diffusion program, Julia Hammer for her introduction to the SHAPE software, Chad Deering for discussions, and Madison Myers for comments on the manuscript. Reviews by J. Ganguly and C.J.N. Wilson were very helpful. TF

acknowledges support from the Atlantis-INVOGE program and a National Science Foundation (NSF) EAPSI fellowship grant. HT is supported by a Royal Society University Research Fellowship. MDH acknowledges support from NSERC (Canada) Discovery grants. This is Laboratory of Excellence ClerVolc contribution number 305.

FIGURE CAPTIONS

Fig. 1. Summary of geological features of Santorini and the Late Bronze Age (LBA) eruption. (a) Map of the Santorini-Amorgos rift zone showing major normal faults (circle on the hanging wall), and the three main subsidence basins. The extent of submarine ignimbrite from the LBA eruption mapped from seismic profiles is shown in yellow (after Sigurdsson et al. 2006). (b) Summary of the activity of Santorini since the 22 ka Cape Riva caldera-forming eruption. (c) Map of Santorini caldera, showing the products of the LBA eruption (including the three main ignimbrite fans from eruptive phase 4), the outline of the shallow Cape Riva caldera that existed prior to the LBA eruption (after Athanassis et al. 2016), the Kameni Line, and the site of the phase 1 vent (after Bond and Sparks 1976). (d) Schematic log of the products of the LBA eruption, with the distributions of the juvenile components and some key cognate components (after Druitt 2014).

Fig. 2. Summary of some previously published petrological features of the Late Bronze Age rhyodacite. (a) Schematic of the different plagioclase types (after Druitt et al. 2012). (b) Histogram of anorthite (An) contents of microprobe analyses, showing that the $An_{40\pm 3}$ rims and $An_{50\pm 7}$ calcic cores of the crystals form two distinct populations (after Druitt et al. 2012). (c) Water contents of plagioclase-hosted melt inclusions as functions of An content, with relationship to the core/rim stratigraphy of a typical core-bearing plagioclase crystal (from Druitt et al. 2016). (d) Water contents of plagioclase-hosted melt inclusions in the rhyodacite as functions of eruptive phase, showing that most H_2O -rich melt inclusions occur in the cores of phenocrysts of phase 1.

Fig. 3. (a,c,e) Back-scattered electron (BSE) images of three representative opx crystals. Crystal S82-30 02 has a magnesian core, whereas crystals S82-34 34 and S82-30 30 each lack one. Numbers in yellow are spot analyses of Mg# (mol %). Numbers in white are diffusion timescales (in years) across the boundaries shown by white lines. (b,d,f) Diffusion profiles for representative zone boundaries (profiles shown by red lines on the BSE images). For reasons of clarity, only the central parts of the profiles appear on the figure. The grey zone around the profile data points shows the analytical uncertainty on the measurements.

Fig. 4. (a) Compositional maps for Mg and Al for a typical weakly zoned opx crystal. The Al map shows clear evidence for sector zoning. (b) Profiles of Mg# (mol %) and Al_2O_3 content (wt %) showing decoupling of the

two and higher frequency variations in Al than in Mg#, suggesting that most of the gradients in Mg# are due to diffusion, not growth. The location of the profile is shown by a yellow line on the Mg map.

Fig. 5. (a,c,e) Back-scattered electron (BSE) images of three representative cpx crystals. Crystals S80-119 04 and S82-34 33 have magnesian cores eaten into by wispy diffusion fronts of more Fe-rich compositions (see insets), whereas crystal S79-101 14 has no core. Numbers in yellow are spot analyses of Mg# (mol %). Numbers in white are diffusion timescales (in years) across the boundaries shown. (b,d,f) Diffusion profiles for representative zone boundaries (profiles shown by red lines on the BSE images). The grey zone around the data points show the analytical uncertainty on the measurements. Orientations of the cpx crystals were determined using the SHAPE software (Dowty 1987), using cpx unit cell parameters a:b:c = 1.097:1:0.596 and $\beta = 106.97^\circ$, by matching measured interfacial angles with those of modelled cpx sections.

Fig. 6. Mg-Fe diffusion timescale spectra for (a) opx and (b) cpx, determined in both cases perpendicular to the c axes of the crystals, as described in the text. Timescales determined on zone boundaries within the crystal rims are colour coded according to fit quality of the diffusion model to the measured Mg# profile. The error bars are calculated from the uncertainty on the $\sqrt{4Dt}$ fit parameter, which takes into account the small analytical error (the error propagation calculation is explained in Online Resource 3). The insets on both figures show diffusion timescale as a function of distance from the crystal edge to the core-rim boundary (or centre of crystal in crystals lacking cores), normalized to one. The opx timescales are calculated using the diffusion coefficient of Dohmen et al. (2016); the small dots show the timescales obtained using Ganguly and Tazzoli (1994) for comparison. The cpx timescales are calculated using the diffusion coefficient of Müller et al. (2013) corrected for diffusion anisotropy, as explained in Online Resource 3.

Fig. 7. (a) Sketches of two endmember scenarios for the growth of pyroxene phenocrysts in the Late Bronze Age magma. In the upper scenario, the crystal grows to its final size, then sits at high temperature until eruption; in the lower scenario the crystal grows progressively. (b,c) Variation of diffusion-derived residence timescales for intra-rim zone boundaries as a function of depth in the crystal. The grey lines depict linear growth from different onset times. Blue dots are timescales from within the crystal rim, and red dots are timescales from the core-rim boundary. Data for the wispy core-rim boundaries in cpx crystals have been omitted, for reasons discussed in the text. Most crystals appear to have grown progressively until the moment of eruption quench, corresponding to the lower case in (a).

Fig. 8. Pressure-temperature diagram showing the pre-eruptive magma chamber (orange) and the main source region for the rhyodacite (green). Crystallinity contours (in wt%, grey lines) and plagioclase anorthite content (An%, black lines) for the LBA rhyodacite are extrapolated from the phase equilibria experiments of

Cadoux et al. (2014) under volatile-saturated conditions at $X_{\text{H}_2\text{O}}=0.85$, where $X_{\text{H}_2\text{O}} = \text{molar H}_2\text{O}/[\text{H}_2\text{O}+\text{CO}_2]$. Green arrows show possible P-T trajectories from the deep source region to the magma chamber. The dashed green arrow depicts ascent slow enough that the magma remains in thermal equilibrium with its surroundings, and involves crystal growth by both decompression and cooling. The solid green arrow depicts very fast, isothermal ascent, and involves crystal growth mostly at shallow depths by cooling and much less by decompression. Note that the paths are schematic and do not take into account adiabatic cooling or release of latent heat of crystallization. The data presented in this paper are more consistent with fast ascent (solid green arrow), as discussed in the text.

Fig. 9. Timeline of events leading up to, and during, the Late Bronze Age (LBA) eruption, following the 22 ka Cape Riva eruption. Timescales are schematic, with solid lines showing best constraints, and dashed lines showing possible extensions. The timelines are coloured according to the three stages, discussed in the main text. Source references for the events are as follows. 1. Druitt et al. (1999), Fabbro et al. (2013). 2 & 9. Druitt (2014), Karátson et al. (2018). 3. Vespa et al. (2006), Vaggelli et al. (2009). 4 & 6. Cottrell et al. (1999), Druitt et al. (2012), Cadoux et al. (2014), Druitt et al. (2016), Andújar et al. (2016), present paper. 5. Satow et al. (2015). 7. Druitt et al. (2016), present paper. 8. Cottrell et al. (1999). 10. Martin et al. (2010). 11. Cioni et al. (2000). 12 & 14. Sparks and Wilson (1990), Druitt (2014). 13. Druitt et al. (2012), present paper.

Fig. 10. Model for the events leading up to the Late Bronze Age (LBA) eruption, showing the three main stages discussed in the text: Stage 1 - prolonged generation of silicic melts in a dioritic/gabbroic region of the subcaldera pluton; stage 2 - instability and compaction-driven transfer of melts to near the top of the pluton, where they accumulate to form a magma chamber; stage 3 - final dyke ascent and the LBA eruption.

ONLINE RESOURCES

- **Online Resource 1.** Compositions of pumices, interstitial glasses and crystal-hosted melt inclusions in the Late Bronze Age (LBA) rhyodacite.
- **Online Resource 2.** Timescale constraints on phenocryst growth and residence from plagioclase.
- **Online Resource 3.** Methods of pyroxene analysis and Mg-Fe diffusion modelling.
- **Online Resource 4.** Results of Mg-Fe diffusion modelling of the Late Bronze Age pyroxenes.
- **Online Resource 5.** Estimation of the sedimentation residence time of a crystal in the Late Bronze Age magma chamber.

REFERENCES

- Allan ASR, Barker SJ, Millet M-A, Morgan DJ, Rooyakkers SJ, Schipper CI, Wilson CJN (2017) A cascade of magmatic events during the assembly and eruption of a super-sized magma body. *Contrib Mineral Petrol* 172:49.
- Allan ASR, Morgan DJ, Wilson CJN, Millet, MA (2013) From mush to eruption in centuries: Assembly of the super-sized Oruanui magma body. *Contrib Mineral Petrol* 166:143–164.
- Andersen DJ, Lindsley DH (1988) Internally consistent solution models for Fe-Mg-Mn-Ti oxides: Fe-Ti oxides. *Am Mineral* 73:714-726.
- Andújar J, Scaillet B, Pichavant M, Druitt TH (2015) Differentiation conditions of a basaltic magma from Santorini, and its bearing on the production of andesite in arc settings. *J Petrol* 56:765–794.
- Andújar J, Scaillet B, Pichavant M, Druitt TH (2016). Generation conditions of dacite and rhyodacite via the crystallization of an andesitic magma. Implications for the plumbing system at Santorini (Greece) and the origin of tholeiitic or calc-alkaline differentiation trends in arc magmas. *J Petrol* 57:1887–1920.
- Annen C (2009) From plutons to magma chambers: Thermal constraints on the accumulation of eruptible silicic magma in the upper crust. *Earth Planet Sci Lett* 284:409–416.
- Annen C, Blundy JD, Sparks RSJ (2006) The genesis of intermediate and silicic magmas in deep crustal hot zones. *J Petrol* 47:505–539.
- Armienti P (2008) Decryption of igneous rock textures: crystal size distribution tools. *Rev Mineral Geochem* 69:623-649.
- Athanassas CD, Bourlès DL, Braucher R, Druitt TH, Nomikou P, Léanni L (2016) Evidence from cosmic ray exposure (CRE) dating for the existence of a pre-Minoan caldera on Santorini, Greece. *Bull Volcanol* 78:35.
- Bachmann O, Bergantz G (2008) The magma reservoirs that feed supereruptions. *Elements* 4:17-21.
- Bachmann O, Bergantz GW (2004) On the origin of crystal-poor rhyolites: Extracted from batholithic crystal mushes. *J Petrol* 45:1565–1582.
- Bachmann O, Huber C (2016) Silicic magma reservoirs in the Earth's crust. *Am Mineral* 101:2377-2404.
- Bacon CR, Druitt TH (1988) Compositional evolution of the zoned calcalkaline magma chamber of Mount Mazama, Crater Lake, Oregon. *Contrib Mineral Petrol* 98:224-256.
- Barboni M, Schoene B (2014) Short eruption window revealed by absolute crystal growth rates in a granitic magma. *Nat Geosci* 7:524–528.
- Barker SJ, Wilson CJN, Morgan DJ, Rowland JV (2016). Rapid priming, accumulation, and recharge of magma driving recent eruptions at a hyperactive caldera volcano. *Geology* 44:323-326.
- Beavan J, Wallace LM, Palmer N, Denys P, Ellis S, Fournier N, Hreinsdottir S, Pearson C, Denham M (2016) New Zealand GPS velocity field: 1995–2013. *New Zealand J Geol Geophys* 59:5-14.
- Bégué F, Deering CD, Gravley DM, Kennedy BM, Chambefort I, Gualda GAR, Bachmann O (2014) Extraction, storage and eruption of multiple isolated magma batches in the paired Mamaku and Ohakuri eruption, Taupo Volcanic Zone, New Zealand. *J Petrol* 55:1653–1684.
- Bergantz GW, Schleicher JM, Burgisser A (2015) Open-system dynamics and mixing in magma mushes. *Nat Geosci* 8: 793–796.
- Bond A, Sparks RSJ (1976) The Minoan eruption of Santorini, Greece. *J Geol Soc Lond* 132:1-16.
- Brophy JG (1991) Composition gaps, critical crystallinity, and fractional crystallization in orogenic (calc-alkaline) magmatic systems. *Contrib Mineral Petrol* 109:173–182.
- Brown M (2013) Granite: From genesis to emplacement. *Geol Soc Am Bull* 125:1079-1113.
- Cadoux A, Scaillet B, Druitt TH, Deloule E (2014) Magma storage conditions of large Plinian eruptions of Santorini Volcano (Greece). *J Petrol* 55:1129-1171.
- Caricchi L, Sheldrake TE, Blundy J (2018) Modulation of magmatic processes by CO₂ flushing. *Earth Planet Sci Lett* 491:160-171
- Cashman KV, Sparks RSJ, Blundy JD (2017) Vertically extensive and unstable magmatic systems: A unified view of igneous processes. *Science* 355:1280.
- Castro JM, Cordonnier B, Schipper CI, Tuffen H, Baumann TS, Feisel Y (2016) Rapid laccolith intrusion driven by explosive volcanic eruption. *Nat Comm* 7: 13585

- Chamberlain KJ, Morgan DJ, Wilson CJN (2014) Timescales of mixing and mobilisation in the Bishop Tuff magma body: Perspectives from diffusion chronometry. *Contrib Mineral Petrol* 168:1034.
- Christopher TE, Blundy J, Cashman K, Cole P, Edmonds M, Smith PJ, Sparks RSJ, Stinton A (2015) Crustal-scale degassing due to magma system destabilization and magma-gas decoupling at Soufriere Hills Volcano, Montserrat. *Geochem Geophys* 16:2797-2811.
- Chu R, Helmlinger DV, Sun D, Jackson JM, Zhu L (2010) Mushy magma beneath Yellowstone. *Geophys Res Lett* 37:L01306.
- Cioni R, Gurioli L, Sbrana, A, Vougioukalakis G (2000) Precursors to the Plinian eruptions of Thera (Late Bronze Age) and Vesuvius (AD 79): Data from archaeological areas. *Phys Chem Earth (A)* 25: 9-11
- Cooper KM (2017) What Does a Magma Reservoir Look Like? The “Crystal's-Eye” View. *Elements* 13:23-28. doi: <https://doi.org/10.2113/gselements.13.1.23>
- Cooper GF, Morgan DJ, Wilson CJN (2017) Rapid assembly and rejuvenation of a large silicic magmatic system: Insights from mineral diffusive profiles in the Kidnappers and Rocky Hill deposits, New Zealand. *Earth Planet Sci Lett* 473:1-13.
- Cooper KM, Kent AJR (2014). Rapid remobilization of magmatic crystals kept in cold storage. *Nature* 506:480–483. doi:10.1038/nature12991
- Costa F, Andreastuti S, Bouvet de Maisonneuve C, Pallister JS (2013) Petrological insights into the storage conditions, and magmatic processes that yielded the centennial 2010 Merapi explosive eruption. *J Volcanol Geotherm Res* 261:209-235.
- Costa F, Dohmen R, Chakraborty S (2008) Time scales of magmatic processes from modelling the zoning patterns of crystals. *Rev Mineral Geochem* 69:545-594.
- Costa F, Morgan D (2010) Time constraints from chemical equilibration in magmatic crystals. In Dosseto A, Turner SP, Van Orman JA (eds), *Timescales of magmatic processes: From core to atmosphere*. Wiley, Chichester, pp 125-159.
- Cottrell E, Gardner JE, Rutherford MJ (1999) Petrologic and experimental evidence for the movement and heating of the pre-eruptive Minoan rhyodacite (Santorini, Greece). *Contrib Mineral Petrol* 135:315–331.
- Cruden AR (1998) On the emplacement of tabular granites. *J Geol Soc Lond* 155:853–862.
- Davidson J, Tepley III F, Palacz Z, Meffan-Main S (2001) Magma recharge, contamination and residence times revealed by in situ laser ablation isotopic analysis of feldspar in volcanic rocks. *Earth Planet Sci Lett* 184:427-442.
- Davis M, Koenders MA, Petford N (2007) Vibro-agitation of chambered magma. *J Volcanol Geotherm Res* 167:24–36.
- Deering CD, Cole JW, Vogel TA (2010) Extraction of crystal-poor rhyolite from a hornblende-bearing intermediate mush: a case study of the caldera-forming Matahina eruption, Okataina volcanic complex. *Contrib Mineral Petrol* 161:129-151.
- Degruyter W, Huber C, Bachmann O, Cooper KM, Kent AJR (2016) Magma reservoir response to transient recharge events: The case of Santorini Volcano (Greece). *Geology* 44:23–26.
- Di Salvo S, Francalanci L, Druitt TH, Braschi E (2013) Short crystal history in the recent magmatic system of Santorini volcano, Greece: inferences from micro-Sr isotope data. *Min Mag* 77:993 (abstract).
- Dohmen R, Ter Heege JH, Becker HW, Chakraborty S (2016) Fe-Mg interdiffusion in orthopyroxene. *Am Mineral*, 101:2210-2221.
- Dowty E (1987) SHAPE Copyright 1994, Shape Software 521 Hidden Valley Road, Kingsport, TN 37663 USA. <http://www.shapesoftware.com>
- Druitt TH (2014) New insights into the initiation and venting of the Bronze-Age eruption of Santorini (Greece), from component analysis. *Bull Volcanol* 76:794.
- Druitt TH, Costa F, Deloule E, Dungan M, Scaillet B (2012). Decadal to monthly timescales of magma transfer and reservoir growth at a caldera volcano. *Nature* 482:77-80.
- Druitt TH, Edwards L, Mellors RM, Pyle DM, Sparks RSJ, Lanphere M, Davies M, Barriero B (1999) Santorini Volcano. *Geol Soc Lond Memoir* 19.
- Druitt TH, Mercier M, Florentin L, Deloule E, Cluzel N, Flaherty T, Médard E, Cadoux A (2016) Magma storage and extraction associated with Plinian and interplinian activity at Santorini caldera (Greece). *J Petrol* 57:461-494.

- Fabbro GN, Druitt TH, Costa F (2017) Storage and eruption of silicic magma across the transition from dominantly effusive to caldera-forming states at an arc caldera (Santorini, Greece). *J Petrol* 58:2429-2464.
- Fabbro GN, Druitt TH, Scaillet S (2013) Evolution of the crustal plumbing system during the build-up to the 22ka caldera-forming eruption of Santorini (Greece). *Bull Volcanol* 75:767.
- Ganguly J, Tazzoli V (1994) Fe²⁺-Mg interdiffusion in orthopyroxene: Retrieval from the data on intracrystalline exchange reaction. *Am Mineral* 79:930-937.
- Gelman SE, Gutiérrez FJ, Bachmann O (2013) On the longevity of large upper crustal silicic magma reservoirs. *Geology* 41:759-762.
- Ghiorso MS, Evans BW (2008) Thermodynamics of rhombohedral oxide solid solutions and a revision of the Fe-Ti two-oxide geothermometer and oxygen-barometer. *Amer J Sci* 308:957-1039
- Glazner AF, Bartley JM, Coleman DS, Taylor RZ (2004) Are plutons assembled over millions of years by amalgamation from small magma chambers? *GSA Today* 14:4-12.
- Gottsmann J, Lavallée Y, Martí J, Aguirre-Díaz G (2009) Magma-tectonic interaction and the eruption of silicic batholiths. *Earth Planet Sci Lett* 284:426-434.
- Grocott J, Arévalo C, Welkner D, Cruden A (2009) Fault-assisted vertical pluton growth: Coastal Cordillera, north Chilean Andes. *J Geol Soc Lond* 166:295-301.
- Gualda GAR, Pamukcu AS, Ghiorso MS, Anderson AT, Sutton SR, Rivers ML (2012) Timescales of quartz crystallization and the longevity of the Bishop giant magma body. *PLoS One* 7:e37492.
- Heiken G, McCoy F, Jr (1984) Caldera development during the Minoan eruption, Thira, Cyclades, Greece. *J Geophys Res* 89:8441-8462.
- Higgins MD (2006) Quantitative textural measurements in igneous and metamorphic petrology. Cambridge University Press, Cambridge, UK.
- Hildreth W, Wilson CJN (2007) Compositional zoning of the Bishop Tuff. *J Petrol* 48:951-999.
- Holness MB (2018) Melt segregation from silicic crystal mushes: a critical appraisal. *Contrib Mineral Petrol* 173:48
- of possible mechanisms and their microstructural record
- Hooft EEE, Nomikou P, Toomey DR, Lampridou D, Getz C, Christopoulou M-E, O'Hara D, Arnoux GM, Bodmer M, Gray M, Heath BA, VanderBeek BP (2017) Backarc tectonism, volcanism, and mass wasting shape seafloor morphology in the Santorini-Christiana-Amorgos region of the Hellenic Volcanic Arc. *Tectonophysics* 712-713:396-414.
- Huang H-H, Lin F-C, Schmandt B, Farrell J, Smith RB, Tsai VC (2015) The Yellowstone magmatic system from the mantle plume to the upper crust. *Science* 15:773-776.
- Huijsmans J (1985) Calc-alkaline lavas from the volcanic complex of Santorini, Aegean Sea, Greece. *Geologica Ultraiectina* 41, 316 pp.
- Huijsmans JPP, Barton M (1989) Polybaric geochemical evolution of two shield volcanoes from Santorini, Aegean Sea, Greece: Evidence for zoned magma chambers from cyclic compositional variations. *J Petrol* 30:583-625.
- Jaxybulatov K, Shapiro NM, Koulakov I, Mordret I, Landès M, Sens-Schönfelder C (2014) A large magmatic sill complex beneath the Toba caldera. *Science* 31:617-619.
- Johnston EN, Sparks RSJ, Phillips JC, Carey S (2014) Revised estimates for the volume of the Late Bronze Age Minoan eruption, Santorini, Greece. *J Geol Soc Lond* 171:583-590.
- Karakas O, Degruyter W, Bachmann O, Dufek J (2017) Lifetime and size of shallow magma bodies controlled by crustal-scale magmatism. *Nat Geosci* 10: 446-450.
- Karátson D, Gertisser R, Telbisz T, Vereb V, Quidelleur X, Druitt TH, Nomikou P, Kósiks (2018) Toward the reconstruction of the disappeared Late Bronze Age intra-caldera island of Santorini, Greece. *Sci Rep* (in press).
- Katz RF, Spiegelman M, Holtzman B (2006) The dynamics of melt and shear localization in partially molten aggregates. *Nature* 442:676-679.
- Kilgour GN, Saunders KE, Blundy JD, Cashman KV, Scott BJ, Miller CA (2014) Timescales of magmatic processes at Ruapehu Volcano from diffusion chronometry and their comparison to monitoring data. *J Volcanol Geotherm Res* 288:62-75.

- Klaver M, Carey S, Nomikou P, Smet I, Godelitsas A, Vroon P (2016) A distinct source and differentiation history for Kolumbo submarine volcano, Santorini volcanic field, Aegean arc. *Geochemistry Geophysics Geosystems* 17:3254–3273.
- Konstantinou KI (2010) Crustal rheology of the Santorini–Amorgos zone: Implications for the nucleation depth and rupture extent of the 9 July 1956 Amorgos earthquake, southern Aegean. *J Geodynam* 50:400-409.
- Koyaguchi T, Kaneko K (1999) A two-stage thermal evolution model of magmas in continental crust. *J Petrol* 40:241-254.
- Krimer D, Costa F (2017) Evaluation of the effects of 3D diffusion, crystal geometry, and initial conditions on retrieved time-scales from Fe-Mg zoning in natural oriented orthopyroxene crystals. *Geochim Cosmochim Acta* 196:271–288.
- Leonard GS, Cole JW, Nairn IA, Self S (2002) Basalt triggering of the c. AD 1305 Kaharoa rhyolite eruption, Tarawera Volcanic Complex, New Zealand. *J Volcanol Geotherm Res* 115:461-486.
- Martin VM, Davidson J, Morgan D, Jerram DA (2010) Using the Sr isotope compositions of feldspars and glass to distinguish magma system components and dynamics. *Geology* 38:539-542.
- Mason BG, Pyle DM, Oppenheimer C (2004) The size and frequency of the largest explosive eruptions on Earth. *Bull Volcanol* 66:735-748.
- Matthews NE, Huber C, Pyle DM, Smith VC (2012) Timescales of magma recharge and reactivation of large silicic systems from Ti diffusion in quartz. *J Petrol* 53:1385-1416.
- Menand T (2011) Physical controls and depth of emplacement of igneous bodies: A review. *Tectonophys* 500:11-19.
- Menand T, Annen C, de Saint Blanquat M (2015) Rates of magma transfer in the crust: Insights into magma reservoir recharge and pluton growth. *Geology* 43:199-202.
- Morgan DJ, Blake S, Rogers NW, De Vivo B, Rolandi G, Davidson JP (2006) Magma chamber recharge at Vesuvius in the century prior to the eruption of AD 79. *Geology* 38:845-848.
- Morgan DJ, Blake S, Rogers NW, DeVivo B, Rolandi G, Macdonald R, Hawkesworth CJ (2004) Time scales of crystal residence and magma chamber volume from modelling of diffusion profiles in phenocrysts: Vesuvius 1944. *Earth Planet Sci Lett* 222:933–946.
- Müller T, Dohmen R, Becker HW, ter Heege JH, Chakraborty S (2013) Fe-Mg interdiffusion rates in clinopyroxene: Experimental data and implications for Fe-Mg exchange geothermometers. *Contrib Mineral Petrol* 166:1563-1576.
- Murphy MD, Sparks RSJ, Barclay J, Carroll MR, Lejeune A-M, Brewer TS, Macdonald R, Black S, Young S (1998) The role of magma mixing in triggering the current eruption at the Soufriere Hills Volcano, Montserrat, West Indies. *Geophys Res Lett* 25:3433-3436. doi: 10.1029/98GL00713
- Nakagawa K, Nagahara H, Ozawa K, Tachibana S, Yasuda A (2005) Experimental determination of AlAl-SiMg interdiffusion coefficient in orthopyroxene. Japan Geoscience Union Meeting, Tokyo, pp. K038-013 (abstract).
- Newman AV, Stiros S, Feng L, Psimoulis P, Moschas F, Saltogianni V, Jiang Y, Papazachos C, Panagiotopoulos D, Karagianni E, Vamvakaris D (2012) Recent geodetic unrest at Santorini Caldera, Greece. *Geophysical Research Letters* 39:L06309.
- Newman, S. & Lowenstern, J. (2002). VolatileCalc: A silicate melt–H₂O–CO₂ solution model written in Visual Basic for excel. *Computers & Geosciences* 28:597-604.
- Nomikou P, Hübscher C, Ruhnau M, Bejelou K (2016) Tectono-stratigraphic evolution through successive extensional events of the Anhydros Basin, hosting Kolumbo volcanic field at the Aegean Sea, Greece. *Tectonophys* 671:202-217.
- Nomikou P, Parks MM, Papanikolaou D, Pyle DM, Mather TA, Carey S, Watts AB, Paulatto M, Kalnins ML, Livanos I, Bejelou K, Simou E, Perros I (2014) The emergence and growth of a submarine volcano: The Kameni islands, Santorini (Greece). *GeoResJ* 1–2:8-18.
- Oppenheimer, C (2013) Nature and impacts of the Minoan eruption of Santorini in 1600. In: Meller H, Bertemes F, Bork HR (eds) *Cultural change in the shadow of the Thera eruption?* pp 49-58.
- Pallister JS, Hoblitt RP, Reyes AG (1992) A basalt trigger for the 1991 eruptions of Pinatubo volcano? *Nature* 356:426-428. doi:10.1038/356426a0

- Pamukcu AS, Gualda GAR, Bégué F, Gravley DM (2015) Melt inclusion shapes: Timekeepers of short-lived giant magma bodies. *Geology* 43:947-950.
- Papale P, Moretti R, Barbato D (2006) The compositional dependence of the saturation surface of H₂O+CO₂ fluids in silicate melts. *Chemical Geology* 229:78-95. doi:10.1016/j.chemgeo.2006.01.013
- Petford N, Kerr RC, Lister JR (1993) Dike transport of granitoid magmas. *Geology* 21:845-848.
- Petrone CM, Bugatti G, Braschi E, Tommasini S (2016) Pre-eruptive magmatic processes re-timed using a non-isothermal approach to magma chamber dynamics. *Nat Comm* 7:12946.
- Putirka KD (2008) Thermometers and barometers for volcanic systems. *Rev Mineral Geochem* 69:61-120.
- Pyle DM (1990) New estimates for the volume of the Minoan eruption. In: Hardy DA, Keller J, Galanopoulos VP, Flemming NC, Druitt TH (eds) *Thera and the Aegean World III*, vol 2. Thera Foundation, London, pp 113-121.
- Pyle DM, Elliott JR (2006) Quantitative morphology, recent evolution, and future activity of the Kameni Islands volcano, Santorini, Greece. *Geosphere* 2:253-268.
- Qin Z, Fangqiong L, Anderson A (1992) Diffusive reequilibration of melt and fluid inclusions. *American Mineralogist* 77: 565–576.
- Rabillard A, Arbaret L, Jolivet L, Le Breton N, Gumiaux C, Augier R, Grasemann B (2015). Interactions between plutonism and detachments during metamorphic core complex formation, Serifos Island (Cyclades, Greece). *Tectonics* 34, doi:10.1002/2014TC003650.
- Rubin AE, Cooper KM, Till CB, Kent AJR, Costa F, Bose M, Gravley D, Deering C, Cole J (2017) Rapid cooling and cold storage in a silicic magma reservoir recorded in individual crystals. *Science* 356:1154-1156. <https://doi.org/10.1126/science.aam8720>
- Ruprecht P, Bachmann O (2010) Pre-eruptive reheating during magma mixing at Quizapu volcano and the implications for the explosiveness of silicic arc volcanoes. *Geology* 38:919-922.
- Satow C, Tomlinson EL, Grant KM, Albert PG, Smith VC, Manning CJ, Ottolini L, Wulf S, Rohling EJ, Lowe JJ, Blockley SPE, Menzies MA (2015). A new contribution to the Late Quaternary tephrostratigraphy of the Mediterranean: Aegean Sea core LC21. *Quat Sci Rev* 117:96-112.
- Saunders K, Blundy J, Dohmen R, Cashman K (2012) Linking petrology and seismology at an active volcano. *Science* 336:1023-1027.
- Schmitt AK, Stockli DF, Song EJ, Storm S (2016) Equilibrium and Disequilibrium of ²³⁰Th-²³⁸U in Zircon from the Minoan Eruption, Santorini, Aegean Sea, Greece. *American Geophysical Union General Assembly* (abstract).
- Schöpa A, Annen C (2013). The effects of magma flux variations on the formation and lifetime of large silicic magma chambers. *J Geophys Res* 118:926–942.
- Schwandt CS, McKay GA (2006) Minor- and trace-element sector zoning in synthetic enstatite. *Am Mineral* 91:1607-1615. doi:10.2138/am.2006.2093.
- Self S (2006) The effects and consequences of very large explosive volcanic eruptions. *Phil Trans Roy Soc* 364:2073-2097.
- Shea T, Costa F, Krimer D, Hammer JE (2015) Accuracy of timescales retrieved from diffusion modeling in olivine: A 3D perspective. *Am Mineral* 100:2026-2042.
- Sigurdsson H, Carey S, Alexandri M, Vougioukalakis G, Croff K, Roman C, Sakellariou D, Anagnostou C, Rousakis G, Ioakim C, Goguo A, Ballas D, Misaridis T, Nomikou P (2006) Marine investigations of Greece's Santorini Volcanic Field. *EOS Trans Am Geophys Union* 87:337-342.
- Sigurdsson H, Carey S, Devine JD (1990) Assessment of the mass, dynamics, and environmental effects of the Minoan eruption of Santorini Volcano. In: Hardy DA, Keller J, Galanopoulos VP, Flemming NC, Druitt TH (eds) *Thera and the Aegean World III*, vol 2. Thera Foundation, London, pp 100-112.
- Singer BS, Costa F, Herrin JS, Hildreth W, Fierstein J (2016) The timing of compositionally-zoned magma reservoirs and mafic “priming” weeks before the 1912 Novarupta-Katmai rhyolite eruption. *Earth Planet Sci Lett* 451:125-137.
- Snyder D (2000) Thermal effects of the intrusion of basaltic magma into a more silicic magma chamber and implications for eruption triggering. *Earth Planet Sci Lett* 175:257-273.
- Solano JMS, Jackson MD, Sparks RSJ, Blundy JD, Annen C (2012) Melt segregation in deep crustal hot zones: A mechanism for chemical differentiation, crustal assimilation and the formation of evolved magmas. *J Petrol* 53:1999–2026.

- Sparks RSJ, Cashman KV (2017) Dynamic magma systems: implications for forecasting volcanic activity. *Elements* 13:35-40.
- Sparks RSJ, Wilson CJN (1990) The Minoan deposits: A review of their characteristics and interpretation In: Hardy DA, Keller J, Galanopoulos VP, Flemming NC, Druitt TH (eds) *Thera and the Aegean World III*, vol 2. Thera Foundation, London, pp 89-99.
- Sparks SRJ, Sigurdsson H, Wilson L (1977) Magma mixing: A mechanism for triggering acid explosive eruptions. *Nature* 267:315-318.
- St. Seymour K, Christanis K, Bouzinos A, Papazisimou S, Papatheodorou G, Moran E, Dénès G (2004) Tephrostratigraphy and tephrochronology in the Philippi peat basin, Macedonia, Northern Hellas (Greece). *Quat Int* 121:53-65.
- Streck MJ (2008) Mineral textures and zoning as evidence for open system processes. *Rev Mineral Geochem* 69:595-622.
- Till CB, Vazquez JA, Boyce JW (2015) Months between rejuvenation and volcanic eruption at Yellowstone caldera, Wyoming. *Geology* 43:695-698.
- Tomiya A, Takahashi E (2005) Evolution of the magma chamber beneath Usu Volcano since 1663: A natural laboratory for observing changing phenocryst compositions and textures. *J Petrol* 46:2395-2426.
- Vaggelli G, Pellegrini M, Vougioukalakis G, Innocenti S, Francalanci L (2009) Highly Sr radiogenic tholeiitic magmas in the latest inter-Plinian activity of Santorini volcano, Greece. *J Geophys Res* 114:B06201.
- Van Orman JA, Cherniak DJ, Kita NT (2014) Magnesium diffusion in plagioclase: Dependence on composition, and implications for thermal resetting of the ^{26}Al – ^{26}Mg early solar system chronometer. *Earth Planet Sci Lett* 385:79-88.
- Vespa M, Keller J, Gertisser R (2006) Interplinian explosive activity of Santorini volcano (Greece) during the past 150,000 years. *J Volcanol Geotherm Res* 153: 262-286.
- Wark DA, Hildreth W, Spear FS, Cherniak DJ, Watson EB (2007) Pre-eruption recharge of the Bishop magma system. *Geology* 35:235-238.
- Weinberg RF, Veveakis E, Regenauer-Lieb K (2015) Compaction-driven melt segregation in migmatites. *Geology* 43:471-474.
- Wiebe RA, Collins WJ (1998) Depositional features and stratigraphic sections in granitic plutons: Implications for the emplacement and crystallization of granitic magma. *J Struct Geol* 20:1273-1289.
- Wilson CJN, Charlier BLA (2009). Rapid rates of magma generation at contemporaneous magma systems, Taupo Volcano, New Zealand: Insights from U-Th model-age spectra in zircons. *J Petrol* 50:875-907.
- Wotzlaw J-F, Bindeman IN, Watts KE, Schmitt AK, Caricchi L, Schaltegger U (2014) Linking rapid magma reservoir assembly and eruption trigger mechanisms at evolved Yellowstone-type supervolcanoes. *Geology* 42:807-810.

Table 1. Summary of timescales of crystal growth and residence in the Late Bronze Age rhyodacite.

Mineral	Method	Time prior to eruption (years)	Geological interpretation of timescale
Orthopyroxene	Mg-Fe diffusion (Dohmen et al. 2016)	1 to 459; outlier at 852	High-T opx phenocryst residence
	Mg-Fe diffusion (Ganguly & Tazzoli 1994)	1 to 110; outlier at 157	High-T opx phenocryst residence
Clinopyroxene	Mg-Fe diffusion	2 to 500; outliers at 1173-2031	High-T cpx phenocryst residence
Plagioclase	Mg diffusion	1.5 to 60	High-T plag phenocryst residence
Plagioclase	Mg diffusion	<1.5	High-T residence of rimless calcic plag crystals
Plagioclase	Crystal size distribution	40-800	Growth time of plag phenocrysts
Plagioclase	H ₂ O diffusion	<2	Arrival of last silicic input into the magma chamber

Opx: orthopyroxene; cpx: clinopyroxene; plag: plagioclase; High-T: high temperature.