

1 **October 31, 2015**

2

3 **Validity and reliability of two field-based leg stiffness devices: implications for**
4 **practical use**

5

6 Luca Ruggiero,^{1,2} Susan Dewhurst,¹ Theodoros M. Bampouras¹

7

8 ¹Department of Medical and Sport Sciences, University of Cumbria, Lancaster,
9 LA1 3JD, United Kingdom; ²School of Health and Exercise Sciences, University of
10 British Columbia, Kelowna, British Columbia, Canada.

11

12 **Funding:** *No external funding were received.*

13 **Conflict of Interest Disclosure:** *The authors have no conflict of interest to disclose.*

14 **Correspondence Address:** *Theodoros M. Bampouras, University of Cumbria, Department of*
15 *Medical and Sport Sciences, Human Performance Laboratory, Bowerham Road, Lancaster LA1*
16 *3JD, United Kingdom. Email: theodoros.bampouras@cumbria.ac.uk. Tel. No.: +44 1524 590837.*

17 **Running Head:** *Field-based leg stiffness measurement.*

18 **Abstract**

19 Leg stiffness is an important performance determinant in several sporting activities. The aim of
20 this study was to evaluate the criterion-related validity and reliability of two field-based leg
21 stiffness devices, Optojump Next® (Optojump) and Myotest Pro® (Myotest) in different testing
22 approaches. Thirty-four males performed, on two separate sessions, three trials of 7 maximal
23 hops, synchronously recorded from a force platform (FP), Optojump and Myotest. Validity
24 (Pearson's correlation coefficient, r ; relative mean bias, bias; 95% limits of agreement, 95%LoA)
25 and reliability (coefficient of variation, CV; standard error of measurement, SEM; intraclass
26 correlation coefficient, ICC) were calculated for first attempt, maximal attempt, and average
27 across three trials. For validity all three methods, Optojump correlated highly to the FP (range $r =$
28 0.98-0.99) with small bias (range 0.91-0.92, 95 LoA 0.86-0.98). Myotest demonstrated high
29 correlation to FP (range $r = 0.81-0.86$) with large bias (range 1.92-1.93, 95% LoA 1.63-2.23).. In
30 terms of reliability, Optojump yielded a low CV (range 5.9%-6.8%), SEM ranging 1.8-2.1 kN/m,
31 and high ICC (range 0.82-0.86). Myotest had a larger CV (range 8.9%-13.0%), SEM ranging
32 from 6.3-8.9 kN/m, and moderate ICC (range 0.64-0.79). The findings present important
33 information for these devices and support the use of a single trial to assess leg stiffness in the
34 field, thus testing in a time-efficient way.

35

36 **Keywords:** hopping test, vertical stiffness, test-retest, sensitivity.

37

38 **Word Count:** 3557

39

Introduction

40 Leg stiffness describes the response of the lower limbs to generate force and resist
41 deformation during rebound activities.^{8,9} Enhanced stiffness is beneficial to reduce metabolic cost
42 of bouncing gait (i.e. running, hopping)¹²⁻¹⁴ as well as to attaining high sprinting speed¹⁵⁻¹⁶,
43 whereas lower leg stiffness may lead to less storage and recoil of elastic energy, placing greater
44 metabolic demand during push-off, and to a reduced ability to sustain impact loads, raising injury
45 risk.^{9,11,17} Thus, leg stiffness evaluation can be important both prior to and during training.

46 Two field-based devices can assess leg stiffness are the Optojump Next[®] (Microgate,
47 Bolzano, Italy; Optojump) and Myotest Pro[®] (Myotest, Sion, Switzerland; Myotest).²¹⁻²²
48 Optojump Next[®] is an optical measurement system consisting of two infrared photocell bars that
49 can derive contact and flight times from the breaking of the transmitted beam, whereas Myotest
50 Pro[®] is a wireless lightweight portable triaxial accelerometer that can be fixed on the athlete.
51 Both are portable and practical, allowing athletes to jump on any given surface, used largely
52 because of their versatility and reasonable cost.²³⁻²⁵

53 Several studies have examined the devices' criterion-related validity and reliability for
54 vertical jump height from squat and countermovement jumps in comparison to a force
55 platform.^{22,25-27} Leg stiffness with the above equipment, however, has either not been examined
56 or has been conducted in a less time-efficient way. For example, in the Choukou et al²² study, the
57 authors processed the data obtained, thus determining the reliability of the processed data rather
58 than the calculated value for Myotest Pro[®], while substantially adding to the analysis time.
59 Moreover, measurement reliability of the criterion-related leg stiffness outcome was not
60 determined, raising uncertainty on interpretation of the results.

61 The aim of the present study was twofold. Criterion-related validity (the force platform as
62 gold standard), reliability and sensitivity of both Optojump Next® and Myotest Pro® (henceforth
63 Optojump and Myotest, respectively) for measuring leg stiffness in hopping was assessed, with
64 no manipulation of the software, hardware or the data obtained, where possible. This approach
65 was deemed to reflect more closely in the field testing conditions while provides realistic
66 information for the equipment (i.e. when used as close to the manufacturer suggestions as
67 possible). These aspects were then examined with three different procedures, namely the first trial
68 executed, the average across three trials, and the maximal stiffness value out of them, to explore
69 whether a single trial was sufficient, offering practical information in terms of timing
70 requirements for leg stiffness testing.

71 **Methods**

72 **Participants**

73 Thirty-four male University students (age 21.8 ± 3.9 years, height 1.83 ± 0.07 m, body
74 mass 79.0 ± 11.4 kg) took part in the study. They were all physically active, free from lower
75 limbs injuries for at least six months prior to the testing sessions, and competing in various team
76 sports. All participants were instructed to refrain from strenuous exercise, alcohol, and caffeine
77 for 2 days, 24 and 2 hours before testing, respectively. Procedures were approved by the
78 University Ethical Committee and informed consent was given by all participants.

79 **Procedures**

80 Participants visited the laboratory on two separate sessions, 1 week apart, at the same time
81 of the day. The same protocol was strictly followed in each session. Following a standardised
82 warm up, participants familiarised themselves with the test. All participants reported to be

83 completely accustomed with the task, and no more than two familiarizing attempts were needed.
84 Following a 5-minute rest, 3 trials of the 7MH were performed, with 2 minutes resting between
85 trials. Participants were instructed to jump as high as possible, with minimal contact time, and
86 with arms akimbo at all times. Hopping was chosen as well-documented functional task, and
87 maximal effort was required as usually performed in field testing.

88 All jumps were performed on a force platform (FP) (AccuPower, AMTI, Watertown, MA,
89 United States; 200 Hz sampling rate). The resulting vertical force-time trace allowed measuring
90 participants' body mass, contact and flight times, used to calculate leg stiffness as $k = \frac{m \times \pi(\text{flight time} + \text{contact time})}{(\text{contact time}^2 \left(\frac{\text{flight time} + \text{contact time}}{\pi} - \frac{\text{contact time}}{4} \right))}$
91 (Eq. 1)¹⁸. Data was synchronously collected by Optojump and Myotest (Figure 1). Optojump 1
92 meter bars (resolution of 96 diodes, sampling rate of 1 kHz) were placed on the lateral border
93 lines of FP. Contact and flight times for all seven jumps of 7MH test and the participant's body
94 mass was used in Eq. 1 to calculate leg stiffness.¹⁸ Myotest (sampling rate of 500 Hz) was fixed
95 on the participants by means of an elastic Velcro waistband, fastened on a line passing on both
96 great trochanters and the medium part of the gluteal region, as per manufacturer instructions.
97 Myotest uses internal algorithms for calculation of leg stiffness taking into account the average of
98 the best three hops from any given trial of 7MH. Leg stiffness values were displayed on the
99 device screen immediately after the trial.
100

101 Data Analysis

102 Leg stiffness was examined for all three devices from a) the 1st trial from each session
103 (K_{First}), b) the average across three trials from session (K_{Avg}), and c) the maximal value from
104 session (K_{Max}).

Commented [TB1]: Luca, doublecheck it is correct (especially that I did not miss a bracket somewhere).

105 For the K_{Max} approach, Wilcoxon signed-rank test was used to check for conformity of the
106 trial number wherein the maximum stiffness value occurred between each device and FP. No
107 significant difference was revealed for any comparison. For the K_{Avg} approach, within-subject
108 variation over the three trials was assessed via 1-way repeated measures ANOVA before
109 averaging, reporting no significant differences. Therefore, stiffness results for each subject were
110 collapsed to a single value per session.

111 **Criterion-related validity assessment procedures**

112 As no significant test-retest differences (examined with paired t -test) between Session 1
113 and Session 2 were reported for any of the equipment, results were collapsed to a single
114 participant value for each of the K_{First} , K_{Max} , and K_{Avg} procedures.²⁸ These single values were
115 then used to investigate for criterion-related validity of the Optojump and Myotest in comparison
116 to the FP. Data was checked for heteroscedasticity by correlating the test score differences
117 between either Optojump or Myotest and the FP to their mean value, for each procedure,
118 following the method by Bland and Altman.²⁹ As significant correlations were found, indicating
119 the presence of heteroscedasticity for the validity investigation, raw data was transformed using
120 the natural logarithm before further analysis occurred.²⁹ Thereby, normality of residuals (log test
121 score differences between either Optojump or Myotest and FP) was examined using the Shapiro-
122 Wilk test, and with normality defined as the ratio of skewness and kurtosis to the respective
123 standard error not exceeding ± 2.0 .³⁰ Normal distribution was confirmed for each procedure and
124 device. Criterion-related validity to the FP was assessed via Pearson's correlation coefficient and
125 relative mean bias. In addition, as suggested by Bland & Altman²⁹, agreement between the
126 measurement devices (either Optojump or Myotest related to FP) was examined, and 95% limits
127 of agreement (95% LoA) were reported. The limits display that, for about 95% of cases, the leg
128 stiffness measurement of the examined device may differ from the one of the FP by the lower

129 limit to the upper limit. Pearson's correlation coefficient (r) was interpreted as indicating high
130 correlation for an r value above 0.8.³¹ Relative mean bias was calculated as the difference
131 between the logarithmic transformed score means of either Optojump or Myotest and FP, and
132 reported as antilog. Because the antilog of the difference between two logarithmic measurements
133 equals to the dimensionless ratio between the same two measurements, the relative mean bias
134 must be interpreted as the ratio between the average outcome of the examined device and that of
135 the FP. Likewise, 95% LoA were calculated on the logarithmic scale, and reported as antilogs as
136 mean difference \pm 1.96 standard deviations of the differences.

137 **Reliability assessment procedures**

138 The residuals (raw 1st – 2nd session score differences) and the respective pair means for
139 each piece of equipment and procedures were correlated, to investigate the presence of
140 heteroscedasticity.²⁹ No significant correlation was found, indicating homoscedastic distribution.
141 Thus, data was further analyzed as raw values. Normality of the residuals was then checked for
142 both each procedure and device, and confirmed.

143 Indices of both absolute and relative reliability were used for the investigation, for each
144 procedure. Absolute intersession reliability was assessed via coefficient of variation and standard
145 error of measurement (CV and SEM, respectively). CV was calculated as the standard deviation
146 (SD) divided by the mean and multiplied by 100 for each participant, and then averaged.³² The
147 threshold was set at 10%, with values below suggesting high consistency.^{33,34} To better represent
148 all individuals, SD of CV was also reported in addition to group mean CV.³³ SEM was calculated
149 as the square root of the mean square error term in a repeated measures ANOVA.³⁰ SEM is of
150 practical importance, as it allows coaches easily determine the minimum difference (MD) needed
151 for a performance change to be considered real (95% confidence) rather than a measurement
152 error^{30,35}, using the following formula:

153
$$MD = SEM \times 1.96 \times \sqrt{2} \quad (\text{Equation 2})$$

154 Finally, relative intersession reliability was assessed by interclass correlation coefficient
155 (ICC), calculated according to Hopkins³⁶ as:

156
$$1 - ((SEM)^2 / (\text{mean of subjects' standard deviation between trials})^2)$$

157 An ICC value above 0.8 was set as a threshold for indicating small measurement error.³⁷ Ninety-
158 five per cent confidence intervals (95 % CI) for ICCs were also calculated using the spreadsheet
159 provided by Hopkins³⁸, representing the likely range of values containing the true population of
160 ICCs in approximately 95% of the cases.

161 Statistical significance level was set for each test at $P < 0.05$. All statistical tests were
162 performed using SPSS software (IBM SPSS Statistics, version 20, Inc., Chicago, IL, USA).

163 **Results**

164 Leg stiffness calculated from Optojump (Table 1), demonstrated high correlation to FP
165 leg stiffness (Table 1) in all analysis procedures (range $r = 0.98-0.99$, $P < .001$) with relative
166 mean bias ranging from 0.91 to 0.92 (Table 2). 95%LoA (Table 2, Figure 2) were not
167 substantially different between procedures. Leg stiffness calculated from Myotest (Table 1) also
168 showed high correlation to FP leg stiffness in all methods (range $r = 0.81 - 0.86$, $P < .001$), with
169 higher measured leg stiffness (relative bias ranging between 1.92 and 1.93, Table 2). 95%LoA
170 reported were wider compared to Optojump (Table 2), evident from different y-axis ranges
171 (Figure 2).

172 FP exhibited low CV, suggesting good absolute reliability (Table 3). However, when
173 relative reliability was considered, only K_{Max} procedure reported an $ICC \geq 0.8$, with K_{First} and
174 K_{Avg} ICCs of 0.74 and 0.79, respectively. Optojump revealed high absolute and relative reliability
175 in all three analysis procedures, shown from relatively low values of group mean CV and high

176 ICC (Table 3). For Myotest, the K_{Avg} procedure was the more consistent one with a low CV but
177 moderate ICC, whereas K_{First} and K_{Max} reported lower consistency (Table 3). For all procedures,
178 Myotest yielded higher SEM than FP and Optojump (Table 3).

179 **Discussion**

180 The aim of this study was to determine criterion-related validity and reliability of two
181 commonly used field-based devices (i.e. Optojump and Myotest) in measuring leg stiffness. In
182 addition, three different analysis procedures were examined (i.e. K_{First} , K_{Max} and K_{Avg}), to provide
183 practical information in terms of timing requirements to assess leg stiffness. Optojump showed a
184 valid leg stiffness measurement compared to FP, with all analysis procedures being reliable.
185 Myotest also showed valid leg stiffness measurement compared to FP, but with moderate
186 reliability for all three procedures.

187 Leg stiffness values measured with Optojump agreed well with the FP values and are
188 within the range reported from previous literature.^{10,18-20} When the three different procedures
189 were considered, all three procedures showed high reliability, with similar indexes to earlier
190 research using the FP.^{39,40} The systematic bias of Optojump was most likely due to the placement
191 of Optojump bars on the FP (Figure 1), meaning the infrared beams were 0.3 cm higher than the
192 FP surface.²⁶ Consequently, increased contact time and reduced flight time compared to those of
193 FP, resulted in lower leg stiffness.^{4,18} Although this height discrepancy may appear as a
194 methodological concern, we opted for this approach as it more closely reflects field testing,
195 where the placement of the Optojump bars on a given surface (e.g. ground, court, track), will be
196 included in the measurement.

197 Leg stiffness values obtained from Myotest were significantly different with the FP and
198 outside the values seen from hopping in previous reports.^{10,18-20} Further, reliability for all three

199 procedures was moderate. Our results contradict the study by Choukou et al.,²² who reported the
200 5 hop test as valid and reliable in measuring leg stiffness using Myotest²². The higher number of
201 total hops considered in Choukou et al.²² (all 5, compared to best 3 in the present investigation)
202 could have reduced within-subject variability³⁶, possibly explaining the discrepancy. The
203 overestimation of leg stiffness and poorer reliability of Myotest in relation to the FP might be
204 attributed to the following reasons. Myotest leg stiffness computation is based on integration of
205 acceleration, with respect to mass and time, and establishes the time interval of integration when
206 the accelerations are null.²² As maximal descending and ascending velocities are not achieved at
207 those exact points, contact time and centre of mass displacement are underestimated, while flight
208 time, force and jump height are overestimated^{22,24}; in turn, magnifying leg stiffness values.
209 Secondly, the fast transition between braking and push-off phase during the maximal hopping
210 task is likely to have caused vibrations of the device and in turn erroneous acceleration
211 detections. Indeed, previous comparisons of the Myotest against FP using single jumps (and,
212 thus, little or no vibrations affecting the measurement) have reported better agreement.²⁷

213 High sensitivity of a device allows for better determining differences resulting from true
214 changes of the physical characteristic evaluated rather than from a measurement error.^{35,42} For
215 this purpose, we calculated SEM, to subsequently determine MD and construct confidence
216 intervals, which can detect with reasonably good confidence (95%) real changes in the variable
217 being measured. The importance of these confidence intervals for each device, the use of MD in
218 assessing changes in performance, and of its magnitude in doing so with small changes can be
219 better illustrated in the following example. Let us suppose that we tested an athlete who in the
220 first testing session achieves a value of 25 kN/m. Following a training intervention, the athlete
221 tests again and achieves a value of 33 kN/m. Replacing the Optojump and Myotest SEM from the

222 K_{FIRST} procedure described in this paper (Table 3) in Eq. 2, the MD representing a true difference
223 will be 5.8 kN/m for Optojump, and 21.1 kN/m for Myotest. As the test-retest difference (33 – 25
224 = 8 kN/m) lies outside the MD for Optojump, we would be certain (more than 95%) of a true
225 change, whereas we would be unable to reach a conclusion using Myotest.

226 Assessing many athletes within the time-restrictions of a training or an assessment
227 session, requires use of scientifically rigorous methods and consideration of the the practical
228 aspects of the assessment (e.g. time availability, set-up and feedback time). Our results showed
229 that leg stiffness assessment can be completed in a valid and reliable manner in the field, with
230 minimal data manipulation (calculation of leg stiffness via Eq. 1). Further, leg stiffness can be
231 confidently assessed with the use of a single trial, allowing time-efficient testing, in particular
232 short time frames are available or large populations are to be tested.

233 References

- 234 1. Blickhan R. The spring-mass model for running and hopping. *J Biomech.* 1989;22(11–
235 12):1217-1227. [http://dx.doi.org/10.1016/0021-9290\(89\)90224-8](http://dx.doi.org/10.1016/0021-9290(89)90224-8)
- 236 2. McMahon TA, Cheng GC. The mechanics of running: how does stiffness couple with
237 speed? *J Biomech.* 1990;23(Suppl. 1):65-78. [http://dx.doi.org/10.1016/0021-
238 9290\(90\)90042-2](http://dx.doi.org/10.1016/0021-9290(90)90042-2)
- 239 3. Farley CT, Gonzales O. Leg stiffness and stride frequency in human running. *J Biomech.*
240 1996;29:181-186. [http://dx.doi.org/10.1016/0021-9290\(95\)00029-1](http://dx.doi.org/10.1016/0021-9290(95)00029-1)
- 241 4. Farley CT, Morgenroth DC. Leg stiffness primarily depends on ankle stiffness during
242 human hopping. *J Biomech.* 1999;32(3):267-273. [http://dx.doi.org/10.1016/S0021-
243 9290\(98\)00170-5](http://dx.doi.org/10.1016/S0021-9290(98)00170-5)
- 244 5. Hobara H, Muraoka T, Omuro K, et al. Knee stiffness is a major determinant of leg

- 245 stiffness during maximal hopping. *J Biomech.* 2009;42(11):1768-1771.
246 <http://dx.doi.org/10.1016/j.jbiomech.2009.04.047>
- 247 6. Kuitunen S, Ogiso K, Komi PV. Leg and joint stiffness in human hopping. *Scand J Med*
248 *Sci Sports.* 2011;21(6):e159-e167. doi: 10.1111/j.1600-0838.2010.01202.x
- 249 7. Farley C, Houdijk H, Van Strien C, Louie M. Mechanism of leg stiffness adjustment for
250 hopping on surfaces of different stiffnesses. *J Appl Physiol.* 1998;85(3):1044-1055.
- 251 8. Bobbert MF, Casius LJR. Spring-like leg behaviour, musculoskeletal mechanics and
252 control in maximum and submaximum height human hopping. *Philos Trans R Soc Lond B*
253 *Biol Sci.* 2011; 366(1570):1516-1529. <http://dx.doi.org/10.1098/rstb.2010.0348>
- 254 9. Oliver JL, Croix MBADS, Lloyd RS, Williams CA. Altered neuromuscular control of leg
255 stiffness following soccer-specific exercise. *Eur J Appl Physiol.* 2014;114(11):2241-2249.
256 doi: 10.1007/s00421-014-2949-z
- 257 10. Hobara H, Kanosue K, Suzuki S. Changes in muscle activity with increase in leg stiffness
258 during hopping. *Neurosci Lett.* 2007;418(1):55-59.
259 <http://dx.doi.org/10.1016/j.neulet.2007.02.064>
- 260 11. Kuitunen S, Kyröläinen H, Avela J, Komi PV. Leg stiffness modulation during exhaustive
261 stretch-shortening cycle exercise. *Scand J Med Sci Sports.* 2007;17(1):67-75. doi:
262 10.1111/j.1600-0838.2005.00506.x.
- 263 12. Oliver JL, Smith PM. Neural control of leg stiffness during hopping in boys and men. *J*
264 *Electromyogr Kinesiol.* 2010;20(5):973-979.
265 <http://dx.doi.org/10.1016/j.jelekin.2010.03.011>
- 266 13. Dalleau G, Belli A, Bourdin M et al. The spring-mass model and the energy cost of
267 treadmill running. *Eur J Appl Physiol.* 1998, 77:257-263. doi: 10.1007/s004210050330.
- 268 14. Barnes KR, Hopkins WG, McGuigan MR, Kilding AE. Warm-up with a weighted vest

269 improves running performance via leg stiffness and running economy. *J Sci Med Sport*.
270 2015;18(1):103-108. <http://dx.doi.org/10.1016/j.jsams.2013.12.005>.

271 15. Chelly SM, Denis C. Leg power and hopping stiffness: relationship with sprint running
272 performance. *Med Sci Sports Exerc*. 2001;33(2):326-33. doi: 10.1097/00005768-
273 200102000-00024.

274 16. Bret C, Rahmani A, Dufour AB, et al. Leg strength and stiffness as ability factors in 100
275 m sprint running. *J Sports Med Phys Fitness*. 2002;42(3):274-81.

276 17. Rabita G, Couturier A, Dorel S, et al. Changes in spring-mass behavior and muscle
277 activity during an exhaustive run at O₂max. *J Biomech*. 2013;46(12):2011-2017.
278 <http://dx.doi.org/10.1016/j.jbiomech.2013.06.011>

279 18. Dalleau G, Belli A, Viale F, et al. A simple method for field measurements of leg stiffness
280 in hopping. *Int J Sports Med*. 2004;25:170-176. doi: 10.1055/s-2003-45252.

281 19. Lloyd RS, Oliver JL, Hughes MG, Williams CA. Reliability and validity of field-based
282 measures of leg stiffness and reactive strength index in youths. *J Sports Sci*.
283 2009;27(14):1565-1573. doi: 10.1080/02640410903311572.

284 20. Lloyd RS, Oliver JL, Hughes MG, Williams, CA. The effect of 4-weeks of plyometric
285 training on reactive strength index and leg stiffness in male youths. *J Strength Cond Res*.
286 2012;26(10):2812-2819. doi:10.1519/JSC.0b013e318242d2ec.

287 21. Maquirriain J. The interaction between the tennis court and the player: how does surface
288 affect leg stiffness?. *Sports Biomech*. 2013;12(1):48-53.
289 doi:10.1080/14763141.2012.725088.

290 22. Choukou MA, Laffaye G, Tairar R. Reliability and validity of an accelerometric system
291 for assessing vertical jumping performance. *Biol Sport*. 2014;31(1):55-62.
292 doi:10.5604/20831862.1086733.

- 293 23. Girard O, Lattier G, Micallef JP, Millet GP. Changes in exercise characteristics, maximal
294 voluntary contraction, and explosive strength during prolonged tennis playing. *Br J Sports*
295 *Med.* 2006;40(6):521-526. doi:10.1136/bjism.2005.023754.
- 296 24. Casartelli N, Müller R, Maffiuletti NA. Validity and reliability of the Myotest
297 accelerometric system for the assessment of vertical jump height. *J Strength Cond Res.*
298 2010;24(11):3186-93. doi: 10.1519/JSC.0b013e3181d8595c.
- 299 25. Castagna C, Ganzetti M, Ditroilo M, et al. Concurrent validity of vertical jump
300 performance assessment systems. *J Strength Cond Res.* 2013;27(3):761-768. doi:
301 10.1519/JSC.0b013e31825dbcc5.
- 302 26. Glatthorn JF, Gouge S, Nussbaumer S, et al. Validity and reliability of Optojump Next
303 photoelectric cells for estimating vertical jump height. *J Strength Cond Res.*
304 2011;27(3):761-8. doi: 10.1519/JSC.0b013e3181ccb18d.
- 305 27. Bampouras TM, Relph NS, Orme D, Esformes JJ. Validity and reliability of the Myotest
306 Pro wireless accelerometer in squat jumps. *Isokinet Exerc Sci.* 2013;21:101-105. doi:
307 10.3233/IES-130484.
- 308 28. Thompson CJ, Bembien MG. Reliability and comparability of the accelerometer as a
309 measure of muscular power. *Med Sci Sports Exerc.* 1999;31(6):897-902.
310 doi: 10.1097/00005768-199906000-00020.
- 311 29. Bland JM, Altman DG. Statistical methods for assessing agreement between two methods
312 of clinical measurement. *Lancet.* 1986;1:307-310. [http://dx.doi.org/10.1016/S0140-](http://dx.doi.org/10.1016/S0140-6736(86)90837-8)
313 [6736\(86\)90837-8](http://dx.doi.org/10.1016/S0140-6736(86)90837-8).
- 314 30. Vincent WJ, Weir JP. *Statistics in kinesiology*. 4th ed. Champaign, IL: Human Kinetics;
315 2012.

- 316 31. Cohen J. *Statistical power analysis for the behavioural sciences*. 2nd ed. Mahwah, NJ:
317 Lawrence Erlbaum; 1988.
- 318 32. Sale DG. Testing strength and power. In: MacDougall JD, Wenger HA, Green HJ, eds.
319 *Physiological testing of the high performance athlete*. 2nd ed. Champaign, IL: Human
320 Kinetics; 1991: 21-106.
- 321 33. Atkinson G, Nevill AM. Statistical method for assessing measurement error (reliability)
322 in variables relevant to sports medicine. *Sports Med*. 1998;26(4):217-238. doi:
323 [10.2165/00007256-199826040-00002](https://doi.org/10.2165/00007256-199826040-00002).
- 324 34. O'Leary TJ, Morris MG, Collett J, Howells K. Reliability of single and paired-pulse
325 transcranial magnetic stimulation in the vastus lateralis muscle. [published online ahead
326 of print January 23, 2015]. *Muscle Nerve*. doi: [10.1002/mus.24584](https://doi.org/10.1002/mus.24584).
- 327 35. Hopkins WG. How to interpret changes in an athletic performance test. *Sportscience*.
328 2004;8:1-7.
- 329 36. Hopkins WG. Measures of reliability in sports medicine and science. *Sports Med*.
330 2000;30(1):1-15. doi: [10.2165/00007256-200030010-00001](https://doi.org/10.2165/00007256-200030010-00001).
- 331 37. Nunnally J, Bernstein I. *Psychometric Theory*. 3rd ed. New York, NY: McGraw Hill;
332 1993.
- 333 38. Hopkins WG. Calculating the reliability intraclass correlation coefficient and its
334 confidence limits (Excel spreadsheet). 2009.
335 <http://www.sportsci.org/resource/stats/xICC.xls>
- 336 39. Joseph CW, Bradshaw EJ, Kemp J, Clark RA. The interday reliability of ankle, knee, leg,
337 and vertical musculoskeletal stiffness during hopping and overground running. *J Appl*
338 *Biomech*. 2013;29:386-394.

- 339 40. McLachlan KA, Murphy AJ, Watsford ML, Rees S. The interday reliability of leg and
340 ankle musculotendinous stiffness measures. *J Appl Biomech.* 2006;22:296-304.
- 341 41. Komi PV, Nicol C. Stretch-shortening cycle of muscle function. In: Komi PV, ed.
342 *Neuromuscular aspects of sport performance.* Oxford, UK: Blackwell Science Ltd;
343 2011:15-31.
- 344 42. Weir JP. Quantifying test-retest reliability using the intraclass correlation coefficient and
345 the SEM. *J Strength Cond Res.* 2005;19(1):231-240.

346

Tables347 **Table1.** Leg stiffness (mean \pm SD) for Session 1 and Session 2.

		Leg Stiffness (kN/m)	
		Session 1	Session 2
K_{First}	FP	26.3 \pm 5.1	26.6 \pm 5.6
	Optojump	24.2 \pm 4.4	24.2 \pm 5.1
	Myotest	53.0 \pm 15.2	50.7 \pm 14.0
K_{Avg}	FP	26.0 \pm 5.2	26.2 \pm 5.0
	Optojump	24.1 \pm 4.6	23.9 \pm 4.4
	Myotest	52.0 \pm 14.3	50.2 \pm 12.4
K_{Max}	FP	27.6 \pm 5.6	27.6 \pm 5.9
	Optojump	25.1 \pm 4.7	24.8 \pm 5.4
	Myotest	55.0 \pm 15.1	51.8 \pm 13.6

348 *Note.* First attempt procedure (**K_{First}**); maximal value procedure (**K_{Max}**); session average value
 349 procedure (**K_{Avg}**); force platform (FP).

350 **Table 2.** Criterion-related validity statistics, compared to FP.

		r	Relative mean bias	95% LoA
K_{First}	Optojump	0.99	0.91	0.86 – 0.96
	Myotest	0.82	1.93	1.63 – 2.23
K_{Avg}	Optojump	0.99	0.92	0.86 – 0.98
	Myotest	0.86	1.92	1.64 – 2.19
K_{Max}	Optojump	0.98	0.92	0.87-0.97
	Myotest	0.81	1.93	1.67 – 2.19

351 *Note.* First attempt procedure (**K_{First}**); maximal value procedure (**K_{Max}**); session average value
 352 procedure (**K_{Avg}**); force platform (FP); Pearson’s product moment correlation coefficient (**r**);
 353 limits of agreement (LoA). All r values were statistically significant at the level of $P < .001$.

354 **Table 3.** Test-retest reliability statistics for every device

		CV ± SD (%)	SEM (kN/m)	ICC (95% CI)
K_{First}	FP	7.7 ± 7.5	2.8	0.74 (0.57 - 0.84)
	Optojump	6.6 ± 5.4	2.1	0.82 (0.70 - 0.90)
	Myotest	12.4 ± 7.0	7.6	0.74 (0.57 - 0.84)
K_{Avg}	FP	6.5 ± 7.7	2.4	0.79 (0.64 - 0.88)
	Optojump	5.9 ± 5.2	1.8	0.86 (0.74 - 0.92)
	Myotest	8.9 ± 7.1	6.3	0.79 (0.64 - 0.88)
K_{Max}	FP	7.3 ± 7.8	2.6	0.80 (0.66 - 0.88)
	Optojump	6.8 ± 6.7	2.1	0.83 (0.71 - 0.90)
	Myotest	13.0 ± 9.4	8.7	0.64 (0.44 - 0.78)

355 *Note.* First attempt procedure (**K_{First}**); maximal value procedure (**K_{Max}**); session average value
 356 procedure (**K_{Avg}**); force platform (**FP**); intraclass correlation coefficient (**ICC**); confidence
 357 intervals (**CI**); coefficient of variation (**CV**); standard deviation (**SD**); standard error of
 358 measurement (**SEM**).

359

360

Figure Captions

361 **Figure 1.** Experimental setup of the devices for synchronous data collection. Note that, custom-
362 made wooden blocks were aligned behind and ahead of the force platform.

363

364 **Figure 2.** Limits of agreement. Ratio of leg stiffness measurements outcome between either
365 Myotest (left side) or Optojump (right side) and Force platform (FP), plotted against their
366 average. The continuous line represents the mean relative bias between the examined device and
367 the FP. Dashed lines represents lower and upper limits with 95 % confidence. A) The 1st trial per
368 session was considered (K_{First}). B) The average across the three trials per session was retained
369 (K_{Avg}). C) The maximal stiffness value per session was considered (K_{Max}).