

1 **Shedding light on the Methylerythritol phosphate (MEP)-pathway: long**
 2 **hypocotyl 5 (HY5)/ phytochrome-interacting factors (PIFs) transcription**
 3 **factors modulating key limiting steps.**

4
 5 Marel Chenge-Espinosa¹, Elizabeth Cordoba¹, Cynthia Romero-Guido¹,
 6 Gabriela Toledo-Ortiz ^{2*} and Patricia León^{1*}

7
 8 ¹ Departamento de Biología Molecular de Plantas, Instituto de Biotecnología,
 9 Universidad Nacional Autónoma de México, Av. Universidad # 2001, Col.
 10 Chamilpa, Cuernavaca, Morelos, México, C.P. 62210.

11 ²Lancaster University, Lancaster Environment Centre, Lancaster LA1 4YQ, UK

12
 13 *Corresponding authors:

14 Patricia León

15 Telephone +52 (55) 5622 7856 Fax: +52 (777) 313 9988

16 email: patricia@ibt.unam.mx

17 Gabriela Toledo-Ortiz

18 Telephone: +44 (0) 1524 595053

19 Email: g.toledo-ortiz@lancaster.ac.uk

20

Total word count	6998
Summary:	219
Introduction:	949
Results:	2673
Discussion:	1338
Materials and Methods:	723
Acknowledgements:	57
Figure Legends	1036
Figures:	7 (Figs. 1 and 7 in color)
Supporting Information	7 (Fig. S1-S6; Table S1)

21

Formatted: Spanish (Spain, Traditional Sort)

Formatted: Spanish (Spain, Traditional Sort)

1 **Significance statement**

2

3

4 Light imposes a direct, rapid and potentially multi-faceted effect that leads to
5 unique protein dynamics to the main flux-limiting steps of the MEP pathway, a key
6 route essential for plants. Through differential direct transcriptional interaction, the
7 key-master integrators of light signals HY5 and PIFs, target the main flux-limiting
8 steps of the pathway. Our work illustrates how light signals can impose contrasting
9 dynamics over a key pathway whose products multi-branch downstream to all
10 chloroplastic isoprenoids.

11

1 **Summary**

2 The plastidial methylerythritol phosphate (MEP) pathway is an essential route for
3 plants as the source of precursors for all plastidial isoprenoids, many of which are
4 of medical and biotechnological importance. The MEP-pathway is highly sensitive
5 to environmental cues as many of these compounds are linked to photosynthesis
6 and growth and light is one of the main regulatory factors. However, the
7 mechanisms coordinating the MEP-pathway with light cues are not fully
8 understood.

9 Here we demonstrate that by a differential direct transcriptional modulation via the
10 key master integrators of light signal transduction HY5 and PIFs that target the
11 genes that encode the rate-controlling *DXS1*, *DXR* and *HDR* enzymes, light
12 imposes a direct, rapid and potentially multi-faceted response that leads to unique
13 protein dynamics of this pathway resulting in up to 10-fold difference in the protein
14 levels. For *DXS1*, PIF1/HY5 act as a direct activation/suppression module. In
15 contrast, *DXR* accumulation in response to light results from HY5 induction with
16 minor contribution of de-repression by PIF1. Finally, *HDR* transcription increases in
17 the light exclusively by suppression of the PIFs repression. This is an example of
18 how light signaling components can differentially multi-target the initial steps of a
19 pathway whose products branch downstream to all chloroplastic isoprenoids.
20 These findings demonstrate the diversity and flexibility of light signaling
21 components that optimize key biochemical pathways essential for plant growth.

22

23 **Keywords:**

24

25 **Isoprenoids, MEP-pathway, Light responses, DXS1, DXR and HDR**
26 **enzymes, Phytochrome interacting factors, Long Hypocotyl 5, Rate-limiting**
27 **enzymes, *Arabidopsis thaliana*.**

28

1 Introduction

2

3 Isoprenoids constitute a family of natural products synthesized in all
4 organisms with diverse function (Chappell, 1995). Isoprenoids are essential for
5 plant development, participating in several key processes such as photosynthesis,
6 respiration and general plant growth (Bouvier *et al.*, 2005). All isoprenoids are
7 produced from the condensation of two universal five-carbon precursors,
8 isopentenyl diphosphate (IPP) and its isomer dimethylallyl diphosphate (DMAPP).
9 Condensation of these basic units gives rise to isoprenoid diversity.

10

11 Plants synthesize IPP and DMAPP by two non-related pathways present in
12 different compartments. The cytoplasmic mevalonic pathway uses acetyl-CoA via
13 mevalonic acid for the synthesis of IPP and DMAPP, and the methyl-D-erythritol 4-
14 phosphate (MEP) pathway takes place in plastids (Croteau *et al.*, 2000, Hemmerlin
15 *et al.*, 2012, Rodriguez-Concepcion and Boronat, 2015). The MEP pathway uses
16 pyruvate and D-glyceraldehyde 3-phosphate (GAP) for IPP and DMAPP synthesis
17 through the activity of seven consecutive enzymes (Phillips *et al.*, 2008). The first
18 step of the MEP pathway is catalyzed by 1-deoxy-D-xylulose 5-phosphate
19 synthase (DXS) that produces 1-deoxy-D-xylulose 5-phosphate (DXP). DXP is
20 rearranged into MEP by the action of the 1-deoxy-D-xylulose 5-phosphate
21 reductoisomerase (DXR), this is the first committed step of the pathway.
22 Subsequently MEP is converted to 1-hydroxy-2-methyl-2-(E)-butenyl 4-
23 diphosphate (HMBPP) by four additional enzymatic steps and in the final step
24 HMBPP is converted into a mixture of IPP and DMAPP via the HMBPP reductase
25 (HDR) enzyme (Eisenreich *et al.*, 2004).

26

27 The MEP pathway is present in eubacteria, plastids and the apicoplast of
28 apicomplexan but is absent in other eukaryotes, including humans (Lange *et al.*,
29 2000). Thus, the MEP pathway is considered an attractive target for development
30 of new antibacterial and antiparasitic drugs, and herbicides (Rodríguez-
31 Concepción, 2004, Rohdich *et al.*, 2005). In plants the MEP pathway is responsible

1 for the production of essential compounds, such as the precursors of
2 photosynthetic pigments (carotenoids and the side chain of chlorophyll),
3 tocopherols and plastoquinones, hormones including gibberellins, abscisic acid
4 and strigolactone and a variety of monoterpenes, diterpenes and some
5 sesquiterpenes (Bouvier *et al.*, 2005, Umehara *et al.*, 2008). Recent studies have
6 reported that the MEP pathway is also essential for the production of stress-
7 specific retrograde signaling molecules (de Souza *et al.*, 2017). Thus, in plants the
8 MEP pathway is essential for plant development and also is an important target for
9 biotechnological manipulation.

10

11 The enzymes in the MEP pathway are subject to modulation at different
12 levels, according to the developmental status of the plant and the fluctuating
13 environmental conditions (Guevara-García *et al.*, 2005, Rodriguez-Concepcion,
14 2006, Cordoba *et al.*, 2009, Banerjee and Sharkey, 2014). This multi-level dynamic
15 is critical to ensure the supply of IPP and DMAPP precursors with the demand of
16 downstream pathways, many of which are dependent of diurnal and light
17 conditions, such as the biosynthesis of chlorophylls and carotenoids (Ruiz-Sola
18 and Rodriguez-Concepcion, 2012). In plants, the plastid-localized enzymes of the
19 MEP pathway come from nuclear-encoded genes. Experimental evidence has
20 demonstrated that the transcript levels of all genes in the pathway are induced
21 during plant development (Guevara-García *et al.*, 2005, Meier *et al.*, 2011), during
22 tomato fruit ripening (Lois *et al.*, 2000) and in response to hormones (Oudin *et al.*,
23 2007). Also, biotic interactions (Walter *et al.*, 2000), circadian clock (Cordoba *et al.*,
24 2009, Vranova *et al.*, 2013) and light modulate the MEP transcripts levels in
25 several plant species (Hsieh and Goodman, 2005, Cordoba *et al.*, 2009).
26 Furthermore, post-transcriptional regulation of DXS, one of the rate-limiting steps
27 of the pathway, leads to changes in its protein accumulation and enzymatic activity
28 (Guevara-García *et al.*, 2005, Wright *et al.*, 2014).

29

30 Light constitutes one of the most critical environmental signals for plant
31 development. From the emergence of the germinating seedling, light changes the

1 developmental program from skotomorphogenesis to photomorphogenesis (Chen
2 *et al.*, 2004). Upon light exposure etioplasts rapidly differentiate into chloroplasts
3 upregulating many genes involved in photosynthesis and other plastid biosynthetic
4 pathways (Jiao *et al.*, 2007). Complex photoreceptor systems allow plants to adjust
5 diverse processes in response to variable light conditions (Franklin and Quail,
6 2010). The phytochrome photoreceptors fine-tune plant photomorphogenesis in
7 response to Red and Far Red light. Mechanistically, light-activated nuclear-
8 phytochromes bind directly to members of the bHLH family of phytochrome-
9 interacting factors (PIFs), promoting their degradation. PIFs act as important
10 repressors of photomorphogenic development in the dark and key signal
11 integrators (Leivar and Quail, 2011, Leivar and Monte, 2014). In addition, active
12 phytochromes prevent the degradation of activators of photomorphogenesis such
13 as ELONGATED HYPOCOTYL 5 (HY5) and its close homolog HYH. These bZIP
14 transcription factors participate in the up-regulation of a variety of genes in
15 response to light and display antagonistic functions to the PIFs in the expression of
16 diverse genes (Tepperman *et al.*, 2001).

17
18 Given the strategic role that light has over isoprenoid production in the
19 present study we investigate the mechanisms coordinating the MEP-pathway with
20 light cues. We examined the role of light signaling components PIFs, HY5 and
21 HYH in modulating the expression and imposing a differential and specific dynamic
22 on *DXS1*, *DXR* and *HDR* genes encoding for flux-controlling enzymes of the MEP
23 pathway (Estévez *et al.*, 2001, Mahmoud and Croteau, 2001, Botella-Pavía *et al.*,
24 2004, Carretero-Paulet *et al.*, 2006). Transcript analyses and CHIP assays
25 confirmed that HY5 and PIFs are direct regulators of the light-modulated
26 expression of these genes. We show that the function of these factors in tuning the
27 MEP-pathway gene expression extends to different developmental stages
28 potentially diversifying the synthesis of multiple compounds of isoprenoid origin,
29 essential for plant growth and development at different stages of the plant life
30 cycle. Furthermore, we demonstrate using protein accumulation studies that the
31 transcriptional regulation mediated by PIF and HY5 impacts the accumulation of

1 DXS1 and HDR enzymes but not DXR. This analysis exemplifies the diverse
2 mechanistic dynamics that the same master light regulators can impose to tune up
3 essential metabolic pathways in response to light.

4

5 **Results**

6

7 ***Cis*-acting elements in the DXS promoter are responsible for the regulation of** 8 **the DXS1 gene.**

9 Previous studies have shown that in developing seedlings, transcripts of the
10 MEP pathway genes accumulate upon light exposure (Botella-Pavía *et al.*, 2004,
11 Hsieh and Goodman, 2005, Cordoba *et al.*, 2009). To analyze if the light induction
12 response is mediated at the transcriptional level, 3 day-old etiolated transgenic
13 lines containing 1510 bp upstream from the *DXS1* gene ATG fused with the GUS
14 gene (Estévez *et al.*, 2000) were exposed to light. GUS activity is detected in the
15 cotyledons of dark grown seedlings (Figure 1a). However, after 6 h of light
16 exposure this staining expands to the hypocotyl (Figure 1d). Quantitative
17 determination of GUS activity confirmed an approximately 2-fold increase after light
18 exposure in comparison to dark control seedlings (Figure 1g). This result confirms
19 that in response to light, *DXS1* transcript levels are transcriptionally up-regulated
20 and also demonstrates that the *cis*-acting elements important for this response are
21 present within the 1.5 Kb upstream regulatory region of this gene.

22

23 To further delimit the region involved in the light response two additional
24 transgenic lines containing 750 (Figure b and e) and 670 bp (Figure c and f)
25 upstream from the *DXS1* ATG of were generated, and their expression in response
26 to light was analyzed. We observed that the GUS expression in these lines is
27 induced upon light exposure at similar levels to the 1.5 Kb original fragment (Figure
28 1g). These results support that the elements responsible for light response in the
29 *DXS1* gene localize within the 670 bp region upstream from the ATG (360 from the
30 transcription initiation site).

31

1 **HY5 and HYH positively regulate transcription of MEP-pathway limiting *DXS1***
2 ***and DXR* genes, but not of *HDR*.**

3 The transcription factor HY5 plays a pivotal role as a positive regulator of
4 photomorphogenesis and greening responses through direct binding to the
5 promoters of diverse light-activated genes (Lee *et al.*, 2007, Zhang *et al.*, 2011). To
6 determine whether HY5 contributes to the light induction of the *DXS1* gene, its
7 transcript level was analyzed after illumination of dark-adapted wild-type and *hy5*
8 mutant seedlings; since a clear accumulation by light for MEP genes was observed
9 under these conditions (Cordoba *et al.*, 2009). Seedlings were grown for 3 days
10 under 16 h light: 8h dark photoperiod and then transferred to darkness for 3
11 additional days (dark-adapted treatment) prior to illumination for 6 h. Northern blot
12 analysis shows that the transcript of *DXS1* is significantly lower in the light *hy5*
13 mutant compared to the Col-0 wild-type counterpart (Figure 2a). In the *hy5hyh*
14 double mutant we observed further reduction of the *DXS1* transcript supporting the
15 notion that while minor, HYH also plays a positive role in the expression of *DXS1* in
16 response to light.

17
18 In addition to DXS, two other enzymes, DXR and HDR, limit the flux through
19 the MEP-pathway (Mahmoud and Croteau, 2001, Botella-Pavía *et al.*, 2004,
20 Carretero-Paulet *et al.*, 2006, Kim *et al.*, 2009). We also analyzed the light
21 responsiveness of *DXR* and *HDR* transcripts in the *hy5* and *hy5hyh* mutants. As
22 shown in Figure 2b the transcript of *DXR* accumulates upon light illumination in the
23 Col-0 wild-type plants. This induction is substantially lower in the *hy5* mutant,
24 demonstrating that HY5 also acts positively on the expression of the *DXR* gene in
25 the light. Relative to *hy5*, the *hy5hyh* double mutant has no additive effect on the
26 level of *DXR* transcript abundance, supporting the lack of involvement of HYH in
27 the light induced up-regulation of this gene. In contrast, the transcript accumulation
28 in response to light for *HDR* is marginal in the wild-type (Figure 2a, b) with no clear
29 role of HY5 and HYH (Figure 2c).

30

1 To further substantiate these results quantitative RT-PCR (qRT-PCR) on the
2 same RNAs samples was performed, confirming the positive role of HY5 on the
3 induction by light of *DXS1* and *DXR* transcripts (Figure 2d, e) but not on *HDR*.
4 qRT-PCR data further corroborated that HYH does not appear to play a major role
5 in the modulation by light of these genes. As monochromatic red light maximizes
6 the light response we analyzed the expression of these genes in the dark-adapted
7 seedlings illuminated with red light (Figure S1). This analysis corroborates an
8 increase of the *HDR* transcript by light and that HY5 does not play a major role in
9 this response. Red-light experiments also demonstrated that in the absence of
10 these bZIP transcription factors there is still some light responsiveness, supporting
11 the notion that additional unknown factors participate in this response.

12

13 **PIFs are negative regulators of *DXS1*, *DXR* and *HDR* genes**

14 For diverse light responses, PIF transcription factors act antagonistically to
15 HY5 and HYH (Chen *et al.*, 2013). To test whether PIFs play an opposing role to
16 HY5 in the light-mediated accumulation of the MEP pathway genes, the transcript
17 level of the *DXS1*, *DXR* and *HDR* were analyzed by northern blots in *pif1*, *pif3*, *pif4*
18 and *pif5* mutants compared to Col-0 wild-type seedlings, using dark-adapted
19 treated seedlings (Leivar *et al.*, 2008). We observed that the transcript levels of the
20 *DXS1* (Figure 3a), *DXR* (Figure 3b) and *HDR* (Figure 3c) genes are elevated in
21 several of the *pif* single mutants in the dark and in the light, supporting a function of
22 these transcription factors as transcriptional repressors of the expression of these
23 rate-limiting MEP pathway genes.

24

25 To further analyze the contribution of the different PIFs we performed qRT-
26 PCR on the same samples illuminated with white light (Figure S2) or with
27 monochromatic red light to maximize response (Figure 3). In the case of *DXS1*,
28 increased transcript accumulation is observed in the *pif1* and *pif3* mutants with a
29 more moderate increment in *pif5*, supporting the repressing function of these
30 factors (Figure 3d and S2a). For *DXR*, transcript levels are higher in the dark and
31 in light-exposed *pif1*, *pif3* and *pif5* mutants demonstrating that these three PIFs

1 contribute to the low accumulation level of this gene in wild-type plants (Figure 3e).
2 Finally, PIF1 appears to have the most significant effect on repressing *HDR*
3 transcript levels, with some contribution of PIF3 (Figure 3f). Collectively our data
4 demonstrate that the PIF factors in a partially redundant form are required to
5 maintain low expression levels of three key genes of the MEP pathway.
6

7 The functional redundancy of PIFs in regulating *DXS1*, *DXR* and *HDR*
8 genes is further illustrated with the quadruple *pifQ* mutant lacking PIF1, PIF3, PIF4
9 and PIF5 (Leivar *et al.*, 2009). Because *pifQ* could suffer from photooxidative
10 damage caused by the overaccumulation of photochlorophyllide in the dark (Chen
11 *et al.*, 2013), *pifQ* analysis was done in 3 day-old seedlings grown in (the reported?)
12 photoperiod and dark-adapted for 36 h before exposure to 6 h of red light. We
13 corroborated that in the *pifQ* mutant the expression level of *DXS1* (Figure 3g), *DXR*
14 (Figure 3h) and *HDR* (Figure 3i) genes are significantly elevated in the dark and
15 after light exposure compared to wild-type seedlings. Over all, our data
16 demonstrates that PIFs are important negative regulators of the three rate-limiting
17 MEP pathway genes.
18

19 **HY5 and PIFs regulators modulate the expression of *DXS*, *DXR* and *HDR*** 20 **genes during de-etiolation**

21 It is known that PIFs display differential expression during distinct
22 developmental responses (Jeong and Choi, 2013). One of the physiologically
23 relevant responses to light is de-etiolation. Previous data demonstrated that during
24 de-etiolation PIF1 represses *PSY* gene expression that plays a limiting role in
25 carotenoid biosynthesis, but under the conditions used (1 h of induction) no
26 regulation by PIF1 was detected for *DXS1* (Toledo-Ortiz *et al.*, 2010). Based on our
27 observations that light induces *DXS*, *DXR* and *HDR* expression at later time points,
28 we re-evaluated the contribution of PIFs and analyzed the role of HY5 by qRT-PCR
29 in 3 day-old etiolated seedlings exposed to red light for 6 h. Similar to the dark-
30 adapted seedlings, the transcription factor HY5 was shown to act as an activator of
31 the *DXS1* and *DXR* genes expression (Figure 4a and b). After light exposure the

1 level of the *DXS1* and *DXR* transcripts is lower in *hy5* compared to the wild-type.
2 There was no significant change in the *HDR* expression level in *hy5* (Figure 4c),
3 indicating that, at this stage, HY5 does not play a major role in the light response of
4 this gene .

5
6 On the other hand, removal of the PIFs results in a significant increase in
7 the transcript levels of the *DXS1*, *DXR* and *HDR* genes at this developmental stage
8 compared to the wild-type seedlings (Figure 4). This result supports and extends
9 the partially redundant, repressive role of PIFs in down-regulating the expression of
10 the *DXS1*, *DXR* and *HDR* genes during de-etiolation. Similar to the dark-adapted
11 seedlings PIF1, PIF3 and PIF5 contribute most significantly to the regulation of
12 these genes and we did not observe any major changes in the relative role of the
13 individual PIFs at this developmental stage. In accordance the *pifQ* mutant
14 accumulates at higher transcript levels than the three genes in the dark and in the
15 light.

17 **PIF1 and HY5 interact with the promoters of the MEP pathway genes *in vivo***

18 To investigate if the changes in the gene expression result from a direct
19 interaction of the PIFs and HY5 transcription factors with the promoters of the
20 *DXS1*, *DXR* and *HDR* genes, we conducted chromatin immunoprecipitation (ChIP)
21 experiments. We selected as a model PIF1, considering that this factor plays a
22 major role in the regulation of the three MEP genes. ChIPs were carried out using
23 seedlings that express PIF1 fused to a myc-tag in a *pif1* background (TAP-PIF1)
24 (Moon *et al.*, 2008) or HY5 with an HA-tag in *hy5* background (HA-HY5) (Lee *et al.*,
25 2007). Lines with a TAP-GFP or HA-GUS in a wild-type background were included
26 as negative controls for nonspecific binding of DNA to the tags used. Transgenic
27 HA-HY5 and TAP-PIF1 are mild-overexpressors, with similar protein expression
28 levels to endogenous HY5 and PIF1 and complement the mutant phenotypes in a
29 wide range of tested-light responses (Lee *et al.*, 2007, Moon *et al.*, 2008).

30

Commented [HG1]: Wasn't sure about this when looking at the plots. Certainly for *DXS1* and *DXR* in the light this is true and also for *DXS1* in the dark but *pif1* looks to be greater than *pifQ* in *DXR* and for *HDR* *pif3* in the dark and *pif1* in the light. I could be missing something though.

1 To address if there is any difference in the interaction of these factors
2 depending on the developmental stage, the ChIP was conducted in dark-adapted
3 seedlings and etiolated seedlings, both exposed to 6 h of red light using myc or HA
4 antibodies. qRT-PCRs were done using specific primers for selected promoter
5 regions of each gene (Table S1, Figures S3, S5 and S6).

6
7 PIFs transcription factors bind to variants of the E boxes (CANNTG),
8 including the G-box (CACGTG) and the PBE-box (CACATG/CATGTG), or to
9 hexameric sequences G-box coupling elements (GCEs) containing the core
10 "ACGT" elements (Toledo-Ortiz *et al.*, 2003, Zhang *et al.*, 2013, Kim *et al.*, 2016).
11 The G-box, together with diverse GCE related motifs, such as the Z box
12 (ATACTGTGT), CA (GACGTA) and CG hybrids (GACGTG), have also been
13 identified as the interaction site of HY5 (Lee *et al.*, 2007, Toledo-Ortiz *et al.*, 2010,
14 Zhang *et al.*, 2011). The analysis of the 1.5 kb *DXS1* promoter sequence using
15 SOGO New PLACE software, showed no-presence of canonical G boxes.
16 However, several E-box related elements, including a PBE-box (CACATG), a CG
17 hybrid box (GACGTG), and a GCE element were found (Figure S3). Four specific
18 oligonucleotide pairs were used to amplify the DNA enriched from the precipitated
19 Protein-DNA complexes from the TAP-PIF1 and HA-HY5 transgenic lines
20 maintained in the dark or exposed to 6 h red light. The qPCR using the
21 oligonucleotide pairs 1 and 2 (Figure S3) showed no significant recovery in
22 comparison to the negative TAP-GFP and HA-GUS controls, indicating no binding
23 of PIF1 or HY5 to these promoter fragments. In contrast, enriched DNA sequences
24 were amplified from the TAP-PIF1 and HA-HY5 immunoprecipitated fractions with
25 the oligonucleotide pair P3 (F3/R3) (Figure S3). PIF1 binding was significant in the
26 extracts from dark grown seedlings and a minor enrichment was seen in the light
27 extracts (Figure 5a). For the HA-HY5 immunoprecipitate we only observed specific
28 amplification in light-exposed seedlings (Figure 5b). For the P4 (F4/R4) primers
29 pair (Figure S3) a minor amplification was detected from both the TAP-PIF1 and
30 HA-HY5 extracts in comparison to the controls, indicating a very weak, and
31 probably not-significant (just slightly above the in-specific GUS-HA background)

1 binding of both factors to this fragment (Figure 5c, d). The fragment that showed
2 preferential binding to PIF1 and HY5 contains a PBE box, a GCE element and CG
3 hybrid box in close proximity to each other (Figure S3). For PIF1, the PBE box
4 (CACATG) is the most likely candidate binding-element and the GCE element
5 (ACGT) for HY5. Very similar results were obtained from the ChIP experiments in
6 de-etiolated seedlings demonstrating that the same sites are involved in the PIF1
7 and HY5 binding in these two light developmental stages (Figure S4).

8

9 We also analyzed the promoter sequences of *DXR* and *HDR* genes for
10 potential PIF1 and HY5 binding sites. Within 1300 pb of the upstream sequence of
11 *DXR* several G box-related sequences were found (Figure S5). Three pairs of
12 primers covering the different elements from the *DXR* promoter were used in ChIP
13 experiments (Table S1 and Figure S5). As shown in Fig. 5f the only significant
14 enrichment detected was for HA-HY5 in the light with the primer pair 1 (F1 and R1)
15 containing two GCE box-related sequences. These data demonstrate that *DXR* is
16 directly upregulated by HY5. In contrast, binding of TAP-PIF1 in the dark was
17 minor, although within statistical significance with the two primer pairs and close
18 with the primer pair 3 (p 0.055) (Figure 5e, g and i). In these fragments, several
19 putative G-box and E-box related sequences are present (Fig. S5). However,
20 considering that the binding of PIF1 to the three sites is so close to the negative
21 control in both photoperiodic and de-etiolation experiments (Figure S4e, g, i), this
22 result probably indicates a non-preferential interaction of PIF1 with the promoter of
23 *DXR*.

24

25 Finally, the *HDR* gene promoter includes 902 bp with only one PBE box and
26 a GCE/ACE motif (Figure S6). Two pairs of primers were designed to cover the
27 potential PIF1/HY5 binding sites and used to amplify the immunoprecipitates from
28 the ChIPs (Table S1). In contrast to the *DXS1* and *DXR* genes, no enrichment was
29 detected with HA-HY5 (Figure 5l), demonstrating that HY5 does not bind to the
30 upstream sequences of the *HDR* gene. For TAP-PIF1 significant binding was
31 observed in dark samples with the primer pair P2 that includes a PBE-box (Figure

1 5k). ChIPs results in de-etiolated seedlings (Figure S4k and l) tightly correlate with
2 those of the photoperiodic-dark-adapted seedlings (Figure 5k, l). In conclusion,
3 these data support the notion that modulation by light of *HDR* transcript levels
4 results from its de-repression from the dark activity of PIF1/PIF3 and not from
5 activation in light by HY5. Whether there is another light regulated activation factor,
6 remains to be investigated. Over all, our ChIP studies likely reflect light imposed
7 changes in the PIFs/HY5 regulators dynamic behavior and differential promoter-
8 binding capacity in response to the light environment.

9

10 **Impact of transcriptional imposed regulation over light modulated levels of** 11 **MEP proteins**

12 The MEP-pathway provides with the intermediaries for the production of
13 multiple compounds essential for photomorphogenic development, including
14 multiple hormones (GA, Cytokinin, ABA, Strigolactones) and photopigments
15 (carotenoids and chlorophylls).

16 A previous study demonstrated that carotenoids and chlorophylls
17 accumulate upon light exposure and this accumulation is affected in the *pifQ* and
18 *hy5* mutants (Toledo-Ortiz *et al.*, 2014). Since the synthesis of these metabolites
19 depends directly on the MEP pathway, the transcriptional regulation of the MEP-
20 transcripts by light can impact the accumulation of the final pathway products. To
21 evaluate the importance that light imposed transcriptional regulation of the *DXS1*,
22 *DXR* and *HDR* transcripts has over the pathway, the levels of the corresponding
23 proteins were analyzed in wild-type, *pifs* and *hy5* mutants. Total protein extracts
24 from dark-adapted seedlings during 3 d (Wt, *pif1*, *pif3*, *pif5* and *hy5*) or 36 h (*pifQ*)
25 were obtained and the level of the *DXS1*, *DXR* and *HDR* proteins compared to
26 dark controls.

27

28 We found that the transcriptional regulation mediated by the HY5/PIF1
29 module results in significant changes in the *DXS1* enzyme level. Compared to the
30 wild-type, the *hy5* mutant has lower accumulation of *DXS1* protein whereas *pif*
31 single mutants and *pifQ* contain up to >10 times higher protein content than wild-

1 type in the dark (Figure 6B). It is worth noticing that although the DXS1 level in the
2 light is higher in the *pifQ* mutant, it does not maintain the same difference observed
3 in the dark (Figure 6b). We hypothesize that this might be the result of post-
4 transcriptional regulation that keeps this protein within certain levels (Flores-Perez
5 *et al.*, 2008). This response correlates well with the transcript trends observed in
6 *pifs*. The accumulation of DXS resembles the model of de-repression by PIFs in
7 the dark /activation by HY5 in the light.

8

9 We also observed accumulation of HDR protein in *pif3* and *pifQ* with a
10 contributed additive effect of primarily PIF3 and PIF1. We did not detect protein
11 differences in *hy5* vs wild-type. The role of PIFs in de-repressing from the dark
12 leads to high protein levels in the light, but no further light-induced upregulation
13 was detected at the time point assessed.

14

15 No major differences were detected in the level of DXR protein in the
16 different mutants analyzed (Figure 6), in agreement with the more moderate
17 transcript differences observed for the *pifs* and *hy5*.

18

19 Discussion

20

21 The MEP pathway is a key biosynthetic route responsible for the synthesis
22 of essential compounds and signals that modulate developmental and stress
23 responses (Bouvier *et al.*, 2005, Umehara *et al.*, 2008, Hemmerlin *et al.*, 2012,
24 Walley *et al.*, 2015, Benn *et al.*, 2016). Also, several of the MEP pathway products
25 have importance for human health and nutrition. Addressing how such a central
26 metabolic pathway is modulated by external and internal cues has big implications
27 for future efforts to regulate its outputs including many compounds of
28 biotechnological or pharmaceutical interest

29

30 Light is one of the most relevant signals that affect plant metabolism,
31 including the production of essential photopigments, growth regulators and stress

1 hormones derived from the MEP-pathway (von Lintig *et al.*, 1997, Rodríguez-
2 Concepción *et al.*, 2004). Previous work on photopigment production demonstrated
3 that light-responsive transcriptional factors HY5 and PIF1 control central carotenoid
4 and chlorophyll biosynthetic genes (Huq *et al.*, 2004, Toledo-Ortiz *et al.*, 2010). In
5 the case of phytoene synthase PIF1 and HY5 act as a module that antagonistically
6 balance the expression of this key carotenogenic gene (Toledo-Ortiz *et al.*, 2014).
7 Our data demonstrate that via the same master-modulators, an additional higher-
8 order layer arises via the light-induced tune up of the MEP-pathway for the
9 coordinated production of the precursors used for multiple compounds involved in
10 photomorphogenesis. Such is the case of carotenoids and the phytol side chain of
11 the chlorophylls.

12

13 The gene products of the MEP pathway accumulate upon light-exposure
14 (Botella-Pavía *et al.*, 2004, Hsieh and Goodman, 2005, Cordoba *et al.*, 2009),
15 however the molecular mechanisms for this upregulation and its impact on the
16 pathway are still not fully understood. The coordinated tuning by light of the MEP
17 pathway genes leads to the possibility that common mechanisms regulate the
18 expression of these genes in response to light. Here we show that the master
19 regulators of light signals transduction HY5 and PIFs directly interact with the up-
20 stream elements of the flux-controlling *DXS1*, *DXR* and *HDR* genes (Estévez *et al.*,
21 2001, Botella-Pavía *et al.*, 2004, Carretero-Paulet *et al.*, 2006) and fine-tune their
22 expression levels in response to light. This regulation involves the interplay and
23 differential contribution of each factor for each gene, leading to unique
24 transcriptional dynamics (Figure 7) that lead to changes in protein accumulation.
25 Considering the flux-controlling capacity of DXS, DXR and HDR (Estévez *et al.*,
26 2001, Botella-Pavía *et al.*, 2004, Enfissi *et al.*, 2005, Carretero-Paulet *et al.*, 2006,
27 Banerjee *et al.*, 2013, Ghirardo *et al.*, 2014), the dynamics observed could
28 importantly modify the flux through the pathway in response to the prevailing light
29 environment.

30 Our findings on the role of PIFs contrast to previous reports that concluded
31 that *DXS1* expression was not regulated by PIF1 (Toledo-Ortiz *et al.*, 2010). This

1 discrepancy probably results from differences in the quantification of transcript
2 levels through a lower-sensitivity microarray analysis compared to qRT-PCR and
3 the kinetics of light induction at earlier or later time points, as all these genes
4 present light/dark oscillatory patterns (Cordoba *et al.*, 2009). In the case of PIF5,
5 using overexpressing PIF5 cell lines it was concluded that this factor was the main
6 positive regulator of all the MEP pathway genes without major participation for
7 PIF1 or PIF3 (Mannen *et al.*, 2014). However, in our study we did not observe
8 major differences of PIF1, PIF3 and PIF5 contribution as negative regulators of the
9 MEP pathway genes during de-etiolation and dark-adaptation. It is possible that
10 these differences result from a dominant negative effect of PIF5-over-expression.
11 Also, we cannot exclude that PIF5 activity could change in other developmental or
12 environmental conditions as a result of interaction with other elements (Mannen *et*
13 *al.*, 2014).

14 Our studies demonstrate that the same key master integrators of light signal
15 signals have the capacity to coordinate the core of the MEP pathway and its
16 multiple outputs through particular mechanisms for each gene (Figure 7). *DXS1*
17 expression is repressed in the dark by the direct binding of PIF1, probably through
18 the PBE box (CACATG), whereas HY5 induces the expression of this gene in the
19 light through direct binding to the nearby GCE element (ACGT). Despite the use of
20 mild-over-expressors of PIF1 or HY5 for ChIP assays, the plant material likely
21 resembles the behavior of the endogenous proteins, as these transgenic lines
22 express comparable protein levels to the endogenous proteins, complement the
23 *pif1* and *hy5* mutant phenotypes and maintain the dark/light dynamics imposed on
24 these proteins by phytochromes and COP1, among others. Since PIFs/HY5
25 targeted *cis*-acting elements are located very close to one another, it is likely that
26 the binding of one regulator results in allosteric interference of binding the second
27 regulator, resulting in an antagonistic mechanism. This regulation resembles the
28 one described for the phytoene synthase (*PSY*) gene (Toledo-Ortiz *et al.*, 2010,
29 Toledo-Ortiz *et al.*, 2014).

30

1 In contrast, the accumulation of *HDR* in the light results exclusively from the
2 degradation of the PIF repressors upon light exposure, without contribution from
3 HY5. PIF1 directly interacts with *cis*-elements in the regulatory region of *HDR*
4 (Figure 7). The strongest candidate as binding-element is a PBE motif present 156
5 bp upstream of the ATG.

6
7 Finally, the expression of *DXR* accumulates in the light as a consequence of the
8 direct interaction of HY5 to elements located around 1.2 Kb upstream of the ATG
9 of this gene (Figure 7). In contrast, PIF1 has a weak interaction in more than one
10 region of the HDR promoter (Figure 7). The weak interaction observed with PIF1
11 could reflect that its binding capacity might depend on other PIFs. In agreement
12 with this possibility our analyses demonstrate that in addition to PIF1, PIF3 and
13 PIF5 also affect the expression level of the *DXS1*, *DXR* and *HDR* genes.
14 Differential affinity of various PIFs for the promoters of MEP-pathway genes could
15 result in modified kinetics of light-responsiveness and modulation by other
16 environmental cues such as photoperiodism and the circadian clock.

17
18 Other studies have demonstrated high affinity of PIF1 and HY5 for G-box elements
19 *in vitro* and *in vivo* (Toledo-Ortiz *et al.*, 2003, Huq *et al.*, 2004, Oh *et al.*, 2009,
20 Zhang *et al.*, 2013). None of the MEP pathway genes analyzed contain canonical
21 G-boxes and our data support that for MEP-genes, the PBE box (CACATG) and
22 the GCE element (ACGT) are the strongest candidates to bind PIF1 for HY5
23 respectively. This differs and extends from the signal integration module
24 established by G-boxes in the case of photopigment related genes. Our analysis
25 from the *DXS1::GUS* transgenic lines carrying different promoter regions of the
26 *DXS1* also supports our ChIP data, positioning the important *cis*-acting elements
27 within 300 bp from the transcription initiation site. It is worth to mention that with the
28 *DXS1::GUS* transgenic constructs although we observed the same response trend,
29 only a 2-fold increase between dark and light conditions was detected with high
30 GUS activity in the dark. This is in contrast to the low *DXS1* transcript levels found
31 in the dark and its increase in response to light detected in our quantification

1 analyzes. This apparent discrepancy probably results from the accumulation of the
2 GUS protein in the dark in the transgenic lines, as the stability of this protein is
3 known to extend for more than 3 days (Kavita and Burma, 2008). Thus, in the case
4 of *DXS1*, GUS is a good marker to identify regulatory *cis* acting elements but not
5 for quantitative analyzes of its expression.

6
7 Hence together, our results indicate that the transcriptional regulation of
8 MEP-pathway genes by PIFs-HY5 results in a unique dynamic behavior for each
9 gene, providing additional flexibility to integrate inputs perceived by these master
10 regulators, such as time keeping or temperature signals. Interestingly, the HY5 and
11 PIF binding motives in *DXS1* gene localize proximal to a potential CCA1 binding
12 site (AAAATCT). CCA1 encodes a MYB-related protein that binds to *Lhcb1*3* and
13 that participates in the phytochrome regulation of this gene (Wang *et al.*, 1997).
14 This factor is also an important component of circadian regulation in coordination
15 of HY5 (Nagel *et al.*, 2015). Since *DXS1*, as well as other genes of the MEP
16 pathway are regulated by the circadian clock (Cordoba *et al.*, 2009), it is possible
17 that the enhanced regulatory dynamic imposed by a PIF/HY5 co-acting module
18 would also bring unique capacity for the light and circadian regulation of *DXS1* via
19 interaction with circadian components such as CCA1.

20
21 Importantly, we provide evidence that the differences in the transcript de-
22 re-pression/activation by light mediated by PIFs/HY5 reflects on changes at the
23 protein level leading to unique protein dynamics. *DXS1* follows a good
24 correspondence with the transcriptional fluctuations including low protein level in
25 the dark and accumulation in the light in a HY5/PIFs dependent manner, resulting
26 in a significant difference in protein levels. These rapid changes in the protein
27 accumulation in response to light supports the possibility of a rapid turnover for this
28 protein and are consistent to its central role as a major flux controller of the
29 pathway in diverse environmental and developmental conditions (Estévez *et al.*,
30 2001, Enfissi *et al.*, 2005, Banerjee *et al.*, 2013, Ghirardo *et al.*, 2014, Wright *et al.*,
31 2014). However, we observed some discrepancies between the levels of the

1 transcript and their corresponding protein such as in the *pifQ* mutant, where only a
2 1.5-fold increase after light exposure is detected. This result probably reflects post-
3 translational regulatory events over the DXS1 that adjust the level of this protein in
4 response to the product demand, as has been previously reported (Pulido *et al.*,
5 2013, Pokhilko *et al.*, 2015, Pulido *et al.*, 2016). Thus, while transcription regulation
6 of *DXS1* plays an important role in control the levels of this enzyme post-
7 translational regulatory events also act as an additional layer of regulation that
8 feedbacks metabolic requirements and impact upon the overall accumulation of
9 this protein.

10 On the other hand, HDR protein levels also reflect on the transcriptional regulation
11 mediated by light, following a different dynamic that results from the de-repression
12 of transcript levels in the dark with no further changes associated with the light.
13 The accumulation of HDR in response to light might be important to fulfill the
14 synthesis requirement of photopigments in coordination to DXS1. This result is
15 consistent with the co-limiting role previously observed for HDR during carotenoid
16 synthesis in dark/light transition and during fruit ripening (Botella-Pavía *et al.*, 2004,
17 Kim *et al.*, 2009). This type of regulation could be particularly important in
18 conditions where different levels of PIFs accumulate, such as photoperiodic
19 conditions, and at the same time could limit the acute responses to the light signal.
20 Finally, changes in the DXR protein accumulation in response to light was reflected
21 in constant protein levels in the light within the time frame analyzed. This result is
22 intriguing and it may reflect on particularities of the half-life of this protein or post-
23 transcriptional events that control protein abundance. Future analyses in this
24 respect are important to address these possibilities.

25

26 Overall, our data support a model where a differential contribution of the
27 master light regulators PIFs and HY5 to light-modulated transcriptional effects
28 reflects in protein changes in MEP-pathway flux-controlling enzymes DXS1 and
29 HDR. The significance of this differential regulation may impact on multiple
30 downstream pathways such as chlorophyll and carotenoid biosynthesis as well as
31 hormone and secondary metabolite synthesis, maintaining a very sensitive

1 responsiveness to the prevailing external conditions. Our studies exemplify how
2 differential multi-targeting of the initial steps of a pathway whose products multi-
3 branch downstream could impose a fine and unique modulation of all chloroplastic
4 isoprenoids.

5

6 **Methods**

7

8 **Plant Material and Growth Conditions.**

9 The *Arabidopsis thaliana* lines used in this work are in Columbia (Col-0)
10 background. Seeds from *pif1-2*, *pif3-3*, *pif4-2*, *pif5-2*, and *pifQ* were kindly provided
11 by P. Quail (University of California Berkeley). Seeds were grown on 1X Murashige
12 and Skoog (MS) media with Gamborg vitamins (Phytotechnology Laboratories,
13 Shawnee Mission, KS) supplemented with 1% (w/v) sucrose and 0.8% (w/v)
14 phytoagar and stratified at 4°C for 4 days. For the light gene expression analysis
15 two treatments were used. For the dark-adapted treatment, seedlings were grown
16 for 3 days in a 16 h light: 8h dark photoperiod at 120 $\mu\text{mol m}^2 \text{sec}^{-1}$, followed by 3
17 day dark adaptation for the *hy5*, *hy5hy5* and *pif* single mutants or 36 h for the *pifQ*.
18 Light treatment was done using 6 h with (100 $\mu\text{mol m}^2 \text{sec}^{-1}$) cool white (Philips
19 F25T8/TL841) or (40 $\mu\text{mol m}^2 \text{sec}^{-1}$) of red (Phillips LED module HF Deep Red
20 177354) lights. For de-etiolation experiments seedlings were exposed to 3 h of
21 white light, transferred to darkness for 3 days and exposed to 6 h of red light (40
22 $\mu\text{mol m}^2 \text{sec}^{-1}$). Control seedlings were maintained in darkness. Growth
23 temperature was maintained between 21- 22°C in all cases.

24

25 **Expression analysis**

26 Total RNA was isolated using TRIzol (Invitrogen, Carlsbad, CA, USA)
27 according to the protocol provided by the manufacturer. For northern-blot analysis,
28 5 μg of total RNA was fractionated and transferred onto a Hybond-N⁺ nylon
29 membrane (GE, Buckinghamshire, UK). Hybridizations and washes were
30 performed under stringent conditions. Probes were ³²P-radiolabeled using the
31 Megaprime DNA labeling system (GE, Buckinghamshire, UK). All probes were

1 obtained by PCR amplification as previously reported (Guevara-García *et al.*,
2 2005). For qRT-PCR experiments seedlings were harvested in the dark for the
3 dark samples and RNA extraction conducted using the RNeasy Plant Mini Kit
4 (Qiagen) or TRIzol. Complementary DNA (cDNA) was obtained from DNase-
5 treated RNA with M-MLV Reverse Transcriptase kit (Invitrogen, Carlsbad, CA). The
6 qRT-PCR experiments were performed using FastStart DNA Master^{PLUS} SYBR
7 Green I (Roche) on an Agilent Technologies Stratagene MX3005P or a Light
8 Cycler 480 Roche. Analyses were done with three independent experiments and
9 technical duplicates were included in each case (n=2). The reference gene used in
10 the qPCR analyses is *ACT7* since the expression of this reference gene has been
11 shown to not have major fluctuations in the conditions analyzed.

12

13 **DXS1 promoter analysis.**

14 From the transcriptional *DXS1::GUS* clone (Estévez *et al.*, 2000), two
15 additional clones were generated containing 750 bp and 660 bp deletions from the
16 ATG of *DXS1*. The fragments were subcloned into the pBin19 binary vector and
17 used to generate transgenic lines through *Agrobacterium tumefaciens*-mediated
18 transformation into the Col-0 ecotype (Clough and Bent, 1998). At least three
19 independent homozygous lines were selected for each construct and analyzed.

20

21 **GUS histochemical and fluorimetric analyses**

22 Three day-old etiolated seedlings exposed to light or dark for 6 h were
23 stained using the GUS histochemical assay (Jefferson *et al.*, 1987). Plants were
24 clarified as reported (Malamy and Benfey, 1997) and visualized using a
25 stereoscopic microscope (Nikon SMZ1500). For the fluorometric analysis the
26 seedlings were homogenized in GUS extraction buffer (50 mM NaHPO₄, pH 7.0; 10
27 mM Na₂EDTA; 1% Triton X-100; 0.1% N-lauroyl sarcosine and 10 mM β-
28 mercaptoetanol). The enzymatic reaction was done using 5 µl of the extracts.
29 Fluorometric quantification was done with TKO 100 fluorimeter (Hoeffer). Specific
30 activity was determined as nmol of methyl-umbelliferone per µg protein⁻¹ per min⁻¹.

31

1 **Protein gel blot analysis**

2 Total protein was obtained from seedlings and 20 µg of the samples was
3 separated by SDS-polyacrylamide gel electrophoresis (SDS-PAGE). The proteins
4 were transferred onto nitrocellulose membrane (Amersham Protan Premium 0.45
5 µm NC GE Healthcare Life Science). To verify equal loading, a parallel gel was run
6 and stained with Coomassie blue. Immunodetection was performed as previously
7 reported (Guevara-García *et al.*, 2005). Detection was done using the Luminata
8 Crescendo Western HRP Substrate (Millipore, USA). Bands from three
9 independent experiments were quantified by densitometric analysis using ImageJ
10 software (Schneider *et al.*, 2012).

11

12 **Chromatin Immunoprecipitation assays**

13 ChIP assays were conducted following the protocol reported previously
14 (Moon *et al.*, 2008) except that in our assays, 2 week old- seedlings were used.
15 Plants were grown as described in (Toledo-Ortiz *et al.*, 2014), and dark adapted for
16 72h before light treatments and sample collection. Samples were collected for dark
17 time points (0h, before lights on at the end of dark adaptation) or after 6 h
18 illumination with red light (40 µmol m⁻² s⁻¹). Plant material used (35S::HA-HY5 in
19 *hy5-215* and 35S::TAP-PIF1 in *pif1-2*) was previously described (Lee *et al.*, 2007,
20 Moon *et al.*, 2008) and kindly provided by the Deng Lab (Yale) and Huq Lab (UT
21 Austin). Both lines are mild-over expressors that complement the mutant
22 phenotypes and show comparative levels and light responses to native *PIF1* and
23 *HY5*. qRT-PCR was conducted on a Roche 480 Light cycler according to standard
24 protocol by the manufacturer. The oligonucleotides sequences used to amplify
25 upstream promoter regions of individual genes are shown in Table S1. Upstream
26 sequences of the *DXS1*, *DXR* and *HDR* genes were analyzed for possible light
27 responsive elements using SOGO New PLACE software
28 (<https://sogo.dna.affrc.go.jp/cgi-bin/sogo.cgi?sid=&lang=en&pj=640&action=page&page=newplace>)
29 and are shown in Figures S2, S3 and S4 (Higo *et al.*, 1999).

30

31

1 **Acknowledgements**

2

3 We thank Dr. Nidia Sánchez and Dr. Gabriela Perez for technical assistance and
4 Geoff Holrooyd and Phoebe Sutton for proof-reading the manuscript. This research
5 was supported by grants from CONACYT (CB 220534 and FC 2016-96) and
6 DGAPA-UNAM (IN204617) for PL and by Royal Society Grant (RG150711) to
7 GTO. MC and CR were supported by PhD fellowship from CONACYT and DGAPA.
8 The authors declare no conflict of interest.

9

10 **Bibliography**

11

12 **Banerjee, A. and Sharkey, T.D.** (2014) Methylerythritol 4-phosphate (MEP) pathway
13 metabolic regulation. *Nat Prod Rep*, **31**, 1043-1055.

14 **Banerjee, A., Wu, Y., Banerjee, R., Li, Y., Yan, H. and Sharkey, T.D.** (2013) Feedback
15 inhibition of deoxy-D-xylulose-5-phosphate synthase regulates the
16 methylerythritol 4-phosphate pathway. *J Biol Chem*, **288**, 16926-16936.

17 **Benn, G., Bjornson, M., Ke, H., De Souza, A., Balmond, E.L., Shaw, J.T. and Dehesh,**
18 **K.** (2016) Plastidial metabolite MEcPP induces a transcriptionally centered
19 stress-response hub via the transcription factor CAMTA3. *Proc Natl Acad Sci U*
20 *S A*, **113**, 8855-8860.

21 **Botella-Pavía, P., Besumbes, O., Phillips, M.A., Carretero-Paulet, L., Boronat, A.**
22 **and Rodríguez-Concepción, M.** (2004) Regulation of carotenoid biosynthesis
23 in plants: evidence for a key role of hydroxymethylbutenyl diphosphate
24 reductase in controlling the supply of plastidial isoprenoid precursors. *Plant J*,
25 **40**, 188-199.

26 **Bouvier, F., Rahier, A. and Camara, B.** (2005) Biogenesis, molecular regulation and
27 function of plant isoprenoids. *Progress in lipid research*, **44**, 357-429.

28 **Carretero-Paulet, L., Cairó, A., Botella-Pavía, P., Besumbes, O., Campos, N.,**
29 **Boronat, A. and Rodríguez-Concepción, M.** (2006) Enhanced flux through
30 the methylerythritol 4-phosphate pathway in *Arabidopsis* plants
31 overexpressing deoxyxylulose 5-phosphate reductoisomerase. *Plant Mol Biol*,
32 **62**, 683-695.

33 **Chappell, J.** (1995) Biochemistry and molecular biology of the isoprenoid pathway in
34 plants. *Ann. Rev. Plant Mol. Biol.*, **46**, 521-547.

35 **Chen, D., Xu, G., Tang, W., Jing, Y., Ji, Q., Fei, Z. and Lin, R.** (2013) Antagonistic basic
36 helix-loop-helix/bZIP transcription factors form transcriptional modules that
37 integrate light and reactive oxygen species signaling in *Arabidopsis*. *Plant Cell*,
38 **25**, 1657-1673.

39 **Chen, M., Chory, J. and Fankhauser, C.** (2004) Light signal transduction in higher
40 plants. *Annual review of genetics*, **38**, 87-117.

- 1 **Clough, S.J. and Bent, A.F.** (1998) Floral dip: a simplified method for Agrobacterium-
2 mediated transformation of *Arabidopsis thaliana*. *Plant J*, **16**, 735-743.
- 3 **Cordoba, E., Salmi, M. and León, P.** (2009) Unravelling the regulatory mechanisms
4 that modulate the MEP pathway in higher plants. *J Exp Bot*, **60**, 2933-2943.
- 5 **Croteau, R., Kutchan, T.M. and Lewis, N.G.** (2000) Natural products (secondary
6 metabolites). In *Biochemistry and Molecular Biology of Plants* (Buchanan, B.B.,
7 Grisse, W. and Jones, R.L. eds). Rockville, MD: American Society of Plant
8 Biology, pp. 1250-1268.
- 9 **de Souza, A., Wang, J.Z. and Dehesh, K.** (2017) Retrograde Signals: Integrators of
10 Interorganellar Communication and Orchestrators of Plant Development. *Annu*
11 *Rev Plant Biol*, **68**, 85-108.
- 12 **Eisenreich, W., Bacher, A., Arigoni, D. and Rohdich, F.** (2004) Biosynthesis of
13 isoprenoids via the non-mevalonate pathway. *Cell Mol Life Sci*, **61**, 1401-1426.
- 14 **Enfissi, E.M.A., Fraser, P.D., Lois, L.M., Boronat, A., Schuch, W. and Bramley, P.M.**
15 (2005) Metabolic engineering of the mevalonate and non-mevalonate
16 isopentenyl diphosphate-forming pathways for the production of health-
17 promoting isoprenoids in tomato. *Plant Biotech J*, **3**, 17-27.
- 18 **Estévez, J.M., Cantero, A., Reindl, A., Reichler, S. and León, P.** (2001) 1-Deoxy-D-
19 xylulose-5-phosphate synthase, a limiting enzyme for plastidic isoprenoid
20 biosynthesis in plants. *J Biol Chem*, **276**, 22901-22909.
- 21 **Estévez, J.M., Cantero, A., Romero, C., Kawaide, H., Jiménez, L.F., Kuzuyama, T.,**
22 **Seto, H., Kamiya, Y. and León, P.** (2000) Analysis of the expression of *CLA1*, a
23 gene that encodes the 1-deoxyxylulose 5-phosphate synthase of the 2-C-
24 methyl-D-erythritol-4-phosphate pathway in *Arabidopsis*. *Plant Physiol*, **124**,
25 95-103.
- 26 **Flores-Perez, U., Sauret-Güeto, S., Gas, E., Jarvis, P. and Rodríguez-Concepción, M.**
27 (2008) A mutant impaired in the production of plastome-encoded proteins
28 uncovers a mechanism for the homeostasis of isoprenoid biosynthetic enzymes
29 in *Arabidopsis* plastids. *Plant Cell*, **20**, 1303-1315.
- 30 **Franklin, K.A. and Quail, P.H.** (2010) Phytochrome functions in *Arabidopsis*
31 development. *J Exp Bot*, **61**, 11-24.
- 32 **Ghirardo, A., Wright, L.P., Bi, Z., Rosenkranz, M., Pulido, P., Rodríguez-**
33 **Concepcion, M., Niinemets, U., Bruggemann, N., Gershenzon, J. and**
34 **Schnitzler, J.P.** (2014) Metabolic flux analysis of plastidic isoprenoid
35 biosynthesis in poplar leaves emitting and nonemitting isoprene. *Plant Physiol*,
36 **165**, 37-51.
- 37 **Guevara-García, A.A., San Roman, C., Arroyo, A., Cortés, M.E., Gutiérrez-Nava, M.L.**
38 **and León, P.** (2005) The characterization of the *Arabidopsis clb6* mutant
39 illustrates the importance of post-transcriptional regulation of the Methyl-D-
40 Erythritol 4-Phosphate Pathway. *Plant Cell*, **17**, 628-643.
- 41 **Hemmerlin, A., Harwood, J.L. and Bach, T.J.** (2012) A raison d'être for two distinct
42 pathways in the early steps of plant isoprenoid biosynthesis? *Progress in lipid*
43 *research*, **51**, 95-148.
- 44 **Higo, K., Ugawa, Y., Iwamoto, M. and Korenaga, T.** (1999) Plant cis-acting
45 regulatory DNA elements (PLACE) database: 1999. *Nucleic Acids Res*, **27**, 297-
46 300.

- 1 **Hsieh, M.H. and Goodman, H.M.** (2005) The Arabidopsis *IspH* homolog is involved in
2 the plastid nonmevalonate pathway of isoprenoid biosynthesis. *Plant Physiol*,
3 **138**, 641-653.
- 4 **Huq, E., Al-Sady, B., Hudson, M., Kim, C., Apel, K. and Quail, P.H.** (2004)
5 Phytochrome-interacting factor 1 is a critical bHLH regulator of chlorophyll
6 biosynthesis. *Science*, **305**, 1937-1941.
- 7 **Jefferson, R.A., Kavanagh, T.A. and Bevan, M.W.** (1987) GUS fusions: beta-
8 glucuronidase as a sensitive and versatile gene fusion marker in higher plants.
9 *Embo J*, **6**, 3901-3907.
- 10 **Jeong, J. and Choi, G.** (2013) Phytochrome-interacting factors have both shared and
11 distinct biological roles. *Mol Cells*, **35**, 371-380.
- 12 **Jiao, Y., Lau, O.S. and Deng, X.W.** (2007) Light-regulated transcriptional networks in
13 higher plants. *Nature reviews. Genetics*, **8**, 217-230.
- 14 **Kavita, P. and Burma, P.K.** (2008) A comparative analysis of green fluorescent
15 protein and beta-glucuronidase protein-encoding genes as a reporter system
16 for studying the temporal expression profiles of promoters. *J Biosci*, **33**, 337-
17 343.
- 18 **Kim, J., Kang, H., Park, J., Kim, W., Yoo, J., Lee, N., Kim, J., Yoon, T.Y. and Choi, G.**
19 (2016) PIF1-Interacting Transcription Factors and Their Binding Sequence
20 Elements Determine the in Vivo Targeting Sites of PIF1. *Plant Cell*, **28**, 1388-
21 1405.
- 22 **Kim, Y.B., Kim, S.M., Kang, M.K., Kuzuyama, T., Lee, J.K., Park, S.C., Shin, S.C. and**
23 **Kim, S.U.** (2009) Regulation of resin acid synthesis in *Pinus densiflora* by
24 differential transcription of genes encoding multiple 1-deoxy-D-xylulose 5-
25 phosphate synthase and 1-hydroxy-2-methyl-2-(*E*)-butenyl 4-diphosphate
26 reductase genes. *Tree Physiol*, **29**, 737-749.
- 27 **Lange, B.M., Rujan, T., Martin, W. and Croteau, R.** (2000) Isoprenoid biosynthesis:
28 the evolution of two ancient and distinct pathways across genomes. *Proc Natl*
29 *Acad Sci USA*, **97**, 13172-13177.
- 30 **Lee, J., He, K., Stolc, V., Lee, H., Figueroa, P., Gao, Y., Tongprasit, W., Zhao, H., Lee,**
31 **I. and Deng, X.W.** (2007) Analysis of transcription factor HY5 genomic binding
32 sites revealed its hierarchical role in light regulation of development. *Plant Cell*,
33 **19**, 731-749.
- 34 **Leivar, P. and Monte, E.** (2014) PIFs: systems integrators in plant development. *Plant*
35 *Cell*, **26**, 56-78.
- 36 **Leivar, P., Monte, E., Oka, Y., Liu, T., Carle, C., Castillon, A., Huq, E. and Quail, P.H.**
37 (2008) Multiple phytochrome-interacting bHLH transcription factors repress
38 premature seedling photomorphogenesis in darkness. *Curr Biol*, **18**, 1815-
39 1823.
- 40 **Leivar, P. and Quail, P.H.** (2011) PIFs: pivotal components in a cellular signaling hub.
41 *Trends Plant Sci*, **16**, 19-28.
- 42 **Leivar, P., Tepperman, J.M., Monte, E., Calderon, R.H., Liu, T.L. and Quail, P.H.**
43 (2009) Definition of early transcriptional circuitry involved in light-induced
44 reversal of PIF-imposed repression of photomorphogenesis in young
45 Arabidopsis seedlings. *Plant Cell*, **21**, 3535-3553.

- 1 **Lois, L.M., Rodríguez-Concepción, M., Gallego, F., Campos, N. and Boronat, A.**
2 (2000) Carotenoid biosynthesis during tomato fruit development: regulatory
3 role of 1-deoxy-D-xylulose 5-phosphate synthase. *Plant J*, **22**, 503-513.
- 4 **Mahmoud, S.S. and Croteau, R.B.** (2001) Metabolic engineering of essential oil yield
5 and composition in mint by altering expression of deoxyxylulose phosphate
6 reductoisomerase and menthofuran synthase. *Proc Natl Acad Sci U S A*, **98**,
7 8915-8920.
- 8 **Malamy, J.E. and Benfey, P.N.** (1997) Organization and cell differentiation in lateral
9 roots of *Arabidopsis thaliana*. *Development*, **124**, 33-44.
- 10 **Mannen, K., Matsumoto, T., Takahashi, S., Yamaguchi, Y., Tsukagoshi, M., Sano,**
11 **R., Suzuki, H., Sakurai, N., Shibata, D., Koyama, T. and Nakayama, T.** (2014)
12 Coordinated transcriptional regulation of isopentenyl diphosphate
13 biosynthetic pathway enzymes in plastids by phytochrome-interacting factor 5.
14 *Biochem Biophys Res Commun*, **443**, 768-774.
- 15 **Meier, S., Tzfadia, O., Vallabhaneni, R., Gehring, C. and Wurtzel, E.T.** (2011) A
16 transcriptional analysis of carotenoid, chlorophyll and plastidial isoprenoid
17 biosynthesis genes during development and osmotic stress responses in
18 *Arabidopsis thaliana*. *BMC Syst Biol*, **5**, 77.
- 19 **Moon, J., Zhu, L., Shen, H. and Huq, E.** (2008) PIF1 directly and indirectly regulates
20 chlorophyll biosynthesis to optimize the greening process in *Arabidopsis*. *Proc*
21 *Natl Acad Sci U S A*, **105**, 9433-9438.
- 22 **Nagel, D.H., Doherty, C.J., Pruneda-Paz, J.L., Schmitz, R.J., Ecker, J.R. and Kay, S.A.**
23 (2015) Genome-wide identification of CCA1 targets uncovers an expanded
24 clock network in *Arabidopsis*. *Proc Natl Acad Sci U S A*, **112**, E4802-4810.
- 25 **Oh, E., Kang, H., Yamaguchi, S., Park, J., Lee, D., Kamiya, Y. and Choi, G.** (2009)
26 Genome-wide analysis of genes targeted by PHYTOCHROME INTERACTING
27 FACTOR 3-LIKE5 during seed germination in *Arabidopsis*. *Plant Cell*, **21**, 403-
28 419.
- 29 **Oudin, A., Mahroug, S., Courdavault, V., Hervouet, N., Zelwer, C., Rodríguez-**
30 **Concepcion, M., St-Pierre, B. and Burlat, V.** (2007) Spatial distribution and
31 hormonal regulation of gene products from methyl erythritol phosphate and
32 monoterpene-secoiridoid pathways in *Catharanthus roseus*. *Plant Mol Biol*, **65**,
33 13-30.
- 34 **Phillips, M.A., León, P., Boronat, A. and Rodríguez-Concepción, M.** (2008) The
35 plastidial MEP pathway: unified nomenclature and resources. *Trends Plant Sci*,
36 **13**, 619-623.
- 37 **Pokhilko, A., Bou-Torrent, J., Pulido, P., Rodríguez-Concepcion, M. and Ebenhoh,**
38 **O.** (2015) Mathematical modelling of the diurnal regulation of the MEP
39 pathway in *Arabidopsis*. *The New phytologist*, **206**, 1075-1085.
- 40 **Pulido, P., Llamas, E., Llorente, B., Ventura, S., Wright, L.P. and Rodríguez-**
41 **Concepcion, M.** (2016) Specific Hsp100 Chaperones Determine the Fate of the
42 First Enzyme of the Plastidial Isoprenoid Pathway for Either Refolding or
43 Degradation by the Stromal Clp Protease in *Arabidopsis*. *PLoS genetics*, **12**,
44 e1005824.
- 45 **Pulido, P., Toledo-Ortiz, G., Phillips, M.A., Wright, L.P. and Rodríguez-**
46 **Concepcion, M.** (2013) *Arabidopsis* J-protein J20 delivers the first enzyme of

1 the plastidial isoprenoid pathway to protein quality control. *Plant Cell*, **25**,
2 4183-4194.

3 **Rodríguez-Concepcion, M.** (2006) Early steps in isoprenoid biosynthesis: multilevel
4 regulation of the supply of common precursors in plant cells. *Phytochem Rev* **5**,
5 1-15.

6 **Rodríguez-Concepción, M.** (2004) The MEP pathway: a new target for the
7 development of herbicides, antibiotics and antimalarial drugs. *Curr Pharm Des*,
8 **10**, 2391-2400.

9 **Rodríguez-Concepcion, M. and Boronat, A.** (2015) Breaking new ground in the
10 regulation of the early steps of plant isoprenoid biosynthesis. *Curr Opin Plant*
11 *Biol*, **25**, 17-22.

12 **Rodríguez-Concepción, M., Forés, O., Martínez-García, J.F., González, V., Phillips,**
13 **M.A., Ferrer, A. and Boronat, A.** (2004) Distinct light-mediated pathways
14 regulate the biosynthesis and exchange of isoprenoid precursors during
15 Arabidopsis seedling development. *Plant Cell*, **16**, 144-156.

16 **Rohdich, F., Bacher, A. and Eisenreich, W.** (2005) Isoprenoid biosynthetic pathways
17 as anti-infective drug targets. *Biochem Soc Trans*, **33**, 785-791.

18 **Ruiz-Sola, M.A. and Rodríguez-Concepcion, M.** (2012) Carotenoid biosynthesis in
19 Arabidopsis: a colorful pathway. *The Arabidopsis book / American Society of*
20 *Plant Biologists*, **10**, e0158.

21 **Schneider, C.A., Rasband, W.S. and Eliceiri, K.W.** (2012) NIH Image to ImageJ: 25
22 years of image analysis. *Nat Methods*, **9**, 671-675.

23 **Tepperman, J.M., Zhu, T., Chang, H.S., Wang, X. and Quail, P.H.** (2001) Multiple
24 transcription-factor genes are early targets of phytochrome A signaling. *Proc*
25 *Natl Acad Sci U S A*, **98**, 9437-9442.

26 **Toledo-Ortiz, G., Huq, E. and Quail, P.H.** (2003) The Arabidopsis basic/helix-loop-
27 helix transcription factor family. *Plant Cell*, **15**, 1749-1770.

28 **Toledo-Ortiz, G., Huq, E. and Rodríguez-Concepcion, M.** (2010) Direct regulation of
29 phytoene synthase gene expression and carotenoid biosynthesis by
30 phytochrome-interacting factors. *Proc Natl Acad Sci U S A*, **107**, 11626-11631.

31 **Toledo-Ortiz, G., Johansson, H., Lee, K.P., Bou-Torrent, J., Stewart, K., Steel, G.,**
32 **Rodríguez-Concepcion, M. and Halliday, K.J.** (2014) The HY5-PIF regulatory
33 module coordinates light and temperature control of photosynthetic gene
34 transcription. *PLoS genetics*, **10**, e1004416.

35 **Umehara, M., Hanada, A., Yoshida, S., Akiyama, K., Arite, T., Takeda-Kamiya, N.,**
36 **Magome, H., Kamiya, Y., Shirasu, K., Yoneyama, K., Kyojuka, J. and**
37 **Yamaguchi, S.** (2008) Inhibition of shoot branching by new terpenoid plant
38 hormones. *Nature*, **455**, 195-201.

39 **von Lintig, J., Welsch, R., Bonk, M., Giuliano, G., Batschauer, A. and Kleinig, H.**
40 (1997) Light-dependent regulation of carotenoid biosynthesis occurs at the
41 level of phytoene synthase expression and is mediated by phytochrome in
42 *Sinapis alba* and *Arabidopsis thaliana* seedlings. *Plant J*, **12**, 625-634.

43 **Vranova, E., Coman, D. and Grusissem, W.** (2013) Network analysis of the MVA and
44 MEP pathways for isoprenoid synthesis. *Annu Rev Plant Biol*, **64**, 665-700.

45 **Walley, J., Xiao, Y., Wang, J.Z., Baidoo, E.E., Keasling, J.D., Shen, Z., Briggs, S.P. and**
46 **Dehesh, K.** (2015) Plastid-produced interorgannellar stress signal MEcPP

1 potentiates induction of the unfolded protein response in endoplasmic
2 reticulum. *Proc Natl Acad Sci U S A*, **112**, 6212-6217.

3 **Walter, M., Fester, T. and Strack, D.** (2000) Arbuscular mycorrhizal fungi induce the
4 non-mevalonate methylerythritol phosphate pathway of isoprenoid
5 biosynthesis correlated with accumulation of the yellow pigment and the other
6 apocarotenoids. *Plant J*, **21**, 571-578.

7 **Wang, Z.Y., Kenigsbuch, D., Sun, L., Harel, E., Ong, M.S. and Tobin, E.M.** (1997) A
8 Myb-related transcription factor is involved in the phytochrome regulation of
9 an Arabidopsis Lhcb gene. *Plant Cell*, **9**, 491-507.

10 **Wright, L.P., Rohwer, J.M., Ghirardo, A., Hammerbacher, A., Ortiz-Alcaide, M.,**
11 **Ragusckke, B., Schnitzler, J.P., Gershenzon, J. and Phillips, M.A.** (2014)
12 Deoxyxylulose 5-Phosphate Synthase Controls Flux through the
13 Methylerythritol 4-Phosphate Pathway in Arabidopsis. *Plant Physiol*, **165**,
14 1488-1504.

15 **Zhang, H., He, H., Wang, X., Wang, X., Yang, X., Li, L. and Deng, X.W.** (2011)
16 Genome-wide mapping of the HY5-mediated gene networks in Arabidopsis that
17 involve both transcriptional and post-transcriptional regulation. *Plant J*, **65**,
18 346-358.

19 **Zhang, Y., Mayba, O., Pfeiffer, A., Shi, H., Tepperman, J.M., Speed, T.P. and Quail,**
20 **P.H.** (2013) A quartet of PIF bHLH factors provides a transcriptionally centered
21 signaling hub that regulates seedling morphogenesis through differential
22 expression-patterning of shared target genes in Arabidopsis. *PLoS genetics*, **9**,
23 e1003244.

24

25 **Supporting information**

26
27 **Supplemental Figure 1.** Expression analysis of the *DXS1*, *DXR* and *HDR* genes
28 in *hy5* mutants in Red light.

29
30 **Supplemental Figure 2.** Expression analysis of the *DXS1*, *DXR* and *HDR* genes
31 in the *pif* single mutants in Red light.

32
33 **Supplemental Figure 3.** *DXS1* upstream regulatory region.

34
35 **Supplemental Figure 4.** Chromatin immunoprecipitation assays for 35S::TAP-PIF1
36 and 35S::HA-HY5 in de-etiolated seedlings.

37
38 **Supplemental Figure 5.** *DXR* upstream regulatory region.

39
40 **Supplemental Figure 6.** *HDR* upstream regulatory region.

41
42 **Supplemental Table I.** Sequence of the oligonucleotides used in the work.

1 **Figure Legends**

2

3 **Figure 1. Transcriptional regulation of *DXS1* by light.** Histochemical expression
4 of 3 day-old transgenic representative lines grown in the dark (a-c) or exposed to 6
5 h of light (d-f) expressing the GUS marker from 1510 bp (a and d), 753 bp (b and e)
6 and 660 bp (c and f) upstream sequences from the ATG of *DXS1*. (g) GUS specific
7 activities from the dark (dark gray) or exposed to light (gray) transgenic lines. The
8 induction level is reported relative to the mean of the specific activity (nmol of
9 methyl-umbelliferone per μg of protein $^{-1}$ min $^{-1}$) of the corresponding dark sample.
10 Each bar is the mean of three independent experiments and error bars represent
11 \pm SD. The numbers above the bars indicate the *P* values according to a Student's *t*
12 test.

13

14 **Figure 2. Expression analysis of the *DXS1*, *DXR* and *HDR* genes in *hy5* and**
15 ***hy5hyh* mutants.** Representative northern blots of *DXS1* (a), *DXR* (b) and *HDR*
16 (c), each lane contains 5 μg of RNA from 3 day-old dark-adapted seedlings
17 maintained in the dark (D) or exposed to light for 6 h (L) from wild-type (Wt), *hy5*
18 and *hy5hyh* mutants and hybridized with probes for the *DXS1* (a), *DXR* (b) and
19 *HDR* (c) genes. The 28S rRNA is shown as loading control. The membrane is
20 representative of three independent biological experiments. qRT-PCR analysis of
21 *DXS1* (d), *DXR* (e) and *HDR* (f) transcript levels from Col-0, *hy5* and *hy5hyh* dark-
22 adapted seedlings maintain in the dark (gray column) or exposed to 6 h of light
23 (white column). Expression is normalized to the Col-0 dark samples and adjusted
24 to *Actin 7 (ACT7)*. Bars are means \pm SE of triplicate biological experiments (each
25 with n=2 technical replicas). The numbers bars indicate P values $p < 0.05$ and the *
26 marks statistical difference ($p < 0.05$) between the light samples from the *hy5* and
27 *hy5hyh* mutants compared to the induction in the Wt according to a Student's *t*
28 test.

29

30 **Figure 3. Expression analysis of the *DXS1*, *DXR* and *HDR* genes in the *pif***
31 **single mutants.** Representative RNA blots of *DXS1* (a), *DXR* (b) and *HDR* (c)

1 from 3 day-old dark-adapted wild-type Col0 (Wt), *pif1*, *pif3*, *pif4* and *pif5* maintained
2 in darkness (D) or exposed to 6 h light (L). The 28S rRNA is shown as a loading
3 control (28S). Membranes are representative of three independent biological
4 experiments. Analysis by qRT-PCR of *DXS1* (d,g), *DXR* (e,h) and *HDR* (f,i)
5 transcripts from Col-0, *pif1*, *pif3*, *pif4* and *pif5* (d-f) dark-adapted for 3 days (gray
6 column) or 36 h for *pifQ* (g-i) and exposed 6 h of red light (white column).
7 Expression is reported relative to the dark Col-0 sample and adjusted relative to
8 *Actin 7 (ACT7)*. Bars are means \pm SE of triplicate biological experiments (each with
9 n=2 technical replicas). The letter above the bars indicate P values $p < 0.05$
10 between dark (a) or light (b) of Wt compared to the mutants (Student's t test).
11

12 **Figure 4. Expression analysis of the *DXS1*, *DXR* and *HDR* genes in the *hy5*,**
13 ***pif* single mutants and *pifQ* during de-etiolation.** qRT-PCR analysis of *DXS1*
14 (a), *DXR* (b) and *HDR* (c) genes from etiolated seedlings of Col-0, *hy5*, *pif1*, *pif3*,
15 *pif4*, *pif5* and *pifQ* maintained in the dark (gray columns) or exposed to 6 h of red
16 light (white columns) relative to the Col-0 dark samples and adjusted relative to
17 *Actin 7 (ACT7)*. Bars are means \pm SE of triplicate biological experiments (each with
18 n=2 technical replicas). The letter above the bars indicate values ($p < 0.05$)
19 decrease (a and b) or increase (c and d) between the dark (a and c) or light (b and
20 d) values between the Col0 and the mutants (Student's t test).
21

22 **Figure 5. Chromatin immunoprecipitation assays for 35S::TAP-PIF1 and**
23 **35S::HA-HY5 in dark-adapted seedlings.** Diagrams of the upstream regions of
24 *DXS1*, *DXR* and *HDR* genes. Primers used for the analyses (arrows) and the
25 potential PIFs (black) and HY5 (gray) binding elements (rectangles). ChIP of three
26 days dark-adapted TAP-PIF and HA-HY5 transgenic seedlings (Lee *et al.*, 2007,
27 Moon *et al.*, 2008) maintained in the dark (grey zone) or exposed to 6h red-light
28 (clear zone). ChIP was conducted using specific antibodies against MYC for PIF1
29 the HA for HY5. 35S::GFP-TAP or 35S::GUS-HA lines were used as controls for
30 unespecific binding. The ChIP/qPCR was done using specific primer pairs (F,
31 forward primer and R, reverse primer) covering the regions containing putative

1 binding elements. ChIP-enriched DNA regions of the TAP-PIF *DXS1* (a and c),
2 *DXR* (e,g and i) and *HDR* (k) or for HA-HY5 *DXS1* (b and d) *DXR* (f,h and j) and
3 *HDR* (l) samples. The bars are the mean \pm SE of triplicate independent
4 experiments (each with technical duplicates n=2). The asterisk indicate the values
5 with significance ($p < 0.05$) between the negative control according to a Student's t
6 test.

Commented [HG2]: In the figure 'f' is incorrectly annotated 'g' so you have two labeled 'g'

7
8 **Figure 6. Protein accumulation of the *DXS1*, *DXR* and *HDR* in mutants of the**
9 ***HY5* and *PIFs*.** (a) Immunoblots with 20 μ g of protein extracts from seedlings dark-
10 adapted for 3 days (*Col-0*, *hy5*, *pif1*, *pif3*, *pif4* and *pif5*) or 36 h (*pifQ*) and
11 maintained in the dark (D) or exposed to 6 h light (L) using specific antibodies for
12 *DXS1*, *DXR* or *HDR* proteins. A Coomassie blue-stained gel (Coo) is shown as a
13 loading control. A representative gel from three independent biological experiments
14 is shown. (b) Densitometric analyses of the *DXS1*, *DXR* and *HDR* protein levels
15 from *Col-0*, *hy5*, *pif1*, *pif3*, *pif5* and *pifQ* immunoblots from the dark (grey bars) or
16 exposed to 6 h light (white bars) samples. The expression level is reported relative
17 to the *Col-0* light samples and adjusted to the corresponding loading control. The
18 bars correspond to the average of three independent biological experiments \pm SD
19 of biological triplicates.

20
21 **Figure 7. Model of the differential light regulation of *DXS1*, *DXR* and *HDR***
22 **gene expression to modulate the MEP pathway.** Light via phytochrome (Pfr)
23 results in the degradation of PIFs and in the accumulation of the HY5. For the
24 *DXS1*, Pfr accumulation inhibits the direct repression of PIF1 (solid arrow), PIF3
25 and PIF5 (dashed arrows) and the activation of HY5 (solid arrow). For *DXR*, Pfr
26 impairs the weak repression of PIF1 in multiple sites (solid arrows), PIF3 and PIF5
27 (dashed arrows). Light via HY5 activate *DXR* (solid line). Finally, Pfr accumulation
28 induces *HDR* expression through the degradation of the PIFs. The dashed arrows
29 mean that a direct interaction was not experimentally demonstrated. The thickness
30 of the arrows reflects the RNA levels and the enrichment detected in the ChIP

- 1 analyses. The orange arrow reflects the dynamic of the light activation observed for
- 2 each gene.
- 3
- 4