

# 1 Using noble gases to trace groundwater evolution and assess helium 2 accumulation in Weihe Basin, central China

3 Wen Zhang <sup>a, b, c, d</sup>, Yuhong Li <sup>b, \*</sup>, Fenghua Zhao <sup>a</sup>, Wei Han <sup>b</sup>, Yan Li <sup>c</sup>, Yunpeng Wang <sup>e</sup>,  
4 Greg Holland <sup>d</sup>, Zheng Zhou <sup>c</sup>

5 a. College of Geoscience and Surveying Engineering, China University of Mining and Technology,  
6 100083, Beijing, China

7 b. Xi'an Center, China Geological Survey, 710054, China

8 c. Lancaster Environment Centre, Lancaster University, LA1 4YQ, UK

9 d. School of Earth and Environmental Sciences, The University of Manchester, M13 9PL, UK

10 e. State Key Laboratory of Organic Geochemistry, Guangzhou Institute of Geochemistry, Chinese  
11 Academy of Sciences, Guangzhou, 510640, China

12

13 **Abstract:** The severe shortage of helium resources is an impending global problem. However, the  
14 helium accumulation processes and conditions favorable for helium enrichment in reservoirs remain  
15 poorly understood, which makes helium exploration challenging. Noble gases are good tracers of  
16 subsurface fluids provenance, migration and storage, as well as indicators of the nature and quantity  
17 of associated phases. In this study the variation of major gases and noble gases data in Weihe Basin  
18 provide us with an excellent opportunity to understand the groundwater evolution and helium  
19 accumulation processes. Twelve gas samples collected from wellheads of geothermal wells can be  
20 classified into three groups, in which Group A has high concentrations of N<sub>2</sub> (58.57% - 91.66%) and  
21 He (0.32% - 2.94%); Group B has high contents of CH<sub>4</sub> (52.94% and 69.50%) and low  
22 concentrations of He (0.057% and 0.062%); Group C has a high content of CH<sub>4</sub> (71.70%) and He  
23 (2.11%). Helium isotopic ratios are predominantly radiogenic in origin and therefore crustally  
24 derived. Measured elemental ratios of noble gases are compared with multiple fractionation models  
25 for Group A and B samples, implying that open system heavy oil-water fractionation with excess  
26 heavy noble gases has occurred in the basin with  $V_{oil}/V_{water}$  ratios of 0.06-0.18. The amount of  
27 helium in Group A and B samples requires the release of all <sup>4</sup>He produced in the crust since

---

\* Corresponding author

E-mail address: [L1763@tom.com](mailto:L1763@tom.com) (Yuhong Li); [wenzhangcn@outlook.com](mailto:wenzhangcn@outlook.com) (Wen Zhang) Tel: +8618734130189 (Yuhong Li)

28 0.30Ma-1.98Ma into the groundwater. The Group C sample requires an additional He flux from  
29 adjacent granitic bodies. The accumulation of helium and hydrocarbon in the Weihe Basin can be  
30 explained by a 4-stage process. Accumulation of commercially viable helium requires high He flux  
31 from source rocks, the existence of a free gas phase of major gas components (CH<sub>4</sub> in most cases,  
32 N<sub>2</sub> or CO<sub>2</sub>) and minimal major gas addition after formation of the free gas phase.

33 **Keywords:** Noble gases; Helium accumulation; Oil-water fractionation; Weihe Basin

34

## 35 1. INTRODUCTION

36 Helium is an essential resource used extensively in nuclear magnetic resonance (NMR),  
37 high-tech industries and scientific research (Cai et al., 2010). Based on present global helium  
38 reserves, a severe shortage of helium is imminent (Nuttall et al., 2012; USGS, 2018), making  
39 helium resource exploration an urgent issue.

40 <sup>4</sup>He production is dominated by  $\alpha$ -decay of <sup>235,238</sup>U and <sup>232</sup>Th in sedimentary strata, basement  
41 or crust (Ballentine and Burnard, 2002; Brown, 2010). The <sup>4</sup>He release process is controlled by two  
42 mechanisms: (1)<sup>4</sup>He is released from crust or aquifer by steady-state flux and accumulated in  
43 groundwater (Torgersen and Clarke, 1985; Torgersen and Ivey, 1985; Castro et al., 1998), making  
44 <sup>4</sup>He an effective tool to date groundwater associated with oil and gas reservoirs (Barry et al., 2017;  
45 Zhou and Ballentine, 2006) and ancient fluid systems (Holland et al., 2013). (2)<sup>4</sup>He can also be  
46 liberated rapidly over a short period of time from ancient rocks by magmatic and tectonic activity,  
47 as shown at Yellowstone in the U.S.A. (Lowenstern et al., 2014) and Tanzania (Danabalan, 2017).  
48 After release, the concentration of <sup>4</sup>He in the subsurface fluid systems can then be altered by  
49 interactions with oil, gas and water. Helium is often associated with natural gas (Ballentine and  
50 Lollar, 2002; Brown, 2010), nitrogen (Danabalan et al., 2016), and carbon dioxide (Gilfillan et al.,  
51 2008) and can be closed linked to groundwater flow, as observed in the Hugoton-Panhandle gas  
52 field (Ballentine and Lollar, 2002; Danabalan, 2017).

53 Noble gases have been widely used in hydrocarbon systems to quantify oil-water-gas  
54 interaction during groundwater movement and hydrocarbon accumulation (Ballentine et al., 1991;

55 Barry et al., 2016; Wen et al., 2015; Zhou et al., 2005) as well as in CO<sub>2</sub> and N<sub>2</sub> systems (Ballentine  
56 and Lollar, 2002; Danabalan, 2017; Gilfillan et al., 2008; Zhou et al., 2012) because of their  
57 chemical inertness. In these case studies, noble gases dissolved in air-saturated water (ASW, water  
58 that has equilibrated with air under recharge conditions) are transported into the subsurface  
59 preserving ASW noble gas signatures, which are subsequently modified by solubility-controlled  
60 equilibration when groundwater contacts oil and/or gas phases (Ballentine et al., 2002; Kipfer et al.,  
61 2002). Thus ASW-derived noble gases dissolved in groundwater or partitioning into gas phases  
62 provide a tool to understand the relationship between groundwater evolution and hydrocarbon and  
63 helium accumulation.

64 In this study, we present the concentrations and isotope ratios of noble gases, major gas  
65 components and carbon isotopes of CH<sub>4</sub> and CO<sub>2</sub> in gas samples collected from geothermal wells in  
66 Weihe Basin, central China. The combination of these data provides an excellent opportunity to  
67 understand the influence of groundwater (traced by ASW-derived noble gases) and major gas  
68 components on helium transportation and accumulation as well as determining the sources of  
69 helium.

70

## 71 **2. GEOLOGICAL SETTING**

72 Weihe Basin, a Cenozoic graben measuring 22000 km<sup>2</sup>, is located in central China (Figure 1).  
73 It is the southern part of the Cenozoic graben system surrounding the Ordos Block, formed during  
74 the continental extension of the Qinling Orogenic Belt (Deng and You, 1985; Li and Ren, 1986).  
75 Geophysical data suggest the crustal thickness is around 35-38km under the Weihe Basin compared  
76 with 40km beneath the Ordos Basin. There is an upper mantle uplift of 5-6 km under the Weihe  
77 Basin (Xu et al., 1993). Neoproterozoic, Paleozoic and Mesozoic granitoid formations occurred  
78 throughout the Qinling Orogenic Belt (Wang et al., 2015b), which are rich in uranium and thorium  
79 (Han et al., 2014). Many Cenozoic faults and fractures are present in the basin and around the  
80 border area, suggesting extension in this region during the Cenozoic (Mercier et al., 2013).  
81 Numerous destructive earthquakes, such as the 1556 M8 Huaxian earthquake (Zhang et al., 1995),

82 demonstrate that this region has been tectonically active until very recently.

83 Two main depressions, the Xi'an Depression and the Gushi Depression, are distributed  
84 throughout the basin. The basin is filled with Cenozoic strata around 7000m thick, which is  
85 comprised of Eocene, Miocene, Pliocene, Pleistocene, and Holocene layers. The major geothermal  
86 reservoir is found within the Pliocene Lantian-Bahe Formation, deposited under fluvial to shallow  
87 lacustrine conditions (Ma et al., 2010). Groundwater is recharged from the Ordos Basin in the north  
88 and the Qinling Orogenic Belt in the south and discharged from the Weihe River in the middle.  
89 Sampling wells are distributed in the area of groundwater circulation from the Qinling Orogenic  
90 Belt to the Weihe River. Depths of sample collection are listed in Table 1.

91

### 92 **3. SAMPLING AND ANALYTICAL METHODS**

#### 93 **3.1 Sampling**

94 Twelve gas samples were collected from the wellheads of different producing geothermal wells  
95 in the Weihe Basin, 7 of which are located in the Xi'an Depression and 5 are located in the Gushi  
96 Depression. A flash tank was equipped at the wellhead of WR-HX to separate gas and water due to  
97 the high gas content. Therefore, the gas sample was collected directly from the exhaust pipe for  
98 WR-HX. Since there were only gas bubbles exsolved from groundwater at other wellheads and  
99 flash tanks are not equipped, a gas-water separator was used to take gas samples (Pinti et al., 2017).  
100 Noble gas samples were collected in standard refrigeration grade 10mm external-diameter copper  
101 tubes (~50cm in length). The gas was allowed to flow freely through the collection system for at  
102 least 10 minutes to remove air contamination before the tubes were sealed by stainless steel  
103 pinch-off clamps (Weiss, 1968; Zhou et al., 2005). Additional samples for analysis of major gas and  
104 carbon isotopes of CH<sub>4</sub> and CO<sub>2</sub> were collected in upside-down glass bottles by displacing saturated  
105 salt water.

106 Gases in Group A and Group B wells are dissolved in groundwater in the subsurface in the  
107 Weihe Basin (Zhang et al., 2014a). Because a gas-water separator was used to separate gas and  
108 water phases during sampling and the gaseous phase has a strong affinity for noble gases (Warr et

109 al., 2018), a complete degassing of noble gases can be assumed. Therefore, the elemental and  
110 isotopic ratios measured in the gas phase of Group A and Group B samples can represent the  
111 associated groundwater from which the gases are exsolved. The much higher gas content at the  
112 wellhead of Group C well (WR-HX) in comparison to other sampling wells, as well as their  
113 associated noble gas signatures suggest that there is a gas reservoir in the subsurface (discussed in  
114 section 5.2.2). The sampled gas at the wellhead of WR-HX is a mixture of free gas and dissolved  
115 gas.

## 116 **3.2 Analytical methods**

### 117 **3.2.1 Major gas components and carbon isotopes**

118 Major gas components were analyzed at the Key Laboratory of Petroleum Resource Research,  
119 Chinese Academy of Sciences, Lanzhou. The analytical procedures were described in detail in  
120 Zhang et al. (2018a). Briefly, the contents of CH<sub>4</sub> and non-hydrocarbon gases were measured by a  
121 MAT271 mass spectrometer. CH<sub>4</sub> and other hydrocarbon gases were analyzed on an on-line  
122 continuous flow gas chromatograph (Agilent 6890 GC), which is equipped with flame ionization  
123 and thermal conductivity detectors. Since CH<sub>4</sub> concentrations were analyzed by both MAT271 and  
124 GC, the non-hydrocarbon and hydrocarbon results were calibrated to 100% in total based on the  
125 CH<sub>4</sub> contents. The analytical errors were less than 1 vol.% for CH<sub>4</sub>, CO<sub>2</sub> and N<sub>2</sub> and less than 10  
126 vol.% for other gases.

127 Carbon isotopes of CH<sub>4</sub> and CO<sub>2</sub> were measured at the State Key Laboratory of Organic  
128 Geochemistry, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou.  
129 The analyses were performed by a Delta Plus II XL GC-IRMS equipped with a CP-Poraplot Q  
130 column using standard techniques as described in Gai et al (2019). Each gas sample was analyzed  
131 twice and the errors were less than  $\pm 0.5\%$ .

### 132 **3.2.2 Noble gases**

133 Noble gas analysis was conducted in the noble gas laboratory at Lancaster University. The  
134 analytical system can be divided into four sections: sample introduction, sample purification, noble  
135 gas separation and noble gas analysis system, which is shown in Figure 2.

136 After connecting a sample copper tube to the gas extraction line, the line was evacuated  
137 through a turbomolecular pump for at least 6 hours to remove air and ensure the preline vacuum  
138 was less than  $2 \times 10^{-7}$  mbar, which was monitored by a compact cold cathode pressure gauge. A  
139 quadrupole mass spectrometer (Hiden Analytical HAL-201) was used to further monitor vacuum  
140 and check for any leaks from the crimped copper tube. After releasing gas from the copper tube,  
141 samples were expanded into a calibrated volume where pressure was recorded by an MKS dual  
142 capacitance manometer. Then the sample was transferred to the purification line where  
143 hydrocarbons and other reactive gases were removed by titanium sponge initially held at 800 °C for  
144 1min and then cooled to room temperature of 21 °C in 20mins. Further purification was carried on a  
145 hot getter (SAES GP-50) held at 250 °C for 15mins. Heavy noble gases (Ar, Kr and Xe) were  
146 trapped in a cold finger by charcoal at 77K (refrigerated by liquid nitrogen) and light noble gases  
147 (He and Ne) were trapped by a Janis cryogenic trap at 9.5K.

148 After trapping all noble gases, temperature of the cryogenic trap was increased to 50K and held  
149 for 5mins to release helium, which was then admitted to the Isotopx NGX mass spectrometer to  
150 measure the helium concentration and isotopic ratio. Neon was subsequently released from the  
151 cryogenic trap at 110K and held for 5mins before admittance to the mass spectrometer.  
152 Doubly-charged  $^{40}\text{Ar}^{++}$  and  $\text{CO}_2^{++}$  were used to correct  $^{20}\text{Ne}$  and  $^{22}\text{Ne}$  following the methods of  
153 Niedermann et al. (1993). After measurements of He and Ne, the charcoal cold finger was heated  
154 from 77K to 450K and held for 10mins to release argon, krypton and xenon completely. Two  
155 aliquots of the released gas were admitted into NGX mass spectrometer successively for separate  
156 analysis of argon and krypton. The remaining Xe in the gas phase was then re-absorbed onto a  
157 charcoal trap held at 210K for 10mins followed by pumping out of the residual unabsorbed gas  
158 (argon and krypton). Finally, the charcoal trap was heated to 450K and held for 10mins again to  
159 release xenon, which was admitted into NGX mass spectrometer for xenon analysis. A cold getter  
160 (SAES GP-50) was held at room temperature (21°C) to further remove hydrogen before allowing  
161 gases into the mass spectrometer.

162 A blank and an air standard were analyzed for each sample. Blank corrections for  $^4\text{He}$ ,  $^{20}\text{Ne}$ ,  
163  $^{40}\text{Ar}$ ,  $^{84}\text{Kr}$  and  $^{130}\text{Xe}$  blanks were 0.37 vol%, 0.04 vol%, 0.67 vol%, 0.16 vol% and 0.11 vol% of the  
164 average sample size, respectively.

165

## 166 4. RESULTS

### 167 4.1 Major gas components and carbon isotopes

168 Sample names, locations, depths, major gas components and carbon isotopes are listed in Table  
169 1. Seven samples and five samples were collected from the Xi'an and Gushi Depressions  
170 respectively. These samples can be classified into three groups based on their  $\text{CH}_4$ ,  $\text{N}_2$  and He  
171 concentrations. Group A includes all samples collected from the Xi'an Depression (XP-1, XP-2,  
172 XP-3, XY-1, XY-2, XY-3 and WG-1) and two samples taken from the Gushi Depression (WR-2 and  
173 WR-051). They have relatively low concentrations of  $\text{CH}_4$  between 2.23% and 25.40% (9.78%  
174 average), high concentrations of  $\text{N}_2$  between 58.57% and 91.66% (76.14% average) and high  
175 concentrations of He between 0.32% and 2.94% (1.62% average). Group B is composed of two  
176 samples from the Gushi Depression (WR-1 and WR-11) with high concentrations of  $\text{CH}_4$  (52.94%  
177 and 69.50%), low concentrations of  $\text{N}_2$  (27.81% and 4.85%) and low concentrations of He (0.057%  
178 and 0.062%). Finally, we classify one sample (WR-HX) as Group C as it is characterized by a high  
179 concentration of He (2.11%) and similar contents of  $\text{CH}_4$  (71.70%) and  $\text{N}_2$  (27.20%) compared to  
180 samples in Group B. The  $\text{CO}_2$  concentrations in the majority of samples are lower than 10% (0.03%  
181 - 9.36%, 4.02% in average) except for WR1 (22.24%), WR11 (13.56%), and XY-2 (31.69%). There  
182 are minor amounts of other hydrocarbon gases ( $\text{C}_{2-6}$ ) in all samples.  $\text{O}_2$  is observed in all samples  
183 except for the Group C sample. The carbon isotopic composition of  $\text{CH}_4$ ,  $\delta^{13}\text{C}(\text{C}_1)$ , shows a large  
184 range from -19.04‰ to -44.46‰ (VPDB).  $\delta^{13}\text{C}(\text{CO}_2)$  values vary between -10.50‰ and -5.47‰,  
185 reported relative to Vienna Pee Dee Belemnite (VPDB), which form a positive correlation ( $R^2 =$   
186 0.63) with  $\text{CO}_2$  concentrations.

### 187 4.2 Noble gases

188 He, Ne, Ar, Kr and Xe abundances and isotopic compositions of the 12 samples are given in

189 Table 2. Helium concentrations in Group A and C samples are higher than those in Group B samples  
190 by one to two orders of magnitude. Other noble gases in Group A samples show higher  
191 concentrations compared with those of Group B and Group C samples. The noble gas signatures are  
192 described from section 4.2.1 to section 4.2.5.

#### 193 4.2.1 Helium

194  $^4\text{He}$  concentrations are between  $32.3$  and  $294 \times 10^{-4} \text{ cm}^3 \text{ STP/cm}^3$  in Group A samples,  $5.74$   
195 and  $6.21 \times 10^{-4} \text{ cm}^3 \text{ STP/cm}^3$  in Group B samples and  $211 \times 10^{-4} \text{ cm}^3 \text{ STP/cm}^3$  in the Group C  
196 sample. The  $^4\text{He}$  concentrations in the Weihe Basin geothermal field are much higher than those in  
197 most natural gas fields around the world. For example,  $^4\text{He}$  concentrations are  $0.04\text{-}116 \times 10^{-6} \text{ cm}^3$   
198  $\text{STP/cm}^3$  in the San Juan Basin, USA (Zhou et al., 2005) and  $5.32\text{-}6.81 \times 10^{-5} \text{ cm}^3 \text{ STP/cm}^3$  in the  
199 Sleipner Vest gas field (Barry et al., 2016) respectively. Helium in the Weihe Basin is dominated by  
200 radiogenic components with helium isotope values ( $^3\text{He}/^4\text{He}$ ) varying between  $0.033\text{Ra} \pm 0.001\text{Ra}$   
201 and  $0.126\text{Ra} \pm 0.002\text{Ra}$ , where Ra is the atmospheric value of  $^3\text{He}/^4\text{He}$  of  $1.4 \times 10^{-6}$  (Mamyrin and  
202 Tolstikhin, 1984). Measured  $^4\text{He}/^{20}\text{Ne}$  values are between 1439 and 37164, which are much higher  
203 than the ratio in ASW (0.288, Kipfer et al., 2002) or air (0.318, Sano et al., 2013). Therefore,  
204 atmospheric or ASW-derived gas has little contribution to  $^4\text{He}$  concentrations. Using a simple  
205 two-component mixing model between a sub-continental lithospheric mantle-like endmember  
206 ( $6.1\text{Ra}$ , Gautheron and Moreira, 2002) and an upper crustal endmember ( $0.008\text{Ra}$ , Ballentine and  
207 Burnard, 2002), the mantle contribution to helium is between 0.40% and 1.94%. The remainder of  
208 the  $^4\text{He}$  is derived from the crust (Fig. 3).

#### 209 4.2.2 Neon

210  $^{20}\text{Ne}$  concentrations are between  $15.9$  and  $39.5 \times 10^{-7} \text{ cm}^3 \text{ STP/cm}^3$  ( $30.0 \times 10^{-7} \text{ cm}^3 \text{ STP/cm}^3$  in  
211 average) in Group A samples,  $0.69$  and  $1.11 \times 10^{-7} \text{ cm}^3 \text{ STP/cm}^3$  in Group B samples, and  $5.50 \times$   
212  $10^{-7} \text{ cm}^3 \text{ STP/cm}^3$  in the Group C sample. Measured  $^{20}\text{Ne}/^{22}\text{Ne}$  ratios vary between  $8.68 \pm 0.15$  and  
213  $10.9 \pm 0.40$ , showing minor deviations from the atmospheric  $^{20}\text{Ne}/^{22}\text{Ne}$  value of 9.80 (Bottomley et  
214 al., 1984, Fig. 4). This may be due to minor mass fractionation as observed previously (Zhou et al.,  
215 2005). Measured  $^{21}\text{Ne}/^{22}\text{Ne}$  ratios vary between  $0.0291 \pm 0.0008$  and  $0.0311 \pm 0.0008$  except for



216 sample WR-HX ( $0.0385 \pm 0.0015$ ), compared with the atmospheric value of  $^{21}\text{Ne}/^{22}\text{Ne}=0.029$   
217 (Bottomley et al., 1984). Ne isotope ratios are indistinguishable between the samples from the two  
218 depressions. Helium isotopes suggest a minor contribution from mantle degassing, therefore the  
219 “excess”  $^{21}\text{Ne}$  can be accounted for by the addition of radiogenic  $^{21}\text{Ne}$  to ASW-derived Ne (Fig. 4).  
220 The crustal Ne endmember is featured by  $^{20}\text{Ne}/^{22}\text{Ne}=0$  and  $^{21}\text{Ne}/^{22}\text{Ne} = 0.47$  (Kennedy et al., 1990).  
221 Based on the simple two-component mixing model between air and crust (Ballentine et al., 2002),  
222 the  $^{21}\text{Ne}$  derived from crust accounts for between 0.0% and 9.5% of the total  $^{21}\text{Ne}$  except for  
223 WR-HX well, which is 33.3%.

#### 224 4.2.3 Argon

225  $^{36}\text{Ar}$  concentrations range between  $7.18$  and  $19.20 \times 10^{-6} \text{ cm}^3 \text{ STP/cm}^3$ , with  $^{40}\text{Ar}/^{36}\text{Ar}$  ratios  
226 varying between  $339 \pm 5$  and  $613 \pm 11$  in Group A samples. Comparatively, Group B samples show  
227 around 10 times lower  $^{36}\text{Ar}$  concentrations ( $2.51$  and  $5.78 \times 10^{-7} \text{ cm}^3 \text{ STP/cm}^3$ ) and slightly higher  
228  $^{40}\text{Ar}/^{36}\text{Ar}$  ratios ( $542 \pm 9$  to  $735 \pm 13$ ). The Group C sample shows a higher  $^{36}\text{Ar}$  concentration  
229 ( $2.88 \times 10^{-6} \text{ cm}^3 \text{ STP/cm}^3$ ) and a higher  $^{40}\text{Ar}/^{36}\text{Ar}$  ratio ( $1142 \pm 19$ ) than Group B samples.  $^{40}\text{Ar}/^{36}\text{Ar}$   
230 ratios show different deviation from the atmospheric  $^{40}\text{Ar}/^{36}\text{Ar}$  ratio of  $298.56 \pm 0.31$  (Lee et al.,  
231 2006; Mark et al., 2011). Resolved excess  $^{40}\text{Ar}^*$  ( $^{40}\text{Ar}^* = ^{40}\text{Ar} - 298.56 \times ^{36}\text{Ar}$ , Battani et al., 2000)  
232 contributes 12.9%-74.1% to the total  $^{40}\text{Ar}$ . Since there is no distinct mantle-derived helium, we  
233 assume the excess  $^{40}\text{Ar}$  is derived from a radiogenic source. Measured  $^{38}\text{Ar}/^{36}\text{Ar}$  ratios are air-like  
234 ( $0.1885 \pm 0.0003$ , Lee et al., 2006; Mark et al., 2011), varying between  $0.180 \pm 0.006$  and  
235  $0.200 \pm 0.005$ .

#### 236 4.2.4 Krypton and Xenon

237  $^{84}\text{Kr}$  concentrations are between  $16.6$  and  $64.8 \times 10^{-8} \text{ cm}^3 \text{ STP/cm}^3$  in Group A samples and  
238 between  $0.869$  and  $2.07 \times 10^{-8} \text{ cm}^3 \text{ STP/cm}^3$  in Group B samples and  $9.95 \times 10^{-8} \text{ cm}^3 \text{ STP/cm}^3$  in the  
239 Group C sample. The isotope ratios of krypton are indistinguishable from corresponding air ratios.  
240 For example,  $^{86}\text{Kr}/^{84}\text{Kr}$  ratios range from  $0.286 \pm 0.014$  to  $0.322 \pm 0.022$  compared to the  
241 atmospheric ratio of  $^{86}\text{Kr}/^{84}\text{Kr} = 0.3035 \pm 0.000008$  (Aregbe et al., 1996).

242  $^{130}\text{Xe}$  concentrations vary from  $14.5$  to  $40.2 \times 10^{-10} \text{ cm}^3 \text{ STP/cm}^3$  in Group A samples, compared

243 with  $0.98 - 2.65 \times 10^{-10} \text{ cm}^3 \text{ STP/cm}^3$  in Group B samples and  $8.17 \times 10^{-10} \text{ cm}^3 \text{ STP/cm}^3$  in the  
244 Group C sample. Samples are featured by air-like Xe ratios, e.g.,  $^{132}\text{Xe}/^{130}\text{Xe}$  ratios vary from  
245  $6.47 \pm 0.27$  to  $7.22 \pm 0.69$  compared to the atmospheric value of  $6.6098 \pm 0.0023$  (Valkiers et al., 1998).

#### 246 4.2.5 Air contamination

247 The relationship between noble gas isotopic ratios and noble gas concentrations is an effective  
248 tool for estimating air contamination (Barry et al., 2016). Fig. 5 shows the relationship between  
249  $^{40}\text{Ar}/^{36}\text{Ar}$  and  $1/^{36}\text{Ar}$  for all samples (Fig. 5a) and samples only in Group A (Fig. 5b) due to large  
250 differences in  $^{36}\text{Ar}$  contents among the groups.  $^{40}\text{Ar}/^{36}\text{Ar}$  ratios show a similar relationship with  
251  $1/^{20}\text{Ne}$  values for all samples (not shown in figures). Given the adjacent sample locations of XP-1,  
252 XP-2 and XP-3 (maximum distance is 2.3 km) and horizontal strata in this area, it is reasonable to  
253 assume similar pristine noble gas signatures for gas samples from the three wells. Since the data of  
254 XP-2 and XP-3 are highly consistent with the mixing line of XP-1 and air in Fig. 5b, the two  
255 samples were deemed to be contaminated by air and regarded as ‘Outlier Samples’. In addition, the  
256  $^{20}\text{Ne}/^{36}\text{Ar}$  ratios of XP-2 and XP-3 (0.405 and 0.374) are also between the atmospheric value (0.542,  
257 Sano et al., 2013) and the ratio of XP-1 (0.295). The concentrations of radiogenic components ( $^4\text{He}$ ,  
258  $^{40}\text{Ar}^*$ ) in XP-2 and XP-3 are also lower than those of XP-1 (Table 2). Therefore, it is possible that  
259 these two samples were contaminated by air. Since the vacuum integrity of the sample preparation  
260 system is monitored by a pressure gauge and a quadrupole mass spectrometer, the air addition  
261 during sample processing in the lab can be excluded. Air contamination may have been acquired  
262 during sample collection. The following discussion excludes these two samples. There is no positive  
263 correlation between  $^{40}\text{Ar}/^{36}\text{Ar}$  ratios and  $1/^{36}\text{Ar}$  values for the remainder of the samples, suggesting  
264 that they are unaffected by air contamination.

265

## 266 5. DISCUSSION

### 267 5.1 Modelling of groundwater evolution

268 Atmosphere-derived noble gases (e.g.,  $^{20}\text{Ne}$ ,  $^{36}\text{Ar}$ ,  $^{84}\text{Kr}$  and  $^{130}\text{Xe}$ ) are dissolved into meteoric  
269 water under surface conditions. They are transported into the subsurface system by aquifer recharge.

270 The absence of other significant subsurface sources makes them ideal tracers for gas-oil-water  
271 fractionation in sedimentary basins (Pinti and Marty, 1995; Wen et al., 2015). In this study, noble  
272 gas data of Group A and B samples are used to identify phase interactions among oil, gas and  
273 groundwater. In addition, the volume ratios of the different phases are quantified as the amount of  
274 oil and gas present in a reservoir is key information for hydrocarbon exploration. In this study we  
275 also test open system versus closed system models, providing a tool to understand the extent of  
276 groundwater circulation and its influence on hydrocarbon and helium accumulation processes.

### 277 **5.1.1 Modelling conditions**

278 Noble gas concentrations and elemental ratios in air saturated water (ASW) were calculated  
279 using Henry's law under recharge conditions of 10°C, 0M NaCl and an average altitude of 1000m in  
280 the recharge area (North Qinling Belt). As a result, the initial  $^{20}\text{Ne}/^{36}\text{Ar}$  ratio,  $^{84}\text{Kr}/^{36}\text{Ar}$  ratio and  
281  $^{130}\text{Xe}/^{36}\text{Ar}$  ratio in ASW are taken to be 0.143,  $4.02 \times 10^{-2}$  and  $4.02 \times 10^{-4}$  respectively. The  
282 geothermal groundwater is extracted from an approximate depth of 2500m, where the temperature  
283 and pressure are taken to be 88°C and 25MPa respectively based on a surface temperature of 13°C,  
284 an average geothermal gradient of 30°C/km and a hydrostatic pressure gradient of 10MPa/km. The  
285 salinity of groundwater is taken to be 0.1 M NaCl equivalent based on the measured value of  
286 2.0-7.8g/L. The geothermal gradient, hydrostatic pressure gradient and groundwater salinity were  
287 measured by Shaanxi Geological Environment Monitoring Station. Henry's constants of noble  
288 gases used for modelling are calculated from empirical equations, corrected with fugacity  
289 coefficients and activity coefficients (Ballentine et al., 2002 and references therein). The calculated  
290 Henry's constants are listed in Table 3.

### 291 **5.1.2 Gas-water fractionation**

292 Solubilities of noble gases in water increase with mass ( $\text{Ne} < \text{Ar} < \text{Kr} < \text{Xe}$ ) (Ballentine et al.,  
293 2002), therefore Ne exsolves into the gas phase most readily whereas progressively heavier noble  
294 gases remain dissolved into groundwater. If the gas-water fractionation occurred in the groundwater  
295 within the Weihe Basin, the  $^{20}\text{Ne}/^{36}\text{Ar}$  ratios in the water phase would be lower than the initial  
296  $^{20}\text{Ne}/^{36}\text{Ar}$  ratio in ASW. However, all measured  $^{20}\text{Ne}/^{36}\text{Ar}$  ratios in Group A and B samples (0.195 -

297 0.382), which represent the noble gas signatures in the water phase, are higher than the ratio of  
298 ASW (0.143). This cannot be explained by the simple gas-water fractionation model.

299 The possibility of multi stage gas-water fractionation (re-solution and effervescence model,  
300 Ballentine et al., 2002) was also considered. A gas phase exsolved from the ASW would have an  
301 elevated  $^{20}\text{Ne}/^{36}\text{Ar}$  ratio. When the exsolved gases dissolve into a small volume of water, this  
302 process will lead to a significant increase of  $^{20}\text{Ne}/^{36}\text{Ar}$  ratio in the groundwater. Iterative exsolving  
303 and dissolving will further increase the  $^{20}\text{Ne}/^{36}\text{Ar}$  ratio in the water phase (Ballentine et al., 2002).  
304 Multi stage exsolution and dissolution models require variation in the pressure/temperature  
305 conditions in the subsurface, presumably due to repeated uplift and subsidence of strata. However,  
306 the Weihe Basin has been mainly subject to extension and settlement continuously since the  
307 Palaeogene (Mercier et al., 2013). This is inconsistent with the complex conditions needed for the  
308 re-solution and effervescence model. Therefore, we consider that the simple and multi stage  
309 gas-water fractionation models are not viable explanations for the noble gas data recorded from the  
310 Weihe Basin.

### 311 **5.1.3 Oil-water fractionation**

312 Oil has been found to adhere in the pipes of geothermal wells in the Gushi Depression and  
313 around 50kg of oil are produced from WR-11 and WR-2 through oil water separating equipment per  
314 day. Therefore, it is reasonable to consider a model in which the groundwater has been equilibrated  
315 with an oil phase earlier during its migration. In addition, the involvement of oil can explain the  
316 high  $^{20}\text{Ne}/^{36}\text{Ar}$  value in groundwater since Ar relative to Ne has higher solubility in oil than water  
317 (Battani et al., 2000; Zhou et al., 2012). If the oil-water equilibrium process occurs in either a closed  
318 system, batch fractionation of noble gases occurs. Conversely, open system equilibration will result  
319 in a Rayleigh fractionation pattern of noble gases. Both cases are modelled in this study assuming  
320 an oil API of 25 and 34 respectively.

#### 321 *Batch fractionation.*

322 When equilibrium has been achieved under closed system conditions, the elemental ratios of  
323 noble gases follow the batch fractionation equation (Ballentine et al., 2002):

$$\left(\frac{A}{B}\right)_{water} = \left(\frac{A}{B}\right)_{ASW} \times \left( \frac{\frac{V_{water} \rho_{water} + (K_B)_{water}}{V_{oil} \rho_{oil} + (K_B)_{oil}}}{\frac{V_{water} \rho_{water} + (K_A)_{water}}{V_{oil} \rho_{oil} + (K_A)_{oil}}} \right) \quad (1)$$

Where subscripts *ASW* and *water* represent the original ASW water and water phase after oil-water equilibrium;  $V_{oil}$  and  $V_{water}$  are the volumes of oil and groundwater involved in the system;  $\rho_{oil}$  is the density of oil, which is taken to be 0.9 g/cm<sup>3</sup> and 0.85 g/cm<sup>3</sup> for API=25 and 34 respectively;  $\rho_{water}$  is the density of groundwater as 1.0g/cm<sup>3</sup>;  $K_{A(water)}$ ,  $K_{B(water)}$  are Henry's constants of species A and B in the groundwater;  $K_{A(oil)}$ ,  $K_{B(oil)}$  are Henry's constants of A and B in the oil. In this study, species A includes <sup>20</sup>Ne, <sup>84</sup>Kr or <sup>130</sup>Xe and species B is <sup>36</sup>Ar.

*Rayleigh fractionation.*

A simple Rayleigh fractionation law is used to model oil-water equilibrium in an open dynamic system (Battani et al., 2000):

$$\left(\frac{A}{B}\right)_{water} = \left(\frac{A}{B}\right)_{ASW} f^{(\alpha-1)} \quad (2)$$

$$\alpha = \frac{\left(\frac{(K_A)_{water}}{(K_B)_{water}}\right)}{\left(\frac{(K_A)_{oil}}{(K_B)_{oil}}\right)} \quad (3)$$

Where  $f$  is the fraction of species B remaining in the water phase;  $\alpha$  is the fractionation coefficient given by Eq. (3); other parameters are the same as those in Eq. (1). Modelling results of both fractionation models are shown in Fig. 6.

Starting with an ASW ratio of <sup>20</sup>Ne/<sup>36</sup>Ar = 0.143, <sup>84</sup>Kr/<sup>36</sup>Ar = 4.02 × 10<sup>-2</sup> and <sup>130</sup>Xe/<sup>36</sup>Ar = 4.02 × 10<sup>-4</sup>, the Batch fractionation show a maximum <sup>20</sup>Ne/<sup>36</sup>Ar ratio in groundwater when the  $V_{oil}/V_{water}$  ratio approaches infinity. The maximum <sup>20</sup>Ne/<sup>36</sup>Ar in groundwater are 0.271 and 0.392 respectively when equilibrating with a light oil phase (API=34) and a heavy oil phase (API=25). From Fig. 6, it is clear that light oil-water fractionation process in a closed system cannot account for all the measured <sup>20</sup>Ne/<sup>36</sup>Ar ratios (0.195 - 0.382). Rayleigh fractionation occurring in a dynamic system, there are no limiting values for elemental ratios when  $V_{oil}/V_{water}$  ratio approaches infinity. Although all three fractionation models (heavy oil-water batch fractionation, heavy oil-water Rayleigh fractionation, and light oil-water Rayleigh fractionation) can explain the measured <sup>20</sup>Ne/<sup>36</sup>Ar ratios,

348 it is apparent that the better fit for the measured ratios is the Rayleigh heavy oil-water fractionation  
 349 with a maximum additional 600% Xe and 100% Kr. Fig. 6 shows that the observed variation in  
 350 measured noble gas concentrations is most consistent with the curvilinear trend of Rayleigh heavy  
 351 oil-water fractionation line. All the  $^{20}\text{Ne}/^{36}\text{Ar}$ ,  $^{84}\text{Kr}/^{36}\text{Ar}$  and  $^{130}\text{Xe}/^{36}\text{Ar}$  ratios are accounted for by  
 352 the Rayleigh heavy oil-water fractionation line and with 600% excess Xe and 100% excess Kr.

353 Previous studies have shown similar Kr and Xe enrichment in natural gas and CO<sub>2</sub> gas fields.  
 354 For example, the  $(^{132}\text{Xe}/^{36}\text{Ar})_{\text{meas.}}/(^{132}\text{Xe}/^{36}\text{Ar})_{\text{air}}$  value is up to 576 in gas separated from  
 355 oil-associated gases from the Elk Hill oil fields, California (Torgersen and Kennedy, 1999) and is  
 356 approximately 100 in coal bed gases from San Juan Basin (Zhou et al., 2005). Similar heavy noble  
 357 gas enrichment is also seen in CO<sub>2</sub> gas from the Colorado Plateau and Rocky Mountain provinces,  
 358 USA (Gilfillan et al., 2008), natural gas from the Sleipner Vest field (Barry et al., 2016) and stray  
 359 gas preserved in the Trinity aquifer, North-Central Texas (Wen et al., 2017). Heavier noble gas  
 360 species have been suggested to preferentially adsorb onto organic-rich minerals (Podosek et al.,  
 361 1981). When buried, these sediments desorb at high temperatures and heavy noble gases are  
 362 released into the groundwater or free gas phase, causing the enrichment of Kr and Xe relative to Ar  
 363 in deep fluids (Zhou et al., 2005; Barry et al., 2016). Therefore, various excess of Xe and Kr  
 364 observed in this study are caused by the variable distribution of organic-rich material.

#### 365 **5.1.4 Oil and groundwater volume ratio in the Weihe Basin**

366 The volume ratio of oil to water is a key parameter to assess oil generation over geological  
 367 time. The  $V_{\text{oil}}/V_{\text{water}}$  ratio can be calculated based on the model of heavy oil-water Rayleigh  
 368 fractionation with heavy noble gas excesses. It is suggested that step-wise gas-water equilibrium  
 369 can approximate Rayleigh fractionation if  $V_{\text{gas}}/V_{\text{water}}$  is small in each stage (Zhou et al., 2005). The  
 370 principle is the same for the oil-water equilibrium.  $V_{\text{oil}}/V_{\text{water}}$  in one stage equilibration between oil  
 371 and water phase is given by Battani et al. (2000):

$$372 \frac{V_{\text{oil}}}{V_{\text{water}}} = \frac{\rho_{\text{water}}}{\rho_{\text{oil}}} \frac{(K_{\text{Ar}})_{\text{oil}}}{(K_{\text{Ar}})_{\text{water}}} \left( \frac{1-f}{f} \right) \quad (4)$$

373 Where  $V_{\text{oil}}$  is the oil volume equilibrating with the groundwater phase;  $V_{\text{water}}$  is the groundwater

374 volume involved in oil-water equilibration;  $\rho_{oil}$  and  $\rho_{water}$  are the density of oil and groundwater,  
375 which are taken to be  $0.9\text{g/cm}^3$  and  $1.0\text{ g/cm}^3$  respectively.  $(K_{Ar})_{water}$  and  $(K_{Ar})_{oil}$  are Henry's  
376 constants of  $^{36}\text{Ar}$  in the groundwater and oil phase respectively.  $f$  is the fraction of  $^{36}\text{Ar}$  remaining in  
377 the residual groundwater.

378 If a small volume of oil ( $V_{oil}/V_{water} = 3 \times 10^{-5}$  is chosen, Zhou et al., 2005) equilibrates with  
379 groundwater, we can calculate the fraction of  $^{36}\text{Ar}$  remaining in the groundwater phase in each  
380 single stage ( $f_{single}$ ). Since excess Kr and Xe have significant impact on the  $^{84}\text{Kr}/^{36}\text{Ar}$  ratio and  
381  $^{130}\text{Xe}/^{36}\text{Ar}$  ratio (Section 5.2.3),  $^{20}\text{Ne}/^{36}\text{Ar}$  ratios are selected to calculate the final fraction of  $^{36}\text{Ar}$  ( $f$ )  
382 remaining in the groundwater after oil-water equilibrium calculated by Eq. (3). The number of  
383 single stages of oil/water equilibration ( $n$ ) can be calculated by  $f = f_{single}^n$  then the total  $V_{oil}/V_{water}$   
384 equals  $n \times 3 \times 10^{-5}$ . The calculated  $V_{oil}/V_{water}$  ratios range between 0.06 and 0.18 (Table 4). Because  
385 total degassing during sampling is assumed and the slight elevation of  $^{20}\text{Ne}/^{36}\text{Ar}$  in sample gas than  
386 that in groundwater is not taken into account, the derived  $V_{oil}/V_{water}$  values are maxima.

387 Based on the total static reserve of geothermal fluids beyond 4000m in depth in the Weihe  
388 Basin ( $14781.20 \times 10^8\text{ m}^3$ , Zhang et al., 2018b and references therein), the oil volume equilibrating  
389 with present groundwater varies from  $887 \times 10^8\text{ m}^3$  to  $2661 \times 10^8\text{ m}^3$ . This suggests that abundant oil  
390 has been generated in Weihe Basin as these values are even higher than that of URR (ultimately  
391 recoverable resources) standard of a mega-giant oil field ( $78.25 \times 10^8\text{ m}^3$ , Ivanhoe and Leckie, 1993).  
392 Given the fact that there are three Palaeogene-Neogene combinations of reservoir and covering  
393 strata (Li et al., 2013), it is likely that oil has accumulated in the lithologic traps in the Weihe Basin.

## 394 5.2 $^4\text{He}$ accumulation

### 395 5.2.1 Relationship between $^4\text{He}$ and groundwater

396  $^{20}\text{Ne}$  in subsurface fluids is derived from ASW without other sources. Since He and Ne have  
397 similar Henry's constants in water and oil (Ballentine et al., 2002), the water-oil-gas fractionation  
398 results in little fractionation between them. Therefore,  $^4\text{He}/^{20}\text{Ne}$  is an excellent parameter to study  
399 the relationship between  $^4\text{He}$  accumulation and groundwater movement.

400 There is a positive correlation between  $^4\text{He}$  and  $^{20}\text{Ne}$  ( $R^2=0.84$ , Fig. 7) for samples in Group A

401 and B with  $^4\text{He}/^{20}\text{Ne}$  ratios varying from 5113 to 9318 except for WR-2 (1439). This is consistent  
 402 with the assumption that  $^4\text{He}$  and  $^{20}\text{Ne}$  were both exsolved from the water phase during sampling.  
 403 As the  $^{20}\text{Ne}/^{36}\text{Ar}$  ratio of WR-2 (0.287) is consistent with those of others (0.195 - 0.382), it excludes  
 404 the air-contamination possibility for the sample (this is also described in section 4.2.5). The lower  
 405  $^4\text{He}/^{20}\text{Ne}$  ratio of WR-2 suggests that some younger groundwater preserved with less or no crustal  
 406  $^4\text{He}$  mixed with the older groundwater. WR-HX has an extremely high  $^4\text{He}/^{20}\text{Ne}$  ratio (37164)  
 407 compared with samples in Group A and B. This ratio is similar to that observed in helium-rich  
 408 Hugoton-Panhandle gas field (24871 - 46023, calculated from Ballentine and Lollar, 2002). The  
 409 data from Hugoton-Panhandle field are shown as squares in Fig. 7. Since groundwater is  
 410 widely distributed in the subsurface and He is unlikely to be transported by advective flow alone  
 411 due to its low abundance in the geological environment (Ballentine and Burnard, 2002), it is  
 412 reasonable to expect that He can migrate in the subsurface together with groundwater movement.  
 413 The high  $^4\text{He}/^{20}\text{Ne}$  ratio of WR-HX suggests a high  $^4\text{He}$  flux or low groundwater flow rate. There is  
 414 no evidence that a low rate of groundwater flow occurs in the area where the WR-HX well is  
 415 located. Also, WR-HX is located near the uranium and thorium-rich Huashan granitic body.  
 416 Therefore, it is reasonable to attribute the high  $^4\text{He}/^{20}\text{Ne}$  ratio of WR-HX to the high  $^4\text{He}$  flux in this  
 417 area.

### 418 5.2.2 $^4\text{He}$ concentrations in groundwater

419 The initial  $^4\text{He}$  concentrations in groundwater are good parameters to quantify the relationship  
 420 between  $^4\text{He}$  and groundwater. For Group A and B samples, when calculating the initial  $^4\text{He}$   
 421 concentrations in groundwater, the volume ratio of gas to groundwater is needed, which can be  
 422 calculated by:

$$423 \frac{V_{gas}}{V_{water}} = \frac{(C_{36Ar})_{water}}{(C_{36Ar})_{gas}} \quad (5)$$

$$424 (C_{36Ar})_{water} = (C_{36Ar})_{ASW} \times f \quad (6)$$

425 Where subscripts *water*, *gas* represent the water phase and the gas phase collected from



426 wellheads during sampling.  $(C_{36Ar})_{ASW}$  is the concentration of  $^{36}Ar$  in original water (ASW).  $f$  is  
427 the fraction of  $^{36}Ar$  remaining in the residual groundwater after oil-water fractionation, which can  
428 be calculated by Eq. (2). The calculated  $V_{gas}/V_{water}$  ratios (under STP conditions) vary from  
429 0.02-0.08 for Group A samples and 1.22-1.38 for Group B samples (Table 4). These vaules provide  
430 maximum ratios especially for Group A samples when considering the slight incomplete degassing  
431 during sampling caused by the low  $V_{gas}/V_{water}$  ratios. The initial helium concentrations in  
432 groundwater before oil-water fractionation for Group A and Group B samples can be calculated  
433 based on the oil-gas fractionation model,  $V_{gas}/V_{water}$  ratios in the subsurface and the  $^4He$   
434 concentrations in sampled gases (Table 2). Although the  $^4He$  concentrations in Group A samples is  
435 higher than those in Group B samples by one order of magnitude, the initial helium concentrations  
436 in groundwater are similar for Group A samples ( $2.36 - 15.4 \times 10^{-4} \text{ cm}^3 \text{ STP/g}$  and  $9.05 \times 10^{-4} \text{ cm}^3$   
437  $\text{STP/g}$  in average) and Group B samples ( $9.06$  and  $12.2 \times 10^{-4} \text{ cm}^3 \text{ STP/g}$ ). The low concentrations of  
438 He measured in Group B gas samples can be caused by addition of  $CH_4$ , i.e., the addition of major  
439 gas has a strong dilution effect on helium concentration.

440 If assuming the gas sample collected at well WR-HX was completely dissolved in groundwater  
441 in the subsurface, based on the  $V_{gas}/V_{water}$  calculation formula (equation 5), the observed 5-12 times  
442 higher  $^{36}Ar$  concentration in the WR-HX sample than those in Group B samples would suggest the  
443 5-12 times lower  $V_{gas}/V_{water}$  ratio at well WR-HX than Group B wells. However, the gas content of  
444 WR-HX well was much larger than in Group B wells, implying that the collected WR-HX gas is not  
445 exsolved from groundwater completely and at least part of gas in the WR-HX well is derived from a  
446 free gas phase in the subsurface. The high  $^{36}Ar$  concentration in the gas reservoir at WR-HX can  
447 then be explained by the exchange of gas contents between groundwater and the gas reservoir. This  
448 process would continuously extract  $^{36}Ar$  and  $^4He$  from the groundwater into the gas phase and  
449 dissolve  $CH_4$  from the reservoir into groundwater during the migration of the groundwater.

450 Since the sampled gas of WR-HX is a mixture of free gas and dissolved gas, two extreme cases  
451 are considered to calculate the initial  $^4He$  concentrations in groundwater: (1) If the WR-HX sample

452 is derived entirely from groundwater, the initial  $^4\text{He}$  concentration in groundwater is calculated to  
453 be  $6.13 \times 10^{-3} \text{ cm}^3 \text{ STP/g}$  following the same calculation procedures as Group A and B samples; (2)  
454 If the WR-HX sample is taken from the gas reservoir completely, the noble gas signatures in the  
455 WR-HX reservoir are the result of a gas-water equilibrium process. Although it is possible that  
456 groundwater in the WR-HX well has equilibrated with an oil phase like Group A and Group B  
457 groundwater before contacting the gas reservoir (section 5.1), the involvement of an oil phase is  
458 neglected since the  $^{20}\text{Ne}/^{36}\text{Ar}$  ratio of WR-HX reservoir (0.198) is only slightly higher than the ratio  
459 of ASW (0.143), which can be account for by a gas-water fractionation process. Because it is an  
460 open system for Group A and Group B groundwater, we assume that the gas reservoir has  
461 undergone gas-water equilibrium in an open system as well. Based on the Rayleigh fractionation  
462 among  $^4\text{He}$ ,  $^{20}\text{Ne}$  and  $^{36}\text{Ar}$  during the gas-water equilibrium under reservoir conditions, the  $^4\text{He}$   
463 concentration in groundwater that is in contact with the gas reservoir for Group C is calculated to be  
464  $6.39 \times 10^{-3} \text{ cm}^3 \text{ STP/g}$ . No matter gas in WR-HX sample is dissolved in groundwater or from free gas  
465 phase underground, there is little difference of  $^4\text{He}$  concentrations in initial groundwater ( $6.13 \times 10^{-3}$   
466 and  $6.39 \times 10^{-3} \text{ cm}^3 \text{ STP/g}$  respectively). In addition, due to the existence of free gas phase in  
467 WR-HX area, the latter is chosen in the following calculation. The helium content in the  
468 groundwater associated with Group C sample is 4-27 times higher than those in groundwater  
469 associated with Group A and B samples. Similarly, radiogenic  $^{40}\text{Ar}^*$  content in groundwater of  
470 Group C is around 3-28 times of those of groundwater of Group A and Group B. Therefore, there  
471 should be a background  $^4\text{He}$  and  $^{40}\text{Ar}^*$  flux for Group A & B samples and an additional  $^4\text{He}$  and  
472  $^{40}\text{Ar}^*$  flux for Group C sample.

### 473 **5.2.3 $^4\text{He}$ flux**

474  $^4\text{He}$  is generated from the radiogenic decay of  $^{235,238}\text{U}$  and  $^{232}\text{Th}$  (Ballentine and Burnard,  
475 2002). The  $^4\text{He}$  in groundwater can be derived from both in-situ production and external flux  
476 (Torgersen, 1980; Torgersen and Clarke, 1985; Zhou and Ballentine, 2006). The in-situ  $^4\text{He}$   
477 production can be expressed as (Torgersen, 1980):

478 
$$[{}^4\text{He}]_{\text{in situ production}} = \frac{\rho \Delta J_4 (1 - \varphi)}{\varphi} t \quad (9)$$

479 Where  $\rho$  is the density of aquifer in g/cm<sup>3</sup>;  $\Delta$  is the <sup>4</sup>He transfer efficiency from rock matrix to  
 480 groundwater;  $\varphi$  is the aquifer porosity;  $t$  is groundwater residence time in year.  $J_4$  is radioactive  
 481 production of <sup>4</sup>He in cm<sup>3</sup> STP <sup>4</sup>He/g<sub>rock</sub> year:

482 
$$J_4 = 0.2355 \times 10^{-12} [U](1 + 0.123[Th]/[U] - 4) \quad (10)$$

483 Where  $[U]$  and  $[Th]$  are the U and Th concentrations in rocks in ppm.

484 The steady-state external flux of <sup>4</sup>He into groundwater can be expressed as (Zhou and  
 485 Ballentine, 2006):

486 
$$[{}^4\text{He}]_{\text{external flux}} = \frac{J_4 \cdot \rho \cdot H}{\varphi h} t \quad (11)$$

487 Where  $J_4$  is radioactive production of <sup>4</sup>He given by Eq. (9) in cm<sup>3</sup> STP <sup>4</sup>He/g<sub>rock</sub> year;  $\rho$  is  
 488 average crust density in g/cm<sup>3</sup>;  $H$  is the average crust thickness in km;  $h$  is the aquifer thickness in  
 489 km;  $\varphi$  is the aquifer porosity;  $t$  is groundwater residence time in year.

490 The parameters used for calculation of <sup>4</sup>He accumulation rates in groundwater are listed in  
 491 Table 5. The values for average crust compositions are taken to be Clark values, cited from Taylor  
 492 and McLennan (1985). U and Th concentrations of the aquifer are the Clark values as well due to  
 493 the absence of values in the local rocks (Table 5). The <sup>4</sup>He accumulation rate in groundwater from  
 494 in-situ production and steady-state external flux (upper crust and lower crust) is  $7.79 \times 10^{-10}$  cm<sup>3</sup>  
 495 STP/g year. The existing <sup>4</sup>He concentrations in groundwater for Group A and B samples ( $2.36 -$   
 496  $15.4 \times 10^{-4}$  cm<sup>3</sup> STP/g) give groundwater ages of 0.30-1.98Ma (assuming 100% releasing and  
 497 transfer efficiency from the crust to water), which are consistent with ages of 0.3-1.3 Ma, derived  
 498 using the <sup>81</sup>Kr dating method (Li et al., 2017). The ages are realistic because the calculated <sup>4</sup>He  
 499 groundwater ages are slightly younger than the age of the main geothermal reservoir (Lantian-Bahe  
 500 Formation), which was deposited during the Pliocene from 2.58Ma to 5.33Ma.

501 Similarly, the accumulation of helium in groundwater in the WR-HX well (Group C) requires  
 502 timescales on the order of 8.2 million years, which is clearly not valid since the groundwater age is  
 503 older than the aquifer age. Therefore, there must be an additional <sup>4</sup>He flux for the Group C sample,

504 which is consistent with the results from the  $^4\text{He}/^{20}\text{Ne}$  ratio in section 5.2.1. The contribution of  
505 Huashan granitic body to the  $^4\text{He}$  accumulation for the Group C sample is considered since it is in  
506 close proximity to the sample location. If the average helium accumulation duration of Group B  
507 (1.37Ma) is chosen for the Group C sample, the total amount of helium accumulation derived from  
508 aquifer and crust is only  $1.07 \times 10^{-3} \text{ cm}^3 \text{ STP/g}$ . This suggests that the additional  $^4\text{He}$  flux from the  
509 Huashan granitic body accounts for 83.3% of the total  $^4\text{He}$  flux.

510 Since we assume 100% releasing and transfer efficiency for  $^4\text{He}$  from crust to groundwater, the  
511 ages are minimum estimates. Some helium is still preserved in rocks (Tolstikhin et al., 1996;  
512 Tolstikhin et al., 2011) or lost during migration. If the maximum groundwater age in Weihe Basin  
513 was the same as the aquifer age (2.58Ma), the maximum release and transfer coefficient of  $^4\text{He}$   
514 from crust and aquifer to groundwater would be calculated to be 76.7%. Alternatively, episodic  
515 release of  $^4\text{He}$  accumulated in ancient rocks over longer time periods can occur, as inferred for  
516 radiogenic He release in Yellowstone (Lowenstern et al., 2014). These authors interpreted high  
517 helium concentrations to be the result of helium accumulated in Archaean cratonic rocks (more than  
518 2.5 billion years old) liberated over the past two million years by intense crustal metamorphism.  
519 The Cenozoic Weihe Basin is a tectonically active basin with many fractures and recorded  
520 destructive earthquakes (Mercier et al., 2013). Therefore, it is possible that a portion of  $^4\text{He}$  was  
521 generated in the crust before the formation of the aquifer, and was released and transferred to the  
522 groundwater. This process would produce a high  $^4\text{He}$  flux and result in maximum estimates for  $^4\text{He}$   
523 accumulation ages.

### 524 **5.3 The concentration and source of $\text{N}_2$ traced by noble gases**

525 Based on the volumetric ratios,  $\text{N}_2$  contents in groundwater associated with Group A samples  
526 and Group B samples are calculated to be 0.020-0.060  $\text{cm}^3\text{STP/g}$  and 0.067-0.340  $\text{cm}^3\text{STP/g}$   
527 respectively. Similarly, the concentrations of  $\text{CH}_4$  in groundwater are calculated to be 0.001-0.009  
528  $\text{cm}^3\text{STP/g}$  for Group A samples and 0.648-0.959  $\text{cm}^3\text{STP/g}$  for Group B samples (Table 4),  
529 suggesting more potential hydrocarbon resources in the Group B area than A. The carbon isotopes  
530 of  $\text{CH}_4$  in samples range between -19.04‰ and -44.46‰ (VPDB) and fall within the expected range

531 for coal-derived methane (Dai et al., 1992).

532 The source of N<sub>2</sub> in groundwater is considered. The N<sub>2</sub>/<sup>36</sup>Ar ratio in original ASW is calculated  
533 to be 1.14×10<sup>4</sup> based on Henry's constants of N<sub>2</sub> and <sup>36</sup>Ar under recharge conditions. After the  
534 oil-water Rayleigh fractionation discussed in section 5.1.3, the residual N<sub>2</sub>/<sup>36</sup>Ar ratios in  
535 groundwater vary from 1.33×10<sup>4</sup> to 1.84×10<sup>4</sup>. The Henry's constants for N<sub>2</sub> in groundwater at  
536 recharge and reservoir conditions are calculated from Wilhelm et al. (1977) and that in oil at  
537 reservoir conditions is calculated based on the solubility of MK-8 oil in Logvinyuk et al. (1970)  
538 (Table 4). The measured N<sub>2</sub>/<sup>36</sup>Ar ratio varies from 5.20×10<sup>4</sup> to 4.62×10<sup>5</sup>, which means that only  
539 13.2-30.3% of N<sub>2</sub> in Group A samples and 2.9% - 8.0% in Group B samples are derived from  
540 air-saturated water. As the helium isotope ratios are clearly crustal, it is reasonable to exclude a  
541 significant contribution from mantle source for N<sub>2</sub>. In addition, N<sub>2</sub> can also be released from  
542 organic matter during thermal decomposition or from sediments and basement during  
543 metamorphism (Danabalan, 2017).

## 544 **5.4 Helium accumulation model and conditions**

### 545 **5.4.1 Helium accumulation model in the Weihe Basin**

546 The relationship between different sample groups and the constraints on fluid evolution history  
547 from noble gas partition modelling are shown in Fig. 8, which illustrates the accumulation of <sup>4</sup>He,  
548 CH<sub>4</sub>, N<sub>2</sub> and ASW-derived noble gases in the Weihe Basin. Specifically, the process includes four  
549 stages:

550 ① <sup>20</sup>Ne, <sup>36</sup>Ar, other ASW-derived noble gases and N<sub>2</sub> (2.9% - 30.3% of total N<sub>2</sub>) from air  
551 dissolved into surface water at recharge conditions of 10°C, 0M NaCl and an average altitude of  
552 1000m (North Qinling Belt) were incorporated.

553 ② Crustal radiogenic noble gases (e.g., <sup>4</sup>He and <sup>40</sup>Ar\*) were transferred to the groundwater on  
554 a time scale between 0.3 and 1.98 million years. This assumes the releasing and transfer efficiency  
555 from source rock to groundwater of <sup>4</sup>He is 100%, therefore the timescales for <sup>4</sup>He accumulation  
556 could be longer if efficiency is lower. On the other hand, <sup>4</sup>He accumulation ages could be shorter if

557 there were episodic release of  $^4\text{He}$  accumulated in ancient rocks, which would produce a high  $^4\text{He}$   
558 flux.

559 ③ Groundwater underwent heavy oil-groundwater Rayleigh fractionation with  $V_{\text{oil}}/V_{\text{water}}$   
560 being 0.06-0.18. This process caused an increase of  $^{20}\text{Ne}/^{36}\text{Ar}$  ratio (0.195-0.382) in groundwater to  
561 values higher than that in ASW (0.143).

562 ④ Addition of different amount of major gases ( $\text{CH}_4$  and  $\text{N}_2$ ) resulted in various signatures in  
563 the three groups of samples. There are three scenarios.

564 Group A: Minor  $\text{CH}_4$  and nitrogen released from organic matter, sediments and/or basement  
565 were dissolved into groundwater, result in  $\text{CH}_4$  and  $\text{N}_2$  concentrations being 0.001-0.009  $\text{cm}^3\text{STP/g}$   
566 and 0.020-0.060  $\text{cm}^3\text{STP/g}$  respectively. The total dissolved gas amount is low ( $V_{\text{gas}}/V_{\text{water}}$  ratios are  
567 0.02-0.08), having a minor dilution effect on noble gases in groundwater and causing the high  
568 concentration of  $^4\text{He}$  (32.3 to  $294 \times 10^{-4} \text{ cm}^3 \text{ STP/cm}^3$ ), high concentration of  $^{20}\text{Ne}$  (15.9 to  $39.5$   
569  $\times 10^{-7} \text{ cm}^3 \text{ STP/cm}^3$ ) in samples.

570 Group B: Compared with Group A, more  $\text{CH}_4$  and  $\text{N}_2$  were generated and dissolved into Group  
571 B groundwater, making  $\text{CH}_4$  and  $\text{N}_2$  concentrations in groundwater being 0.648-0.959  $\text{cm}^3\text{STP/g}$   
572 and 0.067-0.340  $\text{cm}^3\text{STP/g}$  respectively. The gas amount dissolved in groundwater is high  
573 ( $V_{\text{gas}}/V_{\text{water}}$  ratios are 1.22-1.38) caused a major dilution effect on He and other noble gases  
574 concentrations. This process resulted in the low concentrations of  $^4\text{He}$  (5.74 and  $6.21 \times 10^{-4} \text{ cm}^3$   
575  $\text{STP/cm}^3$ ), low concentrations of  $^{20}\text{Ne}$  (0.69 to  $1.11 \times 10^{-7} \text{ cm}^3 \text{ STP/cm}^3$ ) in samples.

576 Group C: Abundant  $\text{CH}_4$  and  $\text{N}_2$  saturated the groundwater and formed a free gas reservoir.  
577 Then continuous gas-water interaction caused partition of less soluble noble gases (e.g.,  $^4\text{He}$ ,  $^{20}\text{Ne}$   
578 and  $^{36}\text{Ar}$ ) into the gas phase. This resulted in the high concentration of He (2.11%) in the WR-HX  
579 sample compared with those in the Group B samples. The high  $^4\text{He}$  flux of the WR-HX sample can  
580 be accounted for by a background  $^4\text{He}$  flux shown in Group A & B samples and an additional  $^4\text{He}$   
581 flux possibly coming from adjacent granitic bodies, which increased the accumulation rate of He.

## 582 **5.4.2 Conditions for helium accumulation**

583 The most fundamental condition for high helium accumulation is high He flux. The primary  
584 reason why the Group C sample has a high helium concentration is the additional helium flux from  
585 the adjacent Huashan granitic body. The granitic bodies and other uranium-rich and thorium-rich  
586 rocks are suitable helium source rocks. Therefore, the wells close to these rocks are more likely to  
587 accumulate helium. In addition, episodic releasing of  $^4\text{He}$  preserved in rocks during its geological  
588 history could increase He flux significantly during tectonic movement. The rift valley area with  
589 good trap conditions could potentially accumulate significant amounts of helium, as observed in the  
590 region surrounding the Tanzanian Craton (Danabalan, 2017).

591 The second condition for economic helium accumulation is the existence of a free gas phase of  
592 the major gas component. If there is a gas reservoir (associated with the Group C sample in this  
593 study), helium and other noble gases dissolved in groundwater would exsolve into a gas phase  
594 continuously rather than migrate elsewhere due to groundwater flow (Group A and B). When  
595 groundwater migrates through the reservoir, the free gas phase works like a “filter screen” to  
596 concentrate noble gases, especially the less soluble helium, into the gas phase. The different helium  
597 concentrations among the groups in our study suggest that the free gas phase plays an important role  
598 in accumulating less soluble helium. Dissolved helium is less commercially viable than free helium  
599 gas since the production of dissolved gases in groundwater from a geothermal field is generally  
600 lower than gas production from a natural gas field.

601 Finally, He accumulation requires little or no major gas (e.g.,  $\text{CH}_4$ ,  $\text{N}_2$  and  $\text{CO}_2$ ) supplement  
602 after the formation of the free gas phase. The early gas phase in contact with groundwater would  
603 acquire most of the helium. If significant major gas migrated along the same pathway into a free gas  
604 reservoir, this would dilute the high helium concentration in the gas reservoir formed in the early  
605 stages (Brown, 2010). Since the total He content is principally controlled by the He flux and  
606 accumulation time, if there was large amount of major gas added, the helium concentration in the  
607 gas reservoir or groundwater would be diluted to below industrial grade (0.05 - 0.1 vol%).

608

609 **6 CONCLUSIONS**

610 Helium is a vital and indispensable resource in the fields of nuclear magnetic resonance  
611 (NMR), high-tech industry and scientific research. Although, there is an impending helium shortage  
612 globally, it has not been widely recognized that helium is an important and valuable by-product in  
613 natural gas and geothermal groundwater extraction. The extent of and controls on helium  
614 accumulation in geothermal or petroliferous basins are still poorly understood.

615 In this contribution, groundwater evolution and helium accumulation processes were modelled  
616 in the Weihe Basin geothermal field, central China using major gas contents, noble gas components  
617 and isotopic ratios, and carbon isotopic ratios. The 12 gas samples can be classified into three  
618 groups with distinctive properties, of which: Group A has high concentrations of N<sub>2</sub> and He; Group  
619 B has high contents of CH<sub>4</sub> and low contents of He; and Group C has a high content of CH<sub>4</sub> and a  
620 low content of He. Noble gas isotopes are predominantly radiogenic and ASW-derived. Modelling  
621 suggests that Group A and B samples have undergone a heavy oil-water fractionation in an open  
622 system with 600% excess Xe and 100% excess Kr with calculations suggesting there are 887 - 2661  
623 × 10<sup>8</sup> m<sup>3</sup> oil generated in the basin during its the geological history.

624 Current average <sup>4</sup>He concentrations dissolved in groundwater for Group A and B samples  
625 require the transfer of all helium produced from crust to groundwater over the last 0.30Ma-1.98Ma.  
626 The free gas phase is deemed to exist in the subsurface of the WR-HX area (Group C), where the  
627 Huashan granitic body provides an additional <sup>4</sup>He flux. Based on the elemental ratios of <sup>4</sup>He/<sup>20</sup>Ne,  
628 N<sub>2</sub>/<sup>36</sup>Ar and contents of ASW-derived noble gases, CH<sub>4</sub> and N<sub>2</sub>, the accumulation processes of  
629 helium and hydrocarbon in the three sample groups are modelled, which can be explained by a  
630 4-stage process.

631 There are 3 major controls on the helium accumulation in general. The most important factor is  
632 a high He flux, which controls the He accumulation rate. The second factor is the existence of a free  
633 gas phase of major gas components (e.g., CH<sub>4</sub>, N<sub>2</sub>, CO<sub>2</sub>), which can strongly apportion helium into  
634 the gas phase. Finally, a minimal amount of major gas should be supplied after the formation of the  
635 free gas phase since they have a dilution effect on the helium concentration.

636



637 **ACKNOWLEDGMENTS**

638 This research has been supported by National Natural Science Foundation in China (No.  
639 41572131). We thank China Scholarship Council (CSC) for offering the opportunity to collaborate  
640 with Lancaster University and University of Manchester. We acknowledge the team of Prof.  
641 Xingyuan Wang of Weihe Energy Company for the assistance during sampling. This work has  
642 benefited from discussions with Junlin Zhou, Qiao Zhang, Chunhui Cao. This paper has been  
643 greatly improved by detailed comments and helpful suggestions from associated editor Dr. Chris M.  
644 Hall and two anonymous reviewers.

645

646 **REFERENCES**

- 647 Ballentine C.J., O'nions R., Oxburgh E., Horvath F. and Deak J. (1991) Rare gas constraints on  
648 hydrocarbon accumulation, crustal degassing and groundwater flow in the Pannonian Basin.  
649 *Earth Planet. Sci. Lett.* **105**, 229-246.
- 650 Ballentine C.J., Burgess R. and Marty B. (2002) Tracing fluid origin, transport and interaction in the  
651 crust. *Rev. Mineral. Geochem.* **47**, 539-614.
- 652 Ballentine C.J. and Burnard P.G. (2002) Production, Release and Transport of Noble Gases in the  
653 Continental Crust. *Rev. Mineral. Geochem.* **47**, 481-538.
- 654 Ballentine C.J. and Lollar B.S. (2002) Regional groundwater focusing of nitrogen and noble gases  
655 into the Hugoton-Panhandle giant gas field, USA. *Geochim. Cosmochim. Acta* **66**, 2483-2497.
- 656 Barry P.H., Lawson M., Meurer W.P., Danabalan D., Byrne D.J., Mabry J.C. and Ballentine C.J.  
657 (2017) Determining fluid migration and isolation times in multiphase crustal domains using  
658 noble gases. *Geology* **45**, 775-778.
- 659 Barry P.H., Lawson M., Meurer W.P., Warr O., Mabry J.C., Byrne D.J. and Ballentine C.J. (2016)  
660 Noble gases solubility models of hydrocarbon charge mechanism in the Sleipner Vest gas field.  
661 *Geochim. Cosmochim. Acta* **194**, 291-309.

- 662 Bottomley D.J., Ross J.D., Clarke W.B. (1984) Helium and neon isotope geochemistry of some  
663 ground waters from the Canadian Precambrian Shield. *Geochim. Cosmochim. Acta* **48**,  
664 1973-1985.
- 665 Brown A.A. (2010) Formation of High Helium Gases: A Guide for Explorationists, AAPG  
666 Conference, April 11-14, New Orleans, Louisiana, USA.
- 667 Cai Z., Clarke R.H., Glowacki B.A., Nuttall, W.J. and Ward, N. (2010) Ongoing ascent to the  
668 helium production plateau—Insights from system dynamics. *Resour. Policy* **35**, 77-89.
- 669 Castro M.C., Goblet P., Ledoux E., Violette S. and Marsily G. (1998) Noble gases as natural tracers  
670 of water circulation in the Paris Basin: 2. Calibration of a groundwater flow model using noble  
671 gas isotope data. *Water Resour. Res.*, **34**(10), 2467-2483.
- 672 Dai, J.X., Pei, X.G., Qi, H.F. (1992). Natural Gas Geology of China. Petroleum Industry Press,  
673 Beijing.
- 674 Danabalan D. (2017) Helium: Exploration Methodology for a Strategic Resource. Ph. D. thesis,  
675 Durham University.
- 676 Danabalan D., Gluyas J.G., Macpherson C.G., Abraham-James T.H., Bluett J.J., Barry P.H. and  
677 Ballentine C.J. (2016) New High-Grade Helium Discoveries in Tanzania, Goldschmidt  
678 Conference, June26 - July 1, Yokohama, Japan.
- 679 Deng Q.D. and You H.C. (1985) The Structural Activity and Formation Mechanism of the  
680 Down-faulted Basins around the Ordos block. Seismological Press, Beijing.
- 681 Gai H.F., Tian H., Cheng P., Zhou Q., Li T.F., Wang X., Xiao X.M. (2019) Influence of retained  
682 bitumen in oil-prone shales on the chemical and carbon isotopic compositions of natural gases:  
683 Implications from pyrolysis experiments. *Mar. Pet. Geol.* **101**, 148-161.
- 684 Gautheron C., Moreira M. (2002) Helium signature of the subcontinental lithospheric mantle. *Earth*  
685 *Planet. Sci. Lett.* **199**, 0-47.
- 686 Gilfillan S.M.V., Ballentine C.J., Holland G., Blagburn D., Lollar B.S., Stevens S., Schoell M. and  
687 Cassidy M. (2008) The noble gas geochemistry of natural CO<sub>2</sub> gas reservoirs from the

688 Colorado Plateau and Rocky Mountain provinces, USA. *Geochim. Cosmochim. Acta* **72**,  
689 1174-1198.

690 Han W., Li Y., Lu J., Ren Z.L., Xu W. and Song B. (2014) The factors responsible for the unusual  
691 content of helium-rich natural gas in the Weihe Basin, Shaanxi Province. *Geol. Bull. China* **33**,  
692 1836-1841 (in Chinese with English abstract).

693 Holland G., Lollar B.S., Li L., Lacrampe-Couloume G., Slater G.F. and Ballentine C.J. (2013) Deep  
694 fracture fluids isolated in the crust since the Precambrian era. *Nature* **497**, 357-360.

695 Ivanhoe L.F. and Leckie G.G. (1993) Global oil, gas fields, sizes tallied, analyzed. *Oil & Gas J.* **91**,  
696 87-91.

697 Kennedy B., Hiyagon H. and Reynolds J. (1990) Crustal neon: a striking uniformity. *Earth Planet.*  
698 *Sci. Lett.* **98**, 277-286.

699 Kipfer R., Aeschbach-Hertig W., Peeters F. and Stute M. (2002) Noble Gases in Lakes and Ground  
700 Waters. *Rev. Mineral. Geochem.* **47**, 615-700.

701 Li J., Pang Z.H., Yang G.M., Tian J., Tong A.L., Zhang X.Y., Hu S.M. (2017) Million-year-old  
702 groundwater revealed by krypton-81 dating in Guanzhong Basin, China. *Sci. Bull.* **62**,  
703 1181-1184.

704 Li X.G. and Ren Y.K. (1986) Effect and characteristics of the neotectonics of the Weihe graben.  
705 Research on Modern Crustal Movements, 2. Seismological Press, Beijing.

706 Logvinyuk V. P., Makarenkov V.V., Malyshev V.V. Panchenkov G.M. (1970) Solubility of gases in  
707 petroleum products. *Chem. Technol. Fuels Oils*, **6**, 353-355.

708 Lowenstern J.B., Evans W.C., Bergfeld D. and Hunt A.G. (2014) Prodigious degassing of a billion  
709 years of accumulated radiogenic helium at Yellowstone. *Nature* **506**, 355.

710 Ma Z., Yu J., Su Y., Xie J., Jia X. and Hu Y. (2010)  $\delta^{18}\text{O}$  shifts of geothermal waters in the central of  
711 Weihe Basin, NW China. *Environ. Earth Sci.* **59**, 995-1008.

712 Mamyrin B.A. and Tolstikhin L.N. (1984) Helium isotopes in nature, Elsevier Sci. Pub.,  
713 Amsterdam.

- 714 Mercier J.L., Vergely P., Zhang Y.Q., Hou M.J., Bellier O. and Wang Y.M. (2013) Structural records  
715 of the Late Cretaceous–Cenozoic extension in Eastern China and the kinematics of the  
716 Southern Tan-Lu and Qinling Fault Zone (Anhui and Shaanxi provinces, PR China).  
717 *Tectonophysics* **582**, 50-75.
- 718 Niedermann S., Graf T. and Marti K. (1993) Mass spectrometric identification of  
719 cosmic-ray-produced neon in terrestrial rocks with multiple neon components. *Earth Planet.*  
720 *Sci. Lett.* **118**, 65-73.
- 721 Nuttall W.J., Clarke R.H. and Glowacki B.A. (2012) The Future of Helium as a Natural Resource.  
722 Routledge, London and New York.
- 723 Pinti D.L., Castro M.C., Lopez-Hernandez A., Han G., Shouakar-Stash O., Hall C.M. and  
724 Ramírez-Montes M. (2017) Fluid circulation and reservoir conditions of the Los Humeros  
725 Geothermal Field (LHGF), Mexico, as revealed by a noble gas survey. *J. Volcanol. Geotherm.*  
726 *Res.* **333-334**, 104-115.
- 727 Podosek F.A., Bernatowicz T.J. and Kramer F.E. (1981) Adsorption of xenon and krypton on shales.  
728 *Geochim. Cosmochim. Acta* **45**, 2401-2415.
- 729 Sano Y., Marty B. and Burnard P. (2013) Noble Gases in the Atmosphere. In The Noble Gases as  
730 Geochemical Tracers (eds. Burnard P.). Springer Berlin, Heidelberg. pp. 17-31.
- 731 Taylor S.R. and McLennan S.M. (1985) The continental crust: its composition and evolution.  
732 United States: Blackwell Scientific, Oxford.
- 733 Tolstikhin I., Lehmann B.E., Loosli H.H. and Gautschi A. (1996) Helium and argon isotopes in  
734 rocks, minerals, and related ground waters: A case study in northern Switzerland. *Geochim.*  
735 *Cosmochim. Acta* **60**, 1497-1514.
- 736 Tolstikhin I., Waber H.N., Kamensky I., Loosli H.H., Skiba V. and Gannibal M. (2011) Production,  
737 redistribution and loss of helium and argon isotopes in a thick sedimentary aquitard-aquifer  
738 system (Molasse Basin, Switzerland). *Chem. Geol.* **286**, 48-58.
- 739 Torgersen T. (1980) Controls on pore-fluid concentration of  $^4\text{He}$  and  $^{222}\text{Rn}$  and the calculation of  
740  $^4\text{He}/^{222}\text{Rn}$  ages. *J. Geochem. Explor.* **13**, 57-75.

741 Torgersen T. and Clarke W.B. (1985) Helium accumulation in groundwater, I: An evaluation of  
742 sources and the continental flux of crustal  $^4\text{He}$  in the Great Artesian Basin, Australia. *Geochim.*  
743 *Cosmochim. Acta* **49**, 1211-1218.

744 Torgersen T. and Ivey G.N. (1985) Helium accumulation in groundwater. II: A model for the  
745 accumulation of the crustal  $^4\text{He}$  degassing flux. *Geochim. Cosmochim. Acta* **49**, 2445-2452.

746 Torgersen T. and Kennedy B. (1999) Air-Xe enrichments in Elk Hills oil field gases: role of water in  
747 migration and storage. *Earth Planet. Sci. Lett.* **167**, 239-253.

748 USGS (United State Geological Survey) (2018) Mineral Commodity Summaries 2018: Helium by J.  
749 E. Hamak. Retrieved from the USGS website:  
750 <https://minerals.usgs.gov/minerals/pubs/commodity/helium/mcs-2018-heliu.pdf> (Accessed 07  
751 August 2018)

752 Wang J., Liu C., Gao F., Zhang D., Li Y. and Li Z. (2015a) Pre-Cenozoic geological characteristics  
753 and oil-gas significance in Weihe basin, Shaanxi Province. *Geol. Bull. China.* **34**(10),  
754 1981-1991 (in Chinese with English abstract).

755 Wang X., Wang T. and Zhang C. (2015b) Granitoid magmatism in the Qinling orogen, central China  
756 and its bearing on orogenic evolution. *Sci. China: Earth Sci.* **58**, 1497-1512.

757 Warr O., Lollar B.S., Fellowes J., Sutcliffe C.N., McDermott J.M., Holland, G., Mabry, J.C.,  
758 Ballentine C.J. (2018) Tracing ancient hydrogeological fracture network age and  
759 compartmentalisation using noble gases. *Geochim. Cosmochim. Acta* **222**, 340-362.

760 Weiss R F. (1968) Piggyback sampler for dissolved gas studies on sealed water samples. *Deep-Sea*  
761 *Res. Oceanogr. Abstr.* **15**, 695-699.

762 Wen T., Castro M.C., Ellis B.R., Hall C.M. and Lohmann K.C. (2015) Assessing compositional  
763 variability and migration of natural gas in the Antrim Shale in the Michigan Basin using noble  
764 gas geochemistry. *Chem. Geol* **417**, 356-370.

765 Wen T., Castro M.C., Nicot J.P., Hall C.M., Pinti D.L., Mickler P., Darvari R. and Larson T. (2017)  
766 Characterizing the Noble Gas Isotopic Composition of the Barnett Shale and Strawn Group

767 and Constraining the Source of Stray Gas in the Trinity Aquifer, North-Central Texas. *Environ.*  
768 *Sci. Technol.* **51**, 6533-6541.

769 Wilhelm E., Battino R. and Wilcock R.J. (1977) Low-pressure solubility of gases in liquid water.  
770 *Chem. Rev.* **77**, 219-262.

771 Xu X., Ma X. and Deng Q. (1993) Neotectonic activity along the Shanxi rift system, China.  
772 *Tectonophysics* **219**, 305-325.

773 Zhang A., Yang Z., Zhong J. and Mi F. (1995) Characteristics of late quaternary activity along the  
774 Southern Border Fault Zone of Weihe Graben Basin. *Quatern. Int.* **25**, 25-31.

775 Zhang L. (2014a) Geothermal and associated water soluble gas resources exploration and prospect  
776 forecast of Weihe Basin. Ph. D. thesis, Chang'an Univ (in Chinese with English abstract).

777 Zhang M., Tang Q., Cao C., Lv Z., Zhang T., Zhang D., Li Z. and Du L. (2018a) Molecular and  
778 carbon isotopic variation in 3.5 years shale gas production from Longmaxi Formation in  
779 Sichuan Basin, China. *Mar. Pet. Geol.* **89**, 27-37.

780 Zhang W., Li Y.H., Wang L., Zhao F.H., Han W., and Song C.G. (2018b) The analysis of helium  
781 accumulation conditions and prediction of helium resource in Weihe Basin. *Nat. Gas Geosci.*  
782 **29**(2), 236-244 (in Chinese with English abstract).

783 Zhou Z. and Ballentine C.J. (2006) <sup>4</sup>He dating of groundwater associated with hydrocarbon  
784 reservoirs. *Chem. Geol.* **226**(3), 309-327.

785 Zhou Z., Ballentine C.J., Kipfer R., Schoell M. and Thibodeaux S. (2005) Noble gas tracing of  
786 groundwater/coalbed methane interaction in the San Juan Basin, USA. *Geochim. Cosmochim.*  
787 *Acta* **69**, 5413-5428.

788 Zhou Z., Ballentine C.J., Schoell M. and Stevens S.H. (2012) Identifying and quantifying natural  
789 CO<sub>2</sub> sequestration processes over geological timescales: The Jackson Dome CO<sub>2</sub> Deposit,  
790 USA. *Geochim. Cosmochim. Acta* **86**, 257-275.

791

## Tables

**Table 1 Major gas concentrations and carbon isotopes of gases from geothermal wells in Weihe Basin.**

Sample	Location	Sampling method <sup>a</sup>	Depth(m) <sup>b</sup>	Group	Compositions (vol%)						Carbon isotopes (‰)	
					C <sub>1</sub>	N <sub>2</sub>	CO <sub>2</sub>	O <sub>2</sub>	C <sub>2</sub>	C <sub>3+</sub>	δ <sup>13</sup> C(CH <sub>4</sub> )	δ <sup>13</sup> C(CO <sub>2</sub> )
<b>Normal samples</b>												
Xi'an Depression												
XP-1	Xingping-Huihao	GWS	-	A	4.35	83.63	4.79	3.11	0.01	0.00	-30.41	-10.50
WG-1	Wugong-Xin1	GWS	-	A	5.12	86.73	3.77	0.12	0.02	0.00	-27.85	-8.06
XY-1	Xianyang-Wen4	GWS	2600-2804	A	3.07	91.66	2.09	0.14	0.01	0.00	-20.60	-8.98
XY-2	Xianyang-Jin1	GWS	2000-4080	A	7.24	58.57	31.69	0.51	0.04	0.01	-25.83	-5.77
XY-3	Xianyang-Sanpu1	GWS	-	A	11.35	73.60	9.36	2.86	0.05	0.01	-32.07	-10.42
Gushi Depression												
WR-051	Huaying-051 well	GWS	1900-2600	A	25.40	66.56	3.82	0.06	0.24	0.08	-	-
WR-2	Weinan-Weire 2	GWS	2000-3800	A	11.90	72.24	7.42	6.85	0.42	0.22	-44.46	-8.23
WR-1	Weinan-Weire 1	GWS	2080-3200	B	69.50	4.85	22.24	0.23	2.57	0.55	-30.23	-5.47
WR-11	Weinan-Weire 11	GWS	2200-3650	B	52.94	27.81	13.56	2.28	2.22	0.91	-19.04	-7.49
WR-HX	Huaxian -Huaxian 2	FT	-	C	71.70	27.20	0.03	0.00	0.57	0.11	-33.47	-
<b>Outlier Samples</b>												
XP-2	Xingping-Xintai	GWS	-	A	2.49	91.53	2.66	0.61	0.02	0.00	-30.59	-9.16
XP-3	Xingping-Jincheng 1	GWS	-	A	2.23	89.48	2.28	3.46	0.01	0.00	-	-

<sup>a</sup> GWS and FT represent that the gas samples are either collected through a gas-water separator (GWS) or a flash tank (FT).

<sup>b</sup> Depth of groundwater reservoir.

**Table 2 Noble gas isotope systematics of gases from Weihe Basin<sup>a</sup>.**

Sample	Group	<sup>4</sup> He×10 <sup>-4</sup>	<sup>20</sup> Ne×10 <sup>-7</sup>	<sup>36</sup> Ar×10 <sup>-6</sup>	<sup>84</sup> Kr×10 <sup>-8</sup>	<sup>130</sup> Xe×10 <sup>-10</sup>	<sup>3</sup> He/ <sup>4</sup> He (R/Ra) <sup>b</sup>	<sup>20</sup> Ne/ <sup>22</sup> Ne	<sup>21</sup> Ne/ <sup>22</sup> Ne	<sup>40</sup> Ar/ <sup>36</sup> Ar	<sup>86</sup> Kr/ <sup>84</sup> Kr	<sup>132</sup> Xe/ <sup>130</sup> Xe
		cm <sup>3</sup> STP/cm <sup>3</sup>										
Air <sup>c</sup>		0.05	165	31.43	65.0	36.6	1	9.80	0.029	298.56	0.305	6.6098
<b>Normal Samples</b>												
Xi'an Depression												
XP-1	A	161±2	30.8±0.4	10.41±0.13	21.7±0.83	19.4±0.6	0.033±0.001	9.61±0.15	0.0296±0.0008	368±6	0.316±0.025	6.57±0.21
WG-1	A	237±2	39.5±0.4	13.78±0.27	44.9±1.2	40.2±1.1	0.036±0.001	9.83±0.16	0.0295±0.0009	427±9	0.315±0.017	6.92±0.21
XY-1	A	294±3	53.2±0.6	19.20±0.44	64.8±1.8	17.8±0.5	0.071±0.001	9.83±0.15	0.0297±0.0008	447±11	0.314±0.018	6.66±0.19
XY-2	A	94±1	15.9±0.2	7.18±0.11	23.2±0.54	21.7±0.5	0.048±0.001	10.1±0.2	0.0296±0.0008	447±8	0.286±0.014	6.47±0.17
XY-3	A	153±2	16.4±0.2	7.96±0.12	23.1±0.63	17.5±0.4	0.066±0.001	9.86±0.15	0.0311±0.0008	613±11	0.322±0.022	6.50±0.18
Gushi Depression												
WR-2	A	32.3±0.3	22.4±0.3	7.84±0.09	17.8±0.41	14.5±0.8	0.052±0.001	8.92±0.16	0.0291±0.0008	339±5	0.322±0.014	7.08±0.41
WR-051	A	161±2	31.5±0.4	8.26±0.10	16.6±1.10	16.7±0.6	0.070±0.001	9.81±0.15	0.0301±0.0009	518±8	0.299±0.032	6.47±0.27
WR-1	B	5.74±0.06	0.69±0.02	0.251±0.004	0.87±0.01	0.98±0.05	0.070±0.002	10.9±0.4	0.0307±0.0021	735±13	0.301±0.007	7.00±0.36
WR-11	B	6.21±0.06	1.11±0.03	0.578±0.008	2.07±0.02	2.65±0.09	0.079±0.001	8.94±0.24	0.0292±0.0015	541±9	0.308±0.005	6.74±0.25
WR-HX	C	211±2	5.50±0.07	2.88±0.04	9.95±0.11	8.17±0.72	0.126±0.002	8.68±0.15	0.0385±0.0015	1142±19	0.307±0.005	7.22±0.69
<b>Outliers Samples</b>												
XP-2	A	82.4±0.8	30.8±0.4	16.82±0.39	41.4±0.9	23.2±0.7	0.038±0.001	9.38±0.16	0.0285±0.0007	321±5	0.305±0.013	6.50±0.26
XP-3	A	85.3±0.9	69.1±0.9	22.94±0.40	72.0±1.2	30.4±1.1	0.022±0.001	9.31±0.14	0.0283±0.0007	312±5	0.307±0.009	6.70±0.18

<sup>a</sup> 1σ errors are shown in the table after the plus and minus signs.

<sup>b</sup> <sup>3</sup>He/<sup>4</sup>He ratios (R) are normalized to the air value Ra = 1.4×10<sup>-6</sup>.

<sup>c</sup> Noble gas concentrations are cited from Sano et al. (2013). <sup>3</sup>He/<sup>4</sup>He ratio in air is cited from Mamyrin and Tolstikhin (1984). <sup>20</sup>Ne/<sup>22</sup>Ne and <sup>21</sup>Ne/<sup>22</sup>Ne ratios in air are cited from Bottomley et al. (1984). <sup>40</sup>Ar/<sup>36</sup>Ar ratio in air is cited from Lee et al. (2006). <sup>86</sup>Kr/<sup>84</sup>Kr ratio in air is cited from Aregbe et al. (1996). <sup>132</sup>Xe/<sup>130</sup>Xe ratio in air is cited from Valkiers et al. (1998).



**Table 3 Henry's constants (atm kg/mol) of noble gases and nitrogen at modeling conditions<sup>a</sup>.**

Condition	Depth (m)	Temperature (°C)	Groundwater Salinity (M NaCl)	Water						Heavy oil (API = 25)					
				Helium	Neon	Argon	Krypton	Xenon	Nitrogen	Helium	Neon	Argon	Krypton	Xenon	Nitrogen
Recharge	0	10	0	2440.12	1975.10	538.11	276.98	156.25	1169.12						
Reservoir	2500	88	0.1	2061.27	2069.94	1206.80	971.72	639.84	1983.84	595.39	585.60	124.34	59.46	25.35	295.74

<sup>a</sup> Henry's constants of noble gases are cited from Ballentine et al. (2002) and references therein; Henry's constants of N<sub>2</sub> in water and oil are calculated from Wilhelm et al. (1977) and Logvinuk et al. (1970) respectively.

**Table 4 Volume ratios of oil to groundwater during fractionation, volume ratios of gas to water in the subsurface, calculated He, N<sub>2</sub> and CH<sub>4</sub> contents in groundwater under modeling conditions and percentages of ASW-derived N<sub>2</sub>.**

Sample	Group	<sup>20</sup> Ne/ <sup>36</sup> Ar	f <sup>36</sup> Ar remaining in groundwater	V <sub>oil</sub> /V <sub>water</sub>	V <sub>gas</sub> /V <sub>water</sub>	He content in groundwater cm <sup>3</sup> STP/g	CH <sub>4</sub> content in groundwater cm <sup>3</sup> STP/g	N <sub>2</sub> content in groundwater cm <sup>3</sup> STP/g	Measured N <sub>2</sub> / <sup>36</sup> Ar	Calculated ASW-derived N <sub>2</sub> / <sup>36</sup> Ar	Percentage of ASW-derived N <sub>2</sub>
XP-1	A	0.295	0.32	0.13	0.04	8.62×10 <sup>-4</sup>	0.002	0.030	1.62×10 <sup>4</sup>	8.87×10 <sup>4</sup>	18.35%
WG-1	A	0.287	0.33	0.13	0.03	9.85×10 <sup>-4</sup>	0.001	0.024	1.60×10 <sup>4</sup>	6.12×10 <sup>4</sup>	26.19%
XY-1	A	0.277	0.35	0.12	0.02	9.09×10 <sup>-4</sup>	0.001	0.020	1.57×10 <sup>4</sup>	5.20×10 <sup>4</sup>	30.33%
XY-2	A	0.222	0.50	0.08	0.08	9.69×10 <sup>-4</sup>	0.006	0.047	1.41×10 <sup>4</sup>	8.17×10 <sup>4</sup>	17.32%
XY-3	A	0.206	0.56	0.07	0.08	1.54×10 <sup>-3</sup>	0.009	0.060	1.36×10 <sup>4</sup>	1.04×10 <sup>5</sup>	13.17%
WR-2	A	0.287	0.33	0.13	0.05	2.36×10 <sup>-4</sup>	0.006	0.036	1.60×10 <sup>4</sup>	9.72×10 <sup>4</sup>	16.50%
WR-051	A	0.382	0.21	0.18	0.03	8.37×10 <sup>-4</sup>	0.008	0.020	1.84×10 <sup>4</sup>	9.05×10 <sup>4</sup>	20.35%
WR-1	B	0.308	0.30	0.14	1.38	1.22×10 <sup>-3</sup>	0.959	0.067	1.65×10 <sup>4</sup>	2.07×10 <sup>5</sup>	8.01%
WR-11	B	0.195	0.61	0.06	1.22	9.06×10 <sup>-4</sup>	0.648	0.340	1.33×10 <sup>4</sup>	4.62×10 <sup>5</sup>	2.88%

**Table 5 Parameters used for calculation of <sup>4</sup>He accumulation rate.**

<sup>4</sup> He source	U (ppm)	Th (ppm)	Density (g/cm <sup>3</sup> )	Porosity (%)	Thickness (km)	Accumulation rate (cm <sup>3</sup> STP <sup>4</sup> He/g H <sub>2</sub> O year)
In situ-aquifer	2.7	9.6	2.6	0.15	0.2	$5.54 \times 10^{-12}$
External-General Upper crust	2.8	10.7	2.6	-	18	$6.86 \times 10^{-10}$
External-Lower crust	0.28	1.07	3.3	-	18	$8.71 \times 10^{-11}$

## Figures and Captions




Figure 1 Map of the Weihe Basin. a) shows sample locations (yellow dots), basin area (white line), granitic body region (orange lines), tectonic unit boundaries (red lines) and locations of cross section lines (black lines). The A-A' and B-B' geological cross sections. b) shows the major geothermal reservoir of Pliocene Lantian-Bahe Formation (N<sub>2</sub>l+b). Groundwater recharges from Ordos Basin and Qinling Orogenic Belt and discharges from the middle Weihe River.




Figure 2 Schematic diagram of the noble gas analytical system at Lancaster University.

This system consists of four major sections including sample introduction, sample purification, noble gas separation and noble gas analysis sub-systems. Details of the noble gas analytical procedures are described in Section 3.2.2.




Figure 3  $R/R_a$  ratios vs.  $^4\text{He}/^{20}\text{Ne}$  ratios. Helium is mainly derived from crust with 0.40%-1.94% mantle contribution. The helium isotope endmembers used in this study are the sub-continental lithospheric mantle-like ratio of  $6.1R_a$  from Gautheron and Moreira (2002) and the upper crustal value of  $0.008R_a$  from Ballentine and Burnard (2002).




Figure 4  $^{20}\text{Ne}/^{22}\text{Ne}$  ratios vs.  $^{21}\text{Ne}/^{22}\text{Ne}$  ratios. The data can be accounted for by a small mass fractionation-related process (MFL) and an addition of crustal  $^{21}\text{Ne}$ . The calculation of two-component mixing model between ASW and crust ( $^{20}\text{Ne}/^{22}\text{Ne}=0$  and  $^{21}\text{Ne}/^{22}\text{Ne} = 0.47$ , [Kennedy et al., 1990](#)) suggests that crust-derived  $^{21}\text{Ne}$  contributes between 0.00% - 9.47% to the total  $^{21}\text{Ne}$  except for WR-HX which is 33.33%.




Figure 5 a)  $^{36}\text{Ar}$  concentrations ( $1/^{36}\text{Ar}$ ) vs. Ar isotopes ( $^{40}\text{Ar}/^{36}\text{Ar}$ ) of samples in the three groups. b) shows the detailed characteristics of samples in Group A, in which the samples XP-2 and XP-3 (open circles) are highly consistent with the mixing line of XP-1 (filled circle) and air values (open rectangle), indicating these two samples are likely contaminated by air. There is no positive correlation between  $^{40}\text{Ar}/^{36}\text{Ar}$  ratios and  $1/^{36}\text{Ar}$  values for the rest of samples.






Figure 6 a)  $^{20}\text{Ne}/^{36}\text{Ar}$  vs.  $^{130}\text{Xe}/^{36}\text{Ar}$  and b)  $^{20}\text{Ne}/^{36}\text{Ar}$  vs.  $^{84}\text{Kr}/^{36}\text{Ar}$  remaining in groundwater phase after oil-water Rayleigh fractionation and batch fractionation. The heavy oil-water batch fractionation can explain data better. All data are in the area restricted by the Rayleigh heavy oil-water fractionation line and that with either 600% excess of Xe or 100% excess of Kr. The excess heavy noble gases may be released from organic-rich minerals (Barry et al., 2016; Zhou et al., 2005).




Figure 7  ${}^4\text{He}$  concentrations vs.  ${}^{20}\text{Ne}$  concentrations in samples from Weihe Basin (filled triangles and circles) and Hugoton-Panhandle gas field (filled squares, data from Ballentine and Lollar, 2002). The agreement of  ${}^4\text{He}$  and  ${}^{20}\text{Ne}$  concentrations in Group A and B samples (all samples except for WR-HX) suggests helium accumulation is closely related with groundwater migration. The high  ${}^4\text{He}/{}^{20}\text{Ne}$  ratio in WR-HX well is similar with those in Hugoton-Panhandle gas field, suggesting a local  ${}^4\text{He}$  flux possibly coming from adjacent granitic bodies overlaying the background  ${}^4\text{He}$  flux.




Figure 8 Cartoon illustrating the accumulation process of <sup>4</sup>He, ASW-derived noble gases, CH<sub>4</sub> and N<sub>2</sub> in Weihe Basin. See detailed discussion and explanation in Section 5.4.1.