





RESEARCH NOTE

Non-invasive visualisation and identification of fluorescent *Leishmania tarentolae* in infected sand flies [version 1; referees: 1 approved, 2 approved with reservations]

Hector M. Diaz-Albiter^{1,2}, Clément Regnault², Edubiel A. Alpizar-Sosa ²,
Dagmara McGuinness ², Michael Barrett², Rod J. Dillon³

¹El Colegio de la Frontera Sur, Villahermosa, Tabasco, 86280, Mexico

²Wellcome Centre for Molecular Parasitology, University of Glasgow, Glasgow, G12 8TA, UK

³Faculty of Health and Medicine, Lancaster University, Lancaster, Lancashire, LA1 4YQ, UK

v1 First published: 13 Dec 2018, 3:160 (<https://doi.org/10.12688/wellcomeopenres.14910.1>)

Latest published: 13 Dec 2018, 3:160 (<https://doi.org/10.12688/wellcomeopenres.14910.1>)

Abstract

Background: The leishmaniasis are neglected diseases that affect some of the most vulnerable populations in the tropical and sub-tropical world. The parasites are transmitted by sand flies and novel strategies to control this neglected vector-borne disease are needed. Blocking transmission by targeting the parasite inside the phlebotomine vector offers potential in this regard. Some experimental approaches can be best performed by longitudinal study of parasites within flies, for which non-destructive methods to identify infected flies and to follow parasite population changes are required.

Methods: *Lutzomyia longipalpis* were reared under standard insectary conditions at the Wellcome Centre for Molecular Parasitology. Flies were artificially infected with *L. tarentolae* expressing green fluorescent protein (GFP). Parasite counts were carried out 5 days post-infection and the percentage of infected flies and survival of infected females was established up to days 5 post-infection. Whole living females were visualised using an epifluorescence inverted microscope to detect the presence parasites inferred by a localised green fluorescent region in the upper thorax. Confirmation of infection was performed by localised-fluorescence of dissected flies and estimates of the parasite population.




Results: *Leishmania tarentolae* was successfully transfected and expressed GFP *in vitro*. *L. tarentolae*-GFP Infected flies showed similar parasite populations when compared to non-transfected parasites (*L. tarentolae*-WT). Survival of non-infected females was higher than *L. tarentolae*-infected groups, (Log-rank (Mantel-Cox) test, $p < 0.05$). *L. tarentolae*-GFP infected females displayed an intense localised fluorescence in the thorax while other specimens from the same infected group did not. Localised fluorescent flies were dissected and showed higher parasite populations compared to those that did not demonstrate high concentrations in this region (t-test, $p < 0.005$).

Conclusion: These results demonstrate the feasibility of establishing a safe non-human infectious fluorescent *Leishmania*-sand fly infection model by allowing non-destructive imaging to signal the establishment of *Leishmania* infections in living sand flies.

Open Peer Review

Referee Status: ? ✓ ?

	Invited Referees		
	1	2	3
version 1 published 13 Dec 2018	? report	✓ report	? report

- 1 **Jesus G Valenzuela** , National Institute of Allergy and Infectious Diseases, National Institutes of Health (NIH), USA
- 2 **Yusuf Özbel** , Ege University, Turkey
- 3 **Antonio Jorge Tempone** , Instituto Oswaldo Cruz - FIOCRUZ, Brazil

Discuss this article

Comments (0)

Keywords

Leishmania, Lutzomyia, sand fly, fluorescence, GFP, parasite-vector interactions

Corresponding author: Hector M. Diaz-Albiter (hector.diazalbiter@glasgow.ac.uk)

Author roles: **Diaz-Albiter HM:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation; **Regnault C:** Methodology, Visualization, Writing – Original Draft Preparation; **Alpizar-Sosa EA:** Investigation; **McGuinness D:** Data Curation, Formal Analysis, Investigation, Methodology, Software, Writing – Original Draft Preparation, Writing – Review & Editing; **Barrett M:** Funding Acquisition, Resources, Writing – Review & Editing; **Dillon RJ:** Resources, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work is supported by the Wellcome Trust [104111]; and the Royal Society, [NF151329].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2018 Diaz-Albiter HM *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Diaz-Albiter HM, Regnault C, Alpizar-Sosa EA *et al.* **Non-invasive visualisation and identification of fluorescent *Leishmania tarentolae* in infected sand flies [version 1; referees: 1 approved, 2 approved with reservations]** Wellcome Open Research 2018, 3:160 (<https://doi.org/10.12688/wellcomeopenres.14910.1>)

First published: 13 Dec 2018, 3:160 (<https://doi.org/10.12688/wellcomeopenres.14910.1>)

Introduction

Leishmania are protozoan parasites belonging to the Order Kinetoplastida. They infect various vertebrate species and are transmitted via the bite of phlebotomine sand flies (Bates, 2007). Approximately 20 species of this genus cause the leishmaniasis in humans, a group of diseases presenting different clinical manifestations, some of which can lead to death (Burza *et al.*, 2018). The disease is endemic in 97 countries reporting approximately 1 million new cases yearly (WHO, 2018). Although, some treatments are available, they are associated with high toxicity and quite often administered too late and treatment failures are common. No vaccine is available and overuse of pesticides to control arthropod populations worldwide has resulted in development of resistance in arthropod vectors, including sandflies that transmit *Leishmania* (Hemingway & Ranson, 2000; Hassan *et al.*, 2012).

Alternative approaches to intervening against the leishmaniasis are sought and novel approaches to transmission control are attractive (Beatty, 2000). Novel strategies aiming to develop new approaches to interrupt the transmission cycle rely on a thorough knowledge of insect-parasite interactions (Kamhawi, 2006). Several studies focusing on *Leishmania* metabolism, drug resistance, and host-parasite interactions led to us establishing a colony of *Lutzomyia longipalpis* sandflies at the Wellcome Centre for Molecular Parasitology (WCMP). *Lutzomyia longipalpis* is permissive to infection with multiple *Leishmania* species (Soares & Turco, 2003; Volf & Myskova, 2007), making it a versatile model for laboratory based work on parasite-vector interactions.

The *Leishmania*-sand fly model established at the WCMP was tested for potential in establishing artificial infections using the non-human pathogenic *Leishmania tarentolae*. This species was first identified in the gecko *Tarentola mauritanica* (Adler & Theodor, 1935; Chatton & Blanc, 1914; Laveran & Franchini, 1921). Both promastigote and (rarely) amastigote forms are found circulating in the blood, whereas promastigotes are located in the intestinal lumen as well as the cloacae of their lizard hosts (Wilson & Southgate, 1979). The natural vectors of this species are *Sergentomyia* spp., and its development in this fly has been described as hypopylarian, i.e. a parasite that establishes in the hindgut of the sand fly (Bates, 2007). Also, DNA of *L. tarentolae* has been also found in *Phlebotomus argentipes* (Gajapathy *et al.*, 2013). However, very little is known about its biology inside the vector. Research on this species has focussed mainly on its use as a model to study antibiotic resistance and metabolism (Aphasizhev *et al.*, 2003; Dey *et al.*, 1994; Ouellette *et al.*, 1990), and more recently, it has become a well-established, commercially available expression system (Breitling *et al.*, 2002; Kushnir *et al.*, 2005; Zhang *et al.*, 1995).

We transfected the parasite to express GFP aiming to perform further experiments visualising parasite dynamics in the sand fly gut, infections with antibiotic-resistant parasites, as well as to provide a safe protocol for training students and staff to perform *Leishmania* infections where containment facilities are not required.

We showed that non-transfected and GFP-expressing *L. tarentolae* were present in the gut 5 days post infection. It was observed that GFP-parasite infected flies presented a strong fluorescence area in the thorax under the microscope before dissecting them, which was the midgut. We confirmed that flies presenting this localised fluorescence were heavily infected. The technique presented here provides, for the first time, the ability to study the *Leishmania*-sandfly relationship in a non-destructive fashion. This non-destructive GFP infection signalling can be used for testing chemicals that might kill parasites within the vector, infected fly behaviour experiments determining microbiome-relationships that might prevent transmission.

Methods

Sand fly rearing

A colony of *Lutzomyia longipalpis* was initiated from flies originating from the colony at Lancaster University and further flies provided from the lab of Paul Kaye (University of York), both derived originally from Jacobina, Brazil. Insects were kept under standard laboratory conditions (Modi & Tesh, 1983). Briefly, all developmental stages were kept at 27°C (±2°C), a relative humidity of ~80% and a photoperiod of 8 hours light/16 hours darkness. Larvae were kept in plastic pots with a plaster substrate and fed a 1:1 rabbit food and droppings mixture that was autoclaved and ground. Adults were fed *ad libitum* with 70% w/v sucrose solution offered in cotton wool. Females were fed on commercially obtained sheep blood (in Alsever's as anticoagulant) via a Hemotek membrane feeder (Discovery Workshops, UK) at 37°C.

Parasite cultures

Leishmania tarentolae (strain p10) promastigotes were kindly donated by Barrie Rooney (University of Kent) and cultured in HOMEM medium (GE Healthcare) supplemented with 10% heat-inactivated foetal bovine serum (HiFBS, Gibco) and 1% penicillin/streptomycin solution (Sigma-Aldrich). Cells were maintained by weekly sub-passage of stationary-phase cultures in fresh complete HOMEM at a density of 10⁵–10⁶ cells/ml. Cultures were kept in non-vented T25 or T75 flasks (Corning) and incubated at 25°C.

Transfections

Transfections were performed on mid-log phase *L. tarentolae* promastigotes. A total of 5 × 10⁷ parasites were pelleted (1200xg, 10 min) and washed once with ice-cold PBS before resuspending in 100 µl of transfection buffer (90 mM NaPO₄, 5 mM KCl, 50 mM HEPES, 0.15 mM CaCl₂, pH7.3). The suspension was transferred to a 1 ml Bio-Rad GenePulser® cuvette, and mixed with 10 µg of plasmid pGL1132 DNA in 20 µl sterile water, or 20 µl of sterile water as a negative control. Parasites were electroporated using an Amaxa Nucleofector™ II and subsequently transferred to 10 ml of complete HOMEM in a T25 non-vented flask (Corning). After an overnight incubation at 25°C, appropriate drugs (G418 or Hygromycin, 25 µg/ml) were added in the culture to select for transfectants. Successful transfection was evaluated by confirmation of fluorescence using a Carl Zeiss Axiovert 40 CFL inverted epifluorescence microscope (488 nm excitation and 509 nm emission).

Sand fly infections

A total of 2×10^6 parasites were pelleted (1200xg, 10 min) and washed once with ice-cold PBS before resuspending in 1ml of blood meal (sheep blood mixed with heat-inactivated serum) and fed to the insects using a chick skin membrane with the Hemotek apparatus at 37°C as described above. Groups of ~80 females were fed for one hour, after which 60 fully-engorged individuals were separated and kept under standard laboratory conditions until required for experimental work. Negative control groups were flies fed only on sheep blood and manipulated as stated above. Sandfly mortality was recorded daily and 14 insects were dissected 5 days post infection (DPI), when defecation had already taken place and parasites have likely established an infection. Parasite counts were performed by dissecting individual midguts in PBS and manually homogenizing them in 50 ul of PBS. Total parasites were counted under the microscope using a haemocytometer. Percentage of infected flies 5 DPI was recorded and sandfly survival was evaluated every day for the duration of the experiment. Three biological replicates were performed.

Assessment of fluorescence *in vivo*

An intense green fluorescence in the thorax area of *L. tarentolae*-GFP infected flies was observed, 14 individuals were then sorted out into 2 groups based on the absence and presence of this localised fluorescence. Midguts were dissected as described above, confirmation of fluorescent parasites inside intact guts was performed using a Carl Zeiss Axiovert 40 CFL inverted epifluorescence microscope (488 nm excitation and 509 nm emission) and parasite counts of each gut were performed using a haemocytometer as described above.

Statistical analysis

Survival analyses were performed using the Kaplan-Meier Log Rank χ^2 test. Groups were compared using unpaired t-test and infection percentages were compared using a Fisher's test. Results are expressed as the group mean \pm SEM. Significance was considered when $p < 0.05$. All data were analysed with GraphPad Prism (version 5.00, Prism Inc).

Results

Sandfly infections with GFP-transfected *L. tarentolae*

Transfection of *L. tarentolae* with plasmid pGL1132 and subsequent expression of GFP was successfully achieved *in vitro* in *L. tarentolae* (Supplementary Figure 1). Fluorescent parasites were tested alongside WT parasites to determine infectivity in flies. Estimates of parasite population from dissected female *L. longipalpis* infected with *L. tarentolae*-GFP showed no significant differences (unpaired t-test, $p = 0.8602$) compared with non-transfected parasites (*L. tarentolae*-WT) 5 days post-infection (Figure 1A). Percentage of *L. tarentolae*-GFP vs. *L. tarentolae*-WT dissected on day 5 post-infection were not statistically different (F test, $p = 0.6298$; Figure 1B). Sand fly mortality was evaluated over the length of the experiments and data are presented as survival curves from *L. tarentolae*-GFP, *L. tarentolae*-WT and blood-fed, non-infected negative control insects (Figure 1C). Survival curves were not statistically different between infected groups (Log-rank (Mantel-Cox) Test, $p = 0.5801$). However, survival of non-infected, negative control group was higher and significantly different compared against *L. tarentolae*-GFP and *L. tarentolae*-WT infected flies (Log-rank (Mantel-Cox) Test, $p < 0.005$, $p < 0.05$, respectively).




Figure 1. Artificial Infection of female *Lutzomyia longipalpis* with *Leishmania tarentolae*. (A) Scatter plot shows estimates of parasite populations of non-transfected parasites (*L. tarentolae*-WT) and GFP transfected parasites (*L. tarentolae*-GFP) infected flies 5 days post infection (DPI). Each circle represent parasite counts in individual flies, bars are mean \pm SEM. (B) Bars represent percentage of infected (black) and non-infected (white) flies 5 DPI. (C) Survival curves of *L. tarentolae*-WT, *L. tarentolae*-GFP and non-infected negative control. Survival curves of WT and GFP were not statistically different, negative control showed significant differences compared with WT and GFP (Log-rank (Mantel-Cox) Test, $p < 0.05$).

Assessment of *L. tarentolae*-GFP fluorescence *in vivo*

Having identified a localised fluorescence in non-dissected *L. tarentolae*-GFP infected flies, 14 flies were sorted under an epifluorescence microscope based on the presence or absence of this feature. **Figure 2A** shows an infected individual under normal light, under fluorescence, and the gut of the same fly also under fluorescence (left to right). **Figure 2B** shows an individual that was part of the same *L. tarentolae*-GFP infected group but without localised fluorescence in the thorax region, under normal light, under fluorescence and the gut of that fly also under fluorescence (left to right). Flies that displayed localised fluorescence in the thorax region, and individuals that did not show this feature, were separated into 2 groups, dissected, and parasite numbers were determined using a Neubauer haemocytometer. **Figure 2C** shows parasite populations of individuals not showing localised fluorescence vs. individuals not showing localised fluorescence. The number of parasites were significantly higher in flies presenting localised fluorescence compared with those that did not (test $p < 0.005$).

Discussion

In this work, we successfully infected female *L. longipalpis* artificially with the not-pathogenic to human *Leishmania tarentolae*. We also transfected *L. tarentolae* to express GFP and showed that parasite populations and infection rates were similar between *L. tarentolae* expressing GFP and non-transfected *L. tarentolae*, five days after artificial infection. We also recorded infected sand fly survival with no significant differences

between *L. tarentolae*-GFP and *L. tarentolae*-WT. However the key finding of this work was the observation of a distinctive area of fluorescence in non-dissected flies, emitted from inside the thorax, where the midgut is located. Further dissections of sand flies presenting this specific fluorescence were performed and showed comparatively large populations of fluorescent parasites, while flies not showing this feature possessed comparatively very low or no parasite populations.

Our results show that it is possible to use a GFP-expressing non-pathogenic species of *Leishmania* to infect *Lutzomyia longipalpis*. The *L. tarentolae*-*Lu. longipalpis* system provides an additional tool to study vector-parasite interactions using a hazard-free system that can be used to train students, technicians and scientists wishing to study host-parasite interactions. The system may also enable experiments analysing *Leishmania*-resistance and susceptibility in sandflies over the life of the fly. It may also expedite gene silencing experiments on infected flies. Insects where the localised fluorescence was found were cold-anaesthetised and this technique allows recovery of living individuals that can then be kept alive for further experiments. Sand flies show different levels of infection ([Pimenta et al., 1994](#)) and the fluorescence-selection process could allow separation of heavily-infected, low-infected flies and non-infected flies, to rear them as separate groups. This could provide a basis to better understand parasite resistance in sand flies. This technique can also be extended for medically relevant species of *Leishmania*.




Figure 2. *L. tarentolae* -GFP fluorescence *in vivo*. (A) *L. tarentolae*-GFP Infected individual with localised green fluorescence in the upper thorax: under normal light, under fluorescence, and dissected gut under fluorescence (left to right). (B) *L. tarentolae*-GFP Infected individual without localised green fluorescence in the upper thorax under normal light, under fluorescence, and dissected gut under fluorescence (left to right). (C) Scatter plot shows parasite numbers of *L. tarentolae*-GFP infected flies without localised thorax fluorescence (non-fluorescent) and *L. tarentolae*-GFP infected flies showing localised thorax fluorescence. Each circle represents individual counts, bars are mean \pm SEM. Asterisk represents significant differences (unpaired t-test, $p < 0.005$).

The first experimental infection of sand flies with *L. tarentolae* was performed with *Phlebotomus minutus* fed on infected geckos (Parrot, 1934). The vectors of this species are *Sergentomyia* spp., and its development in the fly has been described as hypopylarian, this is, a parasite that establishes in the hindgut of the sand fly (Bates, 2007). Vionette, Ginger & Dillon (unpublished data) showed that *L. tarentolae* could infect *L. longipalpis* and establish a hypopylarian infection. In contrast, our results with the *L. tarentolae*-GFP infection in the fly gut indicated diffuse distribution with no specific binding areas in the gut and did not suggest a hypopylarian type of infection. Regarding our data on *L. tarentolae*-infected flies (Figure 1C), we found similar results to Rogers & Bates, (2007) who found that *L. mexicana* infections in *Lu. longipalpis* increased insect mortality.

Fluorescence have been extensively used to study several aspects of parasite-vector interactions. Perhaps the largest body of work come from mosquito-malaria and tsetse-trypanosome models. In mosquitoes, the use of fluorescent *Plasmodium* together with several microscopy and imaging techniques has largely contributed understanding parasite dynamics such as ookinete migration and mosquito midgut invasion as well the behaviour of sporozoites during ejection of saliva. These studies have been performed in dissected guts as well as in intact insects or intact body parts (Frischknecht *et al.*, 2004; Vlachou *et al.*, 2004; Zieler & Dvorak, 2000). Similar experiments performed in tsetse flies have helped to elucidate the establishment of infection by *Trypanosoma brucei*, as well as the dynamics of co-infection with different strains of the parasite in dissected guts (Gibson & Bailey, 2003). Perhaps some of the most elegant recent work include an extremely detailed modelling of Trypanosome motility, including tsetse tissue topology that resulted in the first 3D modelling of the infection (Schuster *et al.*, 2017).

Our finding of localised fluorescence present in the upper thorax of *L. tarentolae*-infected flies, and further confirmation of high parasite numbers in these individuals has not been described before for phlebotomines. Given the extensive work carried out in other vector-parasite models, this is a method that should be explored further, building up from some of the many tools available from other models.

Supplementary material

Supplementary Figure 1. Transfection of *L. tarentolae*. (A) Diagram of plasmid pGL1132 showing insertion site of GFP. (B) GFP-transfected *L. tarentolae* *in vitro* (x400).

[Click here to access the data](#)

Perhaps one of the main limitations of the proposed protocol in this report is the green fluorescence of insect cuticle and Malpighian tubules (Figure 2A and B), which makes imaging analysis challenging. Qualitative selection of fluorescent and non-fluorescent flies was based on knowledge of the sand fly internal anatomy, as the midgut is pushed forward into the thorax as oocytes increase in size and rests where the green intensity in heavily-infected flies was found. This could be addressed in further experiments evaluating different fluorophores such as DsRed or mCherry to enable a quantitative analysis of fluorescence related to infection intensity.

Conclusion

Here we report the successful establishment of a sand fly-*Leishmania* model using a species of *Leishmania* non-pathogenic to mammals using GFP-labelled *L. tarentolae*. We also show that infectivity rates on established infections are similar to those achieved using non-labelled parasites. Finally, we show that it was also possible to detect highly-infected and low-infected individuals without the traditional dissection technique, opening a route to enable identification of infected flies for ongoing experimentation.

Data availability

Figshare: Dataset 1. Non-invasive visualisation and identification of fluorescent *Leishmania tarentolae* in infected sand flies, <https://doi.org/10.6084/m9.figshare.7283546.v1> (Diaz-Albiter, 2018).

Data are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

Grant information

This work is supported by the Wellcome Trust [104111]; and the Royal Society, [NF151329].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We thank Susan Baillie for technical support.

References

- Adler S, Theodor O: Investigation on Mediterranean kala azar X—A note on *Trypanosoma platydictyli* and *Leishmania tarentolae*. *Proc R Soc Lond B Biol Sci*. 1935; **116**(801): 543–544.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Aphasizhev R, Aphasizheva I, Nelson RE, et al.: Isolation of a U-insertion/deletion editing complex from *Leishmania tarentolae* mitochondria. *EMBO J*. 2002; **22**(4): 913–924.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Breitling R, Klingner S, Callewaert N, et al.: Non-pathogenic trypanosomatid protozoa as a platform for protein research and production. *Protein Expr Purif*. 2002; **25**(2): 209–218.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Burza S, Croft SL, Boelaert M: Leishmaniasis. *Lancet*. 2018; **392**(10151): 951–970.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bates PA: Transmission of *Leishmania metacyclic* promastigotes by phlebotomine sand flies. *Int J Parasitol*. 2007; **37**(10): 1097–1106.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Beatty BJ: Genetic manipulation of vectors: A potential novel approach for control of vector-borne diseases. *Proc Natl Acad Sci U S A*. 2000; **97**(19): 10295–10297.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chatton E, Blanc G: Existence de corps leishmaniformes dans les hémato blastes d'un gecko barbaresque *Tarentola mauritanica* L. *Guth. CR Soc Biol*. 1914; **77**(77): 430–433.
- Dey S, Papadopoulou B, Haimeur A, et al.: High level arsenite resistance in *Leishmania tarentolae* is mediated by an active extrusion system. *Mol Biochem Parasitol*. 1994; **67**(1): 49–57.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Diaz-Albiter H: Non-invasive visualisation and identification of fluorescent *Leishmania tarentolae* in infected sand flies. *figshare*. Fileset. 2018.
- Frischknecht F, Baldacci P, Martin B, et al.: Imaging movement of malaria parasites during transmission by *Anopheles* mosquitoes. *Cell Microbiol*. 2004; **6**(7): 687–694.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gajapathy K, Peiris LB, Goodacre SL, et al.: Molecular identification of potential leishmaniasis vector species within the *Phlebotomus (Euphebotomus) argentipes* species complex in Sri Lanka. *Parasit Vectors*. 2013; **6**(1): 302.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gibson W, Bailey M: The development of *Trypanosoma brucei* within the tsetse fly midgut observed using green fluorescent trypanosomes. *Kinetoplastid Biol Dis*. 2003; **2**(1): 1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hassan MM, Widaa SO, Osman OM, et al.: Insecticide resistance in the sand fly, *Phlebotomus papatasi* from Khartoum State, Sudan. *Parasit Vectors*. 2012; **5**(1): 46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hemingway J, Ranson H: Insecticide resistance in insect vectors of human disease. *Annu Rev Entomol*. 2000; **45**(1): 371–391.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kamhawi S: Phlebotomine sand flies and *Leishmania* parasites: friends or foes? *Trends Parasitol*. 2006; **22**(9): 439–445.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kushnir S, Gase K, Breitling R, et al.: Development of an inducible protein expression system based on the protozoan host *Leishmania tarentolae*. *Protein Expr Purif*. 2005; **42**(1): 37–46.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Laveran A, Franchini G: Des hématozoaires du gecko et spécialement de *Herpetomonas tarentolae*, Procédé simple de culture des *Herpetomonas*. *Bull Soc Pathol Exot*. 1921; **14**(14): 323–326.
- Modi GB, Tesh RB: A simple technique for mass rearing *Lutzomyia longipalpis* and *Phlebotomus papatasi* (Diptera: Psychodidae) in the laboratory. *J Med Entomol*. 1983; **20**(5): 568–569.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ouellette M, Fase-Fowler F, Borst P: The amplified H circle of methotrexate-resistant *leishmania tarentolae* contains a novel P-glycoprotein gene. *EMBO J*. 1990; **9**(4): 1027–1033.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Parrot L: Development of *L. tarentolae* in *P. minutus*. *Bulletin de la Société de Pathologie Exotique*. 1934; **27**(9).
- Pimenta PF, Saraiva EM, Rowton E, et al.: Evidence that the vectorial competence of phlebotomine sand flies for different species of *Leishmania* is controlled by structural polymorphisms in the surface lipophosphoglycan. *Proc Natl Acad Sci U S A*. 1994; **91**(19): 9155–9159.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rogers ME, Bates PA: *Leishmania* manipulation of sand fly feeding behavior results in enhanced transmission. *PLoS Pathog*. 2007; **3**(6): e91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Schuster S, Krüger T, Subota I, et al.: Developmental adaptations of trypanosome motility to the tsetse fly host environments unravel a multifaceted *in vivo* microswimmer system. *eLife*. 2017; **6**: pii: e27656.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Soares RP, Turco SJ: *Lutzomyia longipalpis* (Diptera: Psychodidae: Phlebotominae): a review. *An Acad Bras Cienc*. 2003; **75**(3): .301–330.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Vlachou D, Zimmermann T, Cantera R, et al.: Real-time, *in vivo* analysis of malaria ookinete locomotion and mosquito midgut invasion. *Cell Microbiol*. 2004; **6**(7): 671–685.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Volf P, Myskova J: Sand flies and *Leishmania*: specific versus permissive vectors. *Trends Parasitol*. 2007; **23**(3): 91–92.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- WHO: World Health Organization. Geneva: *Leishmaniasis—Fact sheet Updated March*. Accessed: 27 October 2018. 2018.
[Reference Source](#)
- Wilson VC, Southgate BA: Lizard *leishmania*. *Biology of the kinetoplastida*. 1979; **2**: 241–268.
- Zhang WW, Charest H, Matlashewski G: The expression of biologically active human p53 in *Leishmania* cells: a novel eukaryotic system to produce recombinant proteins. *Nucleic Acids Res*. 1995; **23**(20): 4073–4080.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Zieler H, Dvorak JA: Invasion *in vitro* of mosquito midgut cells by the malaria parasite proceeds by a conserved mechanism and results in death of the invaded midgut cells. *Proc Natl Acad Sci U S A*. 2000; **97**(21): 11516–11521.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Referee Status:



Version 1

Referee Report 07 February 2019

<https://doi.org/10.21956/wellcomeopenres.16255.r34472>



Antonio Jorge Tempone 

Laboratório de Biologia Molecular de Parasitos e Vetores-LBMPV, Instituto Oswaldo Cruz - FIOCRUZ, Rio de Janeiro, Brazil

In work titled “Non-invasive visualization and identification of fluorescent *Leishmania tarentolae* in infected sand flies,” the authors developed a system to study interactions between the parasites of *Leishmania* genus and phlebotomine sand flies. The use of a non-human pathogenic *Leishmania tarentolae* favors the students training under better security conditions. Interestingly, this system allows researchers to quickly and inexpensively identify and isolate highly infected individuals from those with low parasite levels, providing greater sample uniformity. Besides, the authors identified a decrease in the life span of infected insects, evidencing the harmful nature of *Leishmania* infection.

Although the authors comment on the use of fluorescent parasites in other parasite-vector interaction models, such as *Plasmodium*-mosquitoes and trypanosomatids-triatomines, there is no citation about the works of Sadlova et al., 2011¹ and Chajbullinova et al., 2012². These authors employed fluorescent *Leishmania* in their works. In the first work, the authors verified the existence of genetic material exchange between *L. donovani* during its development in the gut of *Phlebotomus perniciosus* and *Lutzomyia longipalpis*. In the second study, the authors studied the development of *L. turanica* in three different species of phlebotomines. These works should be included in the discussion and references.

References

1. Sadlova J, Yeo M, Seblova V, Lewis MD, Mauricio I, Volf P, Miles MA: Visualisation of *Leishmania donovani* fluorescent hybrids during early stage development in the sand fly vector. *PLoS One*. 2011; **6** (5): e19851 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Chajbullinova A, Votypka J, Sadlova J, Kvapilova K, Seblova V, Kreisinger J, Jirku M, Sanjoba C, Gantuya S, Matsumoto Y, Volf P: The development of *Leishmania turanica* in sand flies and competition with *L. major*. *Parasit Vectors*. 2012; **5**: 219 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Referee Expertise: Molecular biology of parasite-vector interaction.

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Referee Report 10 January 2019

<https://doi.org/10.21956/wellcomeopenres.16255.r34465>



Yusuf Özbel 

Department of Parasitology, Faculty of Medicine,, Ege University, İzmir, Turkey

In this study, the researchers developed a system for investigating the interactions between sand fly species and Leishmania parasites.

In my opinion, it is very useful, especially for training, and if it is extended to human pathogenic Leishmania species, we can better understand the events happening in the vector sand fly species. This information can give us the explanations why some Leishmania species are specific to some sand fly species and vice versa.

The manuscript is very well written and I do not have any corrections.

It can be accepted as it is.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.**Referee Expertise:** Vector-borne diseases especially sandfly-borne diseases and Leishmaniasis**I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

Referee Report 09 January 2019

<https://doi.org/10.21956/wellcomeopenres.16255.r34466>**Jesus G Valenzuela**

Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health (NIH), Rockville, MD, USA

The work by Diaz-Albiter et al describes the development of a safe system to study *Leishmania*-sand fly interactions that will not need a special containment facility. The group developed a green fluorescent *Leishmania tarentolae* parasite, infected sand flies with these parasites and observed the parasites inside of the gut of the sand fly. They also compared the survival in sand flies of the green fluorescent parasites and the wild type parasites.

The work is relevant particularly because this system will allow vector-parasites studies in a system that does not require stringent biosafety level of containment. There are some questions the authors need to address and some changes to do in the manuscript:

1. In the abstract GFP should be in parenthesis (GFP).
2. Introduction: "argentes" change to "argentipes".
3. Introduction: "well-establish" change to "well-established".
4. Did the authors follow the infection after 5 days post infection?
5. What parasites stages do the authors observe in the sand flies at 5 days post infection?
6. By not extending the days of infection the authors are missing a very important time point, the emergence of metacyclics.
7. This system can be used also to test transmission blocking vaccines.
8. Methods: change "lab" to "laboratory".
9. The authors use sometimes "sandfly" and other times "sand fly" throughout the manuscript, this needs to be consistent for whatever the authors decide to use.
10. Figure 1A. The Y axis numbers do not make much sense. Can the authors write how many parasites were found per insect or per midgut? If the numbers are in the millions please check your calculations.
11. Same for Figure 2C. Can the authors write how many parasites were found per insect or per midgut? If the numbers are in the millions please check your calculations.
12. Figure 1B. I would flip the graph. Show the infected in the bottom and the uninfected in the top. This will match the label of the Y axis. Otherwise it is very confusing.

13. Under “Assessment of *L. tarentolae*-GFP fluorescence in vivo”, change: “did not showed” to “did not show”
14. Discussion. The authors mentioned “This technique can also be extended for medically relevant species of *Leishmania*”. This was actually already shown by Sacks et al using *L. major* fluorescent parasites and by Guevara P et al. (2001)¹. I think the authors should provide the references of *Leishmania* fluorescent parasites being previously used in sand flies. What the presented work emphasizes is actually having a safe system to study vector-parasite interactions using fluorescent parasites. The use of *Leishmania* fluorescent parasite is not novel and it should be mentioned more in this paper, particularly in the paragraph where the authors describe the *Plasmodium* and *Trypanosome* work.

References

1. Guevara P, Pinto-Santini D, Rojas A, Crisante G, Añez N, Ramirez JL: Green fluorescent protein-tagged *Leishmania* in phlebotomine sand flies. *J Med Entomol.* 2001; **38** (1): 39-43 [PubMed Abstract](#)

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Referee Expertise: Vector Biology

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.
