

1 Developing an Intervention for Fall-Related Injuries in Dementia

2 (DIFRID): an integrated, mixed-methods approach

3 Alison Wheatley, Claire Bamford, Caroline Shaw, Elizabeth Flynn, Amy Smith, Fiona Beyer, Chris Fox,

4 Robert Barber, Steve W. Parry, Denise Howel, Tara Homer, Louise Robinson, Louise M. Allan

5

6 Dr Alison Wheatley

7 Research Associate

8 Institute of Health and Society

9 Newcastle University

10 [Alison.wheatley@ncl.ac.uk](mailto:Alison.wheatley@ncl.ac.uk)

11

12 Claire Bamford

13 Senior Research Associate

14 Institute of Health and Society

15 Newcastle University

16 [claire.bamford@ncl.ac.uk](mailto:claire.bamford@ncl.ac.uk)

17

18 Dr Caroline Shaw

19 Research Associate

20 Institute of Health and Society

21 Newcastle University

22 [c.a.shaw@hotmail.co.uk](mailto:c.a.shaw@hotmail.co.uk)

23

24 Elizabeth Flynn

25 Physiotherapy Operational Lead - Rehabilitation Team

26 The Newcastle upon Tyne Hospitals NHS Foundation Trust

27 [Elizabeth.Flynn@nuth.nhs.uk](mailto:Elizabeth.Flynn@nuth.nhs.uk)

28

29 Amy Smith

- 1 Occupational Therapy Clinical Lead
- 2 Tees, Esk and Wear Valleys NHS Foundation Trust
- 3 [amy.smith16@nhs.net](mailto:amy.smith16@nhs.net)
- 4
- 5 Fiona Beyer
- 6 Research Associate
- 7 Institute of Health and Society
- 8 Newcastle University
- 9 [fiona.beyer@ncl.ac.uk](mailto:fiona.beyer@ncl.ac.uk)
- 10
- 11 Professor Chris Fox
- 12 Professor of Old Age Psychiatry
- 13 University of East Anglia
- 14 [Chris.fox@uea.ac.uk](mailto:Chris.fox@uea.ac.uk)
- 15
- 16 Dr Robert Barber
- 17 Consultant Old Age Psychiatrist / Hon Clinical Senior Lecturer
- 18 Northumberland Tyne and Wear NHS Foundation Trust
- 19 [robert.barber@ncl.ac.uk](mailto:robert.barber@ncl.ac.uk)
- 20
- 21 Dr Steve W. Parry
- 22 Clinical Senior Lecturer and Consultant Physician
- 23 Newcastle University
- 24 [Steve.Parry@nuth.nhs.uk](mailto:Steve.Parry@nuth.nhs.uk)
- 25
- 26 Denise Howel
- 27 Senior Lecturer in Epidemiological Studies
- 28 Institute of Health and Society
- 29 Newcastle University
- 30 [denise.howel@ncl.ac.uk](mailto:denise.howel@ncl.ac.uk)
- 31 Tara Homer
- 32 Research Associate

1 Institute of Health and Society  
2 Newcastle University  
3 [tara.homer@ncl.ac.uk](mailto:tara.homer@ncl.ac.uk)  
4  
5 Professor Louise Robinson  
6 Professor of Primary Care and Ageing  
7 Institute of Health and Society  
8 Newcastle University  
9 [a.l.robinson@ncl.ac.uk](mailto:a.l.robinson@ncl.ac.uk)  
10  
11 Professor Louise M Allan (Corresponding author)  
12 Professor of Geriatric Medicine  
13 Institute of Health Research  
14 University of Exeter  
15 South Cloisters  
16 St Luke's Campus  
17 Heavitree Road  
18 Exeter  
19 EX1 2LU  
20 Telephone : +44 1392 726025  
21 [l.allan@exeter.ac.uk](mailto:l.allan@exeter.ac.uk)

## 1 Abstract

### 2 Background

3 Falls in people with dementia can result in a number of physical and psychosocial consequences.  
4 However, there is limited evidence to inform how best to deliver services to people with dementia  
5 following a fall. The aim of the DIFRID study was to determine the feasibility of developing and  
6 implementing a new intervention to improve outcomes for people with dementia with fall-related  
7 injuries; this encompasses both short-term recovery and reducing the likelihood of future falls. This  
8 paper details the development of the DIFRID intervention.

### 9 Methods

10 The intervention was designed using an integrated, mixed-methods approach. This involved a realist  
11 synthesis of the literature and qualitative data gathered through interviews and focus groups with  
12 health and social care professionals (n=81). An effectiveness review and further interviews and  
13 observation were also conducted and are reported elsewhere. A modified Delphi panel approach  
14 with 24 experts was then used to establish a consensus on how the findings should translate into a  
15 new intervention. After feedback from key stakeholders (n=15) on the proposed model, the  
16 intervention was manualised and training developed.

### 17 Results

18 We identified key components of a new intervention covering three broad areas:

- 19 • Ensuring that the circumstances of rehabilitation are optimised for people with dementia
- 20 • Compensating for the reduced ability of people with dementia to self-manage
- 21 • Equipping the workforce with the necessary skills and information to care for this patient  
22 group

23 Consensus was achieved on 54 of 69 statements over two rounds of the Delphi surveys. The  
24 statements were used to model the intervention and finalise the accompanying manual and protocol  
25 for a feasibility study. Stakeholder feedback was generally positive and the majority of suggested  
26 intervention components were approved. The proposed outcome was a 12-week complex  
27 multidisciplinary intervention primarily based at the patient's home.

### 28 Conclusions

29 A new intervention has been developed to improve outcomes for people with dementia following a  
30 fall requiring healthcare attention. The feasibility of this intervention is currently being tested.

### 31 Trial registration

32 ISRCTN41760734 (16/11/2015)

- 1 Keywords

- 2 dementia; falls; intervention development; Delphi consensus; realist synthesis

## 1 Background

2 People with dementia who live in their own home make up 70% of all people living with dementia in  
3 the UK [1], and are ten times more like to fall as people without dementia [2]. The negative  
4 consequences of falls are greater for people with dementia than for other older people [3]. While  
5 even non-injurious falls can result in psychosocial consequences such as loss of confidence and fear  
6 of falling [4], functional decline in people who sustain injuries may be greater than in those who do  
7 not sustain injuries [5, 6]. Despite this, few trials have specifically addressed the management of fall-  
8 related injuries in people with dementia. While multifactorial interventions by specialist falls services  
9 are effective in preventing further falls in older people without dementia [7, 8], evidence of their  
10 effectiveness for people with dementia is inconclusive [9-11]. Similarly, falls-prevention exercise  
11 programmes such as Otago [12] have little evidence of efficacy for people with dementia, though  
12 some work has been done on tailoring the programme for individuals [13, 14]. There is, however,  
13 some evidence that rehabilitation interventions may result in improvements in motor performance  
14 in people with dementia [15] and that motor training can increase physical activity in people with  
15 dementia without increasing the risk of falls [16]. Recently published guidelines acknowledge that  
16 multifactorial falls interventions may not be suitable for a person living with severe dementia, but  
17 provide no recommendations on how to optimise falls interventions for this patient group [17].

18 The brief for this study was therefore to develop a new complex intervention to improve care for  
19 community-dwelling people with dementia with fall-related injuries. In response to calls for a more  
20 systematic approach to, and greater transparency in, intervention development [18-20], this paper  
21 describes the development process in detail. This includes presenting (a) the causal factors and  
22 change mechanisms underpinning falls and rehabilitation care for this patient group; (b) the  
23 outcomes of a consensus-seeking process based on this initial work; (c) the development of a logic  
24 model; and (d) the development of intervention materials.

## 1 Methods

2 The development of the intervention involved qualitative work to map existing care pathways [21]  
3 and explore the views of stakeholders on the content and delivery of a new intervention [22]; an  
4 effectiveness review [5]; a realist synthesis of the literature; a prospective diary study to provide  
5 information on recruitment sources and existing service use; consensus panel meetings of experts; a  
6 Delphi survey; and further qualitative work to elicit stakeholder feedback on the proposed  
7 intervention. The findings were then used to develop a logic model, protocol [23] and intervention  
8 materials. The feasibility and acceptability of the new intervention is currently being evaluated.  
9 Figure 1 illustrates the process of intervention development.

10 [Figure 1 here]

11 *Figure 1: Intervention development*

### 12 Identifying causal factors and change mechanisms

#### 13 Qualitative work and formative realist analysis

14 The initial qualitative work comprised 58 semi-structured interviews and 5 focus groups with health  
15 and social care professionals (full details of qualitative work are reported elsewhere [22]; this also  
16 included observation of care delivery and interviews with patients and carers, although these were  
17 not included in this formative work due to the timescales involved). Professionals were identified  
18 through snowball sampling facilitated by local study investigators. Recruitment continued until data  
19 saturation was reached. Details of participants are provided in additional file 1. Interviews and focus  
20 groups were audio recorded, transcribed, and anonymised prior to analysis.

21 We used realist methodology [24, 25] to identify both causal factors and change mechanisms. This is  
22 an approach to literature review and data analysis which seeks to answer the question 'what works  
23 for whom under what circumstances, how and why', describing mechanisms which produce  
24 particular outcomes in specific contexts [26]. Members of the qualitative team (AW, CB) generated  
25 first "if-then" statements and grouping these according to emerging themes [27]. We refined the if-

1 then statements, looking for data that could be interpreted as a causal factor or a change  
2 mechanism. We expressed these using the realist framework of Context, Mechanism or Outcome  
3 [25], with mechanisms further divided into – ‘resource’ (the intervention component added) and  
4 ‘reasoning’ (what change this resource will produce) [28]. Finally, we presented these initial Context-  
5 Mechanism-Outcome configurations (CMOCs) to a panel of clinicians (LA, RB, CF, SP, LR) and further  
6 refined them based on their feedback. This framework formed the basis for extracting data from the  
7 literature. For examples of finalised CMOCs, see table 1.

#### 8 *Effectiveness review*

9 This has been reported elsewhere [5]. The review could not draw definitive conclusions, since few  
10 interventions were aimed at patients with dementia, and those that were focused mainly on hip  
11 fracture. It therefore indicated that the development of a new intervention was warranted.

#### 12 *Realist synthesis*

13 The protocol for the realist synthesis was registered with PROSPERO (CRD42016040059).

#### 14 *Search strategy*

15 Searches were limited to English. Initially we undertook a comprehensive search (SR). This took place  
16 in November 2015, and was designed to provide a clear understanding of the interaction between  
17 interventions, characteristics of people with dementia and contextual factors around a fall. Iterative  
18 targeted searches aimed to build on that understanding and were completed by March 2017 (FB). As  
19 the aim of the paper is to describe the intervention development process as it occurred, the  
20 searches have not been updated.

21 Comprehensive searches were conducted in MEDLINE, CENTRAL, Health Management Information  
22 Consortium, EMBASE, CINAHL, Web of Science, Allied and Complementary Medicine Database, and  
23 Physiotherapy Evidence Database (PEDro) (see additional file 2 for an example search strategy).  
24 Trials registers were searched, but further grey literature searching was not conducted. Results from  
25 all databases were imported to Endnote. Targeted searches took place in MEDLINE and CINAHL on



1 EBSCO (see additional file 3 for an example targeted search strategy). Additional papers were  
2 identified through citation chaining of included papers and relevant systematic reviews and hand  
3 searches. Figure 2 demonstrates the flow of studies.

4 [Figure 2 here]

5 *Figure 2: Diagram of the search, screening, selection and extraction process*

#### 6 *Data extraction and CMOc refinement*

7 Data were extracted from included papers using a bespoke online form. This included methodology,  
8 appraisal using the Mixed Methods Appraisal Tool [29]; an intervention description, as applicable,  
9 using the TIDieR framework [30]; and evidence describing contexts, mechanisms or outcomes. Data  
10 were extracted by two reviewers independently, one clinician (LA, BB, CF, SP, SL) and one non-  
11 clinician (CB, FB, CS, AW). Data were discussed at a meeting of reviewers and disagreements  
12 resolved. The qualitative team (CB, CS, AW) analysed and summarised the data. Following this  
13 process, the wording of each CMOc and the set of themes were refined (CB, CS, AW). The process  
14 was repeated for the additional papers identified through targeted searches and citation chaining.

#### 15 *Delphi consensus process*

16 We convened a multidisciplinary panel of 24 expert health and social care professionals (see additional  
17 file 1) to review the initial findings and make recommendations regarding the design of the complex  
18 intervention using a modified Delphi panel approach [31-33] (see additional file 4). Panellists were  
19 selected who (1) represented of a range of stakeholder groups identified to be important to the care  
20 of people with dementia who fall; (2) were in contact with people with dementia who fall and/or (3)  
21 had relevant academic expertise; (4) and were able to attend face-to-face meetings. The consensus  
22 panel meetings were audio recorded (with the consent of participants) and transcribed for analysis.

#### 23 *Consensus panel meeting 1*

24 Prior to the first meeting (March 2017), the panel received summaries of the qualitative work,  
25 effectiveness review, and the realist synthesis. At the meeting, members were split into groups to

1 discuss three key aspects of the intervention: feasibility and setting; content; and outcome measures.  
2 Each group discussed all issues. Key points from the discussions were fed back and areas of initial  
3 agreement and dissent were identified.

#### 4 Delphi surveys

5 Following the first consensus panel meeting, a series of statements were identified and sent to panel  
6 members via an online survey tool. Members were asked to respond to specific questions regarding  
7 feasibility of the setting; staffing and training requirements; components of the intervention; and  
8 outcome measures for the feasibility study. A threshold of two-thirds agreement of those completing  
9 the survey was chosen to represent consensus. Responses were received from 14 panel members.

10 Since consensus was not achieved on all items, a second round of the survey was conducted which  
11 included the results of the first round. This gave members the opportunity to revise their responses.  
12 Responses were received from 13 panel members.

13 All respondents completed all items in both rounds of the survey. To facilitate free expression of  
14 opinion, only the independent moderator (BE) could access non-anonymised data. Statements  
15 included in the survey along with consensus results are given in additional file 5.

#### 16 Stakeholder feedback

17 In parallel with the surveys, additional focus groups and interviews were conducted with a range of  
18 stakeholders to explore the feasibility and acceptability of the draft intervention (see additional file 1  
19 for details of participants). These were invited from the pool of participants who took part in WP2,  
20 supplemented by snowball sampling of professionals and additional patients and carers recruited via  
21 the North East and North Cumbria CRN Case Register. Interviews and focus groups were audio-  
22 recorded, transcribed, and anonymised prior to thematic analysis.

## 1 Consensus meeting 2

2 At the second meeting (June 2017), panel members considered the draft protocol for the feasibility  
3 study; results of stakeholder feedback on the proposed intervention; and the proposed roles of  
4 members of the multidisciplinary team (MDT). Small group discussions were facilitated as at the first  
5 meeting.

## 6 Collation of results and development of a logic model

7 We collated the findings of the final round of the Delphi survey, consensus panel discussion, and  
8 stakeholder feedback to finalise the protocol for the feasibility study and model the intervention.

9 The logic model was developed by the qualitative team (CB, AW) informed by existing logic model  
10 templates [34, 35] and was discussed by the Trial Oversight Committee (TOC).

## 11 Preparation of intervention resources

12 Three specific resources needed to implement the intervention were identified from the protocol  
13 and logic modelling process: an assessment document, a manual and a staff training programme.

14 These were developed by the study team (LA, CB, EF, AS, AW) with reference to the final consensus  
15 statements, protocol, and logic model and were further discussed by the TOC and all co-  
16 investigators.

## 17 Results

18 Nine CMOcs were identified as key components of a new intervention for people with dementia  
19 following a fall. These were grouped into three themes:

- 20 • Ensuring that the circumstances of rehabilitation are optimised for people with dementia  
21 (CMOcs 1-3)
- 22 • Compensating for the reduced ability of people with dementia to self-manage (CMOcs 4-6)
- 23 • Equipping the workforce with the necessary skills and information to care for this patient group  
24 (CMOcs 7-9)

1 In presenting each CMOc, we synthesise evidence from the literature, map relevant consensus  
2 statements, describe how intervention components agreed by the consensus process were  
3 operationalised for the DIFRID intervention, and present results of stakeholder feedback. A fourth  
4 theme, covering practicalities relating to intervention delivery and evaluation, is also discussed.  
5 Quotations presented are identified by a unique participant ID. Additionally, role and service type is  
6 provided for professionals. All identifying information, including location, has been removed to  
7 maintain confidentiality.

## 8 Theme 1: Ensuring that the circumstances of rehabilitation are optimised for people 9 with dementia

10 This theme concerns the ways in which broader contextual factors, such as setting and  
11 comorbidities, can affect the engagement of people with dementia in interventions. The outcomes  
12 of the consensus survey and operationalisation of each CMOc are shown in Table 1.

13 [Table 1 here]

### 14 CMOc1: Managing pain

15 Pain is associated with impaired mobility and physical functioning [36-38] and increased agitation  
16 and aggression [39-42] in people with cognitive impairment or dementia. Sleeping and mood  
17 disorders in people with dementia have also been linked to higher pain levels [43]. People with  
18 dementia who are in pain may therefore find it more difficult to engage fully with an intervention.  
19 However, recognising pain in people with dementia can be challenging as they may be unable to  
20 verbally communicate their pain [44].

21 The consensus panel agreed that identifying pain should be part of the DIFRID intervention.  
22 Stakeholders highlighted the complexities of assessing pain in people with dementia:

23 *There are so many different implications. It is not just about us scoring pain. If you*  
24 *are talking about pain assessment, you need to do it properly. That, again, is*

1           *multi-factorial. You need to use the appropriate pain scoring. If you are talking*  
 2           *about people who have got moderate dementia who are cognitively impaired,*  
 3           *you need to be thinking about something like the Abbey Pain Scale or something*  
 4           *like that. It is not verbal. It is behavioural, body language, facial expressions, all*  
 5           *that sort of stuff.*

6           ***(Prof 122, pain nurse, focus group with specialist nurses)***

## 7 **CMOc2: Ensuring a supportive environment**

8       People with dementia may become distressed in an unfamiliar environment, resulting in an  
 9       exacerbation of symptoms [45]. Moreover, since people with dementia may find it difficult to  
 10      articulate basic needs, such as hydration, these may go unrecognised by staff [45]. Carers in one  
 11      qualitative study described negative experiences of hospitalisation, such as a deterioration in  
 12      patients' health, and were keen to avoid readmission [46]. Home-based exercise interventions have  
 13      been shown to be feasible for at least some patients with cognitive impairment and hip fracture [47-  
 14      50], though some studies reported problems with adherence [51]. Literature relating to patients  
 15      with other fall-related injuries was not found.

16      The consensus panel agreed that the home environment would be the most appropriate location for  
 17      the DIFRID intervention. Stakeholder feedback on this aspect of the intervention was generally  
 18      positive, although some stakeholders highlighted the need for flexibility:

19           *I don't know how that would fit in, because we used to enjoy walking, you see, up*  
 20           *in the hills, and I'm not quite sure how that would fit in with physio in the home.*

21           ***(Interview, Patient 15 and Carer 15)***

22      The intervention, therefore, can be delivered in the most appropriate environment for the activities  
 23      and goals identified by participants.

## 24 **CMOc3: Adopting a holistic approach**

25      Holistic assessments to discover and manage falls risk factors emerged as an important theme.

26      Comorbidities that increase mortality risk during and after hospitalisation for hip fracture in older

1 people may go unrecognised and undiagnosed [45, 52, 53]. Psychosocial factors, such as depression  
 2 [53, 54] and social isolation [54], may also be important for the wellbeing and recovery of people  
 3 with dementia following a fall. Holistic assessments, such as Comprehensive Geriatric Assessment  
 4 (CGA), have been shown to improve outcomes for people with cognitive impairment or delirium who  
 5 have fallen [53-55]. Holistic assessment may also aid patient and caregiver understanding of the  
 6 causes of falls [56] and facilitate early intervention for other health issues which might otherwise  
 7 undermine therapy [57].

8 Stakeholders suggested including a social worker in the DIFRID MDT to facilitate assessment of social  
 9 circumstances:

10 *I think it's really important that people get a review of their social circumstances,*  
 11 *especially if they've had a fall. Sometimes [...] the reason that they've fallen is*  
 12 *that they're trying to do something that they would benefit from having a care*  
 13 *package to prevent them having to do themselves.*

14 ***(Prof 71, reablement support worker, focus group at specialist inpatient***  
 15 ***rehabilitation unit)***

16 The consensus panel subsequently agreed that a social worker should be available on referral.  
 17 Additional areas for assessment suggested by stakeholders included: foot assessment; nutrition;  
 18 frailty; existing equipment and aids; and a detailed cognitive profile. Details of the assessment,  
 19 conducted using skilled observation or verbal report from patient and carer, are shown in table 2.

20 **Table 2: Sections of the assessment and intervention document**

<b>Generic assessment (by physiotherapist or occupational therapist)</b>	
Falls history	
Falls risk assessment (including fear of falling, nutrition, fluid intake, pain, urinary incontinence, bowel incontinence, supportive footwear, visual impairment not corrected with glasses)	
Past medical history and comorbidities	
Medication	
Current activity levels	
Challenging behaviour and sleep disturbance	
Assessment of the needs of the informal carer	
Current mobility	
<b>Physiotherapy assessment</b>	<b>Occupational therapy assessment</b>

Posture and general observations of pain, sensation and tone	Details of home environment
Lying and standing BP	Self-care and productivity
Range of movement	Cognition
Muscle power	Task observations
TUG	Functional difficulties relating to spatial awareness, vision and hearing
<b>Intervention planning</b>	
Needs list	
Action planning and patient goals	
MDT record	
Referrals	

1

2 Stakeholders emphasised the need to interpret the results of holistic assessment and identify clear  
3 processes for addressing issues raised:

4 *For example, incontinence, you know, you are not going to engage someone in an*  
5 *exercise programme, or encourage them to stabilise their gait, their balance or*  
6 *posture if actually their real problem is they are retaining urine. They are getting*  
7 *overflow, and when they stand up to go they have a real sense of urgency and*  
8 *they are desperate. You can put in every intervention you like. Unless you address*  
9 *that problem... You need someone who is going to think about that, and*  
10 *understand what is going on. The reason they are in a hurry to get up and go to*  
11 *the loo is not because they are going frequently. They frequency is due to another*  
12 *problem that hasn't been picked up.*

13 ***(Prof 122, pain nurse, focus group with specialist nurses)***

14 In developing the assessment and intervention materials, we therefore added a section dedicated to  
15 referrals for issues identified during assessment, and tasked the MDT with reviewing this. A  
16 substantial component of the DIFRID staff training programme focused on using the assessment  
17 document and managing any problems identified.

1 Theme 2: Compensating for the reduced ability of people with dementia to self-  
2 manage

3 This theme concerns the ways in which intervention delivery can be adapted to compensate for the  
4 symptoms and challenges of dementia. The outcomes of the consensus survey and  
5 operationalisation of each CMOc are shown in table 3.

6 [Table 3 here]

7 **CMOc4: Embedding interventions in day to day life**

8 Individually tailoring exercises to the preferences, interests, and physical and cognitive abilities of  
9 people with dementia has been described as 'vital' to successful interventions for this patient group  
10 [49]. Cognitive impairment may affect the ability of patients to follow instructions and consequently,  
11 rehabilitation success [47, 51, 53-55, 58-62]. However, some people with dementia may have  
12 relatively well preserved procedural memory which may enable them to form new habits [54].  
13 'Embedding' interventions into existing routines could also help make them habitual [63-65].  
14 Effective tailoring requires specialised training for staff and carers involved in intervention delivery  
15 [49, 66]; including a staff training component in the intervention was therefore seen as essential (see  
16 CMOc8).

17 Stakeholders agreed with the consensus panel's recommendation to use the principles underlying  
18 the Otago exercise programme (i.e. individually tailored; progressive; stable and sustainable; and  
19 using walking alongside strength and balance) [12], although they indicated that implementation of  
20 this programme is often inconsistent with the recommended format due to resource limitations.  
21 Stakeholders emphasised that meaningful activities should include mental and social stimulation as  
22 well as physical activity:

23 *This gentleman had really bad dementia. He had sundown so he was up all night.*  
24 *The family came in, and we had a game of dominos. I couldn't communicate with*  
25 *him. You bring out the dominos and he won every time. It was like a different*



1            *person came out in that dominos. [...] Then, by making him stay awake all day*  
 2            *and doing meaningful activities to keep him active, he was more likely to sleep at*  
 3            *night. [...] He is not getting up and falling over.*

4            ***(Prof 121, focus group with specialist nurses)***

5            Music and dance were also considered particularly valuable. Exploring the barriers (including cultural  
 6            barriers) to meaningful activity was identified as one way to increase the likelihood of successfully  
 7            engaging patients in new activities.

8            While setting patient-centred goals achieved a high level of consensus among the panel, some  
 9            stakeholders had reservations about how this might work in practice:

10            *I don't think I could do it. Like, make a cup of tea. I wouldn't trust myself.*

11            ***(Interview, Patient 13)***

12            Professional stakeholders also identified potential problematic elements of goal setting, including  
 13            the difficulty of engaging people with dementia in setting goals, the potential for them to lose  
 14            interest in things they previously enjoyed, and ensuring goals were those of the patient and not only  
 15            the carer.

16            **CMOc5: Providing ongoing support**

17            One quarter (24%) of re-admissions following hip fracture surgery are due to 'failure of  
 18            rehabilitation'—including deterioration, further falls, and inability to cope[57]. This suggests that the  
 19            duration and/or dosage of existing rehabilitation may be insufficient. , As people with dementia  
 20            typically have difficulties with problem solving and self-management, providing only short-term  
 21            interventions may be particularly problematic for this patient group. Professionals in the initial  
 22            qualitative study felt that existing interventions were often too short and lacked continuity in  
 23            content and staffing [21]. They proposed regular follow-up and review to help identify new problems  
 24            or relapses and maintain continuity of care.

1 The intensity and duration of the intervention proved to be the most contentious aspects of the  
2 intervention among the consensus panel and stakeholders. Ultimately, the consensus panel were  
3 constrained by the realities of the project timescale, which could only accommodate a twelve-week  
4 intervention period. Providing on-going support was therefore not feasible. However, the panel  
5 allowed for up to a total of 22 intervention sessions over twelve weeks; this is substantially more  
6 than is provided by many existing services, which our initial qualitative work found were typically  
7 provided for between two and six weeks. The Delphi survey therefore included questions on setting  
8 appropriate boundaries.

9 All groups of stakeholders stressed the need to tailor the number of intervention sessions to the  
10 individual. However, community-based professionals, particularly those in rural areas, raised  
11 concerns over the feasibility of delivering this number of sessions both within and outside the  
12 context of a trial. The duration of individual sessions and the intervention overall were also queried  
13 by some participants:

14 *You need at least, you know, half of that time even strike up a rapport, for them*  
15 *to remember, possibly, who you are, for you to engage with the carer, and that's*  
16 *before you've even done anything and before you've even assessed the person or*  
17 *given them any intervention. That's every time, because every time is like a new*  
18 *time.*

19 ***(Prof 124, physiotherapist, focus group with community health and social care***  
20 ***professionals)***

21 Other participants questioned whether the allotted twelve weeks would be long enough for all  
22 referrals to have been acted upon and for alternative services to have been put in place to provide  
23 ongoing support. Carers also expressed concern about what would happen after the intervention:

24 *That would be my only concern. You're leaving people, then, in limbo. You're*  
25 *offering them something that isn't there anymore. It was there, but 'oh, that's not*  
26 *there now'.*

1           ***(Interview, Carer 12)***

2   The intervention therefore includes mid-point and final review sessions, where intervention staff  
3   check the status of referrals, treat new issues arising during the intervention period, and signpost  
4   participants to other relevant services (such as activity groups) to help maintain progress after the  
5   completion of the twelve-week intervention.

6   **CMOc6: Involving carers in intervention delivery**

7   The involvement of family carers is frequently recommended to improve adherence and outcomes  
8   of interventions [53, 54, 67, 68]. However, this implicitly assumes that carers have capacity and the  
9   skills to assist in intervention delivery. Many family carers report feeling isolated, helpless, and  
10   overstretched by providing care as well as dealing with their own health problems and other  
11   commitments [46, 67]. Having realistic expectations of carers is therefore important [69]. Factors  
12   shown to facilitate carer involvement include exploring concerns about time requirements and  
13   disruption to routines [70], understanding that carers may have difficulty of acknowledging that they  
14   need help [70], and explicitly discussing potential benefits of a rehabilitation intervention to both  
15   people with dementia and carers [70-73]. Carers may also benefit from interventions tailored to  
16   their own needs [66, 74-77]. Carer behaviours, such as preventing the person with dementia from  
17   moving around in order to avoid falls, can negatively influence the relationship between carer and  
18   patient [46] and impede recovery.

19   The consensus panel agreed that educating patients and carers about positive risk and falls  
20   prevention was important. This was also deemed beneficial by stakeholders:

21           *The physios and OTs [...] can assess whether or not that person needs signposting*  
22           *to have some more help. I'm not saying you'd have to have somebody come in*  
23           *with them and do the carer support, but I do think that training them what to*  
24           *look out for, carer fatigue and the stress side of things.*

25           ***(Interview, Carer 12)***

1 However, professionals also emphasised the importance of ensuring that carer needs do not  
2 overshadow those of the patient. To address this concern, the DIFRID training programme includes  
3 advice on managing triadic consultations.

#### 4 Theme 3: Equipping the workforce with the necessary skills and information to care 5 for people with dementia

6 This theme concerns both the training needs of staff and the practical organisation of interventions  
7 to improve information gathering and communication. The outcomes of the consensus survey and  
8 operationalisation of each CMOc are shown in table 4.

9 [Table 4 here]

#### 10 CMOc7: Developing a detailed understanding of the patient

11 A detailed understanding of the patient is fundamental providing tailored, person-centred care in  
12 dementia. As people with dementia may struggle with giving full and accurate medical histories [45,  
13 57, 78], direct observation of the patient in the environment in which they fell was recommended by  
14 professionals in the initial qualitative study [22]. Additional context or confirmation can be provided  
15 by carers [45, 46] or patients' GPs [78]. Drawing on carer expertise to facilitate the care of people  
16 with dementia in hospital has been shown to be effective in reducing agitation and distress and  
17 improving carer satisfaction, though levels of patient satisfaction were not reported [60].

18 Stakeholder feedback about this aspect of the intervention was positive, particularly around using  
19 carers as information sources. Professionals also agreed that assessment by observation was  
20 important, particularly with regard to how participants get around the house.

#### 21 CMOc8: Equipping staff members with appropriate skills

22 Staff members may lack specific training in working with people with dementia and their families,  
23 and negative views about people with dementia and their ability to participate in an intervention  
24 have been reported [66, 75-77]. Several authors recognised the value of providing specialised

1 training of staff to work with older adults and people with dementia, though few provided detailed  
2 information on the content of such training [78-80]. Data from the qualitative study suggests training  
3 should cover dementia-specific adaptation to practice, as well as challenging negative attitudes  
4 towards those with dementia [22]. Training in how best to engage with carers could also be  
5 beneficial [60].

6 Stakeholders identified training as one of the most crucial components of the intervention:

7 *What's jumping out to me is the dependence on the staff training. From a list of*  
8 *interventions none of those are really, hugely, a step away from what we cover.*  
9 *But I know, definitely, still in our organisation staff still need to understand that*  
10 *you can't deliver the same package to someone with a physical condition as to*  
11 *somebody with some challenges, whatever they are.*

12 ***(Prof 35, dementia and falls co-ordinator, focus group with community health***  
13 ***and social care professionals)***

#### 14 **CMOc9: Improving pathways and referral**

15 Collaboration between professionals is an important factor in whether patients receive effective  
16 treatment [52, 81, 82]. A range of social and contextual factors influences decisions to refer to  
17 services, including lack of confidence in the service provided, reluctance to share responsibility for  
18 patient care, or a perception that the patient would not benefit from the service [57, 80]. The initial  
19 qualitative study found staff often lacked knowledge of local services for people with dementia with  
20 fall-related injuries [22]; however, this evidence suggests that a simple lack of knowledge may not be  
21 the only barrier to successful care. The advantages of formalised care pathways include increasing  
22 efficiency of diagnosis and beginning treatment, increasing consistency of care, reducing risk of  
23 errors, reducing costs, and improvements in staff knowledge and team relations [83, 84]. Developing  
24 an evidence-based pathway requires collaboration and input from stakeholders including health  
25 professionals, patients and family members [85]. Ultimately, the consensus panel agreed that  
26 developing a new care pathway for fall-related injuries in dementia was outside of the scope of this

1 study, though the issues of communication and referral were addressed, and the proposed MDT  
2 meetings were seen as a way of maximising use of existing pathways.

3 Stakeholders raised concerns over the feasibility of organising MDT meetings, particularly in rural  
4 areas. While the use of technology could potentially enable virtual MDT meetings, issues were raised  
5 over security and the need for encryption. Overall, professional stakeholders identified a need to  
6 clarify the roles of each member of the DIFRID MDT; this was subsequently discussed at the second  
7 consensus panel meeting. Stakeholders additionally suggested including dietitian/nutritionist;  
8 Alzheimer's Society outreach workers; and advocacy advisers in the MDT. Potential benefits of  
9 including a community psychiatric nurse (CPN) in the MDT were: (i) access to mental health records,  
10 which provided information about dementia, medications and other interventions; (ii) the potential  
11 for rapid referrals and specialist support; and (iii) the potential role of CPNs in reviewing  
12 medications:

13 *[The CPN at our service] can pull up information on where people are at in terms*  
14 *of the support and input that they have had already when they've last been*  
15 *reviewed at memory services. She can review their medications as well which can*  
16 *be really helpful.*

17 ***(Prof 71, reablement support worker, focus group at specialist inpatient***  
18 ***rehabilitation unit)***

19 These additional staff have not been included in the MDT, but intervention materials signpost  
20 therapists to refer to them as required.

#### 21 [Theme 4: Intervention delivery and evaluation](#)

22 The remaining consensus statements concerned issues of practicality and feasibility for the pilot  
23 study (for example, inclusion criteria, recruitment, and outcome measures).

## 1 Design and feasibility

2 In round one of the Delphi survey, 93% of the consensus panel agreed that a complex intervention  
3 was needed. It was deemed feasible to recruit 10 patients from each of three sites to the feasibility  
4 study. Defining the inclusion criteria for the intervention proved more contentious among the panel.  
5 The original brief for this study was to design a new intervention for people with dementia following  
6 a fall-related injury. In the initial interviews and focus groups professionals argued that early  
7 intervention, prior to significant injury, would be more beneficial. The consensus panel also agreed  
8 that the intervention should include patients with non-injurious falls. However, the TOC  
9 subsequently strongly recommended amending this to a fall for which healthcare attention was  
10 sought. Consensus regarding the time period within which patients had to be recruited following a  
11 fall was not reached after two rounds of surveys. Following discussion at the second meeting, it was  
12 agreed that patients could be recruited up to one month after the index fall.

## 13 Outcome measures

14 One aim of the feasibility study is to assess the suitability and acceptability of outcome measures.  
15 While the number of falls was seen as the most appropriate outcome measure by the consensus  
16 panel, other stakeholders expressed reservations about the sensitivity of this measure:

17 *There are maybe subtleties there, from my thinking, that if it was just based on*  
18 *that what might seem like a fall is actually an improvement because the person*  
19 *does feel more confident, is doing more things but is having non-injurious falls as*  
20 *a side-line.*

21 ***(Prof 35, dementia and falls co-ordinator, focus group, community health and***  
22 ***social care professionals)***

23 To address these concerns, a range of outcome measures are being used in the feasibility study,  
24 including measures of function, quality of life and carer burden [23].

## 25 Logic model

26 A logic model (Figure 3) demonstrates the flow of intervention activities to meet project goals.

1 [Figure 3 here]

2 *Figure 3: Logic model*

3

## 4 Discussion

5 We used a mixed-methods approach to develop the DIFRID intervention. We identified causal  
6 factors and change mechanisms through analysis of qualitative data collected in an earlier phase of  
7 the study and a realist synthesis of the literature. This is summarised in three broad themes:

- 8 • Ensuring that the circumstances of rehabilitation are optimised for people with dementia
- 9 • Compensating for the reduced ability of people with dementia to self-manage
- 10 • Equipping the workforce with the necessary skills and information to care for this patient group.

11 An expert panel considered how best to translate these concepts into a new intervention. Consensus  
12 among the panel on which components should be included was achieved through two rounds of a  
13 Delphi survey. This process allowed us to integrate practical, empirical data from experts and  
14 practitioners with evidence from previous studies to create a robust, theoretically-informed design  
15 for a new intervention.

16 Despite the structured approach to intervention development, not all of the CMOcs that emerged  
17 from the initial synthesis were equally present in the consensus surveys. CMOc5, for example, which  
18 concerned ongoing support and follow-up of people with dementia, was deemed beyond the scope  
19 of this study; panel members expressed concerns regarding practicality and feasibility of engaging in  
20 such follow-up when working within constraints such as funding, existing multidisciplinary teams,  
21 existing service provision, and the 12-week limit of the trial. Moreover, for practical reasons relating  
22 to costing the intervention, it was difficult to allow the number of sessions to be open-ended. The  
23 12-week intervention period is quite short in comparison with some trials of exercise in older people  
24 [86]. However, there are a number of trials which have successfully used this intervention period. In  
25 our development work, we found people with dementia received few interventions, often limited to



1 2 or 6 weeks, so a 12-week intervention is a substantial improvement [21]. Additionally, at the end  
2 of the DIFRID intervention therapists are encouraged to refer participants on to community falls  
3 groups or other appropriate ongoing services. It is possible that in future development of this  
4 intervention we could consider extending the intervention beyond 12 weeks but this will not be  
5 possible within the funding for our planned feasibility trial.

6 CMOc9 refers to the creation of a centralised pathway, which was similarly considered beyond the  
7 scope of the study; instead, the consensus process focused on improving communication within and  
8 between staff. Not all components were systematically translated and included in the Delphi survey;  
9 this led to the omission of a statement relating to blanket pain relief as described in CMOc1, for  
10 example. Potential pitfalls associated with this kind of iterative process of intervention development  
11 therefore include ensuring follow-through of ideas at each stage and the potential disconnect  
12 between theoretical ideals and what is considered practical and feasible in everyday practice.  
13 Though we aimed to follow processes for intervention development [19, 20], these were not always  
14 smoothly navigated from one stage to the next. A more rigorous approach to the process of  
15 operationalising CMOcs to Delphi survey to final intervention could help to mitigate some of these  
16 pitfalls.

17 The final intervention agreed is a home-based, tailored therapy intervention delivered by an MDT  
18 that includes physiotherapists, occupational therapists, therapy assistants, and a geriatrician (see  
19 additional file 6). Up to two assessment sessions and 22 intervention sessions will be available. The  
20 resources developed include an intervention manual for staff; a holistic assessment document to  
21 help staff to tailor the intervention; and a staff training programme [23]. This is in concordance with  
22 guidelines that recommend multifactorial interventions for falls in older people [87]. Though some  
23 evidence suggests that such interventions are not effective in people with dementia [9-11], it is  
24 hoped that the individually tailored, embedded approach will help to mitigate some of the factors  
25 affecting intervention success among this patient group [49].

1 The intervention that has been developed is novel in that it is tailored to the needs of people with  
2 dementia and addresses both rehabilitation and the prevention of future falls in people with  
3 dementia. While we are aware of a current study examining enhanced recovery of confused  
4 patients following hip fracture [6], this focuses on a single type of injury. Other current studies are  
5 focusing on falls interventions for people with dementia, but are not targeted at those who have  
6 already had an injurious fall [88, 89]. The DIFRID intervention therefore targets a neglected group,  
7 and could potentially clarify whether the preventive component is effective in patients who have  
8 already fallen.

### 9 [Strengths and limitations](#)

10 A strength of this project lies in the theoretically and empirically-informed intervention development  
11 process. While a response rate of 58% was achieved for the consensus surveys, not all panel  
12 members attended the consensus meetings. Furthermore, the panel did not include patient or lay  
13 representatives. The Delphi approach seemed less accessible for social care professionals, as  
14 evidenced by difficulty recruiting panel members and engaging them in the surveys. These factors  
15 may have implications for the results. However, the iterative nature of our approach to identifying  
16 causal factors and change mechanisms and stakeholder feedback process means that the opinion of  
17 these stakeholders has been considered in other aspects of the development process. While the  
18 effectiveness review highlighted the scarcity of evidence and underpinned the need to develop a  
19 new intervention, it was of limited value in the process of intervention development. In contrast, the  
20 broader, pragmatic realist approach helped to consider underlying mechanisms, and inform  
21 intervention content and delivery.

### 22 [Conclusions](#)

23 A new intervention has been developed to help people with dementia following a fall requiring  
24 healthcare attention. We are currently assessing the feasibility and acceptability of the DIFRID

1 intervention from the perspectives of all stakeholders. If appropriate, the findings will be used to  
2 refine the intervention, and then explore whether it merits rigorous evaluation [19].

### 3 Abbreviations

4	BP	Blood pressure
5	CGA	Comprehensive Geriatric Assessment
6	CMOc	Context-Mechanism-Outcome configuration
7	CPN	Community Psychiatric Nurse
8	CNPI	Checklist of Nonverbal Pain Indicators
9	GAS	Goal Attainment Scaling
10	MDT	Multidisciplinary Team
11	OT	Occupational Therapy/Therapist
12	PEDro	Physiotherapy Evidence Database
13	TOC	Trial Oversight Committee
14	TUG	Timed Up-and-Go test

### 15 Declarations

#### 16 Ethics approval and consent to participate

17 Ethical review for the initial interviews and focus groups with professionals was provided by  
18 Newcastle University Research Ethics Committee (ref BH138009) and any necessary permissions  
19 obtained from research and development departments of participating Trusts. Further approvals for  
20 stakeholder interviews were given by Newcastle and North Tyneside 1 Ethics Committee (reference  
21 15/NE/0397); Newcastle and North Tyneside 2 Ethics Committee (reference 16/NE/0011); and the  
22 Health Research Authority. Additional approvals were received from participating Trusts and Social  
23 Services Departments as required. For non-statutory agencies, approval was sought from senior  
24 managers. All participants gave written or verbal consent; verbal consent was sought for telephone  
25 interviews with professionals and for consensus panel discussions.

## 1 Consent for publication

2 Not applicable

## 3 Availability of data and material

4 The datasets generated during and/or analysed during the current study are available from the  
5 corresponding author (LA) on reasonable request. The data are not publicly available due to them  
6 containing information that could compromise research participant confidentiality.

## 7 Competing interests

8 The authors declare that they have no competing interests.

## 9 Funding

10 This paper presents independent research funded by the National Institute for Health Research  
11 (NIHR) under the Health Technology Appraisal (Grant Reference Number 13/78/02). The views and  
12 opinions expressed by authors in this publication are those of the authors and do not necessarily  
13 reflect those of the NHS, the NIHR, NETSCC, the HTA programme or the Department of Health. The  
14 verbatim quotations included in this publication express the views and opinions of study participants  
15 and do not necessarily reflect those of the authors, those of the NHS, the NIHR, NETSCC, the HTA  
16 programme or the Department of Health.

## 17 Authors' contributions

18 AW drafted the manuscript with contributions from CB. AW, CB, and CS collected and analysed  
19 qualitative data. FB performed literature searches. LA, CB, RB, FB, CF, SP, CS and AW extracted data  
20 and contributed to CMOc development. LA analysed the consensus survey data. LA, CB, EF, AS and  
21 AW developed intervention resources. LA conceived the design of the study. DH, TH, LR and all  
22 authors made substantial contributions to the design of the study, revised the manuscript critically  
23 for important intellectual content, and approved the final manuscript.

## 1 Acknowledgements

2 The DIFRID team would like to acknowledge the contribution of the members of the consensus  
 3 panel: Dr Fiona Shaw, Consultant Geriatrician, Newcastle upon Tyne Hospitals NHS Foundation  
 4 Trust; Dr Frank Sutherland, Consultant, Emergency Medicine, Norfolk and Norwich University  
 5 Hospital; Dr John-Paul Taylor, Senior Clinical Lecturer and Honorary Consultant in Old Age Psychiatry,  
 6 Institute of Neuroscience, Newcastle University and Northumberland, Tyne and Wear NHS Trust; Dr  
 7 Simon Thacker, Lead Consultant Psychiatrist, Royal Derby Hospital; Dr Kate Walters, Director, Centre  
 8 for Ageing & Population Studies and Reader in Primary Care and Epidemiology Dept. Primary Care &  
 9 Population Health, University College London; Danielle Woods, Lead Nurse for Dementia, Bradford  
 10 Teaching Hospitals NHS Foundation Trust; Louise McCarthy, Lead Research Nurse Older Peoples'  
 11 Services, Norfolk and Suffolk Foundation Trust; Professor Pam Dawson, Associate Dean (Strategic  
 12 Workforce Planning & Development), Northumbria University; Dr Pat Chung, Senior Lecturer,  
 13 Occupational Therapy Pathway, Canterbury Christ Church University; Michelle Brumwell, Health and  
 14 Social care coordinator, Newcastle City Council; Zabeen Aslam, Adult Social Services Directorate,  
 15 Newcastle City Council.

16 We are also grateful to all of the study participants, to Dr Miriam Boyles for additional data  
 17 collection, to Shannon Robalino for conducting searches, to Dr Sue Lord for additional data  
 18 extraction, and to Mrs Beth Edgar for administrative support.

## 19 References

- 20 1. Matthews FE, Arthur A, Barnes LE, Bond J, Jagger C, Robinson L, Brayne C, Medical  
 21 Research Council Cognitive F, Ageing C: **A two-decade comparison of prevalence of**  
 22 **dementia in individuals aged 65 years and older from three geographical areas of**  
 23 **England: results of the Cognitive Function and Ageing Study I and II.** *Lancet* 2013,  
 24 **382:1405-1412.**
- 25 2. Allan LM, Ballard CG, Rowan EN, Kenny RA: **Incidence and prediction of falls in**  
 26 **dementia: a prospective study in older people.** *PLoS ONE [Electronic Resource]*  
 27 2009, **4:e5521.**
- 28 3. Shaw FE: **Falls in cognitive impairment and dementia.** *Clinics in Geriatric Medicine*  
 29 2002, **18:159-173.**

- 1 4. Delbaere K, Close J, Brodaty H, Sachdev P, Lord S: **Determinants of disparities**  
2 **between perceived and physiological risk of falling among elderly people: cohort**  
3 **study.** *BMJ* 2010, **341**:c4165.
- 4 5. Robalino S, Nyakang'o SB, Beyer FR, Fox C, Allan LM: **Effectiveness of interventions**  
5 **aimed at improving physical and psychological outcomes of fall-related injuries in**  
6 **people with dementia: a narrative systematic review.** *Systematic Reviews* 2018,  
7 **7**:31.
- 8 6. Hammond SP, Cross JL, Shepstone L, Backhouse T, Henderson C, Poland F, Sims E,  
9 MacLulich A, Penhale B, Howard R, et al: **PERFECTED enhanced recovery (PERFECT-**  
10 **ER) care versus standard acute care for patients admitted to acute settings with hip**  
11 **fracture identified as experiencing confusion: study protocol for a feasibility cluster**  
12 **randomized controlled trial.** *Trials* 2017, **18**:583.
- 13 7. National Institute for Health and Care Excellence: **Falls: Assessment and Prevention**  
14 **of Falls in Older People. NICE Clinical Guideline 161.** In *NICE Clinical Guideline*, vol.  
15 161. Manchester: NICE; 2013.
- 16 8. Gillespie LD, Robertson MC, Gillespie WJ, Sherrington C, Gates S, Clemson LM, Lamb  
17 SE: **Interventions for preventing falls in older people living in the community.**  
18 *Cochrane Database Syst Rev* 2012, **9**:Cd007146.
- 19 9. Hedman AM, Grafstrom M: **Conditions for rehabilitation of older patients with**  
20 **dementia and hip fracture--the perspective of their next of kin.** *Scandinavian*  
21 *Journal of Caring Sciences* 2001, **15**:151-158.
- 22 10. Jurgens FJ, Clissett P, Gladman JR, Harwood RH: **Why are family carers of people**  
23 **with dementia dissatisfied with general hospital care? A qualitative study.** *BMC*  
24 *Geriatrics* 2012, **12**:57.
- 25 11. Vaapio SS, Salminen MJ, Ojanlatva A, Kivela SL: **Quality of life as an outcome of fall**  
26 **prevention interventions among the aged: a systematic review.** *European Journal of*  
27 *Public Health* 2009, **19**:7-15.
- 28 12. Gardner MM, Buchner DM, Robertson MC, Campbell AJ: **Practical implementation**  
29 **of an exercise-based falls prevention programme.** *Age and Ageing* 2001, **30**:77-83.
- 30 13. El-Khoury F, Cassou B, Charles MA, Dargent-Molina P: **The effect of fall prevention**  
31 **exercise programmes on fall induced injuries in community dwelling older adults:**  
32 **systematic review and meta-analysis of randomised controlled trials.** *BMJ* 2013,  
33 **347**:f6234.
- 34 14. Martins AC, Santos C, Silva C, Baltazar D, Moreira J, Tavares N: **Does modified Otago**  
35 **Exercise Program improves balance in older people? A systematic review.**  
36 *Preventive Medicine Reports* 2018, **11**:231-239.
- 37 15. Hauer K, Schwenk M, Zieschang T, Essig M, Becker C, Oster P: **Physical training**  
38 **improves motor performance in people with dementia: a randomized controlled**  
39 **trial.** *J Am Geriatr Soc* 2012, **60**:8-15.
- 40 16. Zieschang T, Schwenk M, Becker C, Uhlmann L, Oster P, Hauer K: **Falls and Physical**  
41 **Activity in Persons With Mild to Moderate Dementia Participating in an Intensive**  
42 **Motor Training: Randomized Controlled Trial.** *Alzheimer Disease & Associated*  
43 *Disorders* 2017, **31**:307-314.
- 44 17. **Dementia - assessment, management and support for people living with dementia**  
45 **and their carers** [<https://www.nice.org.uk/guidance/ng97>]
- 46 18. Hodinott P: **A new era for intervention development studies.** *Pilot and Feasibility*  
47 *Studies* 2015, **1**:36.

- 1 19. Wight D, Wimbush E, Jepson R, Doi L: **Six steps in quality intervention development (6SQuID)**. *J Epidemiol Community Health* 2016, **70**:520-525.
- 2
- 3 20. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M, Medical Research  
4 Council Guidance: **Developing and evaluating complex interventions: the new  
5 Medical Research Council guidance**. *BMJ* 2008, **337**:a1655.
- 6 21. Wheatley A, Bamford C, Shaw C, Boyles M, Fox C, Allan L: **Service organization for  
7 people with dementia after an injurious fall: challenges and opportunities**. *Age and  
8 Ageing* In press.
- 9 22. Bamford C, Wheatley A, Shaw C, Allan LM: **Equipping staff with the skills to  
10 maximise recovery of people with dementia after an injurious fall**. *Aging and  
11 Mental Health* 2018:1-9.
- 12 23. Allan LM, Wheatley A, Flynn E, Smith A, Fox C, Howel D, Barber R, Homer TM,  
13 Robinson L, Parry SW, et al: **Is it feasible to deliver a complex intervention to  
14 improve the outcome of falls in people with dementia? A protocol for the DIFRID  
15 feasibility study**. *Pilot and Feasibility Studies* 2018, **4**:170.
- 16 24. Wong G, Westhorp G, Manzano A, Greenhalgh J, Jagosh J, Greenhalgh T: **RAMESES II  
17 reporting standards for realist evaluations**. *BMC Medicine* 2016, **14**:96.
- 18 25. Pawson R, Greenhalgh T, Harvey G, Walshe K: **Realist review - a new method of  
19 systematic review designed for complex policy interventions**. *J Health Serv Res  
20 Policy* 2005, **10**:S1:21.
- 21 26. Pawson R, Tilley N: *Realistic evaluation*. London: London : Sage; 1997.
- 22 27. Rycroft-Malone J, McCormack B, Hutchinson AM, DeCorby K, Bucknall TK, Kent B,  
23 Schultz A, Snelgrove-Clarke E, Stetler CB, Titler M, et al: **Realist synthesis: illustrating  
24 the method for implementation research**. *Implementation Science* 2012, **7**:1-10.
- 25 28. Dalkin SM, Greenhalgh J, Jones D, Cunningham B, Lhussier M: **What's in a  
26 mechanism? Development of a key concept in realist evaluation**. *Implementation  
27 Science* 2015, **10**:49.
- 28 29. Pluye P, Robert E, Cargo M, Bartlett G, O'Cathain A, Griffiths F, Boardman F, Gagnon  
29 MP, Rousseau MC: **Proposal: A mixed methods appraisal tool for systematic mixed  
30 studies reviews**. 2011.
- 31 30. Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, Altman DG,  
32 Barbour V, Macdonald H, Johnston M, et al: **Better reporting of interventions:  
33 template for intervention description and replication (TIDieR) checklist and guide**.  
34 *BMJ : British Medical Journal* 2014, **348**.
- 35 31. Dalkey N: **The Delphi Method: An Experimental Study of Group Opinion**. Santa  
36 Monica, CA: The Rand Corporation; 1969.
- 37 32. Dalkey N, Helmer O: **An Experimental Application of the DELPHI Method to the Use  
38 of Experts**. *Management Science* 1963, **9**:458-467.
- 39 33. McMillan SS, King M, Tully MP: **How to use the nominal group and Delphi  
40 techniques**. *Int J Clin Pharm* 2016, **38**:655-662.
- 41 34. Kellogg Foundation: **Logic Model Development Guide** Battle Creek, Michigan: W.K.  
42 Kellogg Foundation; 2004.
- 43 35. Dwyer JJ, Makin S: **Using a program logic model that focuses on performance  
44 measurement to develop a program**. *Can J Public Health* 1997, **88**:421-425.
- 45 36. Schepker CA, Leveille SG, Pedersen MM, Ward RE, Kurlinski LA, Grande L, Kiely DK,  
46 Bean JF: **Effect of Pain and Mild Cognitive Impairment on Mobility**. *J Am Geriatr Soc*  
47 2016, **64**:138-143.

- 1 37. Kolanowski A, Mogle J, Fick DM, Hill N, Mulhall P, Nadler J, Colancecco E, Behrens L:  
2 **Pain, delirium, and physical function in skilled nursing home patients with**  
3 **dementia.** *Journal Of The American Medical Directors Association* 2015, **16**:37-40.
- 4 38. Kress HG, Ahlbeck K, Aldington D, Alon E, Coaccioli S, Coluzzi F, Huygen F, Jaksch W,  
5 Kalso E, Kocot-Kepska M, et al: **Managing chronic pain in elderly patients requires a**  
6 **CHANGE of approach.** *Curr Med Res Opin* 2014, **30**:1153-1164.
- 7 39. Flo E, Gulla C, Husebo BS: **Effective pain management in patients with dementia:**  
8 **benefits beyond pain?** *Drugs & Aging* 2014, **31**:863-871.
- 9 40. Husebo BS, Ballard C, Sandvik R, Nilsen OB, Aarsland D: **Efficacy of treating pain to**  
10 **reduce behavioural disturbances in residents of nursing homes with dementia:**  
11 **cluster randomised clinical trial.** *BMJ (Clinical Research Ed)* 2011, **343**:d4065-d4065.
- 12 41. Ahn H, Horgas A: **The relationship between pain and disruptive behaviors in nursing**  
13 **home residents with dementia.** *BMC Geriatrics* 2013, **13**:14-14.
- 14 42. Ahn H, Horgas A: **Does pain mediate or moderate the effect of cognitive**  
15 **impairment on aggression in nursing home residents with dementia?** *Asian Nursing*  
16 *Research* 2014, **8**:105-109.
- 17 43. Husebo BS, Ballard C, Fritze F, Sandvik RK, Aarsland D: **Efficacy of pain treatment on**  
18 **mood syndrome in patients with dementia: a randomized clinical trial.**  
19 *International Journal Of Geriatric Psychiatry* 2014, **29**:828-836.
- 20 44. Herr K, Bjoro K, Decker S: **Tools for assessment of pain in nonverbal older adults**  
21 **with dementia: a state-of-the-science review.** *J Pain Symptom Manage* 2006,  
22 **31**:170-192.
- 23 45. Valeriani L: **Management of demented patients in emergency Department.**  
24 *International Journal of Alzheimer's Disease* 2011, (no pagination).
- 25 46. McIntyre A, Reynolds F: **There's no apprenticeship for Alzheimer's: the caring**  
26 **relationship when an older person experiencing dementia falls.** *Ageing & Society*  
27 2012, **32**:873-896.
- 28 47. Giusti A, Barone A, Pioli G: **Rehabilitation after hip fracture in patients with**  
29 **dementia.** *Journal of the American Geriatrics Society* 2007, **55**:1309-1310.
- 30 48. Wesson J, Clemson L, Brodaty H, Lord S, Taylor M, Gitlin L, Close J: **A feasibility study**  
31 **and pilot randomised trial of a tailored prevention program to reduce falls in older**  
32 **people with mild dementia.** *BMC Geriatrics* 2013, **13**:89.
- 33 49. Taylor ME, Lord SR, Brodaty H, Kurrle SE, Hamilton S, Ramsay E, Webster L, Payne  
34 NL, Close JC: **A home-based, carer-enhanced exercise program improves balance**  
35 **and falls efficacy in community-dwelling older people with dementia.** *Int*  
36 *Psychogeriatr* 2017, **29**:81-91.
- 37 50. Seitz DP, Gill SS, Austin PC, Bell CM, Anderson GM, Gruneir A, Rochon PA:  
38 **Rehabilitation of Older Adults with Dementia after Hip Fracture.** *Journal of the*  
39 *American Geriatrics Society* 2016, **64**:47-54.
- 40 51. Shaw FE, Bond J, Richardson DA, Dawson P, Steen IN, McKeith IG, Kenny RA:  
41 **Multifactorial intervention after a fall in older people with cognitive impairment**  
42 **and dementia presenting to the accident and emergency department: randomised**  
43 **controlled trial.** *BMJ (Clinical research ed)* 2003, **326**:73.
- 44 52. Tarazona-Santabalbina FJ, Domenech-Pascual JR, Belenguier-Varea AA, Rovira Daudi  
45 E: **The approach to patients with cognitive impairment and hip fracture: The role of**  
46 **orthogeriatric care.** *Reviews in Clinical Gerontology* 2014, **24**:219-227.



- 1 53. Aharony L, Sela-Katz P: **Depression, falls and cognitive changes among community-**  
2 **dwelling elderly people.** *Alzheimer's and Dementia* 2011, **1**):S622.
- 3 54. Goldstein FC, Strasser DC, Woodard JL, Roberts VJ: **Functional outcome of**  
4 **cognitively impaired hip fracture patients on a geriatric rehabilitation unit.** *Journal*  
5 *of the American Geriatrics Society* 1997, **45**:35-42.
- 6 55. Marcantonio ER, Flacker JM, Wright RJ, Resnick NM: **Reducing delirium after hip**  
7 **fracture: a randomized trial.** *Journal of the American Geriatrics Society* 2001,  
8 **49**:516-522.
- 9 56. Faes MC, Reelick MF, Banningh L, de Gier M, Esselink RA, Rikkert MGO: **Qualitative**  
10 **study on the impact of falling in frail older persons and family caregivers:**  
11 **Foundations for an intervention to prevent falls.** *Aging & Mental Health* 2010,  
12 **14**:834-842.
- 13 57. Nilsson I, Rogmark C: **Hemiarthroplasty for displaced femoral neck fracture: good**  
14 **clinical outcome but uneven distribution of occupational therapy.** *Disability &*  
15 *Rehabilitation* 2011, **33**:2329-2332.
- 16 58. Deschodt M, Braes T, Broos P, Sermon A, Boonen S, Flamaing J, Milisen K: **Effect of**  
17 **an Inpatient Geriatric Consultation Team on Functional Outcome, Mortality,**  
18 **Institutionalization, and Readmission Rate in Older Adults with Hip Fracture: A**  
19 **Controlled Trial.** *Journal of the American Geriatrics Society* 2011, **59**:1299-1308  
20 1210p.
- 21 59. Gonski PN, Moon I: **Outcomes of a behavioral unit in an acute aged care service.**  
22 *Archives of Gerontology & Geriatrics* 2012, **55**:60-65.
- 23 60. Luxford K, Axam A, Hasnip F, Dobrohotoff J, Strudwick M, Reeve R, Hou C, Viney R:  
24 **Improving clinician-carer communication for safer hospital care: A study of the**  
25 **'TOP 5' strategy in patients with dementia.** *International Journal for Quality in*  
26 *Health Care* 2015, **27**:175-182.
- 27 61. Raivio M, Korkala O, Pitkala K, Tilvis R: **Rehabilitation outcome in hip-fracture:**  
28 **Impact of weight-bearing restriction - A preliminary investigation.** *Physical and*  
29 *Occupational Therapy in Geriatrics* 2004, **22**:1-9.
- 30 62. Toulotte C, Fabre C, Dangremont B, Lensel G, Thevenon A: **Effects of physical**  
31 **training on the physical capacity of frail, demented patients with a history of**  
32 **falling: a randomised controlled trial.** *Age and Ageing* 2003, **32**:67-73.
- 33 63. Judah G, Gardner B, Aunger R: **Forming a flossing habit: An exploratory study of the**  
34 **psychological determinants of habit formation.** *British Journal of Health Psychology*  
35 2012, **18**:338-353.
- 36 64. Fleig L, McAllister MM, Chen P, Iverson J, Milne K, McKay HA, Clemson L, Ashe MC:  
37 **Health behaviour change theory meets falls prevention: Feasibility of a habit-based**  
38 **balance and strength exercise intervention for older adults.** *Psychology of Sport and*  
39 *Exercise* 2016, **22**:114-122.
- 40 65. Lally P, van Jaarsveld Cornelia HM, Potts Henry WW, Wardle J: **How are habits**  
41 **formed: Modelling habit formation in the real world.** *European Journal of Social*  
42 *Psychology* 2009, **40**:998-1009.
- 43 66. de Rotrou J, Cantegreil I, Faucounau V, Wenisch E, Chausson C, Jegou D, Grabar S,  
44 Rigaud AS: **Do patients diagnosed with Alzheimer's disease benefit from a psycho-**  
45 **educational programme for family caregivers? A randomised controlled study.**  
46 *International Journal of Geriatric Psychiatry* 2011, **26**:833-842.

- 1 67. Faes MC, Reelick MF, Joosten-Weyn Banningh LW, Gier M, Esselink RA, Olde Rikkert  
2 MG: **Qualitative study on the impact of falling in frail older persons and family**  
3 **caregivers: Foundations for an intervention to prevent falls.** *Aging & Mental Health*  
4 2010, **14**:834-842.
- 5 68. Ritter MA, Harty LD: **Total joint replacement in patients with dementia syndromes:**  
6 **a report of thirteen cases.** *Orthopedics* 2004, **27**:516-517.
- 7 69. Isbel ST, Jamieson MI: **Views from health professionals on accessing rehabilitation**  
8 **for people with dementia following a hip fracture.** *Dementia (London)* 2016.
- 9 70. Murphy MR, Escamilla MI, Blackwell PH, Lucke KT, Miner-Williams D, Shaw V, Lewis  
10 SL: **Assessment of caregivers' willingness to participate in an intervention research**  
11 **study.** *Research in Nursing & Health* 2007, **30**:347-355.
- 12 71. Suttanon P, Hill KD, Said CM, Byrne KN, Dodd KJ, Suttanon P, Hill KD, Said CM, Byrne  
13 KN, Dodd KJ: **Factors influencing commencement and adherence to a home-based**  
14 **balance exercise program for reducing risk of falls: perceptions of people with**  
15 **Alzheimer's disease and their caregivers.** *International Psychogeriatrics* 2012,  
16 **24**:1172-1182.
- 17 72. Dow B, Moore K, Russel M, Ames D, Cyarto E, Haines T, Hill K, Lautenschlager N,  
18 Mackenzie L, Williams S, Loi S: **Improving mood through physical activity for carers**  
19 **and care recipients (IMPACCT): protocol for a randomised trial.** *Journal Of*  
20 *Physiotherapy* 2013, **59**:125; discussion 125.
- 21 73. Comans TA, Currin ML, Brauer SG, Haines TP: **Factors associated with quality of life**  
22 **and caregiver strain amongst frail older adults referred to a community**  
23 **rehabilitation service: implications for service delivery.** *Disability And Rehabilitation*  
24 2011, **33**:1215-1221.
- 25 74. Cristancho-Lacroix V, Wrobel J, Cantegreil-Kallen I, Dub T, Rouquette A, Rigaud A-S: **A**  
26 **web-based psychoeducational program for informal caregivers of patients with**  
27 **Alzheimer's disease: a pilot randomized controlled trial.** *Journal Of Medical Internet*  
28 *Research* 2015, **17**:e117-e117.
- 29 75. Zarit SH, Lee JE, Barrineau MJ, Whitlatch CJ, Femia EE: **Fidelity and acceptability of**  
30 **an adaptive intervention for caregivers: An exploratory study.** *Aging & Mental*  
31 *Health* 2013, **17**:197-206.
- 32 76. Shim B, Barroso J, Davis LL: **A comparative qualitative analysis of stories of spousal**  
33 **caregivers of people with dementia: Negative, ambivalent, and positive**  
34 **experiences.** *International Journal of Nursing Studies* 2012, **49**:220-229.
- 35 77. Lach HW, Chang Y: **Caregiver perspectives on safety in home dementia care.**  
36 *Western Journal of Nursing Research* 2007, **29**:993-1014.
- 37 78. Rosler A, von Renteln-Kruse W, Muhlhan C, Frilling B: **Treatment of dementia**  
38 **patients with fracture of the proximal femur in a specialized geriatric care unit**  
39 **compared to conventional geriatric care.** *Zeitschrift für Gerontologie und Geriatrie*  
40 2012, **45**:400-403.
- 41 79. Deschodt M, Braes T, Broos P, Sermon A, Boonen S, Flamaing J, Milisen K: **Effect of**  
42 **an inpatient geriatric consultation team on functional outcome, mortality,**  
43 **institutionalization, and readmission rate in older adults with hip fracture: a**  
44 **controlled trial.** *Journal of the American Geriatrics Society* 2011, **59**:1299-1308.
- 45 80. Reuben DB, Ganz DA, Roth CP, McCreath HE, Ramirez KD, Wenger NS: **Effect of nurse**  
46 **practitioner comanagement on the care of geriatric conditions.** *Journal of the*  
47 *American Geriatrics Society* 2013, **61**:857-867.

- 1 81. Ganz DA, Koretz BK, Bail JK, McCreath HE, Wenger NS, Roth CP, Reuben DB: **Nurse**  
2 **practitioner comanagement for patients in an academic geriatric practice.**  
3 *American Journal of Managed Care* 2010, **16**:e343-355.
- 4 82. Lichtenstein BJ, Reuben DB, Karlamangla AS, Han W, Roth CP, Wenger NS: **Effect of**  
5 **Physician Delegation to Other Healthcare Providers on the Quality of Care for**  
6 **Geriatric Conditions.** *Journal of the American Geriatrics Society* 2015, **63**:2164-2170.
- 7 83. Schrijvers G, van Hoorn A, Huiskes N: **The care pathway: concepts and theories: an**  
8 **introduction.** *International Journal of Integrated Care* 2012, **12**:1-7.
- 9 84. Deneckere S, Euwema M, Van Herck P, Lodewijckx C, Panella M, Sermeus W,  
10 Vanhaecht K: **Care pathways lead to better teamwork: results of a systematic**  
11 **review.** *Soc Sci Med* 2012, **75**:264-268.
- 12 85. Vanhaecht K, Panella M, van Zelm R, Sermeus W: *What about care pathways?*; 2011.
- 13 86. Sherrington C, Fairhall NJ, Wallbank GK, Tiedemann A, Michaleff ZA, Howard K,  
14 Clemson L, Hopewell S, Lamb SE: **Exercise for preventing falls in older people living**  
15 **in the community.** *Cochrane Database of Systematic Reviews* 2019.
- 16 87. National Institute for Health and Care Excellence: **Clinical practice guideline for the**  
17 **assessment and prevention of falls in older people.** London; 2004.
- 18 88. Close JCT, Wesson J, Sherrington C, Hill KD, Kurrle S, Lord SR, Brodaty H, Howard K,  
19 Gitlin LN, O'Rourke SD, Clemson L: **Can a tailored exercise and home hazard**  
20 **reduction program reduce the rate of falls in community dwelling older people**  
21 **with cognitive impairment: Protocol paper for the i-FOCIS randomised controlled**  
22 **trial.** *BMC Geriatrics* 2014, **14**.
- 23 89. Harwood RH, van der Wardt V, Goldberg SE, Kearney F, Logan P, Hood-Moore V,  
24 Booth V, Hancox JE, Masud T, Hoare Z, et al: **A development study and randomised**  
25 **feasibility trial of a tailored intervention to improve activity and reduce falls in**  
26 **older adults with mild cognitive impairment and mild dementia.** *Pilot and*  
27 *Feasibility Studies* 2018, **4**:49.
- 28 90. Feldt KS: **The checklist of nonverbal pain indicators (CNPI).** *Pain Management*  
29 *Nursing* 2000, **1**:13-21.
- 30 91. Podsiadlo D, Richardson S: **The Timed "Up & Go": A Test of Basic Functional**  
31 **Mobility for Frail Elderly Persons.** *Journal of the American Geriatrics Society* 1991,  
32 **39**:142-148.
- 33 92. Kiresuk TJ, Sherman RE: **Goal Attainment Scaling: A General Method for Evaluating**  
34 **Comprehensive Community Mental Health Programs.** *Community Mental Health*  
35 *Journal* 1968, **4**:443-453.
- 36 93. Kampe K, Kohler M, Albrecht D, Becker C, Hautzinger M, Lindemann U, Pfeiffer K: **Hip**  
37 **and pelvic fracture patients with fear of falling: development and description of**  
38 **the "Step by Step" treatment protocol.** *Clin Rehabil* 2017, **31**:571-581.
- 39 94. Alzheimer's Association: **Take Care of Yourself: How to recognize and manage**  
40 **caregiver stress.** 2017.
- 41 95. Alzheimer's Society: **Factsheet 523LP: Carers - looking after yourself.** 2016.
- 42

## 1 Figures

2 Figure 1

3 **Legend:** Intervention development

4

5 Figure 2

6 **Legend:** Diagram of the search, screening, selection and extraction process

7

8 Figure 3

9 **Legend:** Logic model

## 10 Additional files

11 **Filename:** Additional file 1.docx

12 **Title of data:** Professional participants

13 **Description of data:** Table showing numbers of participants from various health and social care  
14 professions in each stage of the project.

15

16 **Filename:** Additional file 2.docx

17 **Title of data:** Comprehensive search strategy

18 **Description of data:** Medline literature search strategy for the phase 1 comprehensive search.

19

20 **Filename:** Additional file 3.docx

21 **Title of data:** Targeted search strategy

22 **Description of data:** Example Medline literature search strategy for the phase 2 targeted searches.

23

24 **Filename:** Additional file 4.docx

25 **Title of data:** Delphi survey

26 **Description of data:** Expanded description of rationale and methodology for the Delphi survey.

27

28 **Filename:** Additional file 5.docx

29 **Title of data:** Consensus statements

1 **Description of data:** Full list of statements provided to the consensus panel along with their  
2 outcomes.

3

4 **Filename:** Additional file 6.docx

5 **Title of data:** Final DIFRID intervention

6 **Description of data:** A description of the final intervention using the TIDieR framework.

7

## 1 Tables

<b>Table 1: Optimising the circumstances of rehabilitation for people with dementia: CMOcs, consensus statements and outcomes</b>				
	<b>CMOc</b>	<b>Consensus statements</b>	<b>Outcome</b>	<b>Operationalisation</b>
<b>CMOc1</b>	<p><b>Context:</b> cognitive impairment may limit the ability of people with dementia to articulate pain</p> <p><b>Mechanism (resource):</b> staff use non-verbal pain signifiers and/or give blanket pain relief</p> <p><b>Mechanism (reasoning):</b> people with dementia are not in pain</p> <p><b>Outcome:</b> capacity to engage with an intervention increases</p>	Tools which assess non-verbal signs of pain should be used	Agreed in round 1 (93%)	<ul style="list-style-type: none"> <li>• Checklist of Nonverbal Pain Indicators (CNPI) [44, 90] included in assessment document</li> <li>• Pain management included in staff training</li> </ul>
<b>CMOc2</b>	<p><b>Context:</b> cognitive impairment may limit the ability of people with dementia to adapt to and cope with new environments</p> <p><b>Mechanism (resource):</b> intervention assessment and delivery takes place in appropriate, accessible and familiar environments</p> <p><b>Mechanism (reasoning):</b> people with dementia feel comfortable and less distracted</p> <p><b>Outcome:</b> anxiety and challenging behaviours are reduced</p>	The intervention should primarily take place in the patient's home	Agreed in round 1 (86%)	Intervention delivered mainly in patient's home
<b>CMOc3</b>	<p><b>Context:</b> the role of comorbidities may be underestimated in dementia</p> <p><b>Mechanism (resource):</b> holistic biopsychosocial assessment is employed</p>	<p>A continence assessment is required</p> <p>An assessment of comorbidities is required</p> <p>An osteoporosis risk assessment is required</p> <p>A vision assessment is required</p>	Agreed in round 1 (79% – 100%)	All included in assessment document (see table 2 below)

	<p><b>Mechanism (reasoning):</b> staff understand the range of factors contributing to falls and are able to treat comorbidities more effectively</p> <p><b>Outcome:</b> falls risk may be reduced and recovery enhanced in patients with dementia</p>	A medication review is required		
		An assessment of challenging behaviour is required		
		Formal assessments of gait and balance should be carried out by the Timed Up and Go (TUG) test [91]	No consensus after 2 rounds (54% & 62%)	
		All patients require attendance for a lying and standing blood pressure (BP)		

1

2

3



<b>Table 3: Compensating for the reduced ability of people with dementia to self-manage: CMOcs, consensus statements and outcomes</b>				
	<b>CMOc</b>	<b>Consensus statements</b>	<b>Outcome</b>	<b>Operationalisation</b>
<b>CMOc 4</b>	<b>Context:</b> cognitive impairment may limit the ability of people with dementia to comply with instructions and form habits <b>Mechanism (resource):</b> staff tailor the intervention (e.g. exercises) to the circumstances of people with dementia and embed it in their existing routines <b>Mechanism (reasoning):</b> intervention becomes routine and habitual <b>Outcome:</b> more successful rehabilitation can be achieved	Interventions should be based on goals set by the patient and carer	Agreed in round 1 (86% - 100%)	<ul style="list-style-type: none"> <li>• Goal Attainment Scaling [92] (GAS) implemented</li> <li>• Compass of Life [93] included to assist in goal identification</li> </ul>
		Therapists should work with service users to minimise the risk of falling, as this may improve confidence and enable realistic risk taking.		Falls risk assessment included
		Therapists should help the service user and caregiver to develop a meaningful programme of activities		<ul style="list-style-type: none"> <li>• Assessment records personal preferences, routines, and priorities</li> <li>• Therapists develop programme of meaningful activities based on information gathered</li> </ul>
		Therapists should undertake observed activities with the service user to facilitate new learning		Included in assessment
		Exercise interventions should be informed by evidence based formats such as the Otago programme but tailored to the circumstances of people with dementia and embedded in their daily life	Agreed in round 2 (69%)	<ul style="list-style-type: none"> <li>• During training, staff are encouraged to use evidence-based formats creatively</li> <li>• Training also includes advice on creating programmes and embedding them into routines</li> <li>• Coloured paper provided for embedding strategies</li> </ul>
<b>CMOc 5</b>	<b>Context:</b> cognitive impairment may	The total number of physiotherapy sessions available in the first 3 months (including	No consensus	Implemented 2 assessment sessions and maximum 22 therapy

	<p>limit the ability of people with dementia to self-manage changes in circumstances</p> <p><b>Mechanism (resource):</b> ongoing follow-up is provided</p> <p><b>Mechanism (reasoning):</b> staff are able to reinforce previous interventions and adapt them to meet changing needs</p> <p><b>Outcome:</b> improvements in mobility are sustained and new falls risks reduced</p>	<p>sessions delivered by a support worker) should be 16, 20 or 24</p> <p>The total number of occupational therapy (OT) sessions available in the first 3 months should be 3-4</p>	<p>after 2 rounds (31% - 62%)</p>	<p>sessions delivered by a mix of OT, physiotherapist and support worker</p>
<b>CMOc 6</b>	<p><b>Context:</b> the burden on informal carers is high when caring for relatives or friends with dementia who are at risk of falling</p> <p><b>Mechanism (resource):</b> carer support and education is provided</p> <p><b>Mechanism (reasoning):</b> carer stress is reduced and skills increased</p> <p><b>Outcome:</b> carers' capacity to assist with the delivery of interventions increases</p>	<p>Carer stress should be routinely assessed</p>	<p>Agreed in round 1 (93% - 100%)</p>	<ul style="list-style-type: none"> <li>• Carer stress included in assessment</li> <li>• Training emphasises ensuring carers have capacity to be involved</li> </ul>
		<p>Therapists should facilitate caregivers, family and friends to adopt a positive approach to risk</p>		<ul style="list-style-type: none"> <li>• Training includes advice on carer education, including accepting 'positive risk'</li> </ul>
		<p>Intervention staff should be able to provide basic carer education &amp; support, referring to other agencies as needed</p>	<p>Agreed in round 2 (77%)</p>	<ul style="list-style-type: none"> <li>• Carer education leaflets provided for dissemination [94, 95]</li> </ul>

<b>Table 4: Equipping the workforce with the necessary skills and information to care for people with dementia: CMOcs, consensus statements and outcomes</b>				
	<b>CMOc</b>	<b>Consensus statements</b>	<b>Outcome</b>	<b>Operationalisation</b>
<b>CMOc7</b>	<p><b>Context:</b> cognitive impairment may limit the ability of people with dementia to pass on information</p> <p><b>Mechanism (resource):</b> staff use multiple sources of information including carers and direct observation</p> <p><b>Mechanism (reasoning):</b> staff gain a better understanding of the individual</p> <p><b>Outcome:</b> staff are able to provide appropriate, tailored care</p>	Assessment should involve multiple sources of information including information from carers	Agreed round 1 (93% – 100%)	The assessment (table 2) includes all of these components
		Assessment should include direct observation		
		A home hazard assessment should include a walk around the house to determine where actual falls have occurred and negotiate how these might be reduced		
<b>CMOc8</b>	<p><b>Context:</b> current staff knowledge of, and attitudes to, dementia are variable</p> <p><b>Mechanism (resource):</b> increased dementia training is provided</p> <p><b>Mechanism (reasoning):</b> staff gain skills in and understanding of rehabilitation for people with dementia</p> <p><b>Outcome:</b> staff ability and willingness to engage with people with dementia is enhanced</p>	Tier 2 training is required for intervention staff	Agreed round 2 (85%)	This was deemed unfeasible in the time available. A tailored training programme was developed, including items from tier 2 training.
		Training needs to include how to tailor an intervention for people with dementia.	Agreed round 1 (100%)	Training includes this
		Training needs to include advice on how to engage and motivate people with dementia.		Training includes this
		Training should include on the job role modelling		This was deemed unfeasible in the time available. Training delivered by therapists with experience in working with people with dementia, who

				were available remotely for advice.
<b>CMOc9</b>	<p><b>Context:</b> care pathways are often unclear</p> <p><b>Mechanism (resource):</b> a centralised, collaborative pathway is developed and disseminated</p> <p><b>Mechanism (reasoning):</b> staff are better equipped to refer to the most appropriate services</p> <p><b>Outcome:</b> service users receive better treatment</p>	The setting of the intervention should make use of existing pathways only when referral from the team deems it would be useful for the individual	Agreed round 1 (85.7% - 100%)	Assessment document includes tracking referrals that are decided by MDT
		A multidisciplinary team (MDT) meeting should be available if needed		<ul style="list-style-type: none"> <li>MDT composition agreed as physiotherapist, OT, support workers and geriatrician, with a general nurse available where the team already included this. Community psychiatric nurse (CPN), social workers, reablement workers, old age psychiatrists and podiatrists accessible by referral.</li> <li>MDT meetings available at beginning and middle of intervention period.</li> </ul>
		Therapists should offer service users information on assistive devices and facilitate delivery		This is flagged in the assessment document and available when needed