

Bond University
Research Repository



A systematic review and meta-analysis of the criterion validity of nutrition assessment tools for diagnosing protein-energy malnutrition in the older community setting (the MACRo Study)

Marshall, Skye; Craven, Dana L.; Kelly, Jaimon T; Isenring, Elisabeth

Published in:
Clinical Nutrition

DOI:
[10.1016/j.clnu.2017.09.022](https://doi.org/10.1016/j.clnu.2017.09.022)

Published: 01/12/2018

Document Version:
Peer reviewed version

[Link to publication in Bond University research repository.](#)

Recommended citation(APA):

Marshall, S., Craven, D. L., Kelly, J. T., & Isenring, E. (2018). A systematic review and meta-analysis of the criterion validity of nutrition assessment tools for diagnosing protein-energy malnutrition in the older community setting (the MACRo Study). *Clinical Nutrition*, 37(6), 1902-1912. <https://doi.org/10.1016/j.clnu.2017.09.022>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

A systematic review and meta-analysis of the criterion validity of nutrition assessment tools for diagnosing protein-energy malnutrition in the older community setting (the MACRo Study)

Skye Marshall^{a,b}, Dana Craven^c, Jaimon Kelly^d, Elizabeth Isenring^e

^a BNutr&Diet(Hons), PhD, Accredited Practising Dietitian, Faculty of Health Sciences and Medicine, Bond University, Robina, Queensland, 4226, Australia.

^b Corresponding author. Bond Institute of Health and Sport, Robina, Queensland, 4226, Australia. Telephone: +61 7 5595 5530, Fax: +61 7 5595 3524, skye_marshall@bond.edu.au.

^c BNutr&Diet, PhD Scholar, Accredited Practising Dietitian, School of Health and Sport Sciences, University of the Sunshine Coast, Sippy Downs, Queensland, 4556, Australia. dana.craven@research.usc.edu.au.

^d BHlthSc, MNutr&Diet(Res), PhD Scholar, Accredited Practising Dietitian, Faculty of Health Sciences and Medicine, Bond University, Robina, Queensland, 4226, Australia. jkelly@bond.edu.au.

^e Professor of Nutrition & Dietetics, PhD, Advanced Accredited Practising Dietitian, Faculty of Health Sciences and Medicine, Bond University, Robina, Queensland, 4226, Australia. lisenrin@bond.edu.au.

Abbreviations

CASP: Critical Appraisal Skills Programme

CKD: Chronic kidney disease

COPD: Chronic obstructive pulmonary disease

GRADE: Grading of Recommendations, Assessment, Development and Evaluation

MACRo: Malnutrition in the Aging Community Review

MNA: Mini Nutritional Assessment

SGA: Subjective Global Assessment

PEM: Protein-energy malnutrition

PG-SGA: Patient-Generated Subjective Global Assessment

PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analysis

PROSPERO: International Prospective Register of Systematic Reviews

Accepted: Clinical Nutrition 28/09/2017

1 **Abstract**

2 **Background & aims:** Malnutrition is a significant barrier to healthy and independent ageing
3 in older adults who live in their own homes, and accurate diagnosis is a key step in managing
4 the condition. However, there has not been sufficient systematic review or pooling of existing
5 data regarding malnutrition diagnosis in the geriatric community setting. The current paper was
6 conducted as part of the MACRo (Malnutrition in the Ageing Community Review) Study and
7 seeks to determine the criterion (concurrent and predictive) validity and reliability of nutrition
8 assessment tools in making a diagnosis of protein-energy malnutrition in the general older adult
9 community.

10 **Methods:** A systematic literature review was undertaken using six electronic databases in
11 September 2016. Studies in any language were included which measured malnutrition via a
12 nutrition assessment tool in adults ≥ 65 years living in their own homes. Data relating to the
13 predictive validity of tools were analysed via meta-analyses. GRADE was used to evaluate the
14 body of evidence.

15 **Results:** There were 6,412 records identified, of which 104 potentially eligible records were
16 screened via full text. Eight papers were included; two which evaluated the concurrent validity
17 of the Mini Nutritional Assessment (MNA) and Subjective Global Assessment (SGA) and six
18 which evaluated the predictive validity of the MNA. The quality of the body of evidence for
19 the concurrent validity of both the MNA and SGA was very low. The quality of the body of
20 evidence for the predictive validity of the MNA in detecting risk of death was moderate (RR:
21 1.92 [95%CI: 1.55-2.39]; $P < 0.00001$; $n = 2,013$ participants; $n = 4$ studies; $I^2: 0\%$). The quality
22 of the body of evidence for the predictive validity of the MNA in detecting risk of poor physical
23 function was very low (SMD: 1.02 [95%CI: 0.24-1.80]; $P = 0.01$; $n = 4,046$ participants; $n = 3$
24 studies; $I^2: 89\%$).

25

26 **Conclusions:** Due to the small number of studies identified and no evaluation of the predictive
27 validity of tools other than the MNA, there is insufficient evidence to recommend a particular
28 nutrition assessment tool for diagnosing PEM in older adults in the community. High quality
29 diagnostic accuracy studies are needed for all nutrition assessment tools used in older
30 community samples, including measuring of health outcomes subsequent to nutrition
31 assessment by the SGA and PG-SGA.

32

33 **Keywords:** Protein-energy malnutrition, nutritional status, nutrition assessment, community,
34 aged, systematic review

35

36

37 **Introduction**

38 One of the largest challenges in helping older adults to remain independent in their own homes
39 is protein-energy malnutrition (PEM), a predictor of hospitalisation, institutionalisation and
40 mortality ¹. PEM is the involuntary loss of lean tissues such as muscle, viscera, and blood and
41 immune cells, with or without loss of subcutaneous fat, as a result of inadequate energy, protein
42 and other nutrients over time ^{2,3}. As a result of decreased muscle mass and other lean tissues,
43 PEM results in decreased physical function and quality of life ^{4,5}. Older adults are particularly
44 at risk of PEM due to physiological and social challenges that occur with ageing, such as social
45 isolation, financial strain, multi-morbidities, polypharmacy, and a decreased appetite,
46 frequently referred to as the “anorexia of ageing” ^{5,6}. The first step in improving the nutrition-
47 related independence and wellbeing of older adults living at home is the accurate identification
48 of PEM, so that appropriate intervention may follow ⁷.

49 Nutrition screening is a process whereby a quick and simple validated nutrition screening tool
50 is implemented to identify risk of malnutrition, and should precede diagnostic assessment ⁸.
51 Nutrition assessment tools differ from malnutrition screening tools in that they are a
52 multidimensional and global assessment tool which are applied by a qualified health
53 professional such as a dietitian or a physician ⁹. Owing to the nature of their multidimensional
54 and detailed approach, they may be used to diagnose PEM. There are three accepted nutrition
55 assessment tools used in practice: the Subjective Global Assessment (SGA) ¹⁰, the scored
56 Patient-Generated Subjective Global Assessment (PG-SGA) ¹¹ and the Mini Nutritional
57 Assessment (MNA) ¹². Short versions of the MNA and PG-SGA (the MNA-Short Form and
58 the PG-SGA-Short Form) are available. The intended use of these shorter forms is for
59 screening for malnutrition, not assessment. A review of the validity of nutrition assessment
60 tools was evaluated by Green and Watson in 2006 ¹³ (literature searched up until 2002) and
61 Watterson et. al. in 2009 ¹⁴ (literature searched up until 2006). However, in addition to requiring

62 an update, these reviews did not pool data, used narrow search terms, and did not critically
63 appraise studies nor the body of evidence. Therefore, further investigation of the criterion
64 validity of nutrition assessment tools in older adults in the community-setting is warranted.

65 The MACRo (Malnutrition in the Ageing Community Review) Study was undertaken to
66 systematically review, quantify, and critically appraise all existing epidemiological
67 international literature concerning malnutrition prevalence, methods of risk detection and
68 diagnosis, predictors of community-acquired malnutrition and long-term outcomes of the
69 condition in older community-dwelling adults. Due to the significant amount of research
70 undertaken on this topic, as well as diverse clinical interests in the findings, the results will be
71 reported in a series of articles. This article reports the results of the following research question:
72 What is the criterion (concurrent and predictive) validity and reliability of nutrition assessment
73 tools in making a diagnosis of PEM in the general older adult community?

74 **Materials and methods**

75 This study protocol is reported using the Preferred Reporting Items for Systematic Reviews
76 and Meta-Analysis (PRISMA) 2015 Statement¹⁵ and flow diagram (Figure 1). This study has
77 been registered with the International Prospective Register of Systematic Reviews
78 (PROSPERO number: CRD42016051241).

79




Figure 1: Flowchart of the MACRo Study search and the included studies which evaluate the criterion validity of nutrition assessment tools.

81 Search strategy

82 Peer-reviewed published studies, in any language, were searched for in the electronic databases:
83 The Cochrane Library, CINAHL (via Ebscohost), EMBASE, Health Source:
84 Nursing/Academic Edition (via Ebscohost), MEDLINE (via PubMed) and Web of Science for
85 publications from database inception to the 13 September 2016. The search strategy used a
86 combination of keywords and each databases' controlled vocabulary (appendix). A snowball
87 search was conducted to complement the systematic search using the reference lists of the
88 included studies and studies included in earlier reviews.

89 Eligibility criteria: types of participants and setting

90 Inclusion criteria for types of participants were older adult samples with a mean age of ≥ 65
91 years living independently in the community (including post hospital discharge, outpatients,
92 and medical centres), who were assessed for PEM using a nutrition assessment tool.
93 Participants included in the current study were the general older population living in the
94 community. Results in disease-specific samples will be reported separately. Observational,
95 interventional (baseline or control group only), cross-sectional, retrospective and cohort studies
96 were considered for inclusion. Exclusion criteria for participants were those assessed as
97 inpatients of acute or sub-acute facilities (excepting observational assessment post-discharge),
98 day hospitals, or were living in residential aged care (including nursing homes). Data where
99 community samples were combined with patients in these settings were also excluded;
100 however, studies which used "nationally representative data" where results were not delineated
101 by setting were not excluded. Intervention studies were excluded for evaluation of predictive
102 validity due to the confounding effect of intervention on prediction of outcomes. Exclusion
103 criteria for study types were abstracts, conference papers, qualitative studies, study protocols,
104 opinions, commentaries, news articles and reviews.

105 Eligibility criteria: Criterion validity of nutrition assessment tools

106 To answer the research question, eligible studies were required to evaluate the criterion validity
107 or reliability of a nutrition assessment tool's ability to diagnose PEM (not risk of PEM).
108 Reflecting this, studies in which no participants were malnourished were excluded. For the
109 MNA, malnutrition was considered at an MNA score <17 (score 17-30 at risk/well-nourished)
110 as per the MNA instructions¹⁶; for the SGA and PG-SGA, malnutrition was considered as
111 rating B (moderately malnourished) & C (severely malnourished) as per their instructions^{10,11}.
112 Studies which evaluated the validity and reliability of modified versions of the MNA, SGA and
113 PG-SGA were included and modifications described.

114 Selection of studies

115 Identified citations from all databases were imported into EndNote [Version X7.7, 2016,
116 Thomson Reuters] and duplicates removed by two independent review authors (SM and DC).
117 A two-step screening process was employed for the first phase of study selection. In step 1,
118 two authors independently scanned the titles and abstracts of studies identified by the search
119 for their potential eligibility. At step 2, potentially eligible articles to address each MACRo
120 study research question were separated into participant groups by one author.

121 In the second phase of study selection, full-text articles were screened independently by two
122 review authors to determine eligibility for inclusion. Disagreements regarding eligibility were
123 discussed to reach consensus.

124 Data extraction and management

125 Criterion validity is composed of two types of validity assessment: concurrent and predictive.
126 Concurrent validity is determined by comparing the score of a new measurement to the score
127 of a well-established measurement, or gold standard, for the same construct. Data extracted to
128 reflect the concurrent validity were measures of diagnostic accuracy tests, including sensitivity
129 (malnourished correctly identified as such), specificity (well-nourished correctly identified as
130 such), positive predictive value (correctly identified as malnourished within malnourished

131 sample), negative predictive value (correctly identified as well-nourished within the well-
132 nourished sample), weighted kappa statistics (agreement of categories) and receiver operating
133 characteristics (ROC) curve (discriminative power of a continuous score)¹⁷. For a nutrition
134 assessment tool, there are no generally accepted a-priori values for sensitivity and specificity,
135 though it would be clinically necessary to have a balance between both high sensitivity and
136 specificity. Consideration of the reference standard used was also considered when interpreting
137 concurrent validity, as this may vary considerably due to the absence of a gold standard for
138 PEM diagnosis⁶.

139 For a nutrition assessment tool, predictive validity is usually evaluated by determining a tool's
140 ability to predict health-related outcomes known to be a consequence of PEM, such as
141 hospitalisation and mortality⁶. Outcomes were considered only if they were measured
142 subsequently to the implementation of the nutrition assessment tool, with a timeframe from one
143 week to 10 years considered. For the current study, the following categorical health-related
144 variables were considered: mortality, hospitalisation, institutionalisation, pressure ulcer/injury,
145 and falls; and continuous variables: hospitalisation cumulative length of stay (LOS), cumulative
146 duration of pressure ulcers, depression, physical function, and quality of life. All data was
147 described qualitatively in tables as well as pooled where possible. Where participants were not
148 classified dichotomously as malnourished and well-nourished, or diagnostic accuracy tests
149 were not performed, raw data extracted from the results was used to determine diagnostic test
150 accuracy wherever possible. For studies with missing data, the study authors were contacted.
151 Extracted data from published papers was undertaken by one author (SM), with a random
152 sample of 20% reviewed by a second author (DC) for accuracy.

153 Review of study strength and quality

154 External and internal study quality (including risk of bias) for individual studies was evaluated
155 by two tools depending on the study type. The Critical Appraisal Skills Programme (CASP)

156 Diagnostic Checklist ¹⁸ was chosen to appraise the study quality of studies which evaluate the
157 concurrent validity of nutrition assessment tools. This was chosen as diagnostic studies have
158 unique considerations for internal and external quality. The Academy of Nutrition and
159 Dietetics' Quality Criteria Checklist: Primary Research was chosen to evaluate studies
160 reporting on the predictive validity of nutrition assessment tools, and designates studies as
161 having positive (strong quality), neutral (neither strong nor weak quality) or negative (weak
162 quality) assessment ¹⁹. This tool was chosen to critically appraise study quality as it is
163 applicable for all original research study designs, and evaluates the external validity in respect
164 to nutrition-related conditions. The appraisal of study quality was conducted independently by
165 two authors (SM and DC). Disagreements were discussed and reported.

166 The certainty in the body of evidence for each outcome of interest was classified using the
167 Grading of Recommendations, Assessment, Development and Evaluation (GRADE) approach
168 ²⁰. This approach has four levels of quality: high (very confident the true effect lies close to
169 that estimated), moderate (moderately confident in the effect estimate), low (confidence in the
170 effect estimate is limited) and very low (very little confidence in the effect estimate). The
171 determination of the quality GRADE level was determined independently by two authors (SM
172 and JK), with disagreements managed by consensus.

173 Meta-analysis

174 Pooled data was analysed using Revman [Review Manager 5, Version 5.3, 2014, Cochrane
175 Informatics & Knowledge Management Department]. Pooled outcomes were calculated using
176 nutrition assessment tools as a dichotomous variable of “malnourished” and “well-nourished”,
177 where well-nourished includes the “at risk of malnutrition” category for the MNA.
178 Dichotomous outcome data was expressed as risk ratios (RR) with 95% confidence intervals,
179 using the Mantel-Haenszel test. Effect sizes for continuous outcome data were calculated as
180 mean differences (MD) for studies which used the same assessment tool, and standardised

181 mean differences (SMD) for studies which used different assessment tools for the same
182 construct, with 95% confidence intervals, using the inverse variance test. SMD effect sizes of
183 <0.4 were considered small, 0.4 – 0.7 moderate, and >0.7 large²¹. Where a SMD was used,
184 this was re-expressed into the scale of one the included instruments by multiplying the SMD
185 by the standard deviation of that tool reported in the total sample²². Where two or more tools
186 had scales with opposite directions of physical function (e.g. lower score indicates worse
187 physical function instead of better physical function), one of the directions was multiplied by -
188 1 to ensure scales followed the same direction²³. Acknowledging that malnutrition has
189 significant variance in its presentation between individuals and within sample populations, a
190 random effects model was used for both categorical and continuous variables. If the required
191 data of included studies was not reported, or could not be calculated or obtained, the results of
192 the study were excluded from meta-analysis and described qualitatively. Heterogeneity
193 between studies was assessed using the I² statistic, and was considered substantial if I² was
194 >50%. Where sensitivity analysis was required, analysis was repeated excluding studies with
195 low study quality/high risk of bias, timeframe of the reported outcome, study design or
196 participant characteristics. No subgroup analyses were found to be necessary to answer the
197 research hypothesis.

198 **Results**

199 Search results and included studies

200 The search identified 6,412 records, of which 1,925 were removed as duplicates (Figure 1).

201 The two authors agreed on a total of 104 potentially eligible records evaluating the criterion
202 validity and/or reliability of a nutrition assessment tool in the general older adult community
203 setting. Following full-text review, eight studies were found to be eligible (Figure 1). Studies
204 were included from Europe (n=4 studies), Asia (n=3 studies) and South America (n=1 studies)
205 (Table 1 and Table 2). Most study samples were recruited via home care (n=5 studies); and,
206 two studies were conducted on the same nationally representative sample in the People's

207 Republic of China (Taiwan). Nutrition assessment tools were completed by nurses (n=2),
208 trained researchers (n=2), or personal/domiciliary carers (n=1); none appear to have been
209 completed by dietitians, although the tool was completed by “nutrition scientists” in one study
210 (Table 2). Additionally, the two studies in the People’s Republic of China (Taiwan) using the
211 same nationally representative dataset did not complete any nutrition assessment tool with
212 older adults, but rather constructed the MNA-T2 (MNA Taiwan-version 2) tool based on items
213 from a larger generic health-based questionnaire ^{24,25}. The MNA-T2 differs from the usual
214 MNA by using Taiwanese-specific anthropometric cut-off points. Furthermore, two items of
215 the MNA-T2 could not be obtained by the researchers (pressure ulcers and fluid intake) so the
216 score was proportionately adjusted where a score of 16.5 or less indicated malnutrition, 17-
217 23.5 indicated risk of malnutrition, and 24 or more indicated normal nutrition status ²⁴. No
218 studies were identified which evaluated the reliability of nutrition assessment tools in this
219 setting. No new global and multidimensional nutrition assessment tools were identified.

220

221 **Table 1:** Comparison of concurrent validity of nutrition assessment tools evaluated in the community setting

Study	Nutrition assessment	Population	Sensitivity ^a	Specificity ^a	Positive Predictive Value ^a	Negative Predictive Value ^a	Kappa ^b	ROC AUC ^c	CASP ^d risk of bias
Kozakova 2012. Data pooled: No.	Tool: MNA ^e /SGA ^{f,g} Benchmark: MNA/SGA	n=120, μ age 73.24 years (SD not reported). Country: Czech Republic & Slovakia Setting: Home care Assessed by: Research nurses.	71.7 (56.5-84.0)	86.5 (76.6-93.3)	76.7 (61.4-88.2)	83.1 (72.9-90.7)	Not reported	Not reported	High
Kozakova 2014. Data pooled: No.	Tool: SGA Benchmark: Nutrition assessment not further described ^h	n=470, μ age 77.3 years (SD not reported). Country: Czech Republic Setting: Home care Assessed by: 10 trained nurses.	93.3 (95%CI not reported)	70 (95%CI not reported)	62.6 (95%CI not reported)	98.4 (95%CI not reported)	Not reported	Not reported	High
Kozakova 2014. Data pooled: No.	Tool: SGA Benchmark: MNA ⁱ	As above.	Not reported	Not reported	Not reported	Not reported	0.442 (95%CI not reported)	Not reported	As above.

222 a data presented %, 95% confidence interval.

223 b data presented as kappa coefficient, 95% confidence interval, p-value.

224 c ROC AUC, Receiver Operating Characteristic Area Under the Curve. Data presented as AUC value \pm standard error, 95% confidence interval, p-value.

226 d CASP, Critical Appraisal Skills Programme

227 e MNA, Mini Nutritional Assessment. MNA score of <17 indicates malnutrition.
228 f The comparative validity of the MNA and SGA was assessed by comparing each assessment tool against the other, where the authors
229 considered both tools as the reference standard.
230 g SGA, Subjective Global Assessment. SGA ratings B and C indicate malnutrition.
231 h Authors report that for the reference standard, participants were grouped into two categories: good nutritional status and nutritional risk, based
232 on their nutrition status. The nutrition risk category was created by fusing the risk of malnutrition and malnutrition categories together. However,
233 it is unclear what was used to inform the nutritional status used to create these two categories. It cannot be the MNA, SGA or the Malnutrition
234 Universal Screening Tool, as all these tools were compared against this standard.
235 i SGA (rating B & C) compared against the MNA dichotomised at <24; therefore, including both “at risk of malnutrition” and “malnourished”
236 MNA categories for the kappa coefficient.

237 **Table 2:** The predictive validity of the Malnutrition Screening Tool (MNA) in the community setting

Study	Population	Time-point	Malnourished with outcome ^a	Well-nourished with outcome ^b	Reported results	AND ^c study quality
Mortality						
Ferreira 2011. Data pooled: Yes.	n=1170, μ age not provided. n=675 in 60-74 year age group; n=495 in ≥ 75 year age group. Country: Brazil Setting: Home Assessed by: trained health professionals and nutrition trainees	7 years	17/30 (56.7%)	315/1140 (27.6%)	Compared with well-nourished (MNA score 24-30) adjusted odds of mortality in malnourished (MNA score <17) was: - OR: 6.05 (95%CI 5.75-6.35) for 60-74 years - OR: 2.76 (95%CI 2.51-3.04) for ≥ 75 years	Positive
Inoue 2007. Data pooled: Yes.	n=181, μ age 78.9 \pm 8.7 years. Country: Japan Setting: Home care Assessed by: trained operators	3 years	14/45 (31.1%) ^d	18/136 (13.2%) ^d	Compared with well-nourished (MNA score 24-30) via adjusted hazard ratio of mortality in malnourished (MNA score <17) was: - HR:14.05 (95%CI: 3.171-64.242)	Neutral
Kiesswetter 2014. Data pooled: Yes.	n=353 ^e , μ age 80.9 \pm 7.9 years. Country: Germany Setting: Home care Assessed by: nutrition scientists	1 year	12/42 (28.6%)	33/267 (12.4%)	Compared with well-nourished (MNA score 24-30), adjusted hazard ratio of mortality in malnourished (MNA score <17) was: - HR: 8.75 (95%CI: 2.45-31.18)	Neutral
Lee 2012 and Wang 2013 ^f . Data pooled: Yes.	n=2948, μ age not provided. n=1866 in 65-74 year age group; n=1082 in ≥ 75 year age group. Country: China (Taiwan) Setting: Nationally representative data, not further described.	4 years	70/90 (76.9%)	591/2857 (20.7)	Compared with well-nourished (MNA score 24-30), adjusted hazard ratio of mortality in malnourished (MNA score <17) was: HR: 3.26 (95%CI: 2.31-4.6; P<0.001).	Both studies were Neutral

	Assessed by: Constructed in research setting based on individual data collected during the Taiwan Longitudinal Survey on Aging (TLSA) researchers ^g .					
Saletti 2005. Data pooled: Yes.	n=353, μ age 83.0 \pm 6.8 years. Country: Sweden Setting: Home care Assessed by: Personal / domiciliary carers	3 years	14/29 (50%) ^h	108/324 (33.3%) ^h	Compared with well-nourished (MNA score 24-30), mortality rates in malnourished (MNA score <17) were significantly higher ($P=0.03$).	Positive
Physical function						
Kiesswetter 2014. Data pooled: Yes.	Reported above.	1 year	Not reported.	Not reported.	Barthel Index mean scores (\pm SD) are: Malnourished (MNA score <17): - 32.3 \pm 25.9 (n=30) At risk of malnutrition (MNA score 17 – 23.5) - 53.9 \pm 25.8 (n=148) Well-nourished (MNA score 24-30) - 76.5 \pm 25.8 (n=86). Scores differed significantly across groups ($P<0.05$)	Neutral
Lee 2012 ^e . Data pooled: Yes.	Reported above.	4 years	Became or remained ADL ⁱ -dependent 3/21 (14.3%). Became or remained IADL ^j -dependent	Became or remained ADL-dependent 47/225 (20.9%). Became or remained IADL-dependent	ADL mean scores (\pm SD) are: Malnourished (MNA score <17): - 2.4 \pm 4.9 (n=21) At risk of malnutrition (MNA score 17 – 23.5) - 3.6 \pm 5.9 (n=225) Well-nourished (MNA score 24-30) - 1.2 \pm 3.8 (n=1944).	Neutral

			17/21 (81.0%).	127/225 (56.4%).	Scores differed significantly malnourished and at-risk groups ($P<0.05$) IADL mean scores (\pm SD) are: Malnourished (MNA score <17): - 9.4 \pm 5.3 (n=21) At risk of malnutrition (MNA score 17 – 23.5) - 7.5 \pm 6.6 (n=225) Well-nourished (MNA score 24-30) - 3.7 \pm 5.4 (n=1944). Scores differed significantly malnourished and at-risk groups ($P<0.05$)	
--	--	--	-------------------	---------------------	--	--

- 238 a For categorical/dichotomous outcomes, data reported number with the outcome at the time-point/number malnourished (MNA score <17) in
239 sample at baseline (% with outcome within malnourished sample).
- 240 b For categorical/dichotomous outcomes, data reported number with the outcome at the time-point/number well-nourished (including at risk of
241 malnutrition, MNA score 17-30) in sample at baseline (% with outcome within well-nourished sample).
- 242 c AND, Academy of Nutrition & Dietetics
- 243 d Data was not reported in the study publication but was provided by authors in an email dated 07/03/2017.
- 244 e n=309 (87.5%) were assessed by the MNA
- 245 f The study results on the same sample were reported across two studies, Lee 2012 and Wang 2013; Lee 2012 reported the number of deaths per
246 category. These studies used the Taiwan Version 2 (MNA-T2) as opposed to the traditional English-language MNA. This tool adopts the
247 Taiwanese-specific anthropometric cut-off points and replaced calf circumference with BMI.
- 248 g Data for all items in the long-form MNA (MNA), except items pressure sore/skin ulcers and fluid intake, were available in the survey database.
249 So, the MNA was based on fifteen items with a maximum score of 28 points, rather than seventeen items for 30 points. However, the total
250 score was proportionately adjusted on the full-score basis. A final score of 16.5 or less suggests malnourishment; 17–23.5, at risk of
251 malnutrition; and 24 or more, normal.

252 h Mortality data was reported as a percentage per MNA category for 224 of the 535 who had mortality data available on public registers.
253 However, the number of participants in the 224 subsample each MNA category was not reported. Therefore, mortality data was extrapolated to
254 the large sample size (e.g. 50% died in malnourished group was reported as 14/29 although exact figures are not known).
255 i ADL, Activities of Daily Living. ADL was measured by a questionnaire adapted from the 1984 National Health Interview Survey Supplement
256 on Aging. Becoming or remaining dependent was considered if the participant had 1 or more dependencies.
257 j IADL, Instrumental Activities of Daily Living. IADL was measured by a questionnaire adapted from the 1984 National Health Interview
258 Survey Supplement on Aging. Becoming or remaining dependent was considered if the participant had 1 or more dependencies.

259 The concurrent validity of nutrition assessment tools in the community

260 There were two studies reporting the concurrent validity of the MNA and SGA (Table 1). Two
261 other studies were identified which reported diagnostic accuracy for the MNA; however, one
262 study was excluded as the reference standard was the Fried Frailty Index, a construct which
263 does not represent malnutrition and therefore does not inform on the ability of the MNA to
264 diagnose malnutrition ²⁶. The second study was excluded as the authors did not report which
265 score was used to dichotomise the MNA categories, the reference standard was not reported in
266 the publication and this missing information could not be obtained ²⁷. No studies were
267 identified which evaluated the criterion validity of the Scored PG-SGA.

268 In the 2012 study, the MNA (score <17 indicating malnutrition, score 17-30 indicating well-
269 nourished) and SGA (rating B and C indicating malnutrition, rating A indicating well-
270 nourished) were compared with each other, where neither tool was considered the “reference
271 standard” ²⁸. This study provided the results in a contingency table, and therefore the diagnostic
272 accuracy tests were performed by the current study authors (SM and checked by DC). When
273 compared against each other, the SGA and MNA had good specificity; however, the sensitivity
274 was lower (Table 1). Kozakova et. al. (2014) further compared the MNA and SGA against each
275 other via a kappa coefficient in a larger sample, which revealed moderate agreement. However,
276 the MNA included both the at risk of malnutrition and malnourished categories for this test
277 (score <24 indicating malnutrition, score 24-30 indicating well-nourished) and therefore the
278 two tools would be expected to have less agreement due to inconsistent categorisation ²⁹. In the
279 2014 Kozakova study, the SGA was found to have strong sensitivity but a lower specificity
280 compared to an unknown benchmark which represents both risk of malnutrition and
281 malnutrition. Both studies were considered to have high risk of bias (Online Supplementary
282 Material). The quality of the evidence (GRADE) for the concurrent validity of both the MNA
283 and SGA was “very low” (Table 3).

284 The predictive validity of nutrition assessment tools in the community

285 Studies which report the predictive validity of nutrition assessment tools were only found for
286 the MNA (n=6 studies). Mortality was reported by five studies, and physical function (using
287 three different measurement tools) was reported by two studies (Table 2). Although Lee and
288 Tsai²⁴ and Wang and Tsai²⁵ were both included, their results were on the same study sample,
289 overseen by the same senior author, and both used the MNA-T2 so were reported together
290 (Table 2). No other outcomes were reported to evaluate the predictive validity of nutrition
291 assessment tools in the community.

292 Regarding mortality, the time to follow-up ranged from 1 – 7 years, and included samples from
293 Asia, Europe and South America. The number of deaths per MNA category were not provided
294 in the study reported by Inoue and Kato³⁰; however, the study authors provided this data by
295 email. There was high heterogeneity in the meta-analysis of mortality when all five studies
296 were included (RR: 2.30 [95%CI: 1.43 – 3.70]; $P < 0.0006$; n=6,152 participants; n=5 studies;
297 I^2 : 89%). However, sensitivity analysis reduced the heterogeneity to I^2 : 0% when Lee and Tsai
298²⁴, which used the MNA-T2, was deselected, as this version differs to the usual MNA in several
299 ways. This result showed that the MNA categorisation of malnutrition (MNA score <17) was
300 able to predict a two-fold risk of death compared to community dwelling older adults
301 categorised as at risk of malnutrition or well-nourished (MNA score 17-30) (RR: 1.92 [95%CI:
302 1.55-2.39]; $P < 0.00001$; n=2,013 participants; n=4 studies; I^2 : 0%) (Figure 2). Two included
303 studies were considered to have positive quality, two to have neutral quality (Online
304 Supplementary Material). The quality of the evidence (GRADE) for the predictive validity of
305 the MNA in detecting risk of death was “moderate” (Table 3).

306


307 **Table 3:** Quality of the body of evidence for each outcome of interest reflecting the Grading of Recommendations, Assessment, Development
 308 and Evaluation (GRADE) approach

Outcome	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication bias^a	Quality of evidence (GRADE)^b
Concurrent validity of the MNA	Very serious ^c	Not applicable	Serious ^d	Not serious	Could not be assessed	⊕○○○ Very Low
Concurrent validity of the SGA	Very serious ^c	Not applicable	Serious ^d	Serious (data not reported)	Could not be assessed	⊕○○○ Very low
Predictive validity of the MNA (mortality)	Serious ^e	Not serious	Not serious	Not serious	Could not be assessed	⊕⊕⊕○ Moderate
Predictive validity of the MNA (physical function)	Serious ^f	Serious ^g	Not serious	Serious ^h	Could not be assessed	⊕○ ○○ Very Low

- 309 a. Could not be assessed for any outcome due to the small number of included studies.
- 310 b. Graded on a scale of high, moderate, low to very low quality of evidence. Each study was downgraded one level for having a “serious” limitation,
 311 and downgraded two levels for a “very serious” limitation to the quality of all studies informing the outcome.
- 312 c. Found to have a high risk of bias when evaluated using the CASP diagnostic checklist (Online Supplementary Material)
- 313 d. Low generalisaibility due to poor description of the persons who undertook the nutrition assessment, their level of training, how the nutrition
 314 assessment was completed, and representing only one study sample.
- 315 e. Two were found to have positive study quality and three neutral study quality by the AND tool (Online Supplementary Material).
- 316 f. Both studies were found to have neutral study quality by the AND tool (Online Supplementary Material).
- 317 g. The meta-analysis of this outcome variable showed substantial heterogeneity.
- 318 h. The meta-analysis of this outcome variable showed a substantial confidence interval.

319 Physical function was measured 1-year and 4-years following nutritional assessment. It was
320 not possible to compare the malnourished participants to the combined well-nourished and at
321 risk of malnutrition groups, so analysis was repeated comparing malnutrition to each MNA
322 category respectively. The results by Lee and Tsai ²⁴ were entered twice as they presented data
323 using two measures of physical function (Table 2). There were significant results when
324 participants in the malnourished category (MNA score <17) were compared to the well-
325 nourished category (MNA score 24-30), revealing a large but imprecise effect size of physical
326 dysfunction in the malnourished group (SMD: 1.02 [95%CI: 0.24-1.80]; $P=0.01$; $n=4,046$
327 participants; $n=3$ studies; $I^2:89\%$) (Figure 3). When transformed back into the Barthel Index
328 on a scale of 0 – 100, where a higher score indicates better physical function, the MNA
329 predicted a difference of 29.4 points between the MNA malnourished and well-nourished
330 categories. The Barthel Index was chosen to represent the difference in physical function as
331 this was the only tool represented in the meta-analysis which has been well described and
332 validated for use in older adults ³¹. The high heterogeneity, which did not significantly improve
333 with sensitivity analysis, is likely due to the differences in the MNA tool used between Lee and
334 Tsai ²⁴ and Kiesswetter ³², as well as the use of three different physical function assessment
335 tools, which may represent different constructs of physical function. The meta-analysis found
336 no significant result when malnutrition (MNA score <17) was compared to at risk of
337 malnutrition (MNA score 17-23.5), and this did not improve with sensitivity analysis (SMD:
338 0.32 [95%CI: -0.28-0.91]; $P=0.30$; $n=670$ participants; $n=3$ studies; $I^2: 82\%$). The two studies
339 which reported the physical function were both rated as having neutral quality (Online
340 Supplementary Material). The quality of the evidence (GRADE) for the predictive validity of
341 the MNA in detecting risk of poor physical function was “very low” (Table 3).

342




343

344 **Figure 2:** The relative risk of death in malnourished (MNA score <17) compared to risk of malnutrition/well-nourished (MNA score 17-30)
 345 community-dwelling older adults as determined by the Mini Nutritional Assessment (MNA).

346

347



348

349 **Figure 3:** The standardised mean difference in physical function between malnourished (MNA score <17) compared to well-nourished (MNA
 350 score 24-30) community-dwelling older adults as determined by the Mini Nutritional Assessment (MNA).

351

352 **Discussion**

353 This is a comprehensive systematic literature review and meta-analysis of the criterion validity
354 of nutrition assessment tools in the community for the diagnosis of PEM. Overall, few studies
355 have evaluated the criterion validity and no studies have evaluated the reliability of nutrition
356 assessment tools in this setting. The results presented in this study reveal that although nutrition
357 assessment tools are frequently used by clinicians and researchers in the older community
358 setting, the current body of evidence provides very little confidence in their diagnostic accuracy
359 indicated by having a “very low” quality of evidence across all studies (Table 3). The 2012
360 study by Kozakova et. al.³³ found that when the MNA and SGA are compared against each
361 other, there is adequate specificity (86.5%) but inadequate sensitivity for a nutrition assessment
362 tool (71.7%). The poor sensitivity between the SGA and MNA agrees with previous research
363 in other settings which has found that the MNA and SGA do not consider the same patients as
364 malnourished, where the MNA has been considered to underestimate malnutrition (MNA score
365 <17) when compared to various reference standards⁶.

366 In the 2014 study by Kozakova et. al.²⁹, the SGA was reported to have excellent sensitivity
367 (93.3%) but inadequate specificity (70%); however, it is likely that the true specificity is higher
368 as the unknown reference standard used was reported to include both “malnourished” and “at
369 risk of malnutrition” individuals, which would lead to a lower reported specificity. Overall,
370 these two studies contribute little to the understanding of the concurrent validity of the MNA
371 and SGA in the older adult community. Both were found to have a high risk of bias due to
372 both studies being completed by non-blinded researchers who undertook all data collection, a
373 lack of appropriate diagnostic accuracy statistics, no description of the training of the
374 researchers who do not have backgrounds in nutrition, and reference standard used to evaluate
375 the SGA was unknown (Online Supplementary Material). Although it must be acknowledged
376 that the lack of a gold standard in diagnosing PEM lends to difficulties in identifying an

377 appropriate reference standard to evaluate the concurrent validity of nutrition assessment tools,
378 the reference standard should be multidimensional, represent PEM, and be well described.
379 Although the current study revealed a poor quality of evidence regarding the diagnostic
380 accuracy of nutrition assessment tools in the community setting, the MNA, SGA and PG-SGA
381 have undergone more rigorous evaluation in acute, subacute and disease-specific populations
382 ^{6,12,34-37}.

383 Only the MNA could be evaluated for predictive validity. This study found that the current
384 body of evidence provides moderate confidence in the ability of the MNA category of
385 malnutrition to predict the risk of death 1 to 7 years following the diagnosis of malnutrition.
386 However, the body of evidence provides only very limited confidence for the ability of the
387 MNA to predict physical dysfunction. Although the MNA has not been evaluated appropriately
388 for concurrent validity, the finding that it has good predictive validity for risk of death is
389 clinically relevant, as prediction of poor health outcomes may be of more clinical significance
390 by indicating the need for intervention, than diagnostic accuracy in the community setting.

391 Further diagnostic accuracy, reliability and prognostic studies in the general older community
392 will help guide which nutrition assessment tool is best suited to this setting. However, until
393 further research is undertaken to guide tool selection, nutrition assessment should continue to
394 be done to identify patients that may be malnourished; however, monitoring response to
395 intervention is of high importance in the absence of evidence for accurate and reliable
396 diagnostic tools ⁷. Additionally, poor sensitivity in the nutrition assessment tools suggests that
397 intervention may be necessary for some individuals identified as at risk of malnutrition or with
398 borderline results, either to prevent malnutrition from developing or to provide treatment to an
399 individual inaccurately identified as “well-nourished”. As per best practice guidelines, such
400 treatment should be individualised ¹⁴.

401 Limitations

402 This systematic literature review is limited in that it did not include grey literature and was
403 unable to obtain complete results for all studies. This was due to poor reporting in some original
404 studies and because most authors were unable to be contacted or they no longer had access to
405 the raw data to generate the results needed for this review. Although the literature search
406 conducted for this study was comprehensive, there remains the possibility that important
407 studies were missed due to not being included in the search or mistakenly excluded by review
408 authors. The results of the criterion validity of nutrition assessment tools were narrowed by
409 excluding studies undertaken with samples combining community-dwelling participants with
410 inpatient or residential aged care participants, as this led to the exclusion of otherwise eligible
411 studies. This study did not evaluate nutrition assessment tool translation or discriminant
412 validity, or responsiveness. Therefore, future systematic reviews are needed to evaluate these
413 important aspects of nutrition assessment.

414 Conclusions

415 This review found that no nutrition assessment tool has undergone sufficient evaluation for
416 concurrent validity in community-dwelling older adults. There is moderate confidence in the
417 ability of the MNA to predict a two-fold risk of death and very limited confidence in its ability
418 to predict physical dysfunction following nutrition assessment. Due to the small number of
419 studies identified and no evaluation of the predictive validity of tools other than the MNA,
420 there is insufficient evidence to recommend a particular nutrition assessment tool for
421 diagnosing PEM in older adults in the community; however, nutrition assessment should
422 continue to be undertaken to ensure malnourished patients are managed and supported. High
423 quality diagnostic accuracy studies are needed for all nutrition assessment tools in non-disease
424 specific older community samples; and studies are needed which measure health outcomes
425 subsequent to nutrition assessment by the SGA and PG-SGA.

426

427 **Highlights**

- 428 • Quality of the evidence for the concurrent validity of the MNA and SGA was very
429 low
- 430 • Quality of the evidence for the MNA to predict risk of death was moderate
- 431 • Quality of the evidence for the MNA to predict risk of physical dysfunction was very
432 low
- 433 • There is insufficient evidence to recommend a particular nutrition assessment tool
- 434 • High quality diagnostic studies are needed for all nutrition assessment tools

435

436 **Declarations**

437 Competing interests

438 The authors declare that they have no financial, personal or potential competing interests.

439 Funding

440 This research did not receive any specific grant from funding agencies in the public,
441 commercial or not-for-profit sectors.

442 Authors' contributions

443 SM and DC carried out the literature search, record screening, data extraction and critical
444 appraisal of individual studies. SM and JK completed the GRADE assessment. SM drafted and
445 revised the manuscript and undertook the meta-analyses. JK provided advice, guidance in the
446 planning of the meta-analysis, and assisted in interpretation of pooled results. DC, JK and EI
447 provided guidance and revision of the manuscript.

448 Acknowledgements

449 The authors gratefully acknowledge the assistance of D. Honeyman, Librarian at Bond
450 University, for contributing to the search strategy. The authors also thank Professor Keiko
451 Inoue for providing additional data beyond that included in her publication upon request.

References

1. Marshall S, Bauer J, Isenring E. The consequences of malnutrition following discharge from rehabilitation to the community: a systematic review of current evidence in older adults. *J Hum Nutr Diet.* 2014;27(2):133-141.
2. Skipper A. Agreement on defining malnutrition. *Journal of Parenteral and Enteral Nutrition.* 2012;36(3):261-262.
3. Pleuss J. Alterations in nutritional status. In: Porth CM, ed. *Pathophysiology, concepts of altered health states.* Vol 7th e.d. Philadelphia: Lippincott Williams & Wilkins; 2005:217 - 238.
4. Agarwal E, Marshall S, Miller M, Isenring E. Optimising nutrition in residential aged care: a narrative review. *Maturitas.* 2016;92:70-78.
5. Marshall S. Why is the skeleton still in the hospital closet? A look at the complex aetiology of malnutrition and its implications for the nutrition care team. *J Nutr Health Aging.* 2017;In Press.
6. Marshall S. Protein-energy malnutrition in the rehabilitation setting: evidence to improve identification. *Maturitas.* 2016;86:77-85.
7. Lacey K, Prichett E. Nutrition Care Process and Model: ADA adopts road map to quality care and outcomes management. *J Am Diet Assoc.* 2003;103(8):1061-1072.
8. Cederholm T, Barazzoni R, Austin P, et al. ESPEN guidelines on definitions and terminology of clinical nutrition. *Clin Nutr.* 2017;36(1):49-64.
9. Field L, Hand R. Differentiating Malnutrition Screening and Assessment: A Nutrition Care Process Perspective. *Journal of the Academy of Nutrition and Dietetics.* 2015;115(5):824-828.
10. Detsky AS, McLaughlin JR, Paker JP, et al. What is subjective global assessment of nutritional status. *JPEN.* 1987;11:8 - 11.
11. Ottery FD. Patient-Generated Subjective Global Assessment. In: McCallum PD, Polisena CG, eds. *The Clinical Guide to Oncology Nutrition.* Chicago: The American Dietetic Association; 2000:11-23.
12. Vellas B, Guigoz Y, Garry PJ, et al. The Mini Nutritional Assessment (MNA) and its use in grading the nutritional state of elderly patients. *Nutrition.* 1999;15(2):116-122.
13. Green SM, Watson R. Nutritional screening and assessment tools for older adults: Literature review. *J Adv Nurs.* 2006;54(54):477 - 490.
14. Watterson C, Fraser A, Banks M, et al. Evidence based practice guidelines for the nutritional management of malnutrition in patients across the continuum of care. *Nutrition & Dietetics.* 2009;66(Suppl 3):S1 - S34.
15. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg.* 2010;8(5):336-341.
16. *Nutrition screening as easy as mna. A guide to completing the Mini Nutritional Assessment (MNA).* Switzerland Nestle Nutrition Institute.
17. Šimundić A-M. Measures of diagnostic accuracy: basic definitions. *Med Biol Sci.* 2008;22(4):61-65.
18. Critical Appraisal Skills Programme (CASP) Diagnostic Test Study Checklist. Critical Appraisal Skills Programme; 2013.
19. *Evidence analysis manual: Steps in the Academy Evidence Analysis Process.* www.andevidencelibrary.com: Academy of Nutrition and Dietetics;2012.
20. Guyatt GH, Oxman AD, Schünemann HJ, Tugwell P, Knottnerus A. GRADE guidelines: a new series of articles in the Journal of Clinical Epidemiology. *J Clin Epidemiol.* 2011;64(4):380-382.

21. Higgins, Julian, Green. 17.8.2 Study summaries using more than one patient-reported outcome. *Cochrane handbook for systematic reviews of interventions*2011.
22. Higgins, Julian, Green. 12.6.4 Re-expressing SMD using a familiar instrument. *Cochrane handbook for systematic reviews of interventions*2011.
23. Higgins, Julian, Green. 9.2.3.2 The standardized mean difference. *Cochrane handbook for systematic reviews of interventions*2011.
24. Lee LC, Tsai AC. Mini-Nutritional Assessment predicts functional decline of elderly Taiwanese: Result of a population-representative sample. *Br J Nutr*. 2012;107(11):1707-1713.
25. Wang JY, Tsai AC. The short-form mini-nutritional assessment is as effective as the full-mini nutritional assessment in predicting follow-up 4-year mortality in elderly Taiwanese. *Journal of Nutrition Health & Aging*. 2013;17(7):594-598.
26. Jurschik P, Botigue T, Nuin C, Lavedan A. Association between Mini Nutritional Assessment and the Fried frailty index in older people living in the community. *Med Clin (Barc)*. 2014;143(5):191-195.
27. Calderon Reyes ME, Ibarra Ramirez F, Garcia J, Gomez Alonso C, Rodriguez-Orozco AR. [Compared nutritional assessment for older adults at family medicine settings]. *Nutr Hosp*. 2010;25(4):669-675.
28. Kozakova R, Jarosova D, Zelenikova R. Comparison of three screening tools for nutritional status assessment of the elderly in their homes. *Biomedical Papers-Olomouc*. 2012;156(4):371-376.
29. Kozakova R, Zelenikova R. Assessing the nutritional status of the elderly living at home. *European Geriatric Medicine*. 2014;5(6):377-381.
30. Inoue K, Kato M. Usefulness of the Mini-Nutritional Assessment (MNA) to evaluate the nutritional status of Japanese frail elderly under home care. *Geriatrics & Gerontology International*. 2007;7(3):238-244.
31. Mahoney FI, Barthel D. Functional evaluation: the Barthel Index. *Md State Med J*. 1965;14:51-61. Used with permission.
32. Kiesswetter E, Pohlhausen S, Uhlig K, et al. Prognostic differences of the mini nutritional assessment short form and long form in relation to 1-year functional decline and mortality in community-dwelling older adults receiving home care. *J Am Geriatr Soc*. 2014;62(3):512-517.
33. Kozakova R, Jarosova D, Zelenikova R. Comparison of three screening tools for nutritional status assessment of the elderly in their homes. *Biomedical Papers*. 2012;156(4):371-376.
34. Marshall S, Young A, Bauer J, Isenring E. Malnutrition in geriatric rehabilitation: prevalence, patient outcomes and criterion validity of the Scored Patient-Generated Subjective Global Assessment (PG-SGA) and the Mini Nutritional Assessment (MNA) *Journal of the Academy of Nutrition and Dietetics*. 2016;116(5):785-794.
35. Isenring E, Bauer J, Capra S. The scored Patient-generated Subjective Global Assessment (PG-SGA) and its association with quality of life in ambulatory patients receiving radiotherapy. *Eur J Clin Nutr*. 2003;57(2):305-309.
36. Bauer J, Capra S, Ferguson M. Use of the scored Patient-Generated Subjective Global Assessment (PG-SGA) as a nutrition assessment tool in patients with cancer. *Eur J Clin Nutr*. 2002;56:779-785.
37. Bauer JM, Vogl T, Wicklein S, Trögner J, Mühlberg W, Sieber C. Comparison of the Mini Nutritional Assessment, Subjective Global Assessment, and Nutritional Risk Screening (NRS 2002) for nutritional screening and assessment in geriatric hospital patients. *Z Gerontol Geriatr*. 2005;38(5):322-327.

Appendix: Search strategy implemented across six electronic databases and results of total records retrieved

Set	Search Terms
<i>MEDLINE (via PubMed) - searched 13 September 2016 using keywords (text word) and MeSH Terms. Result = 1,766 records</i>	
#1	PGSGA [Text Word] OR SGA[Text Word] OR MNA[Text Word] OR "Patient generated subjective global assessment"[Text Word] OR "subjective global assessment"[Text Word] OR "mini nutritional assessment"[Text Word]
#2	Nutrition* [Text Word] OR malnutrition [Text Word] OR “nutrition* status” [Text Word] OR undernutrition [Text Word] OR emaciation [Text Word] OR undernourish* [Text Word] OR protein energy malnutrition [MeSH term] OR malnutrition [MeSH term] OR nutritional status [MeSH term] OR undernutrition [MeSH term] OR nutritional deficiency [MeSH term] OR protein calorie malnutrition [MeSH term] OR emaciation [MeSH term] OR nutrition status [MeSH term] OR protein deficiency [MeSH term]
#3	Screen* [Text Word] OR mass screening [MeSH Terms]
#4	2 AND 3
#5	Diagnos* [Text Word] OR evaluat* [Text Word] OR valid* [Text Word] OR compar* [Text Word] OR “outcome assessment” [Text Word] OR “outcome measure*” [Text Word] OR agreement [Text Word] OR precision [Text Word] OR kappa* [Text Word] OR specificit* [Text Word] OR sensitiv* [keyword] OR accur* [Text Word] OR outcome assessment health care [MeSH term] OR diagnostic related groups [MeSH term] OR diagnosis [MeSH term] OR diagnoses and examinations [MeSH term] OR examinations and diagnoses [MeSH term] OR validity of results [MeSH term]
#6	4 AND 5
#7	Community [Text Word] “community dwelling” [Text Word] OR “community living” [Text Word] OR “community based” [Text Word] OR “community setting” [Text Word] OR “free living” [Text Word] OR “independent living” [Text Word] OR “home” [Text Word] OR “general practice” [Text Word] OR “primary health care” [Text Word] OR “primary care” [Text Word] OR “primary healthcare” [Text Word] OR “primary nursing” [Text Word] OR [Text Word] OR “primary nursing care” [Text Word] OR general practice [MeSH term] OR primary health care [MeSH term] OR primary care nursing [MeSH term] OR primary healthcare [MeSH term] OR primary nursing [MeSH term] OR care, primary nursing [MeSH term] OR primary nursing care [MeSH term] OR agencies, home care [MeSH term] OR home care services [MeSH term] OR home nursing [MeSH term] OR independent living [MeSH term]
#8	(1 OR 6) AND 7

<i>CINAHL (via Ebscohost) was searched on 13 September 2016 using keywords and CINAHL Headings. Results = 1,068 records</i>	
#1	“PGSGA” [keyword] OR “SGA” [keyword] OR “MNA” [keyword] OR "Patient generated subjective global assessment"[keyword] OR "subjective global assessment"[keyword] OR "mini nutritional assessment"[keyword]
#2	(MH "Geriatric Nutrition") OR (MH "Malnutrition") OR "malnutrition" OR (MH "Protein-Energy Malnutrition+") OR (MH "Nutritional Status") OR "nutrition status" OR "undernutrition" OR "nutritional deficiency" OR (MH "Nutrition") OR "Nutrition" OR (MH "Nutritional Assessment") OR "nutritional assessment"
#3	(MH "Health Screening+") OR (MH "Rescreening")
#4	2 AND 3
#5	"(MH "Diagnosis+") OR "DIAGNOSIS" OR (MH "Diagnosis, Differential") OR (MH "Predictive Validity") OR (MH "Criterion-Related Validity+") OR (MH "Concurrent Validity") OR (MH "Validity+") OR "VALIDITY" OR (MH "Reliability and Validity+") OR (MH "External Validity") OR (MH "Internal Validity") OR (MH "Sensitivity and Specificity") OR (MH "Outcome Assessment") OR "OUTCOME MEASURE" OR (MH "Kappa Statistic") OR "KAPPA" OR (MH "Intrarater Reliability") OR (MH "Interrater Reliability") OR (MH "Consensus")
#6	4 AND 5
#7	"(MH "Community Living+") OR (MH "Communities+") OR "community" OR "community dwelling" OR (MH "Community Health Nursing+") OR "community health nursing" OR (MH "Community Health Services+") OR (MH "Home Nursing, Professional") OR (MH "Home Nutritional Support") OR (MH "Primary Nursing") OR "primary nursing" OR "free living" OR "independent living" OR (MH "Family Practice") OR "general practice" OR (MH "Home Health Care+") OR "meals on wheels" OR "community dietitian" OR "community dietician" OR (MH "Rehabilitation, Community-Based")
#8	(1 OR 6) AND 7
<i>The Cochrane Library was searched on 13 September 2013 using keywords and MeSH Headings. Results = 885 records</i>	
#1	“PGSGA” [keyword] OR “SGA” [keyword] OR “MNA” [keyword] OR "Patient generated subjective global assessment"[keyword] OR "subjective global assessment"[keyword] OR "mini nutritional assessment"[keyword]
#2	Nutrition* [Text Word] OR malnutrition [Text Word] OR “nutrition* status” [Text Word] OR “nutrition risk” [Text Word] OR undernutrition [Text Word] OR “nutrition* defici*” [Text Word] OR emaciation [Text Word] OR undernourish* [Text Word] OR protein-energy malnutrition [exp] [MeSH term] OR malnutrition [exp] [MeSH term] OR nutritional status [exp] [MeSH term] OR emaciation [exp] [MeSH term] OR nutrition status [MeSH term] OR protein deficiency [MeSH term]

#3	Screen* [keyword] OR Mass Screening [exp] [Mesh term]
#4	2 AND 3
#5	Diagnos* [Text Word] OR evaluat* [Text Word] OR valid* [Text Word] OR compar* [Text Word] OR “outcome assessment” [Text Word] OR “outcome measure*” [Text Word] OR agreement [Text Word] OR precision [Text Word] OR kappa [Text Word] OR specificit* [Text Word] OR sensitiv* [keyword] OR accura* [Text Word] OR Outcome Assessment (Health Care) [exp] [MeSH term] OR “Diagnosis-Related Groups” [exp] [MeSH term] OR Diagnosis [exp] [MeSH term] OR Reproducibility of Results [exp] [MeSH term]
#6	4 AND 5
#7	Community [Text Word] OR “free living” [Text Word] OR “independent living” [Text Word] OR “home” [Text Word] OR “general practice” [Text Word] OR “primary health care” [Text Word] OR “primary healthcare” [Text Word] OR “primary nursing” [Text Word] OR “home nursing” [Text Word] OR General Practice [exp] [MeSH term] OR Primary Health Care [exp] [MeSH term] “Primary Nursing” [exp] [MeSH term] OR “Home Care Services [exp] [MeSH term] OR Home Care Agencies [exp] [MeSH term] OR Independent Living [exp] [MeSH term]
#8	(1 OR 6) AND 7
<i>Health Source: Nursing/Academic Edition (via Ebscohost) was searched 2 September 2016 using keywords (all text for #1 keywords, title only for other keywords) and Health Source Subject Terms. Results = 128 records</i>	
#1	PGSGA [keyword] OR SGA [keyword] OR MNA [keyword] OR “patient generated subjective global assessment” [keyword] OR “subjective global assessment” [keyword] OR “mini nutritional assessment” [keyword]
#2	Nutrition* [keyword] OR malnutrition [keyword] OR “nutrition* status” [keyword] OR undernutrition [keyword] OR undernourish* [keyword] OR malnutrition [exp] [subject term] OR nutritional status [exp] [subject term] OR nutrition evaluation [exp] [subject term] OR deficiency diseases [exp] [subject term] OR protein-energy malnutrition [exp] [subject term] OR malnutrition diagnosis [exp] [subject term]
#3	Screen* [keyword] OR medical screening [exp] [subject term]
#4	2 AND 3
#5	Community [keyword] OR “free living” [keyword] OR “independent living” [keyword] OR “home” [keyword] OR general practice [keyword] OR “primary care” [keyword] OR home care services [exp] [subject term] OR Home Nursing [exp] [subject term] OR Independent Living [exp] [subject term] OR family medicine [exp] [subject term] OR primary health care [exp] [subject term]

#6	(1 OR 4) AND 5
EMBASE was searched 2 September 2016 for citations from both Embase and MEDLINE using keywords (abstract and title) and Emtree terms (limits: human studies, adults, middle aged, aged, very elderly). Results = 1,187 records	
#1	PGSGA [keyword] OR SGA [keyword] OR MNA [keyword] OR “patient generated subjective global assessment” [keyword] OR “subjective global assessment” [keyword] OR “mini nutritional assessment” [keyword]
#2	Nutrition* [keyword] OR malnutrition [keyword] OR “nutrition* status” [keyword] OR undernutrition [keyword] OR “nutrition* deficien*” [keyword] OR emaciation [keyword] OR undernourish* [keyword] OR malnutrition [exp] [Emtree term] OR protein deficiency [exp] [Emtree term] OR protein calorie malnutrition [exp] [Emtree term] OR nutritional status [exp] [Emtree term] OR nutritional assessment [exp] [Emtree term]
#3	Screen* [keyword] OR screening [exp] [Emtree term] OR screening test [exp] [Emtree term] OR mass screening [exp] [Emtree term]
#4	2 AND 3
#5	Diagnos* [keyword] OR evaluat* [keyword] OR valid* [keyword] OR compar* [keyword] OR “outcome assessment” [keyword] OR “outcome measure*” [keyword] OR agreement [keyword] OR precision [keyword] OR kappa* [keyword] OR specificit* [keyword] OR sensitiv* [keyword] OR accur* [keyword] OR diagnostic accuracy [exp] [Emtree term] OR diagnostic test [exp] [Emtree term] OR diagnostic test accuracy study [exp] [Emtree term] OR diagnostic value [exp] [Emtree term] OR diagnosis [exp] [Emtree term] OR differential diagnosis [exp] [Emtree term] OR quantitative diagnosis [exp] [Emtree term] OR qualitative diagnosis [exp] [Emtree term] OR validity [exp] [Emtree term] OR concurrent validity [exp] [Emtree term] OR criterion related validity [exp] [Emtree term] OR discriminant validity [exp] [Emtree term] OR external validity [exp] [Emtree term] OR predictive validity [exp] [Emtree term]
#6	4 AND 5
#7	Community [keyword] OR “free living” [keyword] OR “independent living” [keyword] OR “home” [keyword] OR general practice [keyword] OR “primary health care” [keyword] OR “primary care” [keyword] OR ‘independent living’ [exp] [Emtree term] OR ‘community care’ [exp] [Emtree term] OR ‘community living’ [exp] [Emtree term] OR ‘home care’ [exp] [Emtree term] OR ‘home health agency’ [exp] [Emtree term] OR general practice [exp] [Emtree term] OR primary medical care [exp] [Emtree term] OR primary health care [exp] [Emtree term] OR general practitioner [exp] [Emtree term]
#8	(1 OR 6) AND 7
Web of Science was searched 2 September 2016 for the following keywords in topic or title (limits: article, editorial material). Results = 1,377 records	

#1	PGSGA OR SGA OR MNA OR “patient generated subjective global assessment” OR “subjective global assessment” OR “mini nutritional assessment”
#2	Nutrition* OR malnutrition OR “nutrition* status” OR undernutrition OR “nutrition* deficien*” OR emaciation OR undernourish* OR protein deficien*
#3	Screen*
#4	2 AND 3
#5	Diagnos* OR evaluat* OR valid* OR compar* OR “outcome assessment” OR “outcome measure*” OR agreement OR precision OR kappa* OR specificit* OR sensitiv* OR accura*
#6	4 AND 5
#7	Community OR “free living” OR “independent living” OR “home” OR general practice OR “primary care”
#8	(1 OR 6) AND 7
Total	<i>6,412 records</i>