

## SUBSOLUTION THEOREM FOR THE COMPLEX HESSIAN EQUATION

BY NGOC CUONG NGUYEN

**Abstract.** We prove the subsolution theorem for a complex Hessian equation in a smoothly bounded strongly  $m$ -pseudoconvex domain in  $\mathbb{C}^n$ .

**Introduction.** Let  $\Omega$  be a bounded domain in  $\mathbb{C}^n$  with the canonical Kähler form  $\beta = dd^c\|z\|^2$ , where  $d = \partial + \bar{\partial}$ ,  $d^c = i(\bar{\partial} - \partial)$ . For  $1 \leq m \leq n$ , we denote  $\mathbb{C}_{(1,1)}$  the space of  $(1,1)$ -forms with constant coefficients. One defines the positive cone

$$(0.1) \quad \Gamma_m = \{\eta \in \mathbb{C}_{(1,1)} : \eta \wedge \beta^{n-1} \geq 0, \dots, \eta^m \wedge \beta^{n-m} \geq 0\}.$$

A  $C^2$  smooth function  $u$  is called  $m$ -subharmonic in  $\Omega$  if at every point  $z \in \Omega$  the  $(1,1)$ -form associated to its complex Hessian belongs to  $\Gamma_m$ , i.e.

$$(0.2) \quad \sum_{j,k=1}^n \frac{\partial^2 u(z)}{\partial z_j \partial \bar{z}_k} i dz_j \wedge d\bar{z}_k \in \Gamma_m.$$

It was observed by Błocki (see [3]) that one may relax the smoothness condition in definition (0.2) and consider this inequality in the sense of distributions to obtain a class, denoted by  $SH_m(\Omega)$  (see preliminaries). When functions  $u_1, \dots, u_k$ ,  $1 \leq k \leq m$ , are in  $SH_m(\Omega)$  and are locally bounded, one may still define  $dd^c u_1 \wedge dd^c u_2 \wedge \dots \wedge dd^c u_k \wedge \beta^{n-m}$  as a closed positive current of bidegree  $(n-m+k, n-m+k)$ . In particular  $(dd^c u)^m \wedge \beta^{n-m}$  is a positive measure for  $u$  bounded  $m$ -subharmonic. Thus, it is possible to study bounded solutions of the Dirichlet problem with positive Borel measures  $\mu$  in  $\Omega$  and continuous

---

2000 *Mathematics Subject Classification.* 32U05, 32U40, 35J99.

*Key words and phrases.* Hessian equation, subsolution theorem, capacity.

boundary data  $\varphi \in C(\partial\Omega)$ :

$$(0.3) \quad \begin{cases} u \in SH_m(\Omega) \cap L^\infty(\Omega), \\ (dd^c u)^m \wedge \beta^{n-m} = d\mu, \\ u(z) = \varphi(z) \quad \text{on } \partial\Omega. \end{cases}$$

The Dirichlet problem for complex Hessian equation (0.3) in smooth cases was first considered by S.Y. Li (see [15]). His main result says that if  $\Omega$  is smoothly bounded and strongly  $m$ -pseudoconvex (see Definition 1.5) then, for a smooth boundary data and for a smooth positive measure, i.e.  $d\mu = f\beta^n$  and  $f > 0$  smooth, there exists a unique smooth solution of the Dirichlet problem for the Hessian equation.

The weak solutions of the equation (0.3), when the measure  $d\mu$  is possibly degenerate, were first considered by Błocki [3]. More precisely, he proved that there exists a unique continuous solution of the homogeneous Dirichlet problem in the unit ball in  $\mathbb{C}^n$ .

Very recently, in [10] Dinew and Kołodziej investigated weak solutions of complex Hessian equations (0.3) with the right hand side more general, namely  $d\mu = f\beta^n$  where  $f \in L^p$ , for  $p > n/m$ . One of their results extended Li's theorem: they proved that the Dirichlet problem still has a unique continuous solution with continuous boundary data and  $d\mu$  in  $L^p$  as above. Their method exploited the new counterpart of pluripotential theory for  $m$ -subharmonic functions, after showing a crucial inequality between the usual volume and  $m$ -capacity which is a version of the relative capacity for  $m$ -subharmonic functions.

In the case  $m = n$ , the subsolution theorem due to Kołodziej [13] (see [14] for a simpler proof) says that Dirichlet problem (0.3) in a strongly pseudoconvex domain is solvable if there is a subsolution. Thus, one may ask the same question when  $m < n$ . In this note we show that the subsolution theorem, Theorem 2.2, for the complex Hessian equation is still true by combining Dinew and Kołodziej's new results for weak solutions of the complex Hessian equations and the method used to prove the subsolution theorem in the pluripotential case.

**Acknowledgements.** I am indebted to my advisor, professor Sławomir Kołodziej, for suggesting the problem and for many stimulating discussions. I would also like to thank the referee whose suggestions and remarks helped me to improve the exposition of the paper. This work is supported by the International Ph.D. Programme *Geometry and Topology in Physical Models*.

## 1. Preliminaries.

1.1. *m-subharmonic functions.* We recall basic notions and results which are adapted from pluripotential theory. The main sources are [1, 2, 5–8, 14] for plurisubharmonic functions and [3, 10] for  $m$ -subharmonic functions. Since a major part of pluripotential theory can be easily adapted to  $m$ -subharmonic case, when the proof is only a copy of the original one with obvious changes of notations, for the proofs we refer the reader to the above references. Let  $\mathbb{C}_{(k,k)}$  be the space of  $(k, k)$ -forms with constant coefficients, and

$$\Gamma_m = \{\eta \in \mathbb{C}_{(1,1)} : \eta \wedge \beta^{n-1} \geq 0, \dots, \eta^m \wedge \beta^{n-m} \geq 0\}.$$

We denote by  $\Gamma_m^*$  its dual cone

$$(1.1) \quad \Gamma_m^* = \{\gamma \in \mathbb{C}_{(n-1,n-1)} : \gamma \wedge \eta \geq 0 \text{ for every } \eta \in \Gamma_m\}.$$

By Proposition 2.1 in [3] we know that  $\{\eta_1 \wedge \dots \wedge \eta_{m-1} \wedge \beta^{n-m}; \eta_1, \dots, \eta_{m-1} \in \Gamma_m\} \subset \Gamma_m^*$ . Moreover if we consider  $\Gamma_m^{**} = \{\eta \in \mathbb{C}_{(1,1)} : \eta \wedge \gamma \geq 0 \text{ for every } \gamma \in \Gamma_m^*\}$  then we have

$$\Gamma_m = \Gamma_m^{**}$$

as  $\{\eta_1 \wedge \dots \wedge \eta_{m-1} \wedge \beta^{n-m}; \eta_1, \dots, \eta_{m-1} \in \Gamma_m\}^* \subset \Gamma_m$ . Therefore

$$(1.2) \quad \Gamma_m^* = \{\eta_1 \wedge \dots \wedge \eta_{m-1} \wedge \beta^{n-m}; \eta_1, \dots, \eta_{m-1} \in \Gamma_m\}.$$

Since  $\Gamma_n \subset \Gamma_{n-1} \subset \dots \subset \Gamma_1$ , we thus obtain

$$\Gamma_n^* \supset \Gamma_{n-1}^* \supset \dots \supset \Gamma_1^* = \{t\beta^{n-1}; t \geq 0\}.$$

In particular, when  $\eta \in \Gamma_m^*$ , and it has a representation

$$\sum a^{j\bar{k}} i^{(n-1)^2} \hat{d}z_j \wedge \hat{d}\bar{z}_k$$

(this notation means that in the  $(n-1, n-1)$ -form only  $dz_j$  and  $d\bar{z}_k$  vanish in the complete form  $dz \wedge d\bar{z}$  at positions  $j$ -th and  $k$ -th) then the Hermitian matrix  $(a^{j\bar{k}})$  is nonnegative definite. In the language of differential forms, a  $C^2$  smooth function  $u$  is  $m$ -subharmonic ( $m$ -sh for short) if

$$dd^c u \wedge \beta^{n-1} \geq 0, \dots, (dd^c u)^m \wedge \beta^{n-m} \geq 0 \text{ at every point in } \Omega.$$

DEFINITION 1.1. Let  $u$  be a subharmonic function on an open subset  $\Omega \subset \mathbb{C}^n$ . Then  $u$  is called  $m$ -subharmonic if for any collection of  $\eta_1, \dots, \eta_{m-1}$  in  $\Gamma_m$ , the inequality

$$dd^c u \wedge \eta_1 \wedge \dots \wedge \eta_{m-1} \wedge \beta^{n-m} \geq 0$$

holds in the sense of currents. Let  $SH_m(\Omega)$  stands for the set of all  $m$ -sh functions in  $\Omega$ .

REMARK 1.2. **(a)** Condition (1.1) is equivalent to  $dd^c u \wedge \eta \geq 0$  for every  $\eta \in \Gamma_m^*$  by (1.2). Hence, a subharmonic function  $u$  is  $m$ -subharmonic if

$$(1.3) \quad \int_{\Omega} u dd^c \phi \wedge \eta = \int_{\Omega} u \sum_{j,k=1}^n a^{j\bar{k}} \frac{\partial^2 \phi}{\partial z_j \partial \bar{z}_k} \beta^n \geq 0$$

for every non-negative test function  $0 \leq \phi$  in  $\Omega$  and for every nonnegative definite Hermitian matrix  $(a^{j\bar{k}})$  with constant coefficients such that  $\eta = \sum_{j,k=1}^n a^{j\bar{k}} i^{(n-1)^2} dz_1 \wedge \dots \wedge \hat{d}z_j \wedge \dots \wedge dz_n \wedge d\bar{z}_1 \wedge \dots \wedge \hat{d}\bar{z}_k \wedge \dots \wedge d\bar{z}_n$  belongs to  $\Gamma_m^*$ . This means that  $u$  is subharmonic with respect to a family of elliptic operators with constant coefficients.

**(b)** A  $C^2$  function  $v$  is  $m$ -subharmonic iff  $dd^c v(z)$  belongs to  $\Gamma_m$  at every  $z \in \Omega$ . Hence

$$dd^c u \wedge dd^c v_1 \wedge \dots \wedge dd^c v_{m-1} \wedge \beta^{n-m} \geq 0$$

holds in  $\Omega$  in the weak sense of currents, for every collection  $v_1, \dots, v_{m-1} \in SH_m \cap C^2(\Omega)$  and any  $u \in SH_m(\Omega)$ .

PROPOSITION 1.3. *Let  $\Omega \subset \mathbb{C}^n$  be a bounded open subset. Then*

1.  $PSH(\Omega) = SH_n(\Omega) \subset SH_{n-1}(\Omega) \subset \dots \subset SH_1(\Omega) = SH(\Omega)$ .
2.  $SH_m(\Omega)$  is a convex cone.
3. *The limit of a decreasing sequence in  $SH_m(\Omega)$  belongs to  $SH_m(\Omega)$ . Moreover, the standard regularization  $u * \rho_\varepsilon$  of a  $m$ -sh function is again a  $m$ -sh function. Here  $\rho_\varepsilon(z) = \frac{1}{\varepsilon^{2n}} \rho(\frac{z}{\varepsilon})$ ,  $\rho(z) = \rho(\|z\|^2)$  is a smoothing kernel, with  $\rho : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  defined by*

$$\rho(t) = \begin{cases} \frac{C}{(1-t)^2} \exp(\frac{1}{t-1}) & \text{if } 0 \leq t \leq 1, \\ 0 & \text{if } t > 1, \end{cases}$$

for a constant  $C$  such that

$$\int_{\mathbb{C}^n} \rho(\|z\|^2) \beta^n = 1.$$

4. *If  $u \in SH_m(\Omega)$  and  $\gamma : \mathbb{R} \rightarrow \mathbb{R}$  is a convex, nondecreasing function then  $\gamma \circ u \in SH_m(\Omega)$ .*
5. *If  $u, v \in SH_m(\Omega)$  then  $\max\{u, v\} \in SH_m(\Omega)$ .*
6. *Let  $\{u_\alpha\} \subset SH_m(\Omega)$  be a locally uniformly bounded from above and  $u = \sup u_\alpha$ . Then the upper semi-continuous regularization  $u^*$  is  $m$ -sh and is equal to  $u$  almost everywhere.*

PROOF. (1) and (2) and the first part of (3) easily follow from the definition of  $m$ -sh functions. From formula (1.3), for  $\eta \in \Gamma_m^*$ , there follows:

$$\int (u * \rho_\varepsilon) dd^c \phi \wedge \eta = \int u dd^c(\phi * \rho_\varepsilon) \wedge \eta \geq 0,$$

since  $\phi * \rho_\varepsilon$  is again a nonnegative test function. Thus (3) is proved. For (4), the smooth function  $\gamma * \rho_\varepsilon$  (the standard regularization on  $\mathbb{R}$ ) is convex and increasing, therefore  $(\gamma * \rho_\varepsilon) \circ u \in SH_m(\Omega)$ . Since  $(\gamma * \rho_\varepsilon) \circ u$  decreases to  $\gamma \circ u$  as  $\varepsilon \rightarrow 0$ , applying the first part of (3) we have  $\gamma \circ u \in SH_m(\Omega)$ . In order to prove (5), note that by using (3) it is enough to show that  $w = \max\{u_\varepsilon, v_\varepsilon\}$  is  $m$ -sh, where  $u_\varepsilon := u * \rho_\varepsilon, v_\varepsilon := v * \rho_\varepsilon$ . Since  $w$  is semi-convex, i.e. there is a constant  $C = C_\varepsilon > 0$  large enough for  $w + C\|z\|^2 = \max\{u_\varepsilon + C\|z\|^2, v_\varepsilon + C\|z\|^2\}$  to be a convex function in  $\mathbb{R}^{2n}$ , hence it has the second derivative almost everywhere and  $dd^c w(x) \in \Gamma_m$  for almost every  $x$  in  $\Omega$ . Let  $w_\varepsilon$  be a regularization of  $w$ , by the formula of the convolution  $w_\varepsilon(x) = \int_\Omega w(x - \varepsilon y) \rho(y) \beta^n(y)$  we have

$$dd^c w_\varepsilon(x) = \int_\Omega dd^c w(x - \varepsilon y) \rho(y) \beta^n(y).$$

Thus, for  $\eta \in \Gamma_m^*$

$$dd^c w_\varepsilon(x) \wedge \eta = \int_\Omega [dd^c w(x - \varepsilon y) \wedge \eta] \rho(y) \beta^n(y) \geq 0.$$

(6) is a consequence of (5) and Choquet's Lemma.  $\square$

1.2. *The complex Hessian operator.* For  $1 \leq k \leq m, u_1, \dots, u_k \in SH_m \cap L_{loc}^\infty(\Omega)$  the operator  $dd^c u_k \wedge dd^c u_{k-1} \wedge \dots \wedge dd^c u_1 \wedge \beta^{n-m}$  is defined inductively by (see [3], [10])

$$(H_k) \quad dd^c u_k \wedge dd^c u_{k-1} \wedge \dots \wedge dd^c u_1 \wedge \beta^{n-m} := dd^c(u_k dd^c u_{k-1} \wedge \dots \wedge dd^c u_1 \wedge \beta^{n-m})$$

which is a closed positive current of bidegree  $(n - m + k, n - m + k)$ . This operator is also continuous under decreasing sequences and symmetric (see Remark 1.10). In the case  $k = m, dd^c u_1 \wedge dd^c u_2 \wedge \dots \wedge dd^c u_m \wedge \beta^{n-m}$  is a nonnegative Borel measure, in particular, when  $u = u_1 = \dots = u_m$  currents (measures)  $(dd^c u)^m \wedge \beta^{n-m}$  are well-defined for  $u \in L_{loc}^\infty(\Omega)$ . The above definitions essentially follow from the analogous definitions due to Bedford and Taylor [1, 2] for plurisubharmonic functions.

PROPOSITION 1.4 (Chern–Levine–Nirenberg inequalities). *Let  $K \subset\subset U \subset\subset \Omega$ , where  $K$  is compact,  $U$  is open. Let  $u_1, \dots, u_k \in SH_m \cap L^\infty(\Omega), 1 \leq k \leq m$  and  $v \in SH_m(\Omega)$ . Then there exists a constant  $C = C_{K,U,\Omega} \geq 0$  such that*

- (i)  $\|dd^c u_1 \wedge \dots \wedge dd^c u_k \wedge \beta^{n-m}\|_K \leq C \|u_1\|_{L^\infty(U)} \dots \|u_k\|_{L^\infty(U)},$
- (ii)  $\|dd^c u_1 \wedge \dots \wedge dd^c u_k \wedge \beta^{n-m}\|_K \leq C \|u_1\|_{L^1(\Omega)} \cdot \|u_2\|_{L^\infty(\Omega)} \dots \|u_k\|_{L^\infty(\Omega)},$
- (iii)  $\|v dd^c u_1 \wedge \dots \wedge dd^c u_k \wedge \beta^{n-m}\|_K \leq C \|v\|_{L^1(\Omega)} \cdot \|u_1\|_{L^\infty(\Omega)} \dots \|u_k\|_{L^\infty(\Omega)}.$

PROOF. (i) By induction we only need to prove that

$$\|dd^c u_1 \wedge \dots \wedge dd^c u_k \wedge \beta^{n-m}\|_K \leq C \|u_1\|_{L^\infty(U)} \|dd^c u_2 \wedge \dots \wedge dd^c u_k \wedge \beta^{n-m}\|_U.$$

In fact, let  $\chi \geq 0$  be a test function equal to 1 on  $K$ . Then an integration by parts yields

$$\begin{aligned} \|dd^c u_1 \wedge \dots \wedge dd^c u_k \wedge \beta^{n-m}\|_K &\leq C \int_U \chi dd^c u_1 \wedge \dots \wedge dd^c u_k \wedge \beta^{n-k} \\ &= C \int_U u_1 dd^c \chi \wedge \dots \wedge dd^c u_k \wedge \beta^{n-k}. \end{aligned}$$

Thus,

$$\|dd^c u_1 \wedge \dots \wedge dd^c u_k \wedge \beta^{n-m}\|_K \leq C' \|u_1\|_{L^\infty(U)} \|dd^c u_2 \wedge \dots \wedge dd^c u_k \wedge \beta^{n-m}\|_U,$$

where  $C'$  depends only on bounds of coefficients of  $dd^c \chi$  and on the set  $U$ .

(ii) It is a simple consequence of (i), and the result  $\|dd^c w \wedge \beta^{n-1}\|_K \leq C_{K,U} \|w\|_{L^1(U)}$  for every  $w \in SH_m(\Omega)$  (see [8], Remark 3.4).

(iii) See [8] Proposition 3.11.  $\square$

1.3. *m-pseudoconvex domains.* Let  $\Omega$  be a bounded domain with  $\partial\Omega$  in the class  $C^2$ . Let  $\rho \in C^2$  in a neighborhood of  $\bar{\Omega}$  be a defining function of  $\Omega$ , i.e. a function such that

$$\rho < 0 \text{ on } \Omega, \quad \rho = 0 \text{ and } d\rho \neq 0 \text{ on } \partial\Omega.$$

DEFINITION 1.5. A  $C^2$  bounded domain is called strongly  $m$ -pseudoconvex if there are a defining function  $\rho$  and some  $\varepsilon > 0$  such that  $(dd^c \rho)^k \wedge \beta^{n-k} \geq \varepsilon \beta^n$  in  $\bar{\Omega}$  for every  $1 \leq k \leq m$ .

It is obvious that a strongly pseudoconvex domain is a strongly  $m$ -pseudoconvex domain. The properties of strongly  $m$ -pseudoconvex domains are similar to those of strongly pseudoconvex domains, e.g. it can be shown that strongly  $m$ -pseudoconvexity is characterized by a condition on its boundary (see [15], Theorem 3.1). We also have the criterion that if the Levi form of  $\Omega$  corresponding to  $\rho$  belongs to the interior of  $\Gamma_{m-1}$  then  $\Omega$  is strongly  $m$ -pseudoconvex (see [15], Proposition 3.3).

1.4. *Cegrell's inequalities for the complex Hessian operator.* It is sufficient for our purpose in this section to work within the class of  $m$ -sh functions which are continuous up to the boundary and equal to 0 on the boundary. Let  $\Omega$  be a strongly  $m$ -pseudoconvex domain in  $\mathbb{C}^n$ . We denote

$$\mathcal{E}_0(m) = \{u \in SH_m(\Omega) \cap C(\bar{\Omega}); u|_{\partial\Omega} = 0, \int_{\Omega} (dd^c u)^m \wedge \beta^{n-m} < +\infty\}.$$

For the case  $m = n$ , this class was introduced by Cegrell in [5]. It is a convex cone for  $1 \leq m \leq n$  (see [5], p. 188). Our goal is to establish inequalities very similar to the one due to Cegrell (see [6], Lemma 5.4, Theorem 5.5) for the Monge–Ampère operator. In order to avoid confusions and trivial statements, we only consider  $2 \leq m \leq n - 1$ .

PROPOSITION 1.6. *Let  $u, v, h \in \mathcal{E}_0(m)$ , and  $1 \leq p, q \leq m$ ,  $p + q \leq m$ . Set  $T = -hS$  where  $S = dd^c h_1 \wedge \dots \wedge dd^c h_{m-p-q} \wedge \beta^{n-m}$  with  $h_1, \dots, h_{m-p-q}$  are also in  $\mathcal{E}_0(m)$ , then*

$$\int_{\Omega} (dd^c u)^p \wedge (dd^c v)^q \wedge T \leq \left[ \int_{\Omega} (dd^c u)^{p+q} \wedge T \right]^{\frac{p}{p+q}} \left[ \int_{\Omega} (dd^c v)^{p+q} \wedge T \right]^{\frac{q}{p+q}}.$$

PROOF. See Lemma 5.4 in [6]. We only here remark that both sides of the inequality are finite because of the convexity of the cone  $\mathcal{E}_0(m)$ .  $\square$

REMARK 1.7. The statement in Proposition 1.5 is still true when  $h \in SH_m \cap L^\infty(\Omega)$ ,  $\lim_{\zeta \rightarrow \partial\Omega} h(\zeta) = 0$  and  $\int_{\Omega} (dd^c h)^m \wedge \beta^{n-m} < +\infty$  since the integration by parts formula is valid as in the case of the continuous case (see [6], Corollary 3.4).

Applying Proposition 1.6 for some special cases of  $m$ -sh functions in  $\mathcal{E}_0(m)$ , we obtain

COROLLARY 1.8. *If  $u, v, h \in \mathcal{E}_0(m)$ ,  $1 \leq p \leq m - 1$ , then*

(i)

$$\begin{aligned} & \int_{\Omega} -h(dd^c u)^p \wedge (dd^c v)^{m-p} \wedge \beta^{n-m} \\ & \leq \left[ \int_{\Omega} -h(dd^c u)^m \wedge \beta^{n-m} \right]^{\frac{p}{m}} \left[ \int_{\Omega} -h(dd^c v)^m \wedge \beta^{n-m} \right]^{\frac{m-p}{m}}, \end{aligned}$$

(ii)

$$\begin{aligned} & \int_{\Omega} (dd^c u)^p \wedge (dd^c v)^{m-p} \wedge \beta^{n-m} \\ & \leq \left[ \int_{\Omega} (dd^c u)^m \wedge \beta^{n-m} \right]^{\frac{p}{m}} \left[ \int_{\Omega} (dd^c v)^m \wedge \beta^{n-m} \right]^{\frac{m-p}{m}}. \end{aligned}$$

PROOF. (i) follows from Proposition 1.6 when  $u = u_1 = \dots = u_p$ ,  $v = v_1 = \dots = v_q$ . (ii) comes from the fact that for  $\rho$  a defining function of  $\Omega$  we have

$$\int_{\Omega} (dd^c u)^p \wedge (dd^c v)^{m-p} \wedge \beta^{n-m} = \lim_{\varepsilon \rightarrow 0} \int_{\{\rho < -\varepsilon\}} (dd^c u)^p \wedge (dd^c v)^{m-p} \wedge \beta^{n-m},$$

and

$$\begin{aligned}
& \int_{U_\varepsilon} (dd^c u)^p \wedge (dd^c v)^{m-p} \wedge \beta^{n-m} \\
& \leq \int_{\Omega} -h_{U_\varepsilon, \Omega}^* (dd^c u)^p \wedge (dd^c v)^{m-p} \wedge \beta^{n-m} \\
& \leq \left[ \int_{\Omega} -h_{U_\varepsilon, \Omega}^* (dd^c u)^m \wedge \beta^{n-m} \right]^{\frac{p}{m}} \left[ \int_{\Omega} -h_{U_\varepsilon, \Omega}^* (dd^c v)^m \wedge \beta^{n-m} \right]^{\frac{m-p}{m}} \\
& \leq \left[ \int_{\Omega} (dd^c u)^m \wedge \beta^{n-m} \right]^{\frac{p}{m}} \left[ \int_{\Omega} (dd^c v)^m \wedge \beta^{n-m} \right]^{\frac{m-p}{m}},
\end{aligned}$$

where  $U_\varepsilon = \{\rho < -\varepsilon\}$  and  $h_{U_\varepsilon, \Omega} = \sup\{u \in SH_m(\Omega); u \leq 0; u|_{U_\varepsilon} \leq -1\}$ . It is clear that  $-1 \leq h_{U_\varepsilon, \Omega}^* \leq 0$ ,  $\lim_{\zeta \rightarrow \partial\Omega} h_{U_\varepsilon, \Omega}^*(\zeta) = 0$  and  $\int_{\Omega} (dd^c h_{U_\varepsilon, \Omega}^*)^m \wedge \beta^{n-m} < +\infty$ . Hence inequality (i) is still applicable by Remark 1.7.  $\square$

1.5. *m-capacity, convergence theorems, the comparison principle.* For  $E$  a Borel set in  $\Omega$  we define

$$cap_m(E, \Omega) = \sup\left\{ \int_E (dd^c u)^m \wedge \beta^{n-m}, u \in SH_m(\Omega), 0 \leq u \leq 1 \right\}.$$

In view of Proposition 1.4, it is finite if  $E$  is relatively compact in  $\Omega$ . This is the version of the relative capacity in the case of  $m$ -subharmonic functions. It is an useful tool to establish convergence properties, especially the comparison principle.

**THEOREM 1.9** (Convergence theorem). *Let  $\{u_k^j\}_{j=1}^\infty$ ,  $k = 1, \dots, m$  be locally uniformly bounded sequences of  $m$ -subharmonic functions in  $\Omega$ ,  $u_k^j \rightarrow u_k \in SH_m \cap L^\infty(\Omega)$  in  $cap_m$  as  $j \rightarrow \infty$ . Then*

$$\lim_{j \rightarrow \infty} dd^c u_1^j \wedge \dots \wedge dd^c u_m^j \wedge \beta^{n-m} = dd^c u_1 \wedge \dots \wedge dd^c u_m \wedge \beta^{n-m}$$

*in the topology of currents.*

**PROOF.** See the proof of Theorem 1.11 in [14].  $\square$

**REMARK 1.10.** As in Theorem 2.1 of [2], one may prove that for  $1 \leq k \leq m$ , if  $u_1^j, \dots, u_k^j$  are decreasing sequences of locally bounded  $m$ -sh functions such that  $\lim_{j \rightarrow \infty} u_l^j(z) = u_l(z) \in SH_m \cap L_{loc}^\infty(\Omega)$  for all  $z \in \Omega$  and  $1 \leq l \leq k$ , then

$$\lim_{j \rightarrow \infty} dd^c u_1^j \wedge \dots \wedge dd^c u_k^j \wedge \beta^{n-m} = dd^c u_1 \wedge \dots \wedge dd^c u_k \wedge \beta^{n-m}$$

in the sense of currents. Thus, the currents obtained in inductive definition  $(H_k)$  of the wedge product of currents associated to locally bounded  $m$ -sh functions are closed positive currents.



PROPOSITION 1.11. *If  $u_j \in SH_m \cap L^\infty(\Omega)$  is a sequence decreasing to a bounded function  $u$  in  $\Omega$  then it converges to  $u \in SH_m \cap L^\infty(\Omega)$  with respect to  $cap_m$ . In particular, Theorem 1.9 holds in this case.*

PROOF. See Proposition 1.12 in [14].  $\square$

THEOREM 1.12 (Quasi-continuity). *For an  $m$ -subharmonic function  $u$  defined in  $\Omega$  and for each  $\varepsilon > 0$ , there is an open subset  $U$  such that  $cap_m(U, \Omega) < \varepsilon$  and  $u$  is continuous in  $\Omega \setminus U$ .*

PROOF. See Theorem 1.13 in [14].  $\square$

From the quasi-continuity of  $m$ -subharmonic functions one can derive several important results.

THEOREM 1.13. *Let  $u, v$  be locally bounded  $m$ -sh functions on  $\Omega$ . Then the following inequality holds:*

$$(dd^c \max\{u, v\})^m \wedge \beta^{n-m} \geq \mathbf{1}_{\{u \geq v\}} (dd^c u)^m \wedge \beta^{n-m} + \mathbf{1}_{\{u < v\}} (dd^c v)^m \wedge \beta^{n-m}.$$

PROOF. See Theorem 6.11 in [7].  $\square$

THEOREM 1.14 (Comparison principle). *Let  $\Omega$  be an open bounded subset of  $\mathbb{C}^n$ . For  $u, v \in SH_m \cap L^\infty(\Omega)$  satisfying  $\liminf_{\zeta \rightarrow z} (u - v)(\zeta) \geq 0$  for any  $z \in \partial\Omega$ , we have*

$$\int_{\{u < v\}} (dd^c v)^m \wedge \beta^{n-m} \leq \int_{\{u < v\}} (dd^c u)^m \wedge \beta^{n-m}.$$

PROOF. The proof follows the lines of the proof of Theorem 1.16 in [14]. First consider  $u, v \in C^\infty(\Omega)$ ,  $E = \{u < v\} \subset\subset \Omega$ , and smooth  $\partial\Omega$ . In this case, put  $u_\varepsilon = \max\{u + \varepsilon, v\}$  and use Stokes' theorem to get

$$(1.4) \quad \begin{aligned} \int_E (dd^c u_\varepsilon)^m \wedge \beta^{n-m} &= \int_{\partial E} d^c u_\varepsilon \wedge (dd^c u_\varepsilon)^{m-1} \wedge \beta^{n-m} \\ &= \int_{\partial E} d^c u \wedge (dd^c u)^{m-1} \wedge \beta^{n-m} = \int_E (dd^c u)^m \wedge \beta^{n-m} \end{aligned}$$

(since  $u_\varepsilon = u + \varepsilon$  in a neighborhood of  $\partial E$ ). By Theorem 1.9,  $(dd^c u_\varepsilon)^m \wedge \beta^{n-m}$  converges weakly\* to  $(dd^c v)^m \wedge \beta^{n-m}$  as  $\varepsilon \rightarrow 0$  on the open set  $E$ , which implies that

$$\int_E (dd^c v)^m \wedge \beta^{n-m} \leq \liminf_{\varepsilon \rightarrow \infty} \int_E (dd^c u_\varepsilon)^m \wedge \beta^{n-m}.$$

This combined with (1.4) implies the statement.

For the general case, suppose  $\|u\|, \|v\| < 1$ , fix  $\varepsilon > 0$  and  $\delta > 0$ . By the quasi-continuity, there is an open set  $U$  such that  $cap_m(U, \Omega) < \varepsilon$  and  $u = \tilde{u}$ ,  $v = \tilde{v}$  on  $\Omega \setminus U$  for some continuous functions  $\tilde{u}, \tilde{v}$  in  $\Omega$ . Let  $u_k, v_k$  be the standard regularizations of  $u$  and  $v$ . By Dini's theorem  $u_k$  and  $v_k$  uniformly

converge to  $u$  and to  $v$ , respectively, on  $\Omega \setminus U$ . Then for  $k > k_0$  big enough, subsets  $E(\delta) := \{\tilde{u} + \delta < \tilde{v}\}$  and  $E_k(\delta) := \{u_k + \delta < v_k\}$  satisfy

$$(1.5) \quad E(2\delta) \setminus U \subset \subset \bigcap_k E_k(\delta) \setminus U \quad \text{and} \quad \bigcup_k E_k(\delta) \setminus U \subset \subset \{\tilde{u} < \tilde{v}\}.$$

In what follows we shall often use the estimate

$$\int_U (dd^c w)^m \wedge \beta^{n-m} \leq \text{cap}_m(U, \Omega) < \varepsilon \quad \text{where} \quad 0 \leq w \leq 1,$$

not mentioning this any more. Since  $\{u + 2\delta < v\} = \{\tilde{u} + 2\delta < \tilde{v}\}$  on  $\Omega \setminus U$ ,

$$(1.6) \quad \int_{\{u+2\delta < v\}} (dd^c v)^m \wedge \beta^{n-m} \leq \int_{\{\tilde{u}+2\delta < \tilde{v}\} \setminus U} (dd^c v)^m \wedge \beta^{n-m} + \varepsilon \\ = \int_{E(2\delta) \setminus U} (dd^c v)^m \wedge \beta^{n-m} + \varepsilon.$$

Since  $(dd^c v_k)^m \wedge \beta^{n-m}$  weakly\* converges to  $(dd^c v)^m \wedge \beta^{n-m}$  and  $E(2\delta)$  is open, by (1.5) we get

$$(1.7) \quad \int_{E(2\delta)} (dd^c v)^m \wedge \beta^{n-m} \leq \liminf_{k \rightarrow \infty} \int_{E(2\delta)} (dd^c v_k)^m \wedge \beta^{n-m} \\ \leq \liminf_{k \rightarrow \infty} \int_{E_k(\delta)} (dd^c v_k)^m \wedge \beta^{n-m} + \varepsilon.$$

Now, by Sard's theorem, we may assume that  $E_k(\delta)$  has smooth boundary (changing  $\delta$  if needed), thus using the argument of the smooth case we have

$$(1.8) \quad \int_{E_k(\delta)} (dd^c v_k)^m \wedge \beta^{n-m} \leq \int_{E_k(\delta)} (dd^c u_k)^m \wedge \beta^{n-m}.$$

Therefore, by (1.6), (1.7) and (1.8), we have

$$(1.9) \quad \int_{\{u+2\delta < v\}} (dd^c v)^m \wedge \beta^{n-m} \leq \liminf_{k \rightarrow \infty} \int_{E_k(\delta)} (dd^c u_k)^m \wedge \beta^{n-m} + 2\varepsilon.$$

Furthermore, using (1.5) and the fact that  $(dd^c u_k)^m \wedge \beta^{n-m}$  weakly\* converges to  $(dd^c u)^m \wedge \beta^{n-m}$  we obtain

$$(1.10) \quad \limsup_{k \rightarrow \infty} \int_{\bigcup_k E_k(\delta) \setminus U} (dd^c u_k)^m \wedge \beta^{n-m} \leq \int_{\bigcup_k E_k(\delta) \setminus U} (dd^c u)^m \wedge \beta^{n-m}.$$

Thus, from (1.5), (1.9) and (1.10) there follows

$$(1.11) \quad \int_{\{u+2\delta < v\}} (dd^c v)^m \wedge \beta^{n-m} \leq \int_{\{\tilde{u} < \tilde{v}\}} (dd^c u)^m \wedge \beta^{n-m} + 3\varepsilon \\ \leq \int_{\{u < v\}} (dd^c u)^m \wedge \beta^{n-m} + 4\varepsilon.$$

Finally, letting  $\delta$  and  $\varepsilon$  tend to 0 in (1.11) the statement is proved.  $\square$

COROLLARY 1.15. *Under the assumption of Theorem 1.14 we have*

- (a) *If  $(dd^c u)^m \wedge \beta^{n-m} \leq (dd^c v)^m \wedge \beta^{n-m}$  then  $v \leq u$ ,*
- (b) *If  $(dd^c u)^m \wedge \beta^{n-m} = (dd^c v)^m \wedge \beta^{n-m}$  and  $\lim_{\zeta \rightarrow z} (u - v)(\zeta) = 0$  for  $z \in \partial\Omega$  then  $u = v$ ,*
- (c) *If  $\lim_{\zeta \rightarrow \partial\Omega} u(\zeta) = \lim_{\zeta \rightarrow \partial\Omega} v(\zeta) = 0$  and  $u \leq v$  in  $\Omega$ , then*

$$\int_{\Omega} (dd^c v)^m \wedge \beta^{n-m} \leq \int_{\Omega} (dd^c u)^m \wedge \beta^{n-m}.$$

PROOF. For (a) and (b) see Corollary 1.17 in [14]. For (c), let  $\varepsilon > 0$ . Applying Theorem 1.14 we have

$$\int_{\Omega} (dd^c v)^m \wedge \beta^{n-m} \leq (1 + \varepsilon)^n \int_{\Omega} (dd^c u)^m \wedge \beta^{n-m}.$$

Then, letting  $\varepsilon \rightarrow 0$ , we obtain the result.  $\square$

**2. Subsolution theorem.** In this section we will prove our main theorem. The method we use here is similar to the one from the proof of the plurisubharmonic case (see [14], Theorem 4.7). We first recall the theorem due to Dinew and Kołodziej about the weak solution of the complex Hessian equation with the right hand side in  $L^p$  (see [10], Theorem 2.10). From now on we only consider  $1 < m < n$ .

THEOREM 2.1 ([10]). *Let  $\Omega$  be a smoothly strongly  $m$ -pseudoconvex domain. Then for  $p > n/m$ ,  $f \in L^p(\Omega)$  and a continuous function  $\varphi$  on  $\partial\Omega$  there exists  $u \in SH_m(\Omega) \cap C(\bar{\Omega})$  satisfying*

$$(dd^c u)^m \wedge \beta^{n-m} = f \beta^n,$$

and  $u = \varphi$  on  $\partial\Omega$ .

Let us state the subsolution theorem.

THEOREM 2.2. *Let  $\Omega$  be a smoothly strongly  $m$ -pseudoconvex domain in  $\mathbb{C}^n$ , and let  $\mu$  be a finite positive Borel measure in  $\Omega$ . If there is a subsolution  $v$ , i.e.*

$$(2.1) \quad \begin{cases} v \in SH_m \cap L^\infty(\Omega), \\ (dd^c v)^m \wedge \beta^{n-m} \geq d\mu, \\ \lim_{\zeta \rightarrow z} v(\zeta) = \varphi(z) \text{ for any } z \in \partial\Omega, \end{cases}$$

then there is a solution  $u$  of the following Dirichlet problem

$$(2.2) \quad \begin{cases} u \in SH_m \cap L^\infty(\Omega), \\ (dd^c u)^m \wedge \beta^{n-m} = d\mu, \\ \lim_{\zeta \rightarrow z} u(\zeta) = \varphi(z) \text{ for any } z \in \partial\Omega. \end{cases}$$

PROOF. We first prove Theorem 2.1 under two additional assumptions:

- 1) the measure  $\mu$  has compact support in  $\Omega$ ;
- 2) the function  $\varphi$  is in the class  $C^2$ .

Using the first of those conditions we can modify  $v$  so that  $v$  is  $m$ -subharmonic in a neighborhood of  $\Omega$  (and still is a subsolution). To do this take an open subset  $U$  such that  $\text{supp } \mu \subset\subset U \subset\subset \Omega$  and consider the envelope

$$\hat{v} = \sup\{w \in SH_m(\Omega) : w \leq 0, w \leq v \text{ on } U\}.$$

Then from Proposition 1.3-(6)  $\hat{v}^*$  is a competitor in the definition of the envelope, hence  $\hat{v} = \hat{v}^* \in SH_m(\Omega)$ . The balayage procedure implies that  $\hat{v} = v$  on  $U$  and  $\lim_{\zeta \rightarrow z} \hat{v}(\zeta) = 0$  for any  $z \in \partial\Omega$  (the balayage still works as in the case of plurisubharmonic functions by results in [3], Theorem 1.2, Theorem 3.7). Thus,  $(dd^c \hat{v})^m \wedge \beta^{n-m} \geq d\mu$  as  $\text{supp } \mu \subset\subset U$ . Next, take  $\rho$  a defining function of  $\Omega$  which is smooth on a neighborhood  $\Omega_1$  of  $\bar{\Omega}$  and  $(dd^c \rho)^k \wedge \beta^{n-k} \geq \varepsilon \beta^n$ ,  $1 \leq k \leq m$ , in  $\bar{\Omega}$  for some  $\varepsilon > 0$ . Since  $\hat{v}$  is bounded, we can further choose  $\rho$  satisfying  $\rho \leq \hat{v}$  on  $\bar{U}$ . Put

$$v_1(z) := \begin{cases} \max\{\rho(z), \hat{v}(z)\} & \text{on } \bar{\Omega}, \\ \rho(z) & \text{on } \Omega_1 \setminus \bar{\Omega}. \end{cases}$$

Hence  $v_1$  is a subsolution which is defined and  $m$ -subharmonic in a neighborhood of  $\bar{\Omega}$ . We still write  $v$  instead of  $v_1$  in what follows. Furthermore, using the balayage procedure (as above) one can make the support of  $d\nu := (dd^c v)^m \wedge \beta^{n-m}$  compact in  $\Omega$ .

Now, we can sketch the rest of the proof of the theorem. We will approximate  $d\mu$  by a sequence of measures  $\mu_j$  for which the Dirichlet problem is solvable (using Theorem 2.1) obtaining a sequence of solutions  $\{u_j\}$  corresponding to  $\mu_j$ . Then we take a limit point  $u$  of  $\{u_j\}$  in  $L^1(\Omega)$ . Finally we show that  $u_j \rightarrow u$  with respect to  $\text{cap}_m$  in order to conclude that  $u$  is a solution of (2.2).

By the Radon–Nikodym theorem,  $d\mu = h d\nu$ ,  $0 \leq h \leq 1$ . For the subsolution  $v$  one can define the regularizing sequence  $w_j \downarrow v$  in a neighborhood of the closure of  $\Omega$ . Let us write  $(dd^c w_j)^m \wedge \beta^{n-m} = g_j \beta^n$ ,  $\mu_j := h g_j \beta^n$ . Then by Proposition 1.11  $\lim_{j \rightarrow \infty} \mu_j = \mu$ . As  $\mu$  has the compact support, so  $\mu_j$ 's does. In particular,  $h g_j \in L^p(\Omega)$  for every  $p > 0$ . Therefore, applying Theorem 2.1 we have  $u_j$  solving

$$(2.3) \quad \begin{cases} u_j \in SH_m(\Omega) \cap C(\bar{\Omega}), \\ (dd^c u_j)^m \wedge \beta^{n-m} = \mu_j, \\ u_j(z) = \varphi(z) \text{ for } z \in \partial\Omega. \end{cases}$$

Now we set  $u = (\limsup u_j)^*$ , and passing to a subsequence we assume that  $u_j$  converges to  $u$  in  $L^1(\Omega)$ . By the definition of  $w_j$  they are uniformly bounded.

Choosing a uniform constant  $C$  such that  $w_j - C < \varphi$  on  $\partial\Omega$ , by Corollary 1.15-(a),  $w_j - C \leq u_j \leq \sup_{\bar{\Omega}} \varphi$ . Thus,  $\{u_j\}$  is uniformly bounded. In particular,  $u$  also is bounded and now we shall check that  $\lim_{\Omega \ni \zeta \rightarrow z} u(\zeta) = \varphi(z)$  for every  $z \in \partial\Omega$ . For this we only need  $\varphi$  to be continuous.

Since  $w_j$  converges uniformly to  $v$  on  $\partial\Omega$  and  $\partial\Omega$  is compact, given  $\varepsilon > 0$  we have  $|w_j - v| < \varepsilon$  on a small neighborhood of  $\partial\Omega$  when  $j$  big enough. Since  $\varphi$  is continuous on  $\partial\Omega$ , there is an approximant  $g \in C^2(\bar{\Omega})$  of the continuous extension of  $\varphi$  such that  $|g - \varphi| < \varepsilon$  on  $\partial\Omega$ . For  $A > 0$  big enough,  $A\rho + g$  is a  $m$ -sh function. By the comparison principle, it implies that  $w_j + A\rho + g - 2\varepsilon \leq u_j$  on  $\Omega$ . Then  $v + A\rho + \varphi - 4\varepsilon \leq u_j$  on a small neighborhood of  $\partial\Omega$  for  $j$  big enough. Hence,  $v + A\rho + \varphi - 4\varepsilon \leq \liminf_{j \rightarrow \infty} u_j \leq u$  on a small neighborhood of  $\partial\Omega$ . Because this is true for an arbitrary  $\varepsilon > 0$ , we obtain  $\lim_{\zeta \rightarrow z} u(\zeta) = \varphi(z)$  for any  $z \in \partial\Omega$ .

The difficult part is to show that  $u_j$  converges in  $cap_m$  to  $u$ .

LEMMA 2.3. *The function  $u$  defined above solves Dirichlet problem (0.3) provided that for any  $a > 0$  and any compact  $K \subset \Omega$  we have*

$$(2.4) \quad \lim_{j \rightarrow \infty} \int_{K \cap \{u - u_j \geq a\}} (dd^c u_j)^m \wedge \beta^{n-m} = \lim_{j \rightarrow \infty} \mu_j(K \cap \{u - u_j \geq a\}) = 0.$$

PROOF OF LEMMA 2.3. Using Theorem 1.13 we have

$$\begin{aligned} (dd^c u_j)^m \wedge \beta^{n-m} &= 1_{\{u - u_j \geq a\}} (dd^c u_j)^m \wedge \beta^{n-m} + 1_{\{u - u_j < a\}} (dd^c u_j)^m \wedge \beta^{n-m} \\ &\leq 1_{\{u - u_j \geq a\}} \mu_j + (dd^c \max\{u, u_j + a\})^m \wedge \beta^{n-m}. \end{aligned}$$

It follows that

$$(2.5) \quad \mu_j \leq 1_{\{u - u_j \geq a\}} \mu_j + (dd^c \max\{u - a, u_j\})^m \wedge \beta^{n-m}.$$

Now, for any integer  $s$  we may choose  $j(s)$  such that  $\mu_{j(s)}(\{u - u_{j(s)} \geq 1/s\}) < 1/s$ . From (2.4) and (2.5) we infer that

$$(2.6) \quad \mu \leq \liminf_{s \rightarrow \infty} (dd^c \rho_s)^m \wedge \beta^{n-m},$$

which means that  $\mu$  is less than any limit point of the right hand side, where  $\rho_s = \max\{u - 1/s, u_{j(s)}\}$ . By the Hartogs lemma,  $\rho_s \rightarrow u$  uniformly on any compact  $E$  such that  $u|_E$  is continuous. So it follows from the quasi-continuity of  $m$ -sh functions that  $\rho_s$  converges to  $u$  in  $cap_m$ . Therefore, by Theorem 1.9  $(dd^c \rho_s)^m \wedge \beta^{n-m} \rightarrow (dd^c u)^m \wedge \beta^{n-m}$  as measures. This combined with (2.6) implies

$$(2.7) \quad \mu \leq (dd^c u)^m \wedge \beta^{n-m}.$$

For the reverse inequality, let  $\Omega_\varepsilon = \{z \in \Omega ; \text{dist}(z, \partial\Omega) < \varepsilon\}$ . We will show that for  $\varepsilon > 0$

$$(2.8) \quad \mu(\Omega) \geq \int_{\Omega_\varepsilon} (dd^c u)^m \wedge \beta^{n-m}.$$

Indeed, firstly we note that  $\rho_s = u_{j(s)}$  on a neighborhood of  $\partial\Omega_\varepsilon$  for  $\varepsilon$  small enough since  $u - u_{j(s)} < 1/s$  on  $\partial\Omega$ ,  $u - u_{j(s)}$  is upper semi-continuous on  $\Omega$  and  $\partial\Omega$  is compact. Hence, by the weak\* convergence  $\mu_{j(s)} \rightarrow \mu$  and Stokes' theorem,

$$\begin{aligned} \mu(\Omega) &\geq \mu(\overline{\Omega_\varepsilon}) \geq \limsup_{j(s) \rightarrow \infty} \mu_{j(s)}(\overline{\Omega_\varepsilon}) \\ &\geq \liminf_{j(s) \rightarrow \infty} \mu_{j(s)}(\Omega_\varepsilon) \\ &= \liminf_{j(s) \rightarrow \infty} \int_{\Omega_\varepsilon} (dd^c u_{j(s)})^m \wedge \beta^{n-m} = \liminf_{j(s) \rightarrow \infty} \int_{\Omega_\varepsilon} (dd^c \rho_s)^m \wedge \beta^{n-m} \\ &\geq \int_{\Omega_\varepsilon} (dd^c u)^m \wedge \beta^{n-m}, \end{aligned}$$

where in the last inequality we use the weak\* convergence  $(dd^c \rho_s)^m \wedge \beta^{n-m} \rightarrow (dd^c u)^m \wedge \beta^{n-m}$ . Therefore, (2.8) is proved. Let  $\varepsilon \rightarrow 0$ ; there follows  $\mu(\Omega) \geq (dd^c u)^m \wedge \beta^{n-m}(\Omega)$ . Thus the measures in (2.7) are equal. The lemma follows.  $\square$

It remains to prove (2.4) in Lemma 2.2 above. It is a consequence of the following lemma.

LEMMA 2.4. *Suppose that there is a subsequence of  $\{u_j\}$ , still denoted by  $\{u_j\}$ , such that*

$$\int_{\{u-u_j \geq a_0\}} (dd^c u_j)^m \wedge \beta^{n-m} > A_0, \quad A_0 > 0, a_0 > 0.$$

*Then, for  $0 \leq p \leq m$ , there exist  $a_p, A_p, k_1 > 0$  such that*

$$(2.9) \quad \int_{\{u-u_j \geq a_p\}} (dd^c v_j)^{m-p} \wedge (dd^c v_k)^p \wedge \beta^{n-m} > A_p, \quad j > k > k_1,$$

*for  $v_j$  being the solution of the following of Dirichlet problem (cf. Theorem 2.1):*

$$(2.10) \quad \begin{cases} v_j \in SH_m(\Omega) \cap C(\overline{\Omega}), \\ (dd^c v_j)^m \wedge \beta^{n-m} = \nu_j (= g_j \beta^n), \\ v_j(z) = 0 \quad \text{on} \quad \partial\Omega. \end{cases}$$

*Note that  $\{v_j\}$  is uniformly bounded as a consequence of the uniform boundedness of  $\{w_j\}$  and Corollary 1.15-(a).*

PROOF OF LEMMA 2.4. We will prove it by induction over  $p$ . For  $p = 0$  the statement holds by the hypothesis. Suppose that (2.9) is true for  $p < m$ , we need to prove it for  $p + 1$ . The first observation is that if  $T(r, s) := (dd^c u_r)^q \wedge (dd^c v_s)^{m-q} \wedge \beta^{n-m}$  then there is a constant  $C$  independent of  $r, s$  such that

$$(2.11) \quad \int_{\Omega} T(r, s) \leq C.$$

Indeed, fix a  $C^2$  extension of  $\varphi$  to a neighborhood of the closure of  $\Omega$ . If  $\rho$  is a defining function of  $\Omega$ , then there is a constant  $A > 0$  such that  $A\rho \pm \varphi \in SH_m(\Omega)$ . We shall check that  $u_r + A\rho - \varphi$  belongs to  $\mathcal{E}_0(m)$ . It is enough to verify

$$\int_{\Omega} (dd^c(u_r + A\rho - \varphi))^m \wedge \beta^{n-m} < +\infty.$$

In fact, from  $(dd^c u_r)^m \wedge \beta^{n-m} = hg_r \beta^n \leq (dd^c(M_r \rho + \varphi))^m \wedge \beta^{n-m}$  for some  $M_r > 0$  and Corollary (1.15)-(a) we have  $u_r \geq M_r \rho + \varphi$  in  $\Omega$ . Hence,  $u_r + A\rho - \varphi \geq (M_r + A)\rho$  in  $\Omega$ . Thus, by Corollary 1.15-(c)

$$\int_{\Omega} (dd^c u_r + A\rho - \varphi)^m \wedge \beta^{n-m} \leq \int_{\Omega} (dd^c(M_r + A)\rho)^m \wedge \beta^{n-m} < +\infty.$$

Now, we note that  $\mu_r(\Omega)$  and  $\nu_s(\Omega)$  are bounded as  $\mu$  and  $\nu$  have compact supports. Next, from Cegrell's inequalities, Corollary 1.8-(ii), for  $1 \leq k \leq m - 1$ , there follows

$$\begin{aligned} & \int_{\Omega} (dd^c(u_r + A\rho - \varphi))^k \wedge (dd^c \rho)^{m-k} \wedge \beta^{n-m} \\ & \leq \left[ \int_{\Omega} (dd^c(u_r + A\rho - \varphi))^m \wedge \beta^{n-m} \right]^{\frac{k}{m}} \left[ \int_{\Omega} (dd^c \rho)^m \wedge \beta^{n-m} \right]^{\frac{m-k}{m}}. \end{aligned}$$

Hence,

$$\begin{aligned} (2.12) \quad I(r) &= \int_{\Omega} (dd^c(u_r + A\rho - \varphi))^m \wedge \beta^{n-m} \\ &\leq \int_{\Omega} (dd^c u_r)^m \wedge \beta^{n-m} + \int_{\Omega} (dd^c(A\rho - \varphi))^m \wedge \beta^{n-m} \\ &\quad + C(A, \varphi) \sum_{k=1}^{m-1} \int_{\Omega} (dd^c u_r + A\rho - \varphi)^k \wedge (dd^c \rho)^{m-k} \wedge \beta^{n-m} \\ &\leq \mu_r(\Omega) + C(A, \rho, \varphi) + C(A, \varphi) \cdot \\ &\quad \cdot \sum_{k=1}^{m-1} \left[ \int_{\Omega} (dd^c(u_r + A\rho - \varphi))^m \wedge \beta^{n-m} \right]^{\frac{k}{m}} \left[ \int_{\Omega} (dd^c \rho)^m \wedge \beta^{n-m} \right]^{\frac{m-k}{m}} \\ &\leq \mu_r(\Omega) + C(A, \rho, \varphi) + C'(A, \varphi, \rho) \sum_{k=1}^{m-1} [I(r)]^{\frac{k}{m}}. \end{aligned}$$

Consider the two sides of inequality (2.12) as two positive functions in  $r$ .  $\mu_r(\Omega)$ 's are bounded, and the degree of  $I(r)$  on the right hand side is strictly less than the degree of  $I(r)$  on the left hand side; therefore,  $I(r)$  are bounded by a constant independent of  $r$ . Again, by Cegrell's inequalities, Corollary 1.8-(ii), as  $v_s$  obviously belongs to  $\mathcal{E}_0(m)$ ,

$$\begin{aligned} \int_{\Omega} T(r, s) &\leq \int_{\Omega} (dd^c(u_r + A\rho - \varphi))^q \wedge (dd^c v_s)^{m-q} \wedge \beta^{n-m} \\ &\leq \left[ \int_{\Omega} (dd^c(u_r + A\rho - \varphi))^m \wedge \beta^{n-m} \right]^{\frac{q}{m}} \left[ \int_{\Omega} (dd^c v_s)^m \wedge \beta^{n-m} \right]^{\frac{m-q}{m}} \\ &\leq [I(r)]^{\frac{q}{m}} [\nu_s(\Omega)]^{\frac{m-q}{m}} \\ &\leq C''(A, \varphi, \rho), \end{aligned}$$

because  $I(r)$  and  $\nu_s(\Omega)$  are bounded. Thus we have proved (2.11). We may assume that  $-1 < u_j, v_j < 0$ , because all functions  $u_j, v_j$  are uniformly bounded by a constant independent of  $j$ . Then the estimates in the statement of Lemma 2.4 will only be changed by a positive constant. To simplify notations we set  $S(j, k) := (dd^c v_j)^{m-p-1} \wedge (dd^c v_k)^p \wedge \beta^{n-m}$ . Fix a positive number  $d > 0$  (specified later in (2.19)) and recall that we need a uniform estimate from below for  $\int_{\{u-u_j \geq d\}} dd^c v_k \wedge S(j, k)$ . From the assumption on  $u_j, v_j$ , we have  $u - u_j \leq \mathbf{1}_{\{u-u_j \geq d\}} + d$ . It follows that

$$\begin{aligned} J(j, k) &:= \int_{\Omega} (u - u_j) (dd^c v_k) \wedge S(j, k) \\ &\leq \int_{\Omega} \mathbf{1}_{\{u-u_j \leq d\}} dd^c v_k \wedge S(j, k) + d \int_{\Omega} dd^c v_k \wedge S(j, k) \\ &\leq \int_{\{u-u_j \geq d\}} dd^c v_k \wedge S(j, k) + dC, \end{aligned}$$

where  $C$  is as in (2.11). Therefore

$$(2.13) \quad \int_{\{u-u_j \geq d\}} dd^c v_k \wedge S(j, k) \geq J(j, k) - dC.$$

The induction hypothesis says that there exist  $a_p, A_p > 0$  and  $k_1 > 0$  such that

$$(2.14) \quad \int_{\{u-u_j \geq a_p\}} (dd^c v_j)^{m-p} \wedge (dd^c v_k)^p \wedge \beta^{n-m} > A_p, \quad j > k > k_1.$$

We fix another small positive constant  $\varepsilon > 0$  and put  $J'(j, k) := \int_{\Omega} (u - u_j) dd^c v_j \wedge S(j, k)$ .



CLAIM.

- (a)  $J'(j, k) - J(j, k) \leq \varepsilon$ ,  
 (b)  $J'(j, k) \geq a_p A_p - \varepsilon(1 + C)$  for  $j > k > k_2$ .

PROOF OF CLAIM. **(a)** By the quasi-continuity, we can choose an open set  $U$  such that functions  $u, v$  are continuous off the set  $U$  and  $\text{cap}_m(U, \Omega) < \varepsilon/2^{m+1}$ . Then

$$(2.15) \quad \int_U (dd^c(v_j + v_k))^m \wedge \beta^{n-m} < 2^m \text{cap}_m(U, \Omega) < \varepsilon/2,$$

$$(2.16) \quad \int_U (dd^c(u_j + v_k))^m \wedge \beta^{n-m} < \varepsilon/2.$$

Therefore

$$J'(j, k) - J(j, k)$$

$$(2.17) \quad \begin{aligned} &= \int_{\Omega} (u - u_j) dd^c v_j \wedge S(j, k) - \int_{\Omega} (u - u_j) dd^c v_k \wedge S(j, k) \\ &= \int_{\Omega} v_j dd^c(u - u_j) \wedge S(j, k) - \int_{\Omega} v_k dd^c(u - u_j) \wedge S(j, k) \\ &= \int_{\Omega} (v_j - v_k) dd^c(u - u_j) \wedge S(j, k) \\ &= \int_{\Omega \setminus U} (v_j - v_k) dd^c(u - u_j) \wedge S(j, k) + \int_U (v_j - v_k) dd^c(u - u_j) \wedge S(j, k) \\ &\leq \int_{\Omega \setminus U} \|v_j - v_k\| dd^c(u + u_j) \wedge S(j, k) + \int_U dd^c(u + u_j) \wedge S(j, k), \end{aligned}$$

where in the second equality we used the integration by parts formula twice with  $u = u_j = \varphi$ ,  $v_j = 0$  on the boundary, and in the last estimate we used the fact  $-1 < u_j, v_j < 0$ . Since the sequence  $(v_j)$  converges uniformly to  $v$  on  $\Omega \setminus U$ , one can find  $l > k_1$  such that  $\|v_j - v_k\| < \varepsilon/2C$  on  $\Omega \setminus U$  for  $j > k > l > k_1$ . From (2.11) and  $u_s$  converges in  $L_{loc}^1$  to  $u$ , we have

$$\int_{\Omega} dd^c u \wedge S(j, k) \leq \liminf_{s \rightarrow +\infty} \int_{\Omega} dd^c u_s \wedge S(j, k) < C.$$

This combined with (2.15) and (2.16) implies that each of the integrals in the last line of (2.17) is at most  $\varepsilon/2$ . The first part of the claim follows.

**(b)** We first observe that since  $\sup \varphi$  (resp. 0) is the upper bound of all  $u_j$  (resp.  $v_j$ ) on the boundary, then, for  $k > k_2 > l$ , in a neighborhood of  $\partial\Omega$ , there is

$$(2.18) \quad v_k \leq v + \varepsilon \quad \text{and} \quad u_k \leq u + \varepsilon.$$

Those inequalities are still valid (after increasing  $k_2$ ) on  $\Omega \setminus U$  thanks to the Hartogs lemma. Hence, using (2.11), (2.15) and (2.18) for  $j > k > k_2$  there is

$$\begin{aligned}
J'(j, k) &= \int_{\Omega} (u - u_j) dd^c v_j \wedge S(j, k) \\
&\geq a_p \int_{\{u - u_j \geq a_p\}} dd^c v_j \wedge S(j, k) + \int_{\{u - u_j < a_p\}} (u - u_j) dd^c v_j \wedge S(j, k) \\
&= a_p \int_{\{u - u_j \geq a_p\}} dd^c v_j \wedge S(j, k) + \int_{\{u - u_j < a_p\} \cap (\Omega \setminus U)} (u - u_j) dd^c v_j \wedge S(j, k) \\
&\quad + \int_{\{u - u_j < a_p\} \cap U} (u - u_j) dd^c v_j \wedge S(j, k) \\
&\geq a_p \int_{\{u - u_j \geq a_p\}} dd^c v_j \wedge S(j, k) - \varepsilon \int_{\Omega \setminus U} dd^c v_j \wedge S(j, k) \\
&\quad - \int_U dd^c v_j \wedge S(j, k) \\
&\geq a_p A_p - \varepsilon(1 + C).
\end{aligned}$$

Thus the proof of the claim is finished.  $\square$

From *Claim* and (2.13) we get

$$\begin{aligned}
\int_{\{u - u_j \geq d\}} dd^c v_k \wedge S(j, k) &\geq J(j, k) - dC \geq J'(j, k) - \varepsilon - dC \\
&\geq a_p A_p - \varepsilon(1 + C) - \varepsilon - dC.
\end{aligned}$$

If we take

$$(2.19) \quad a_{m+1} := d = \frac{a_p A_p}{4C} \quad \text{and} \quad \varepsilon \leq \frac{a_p A_p}{2(2 + C)},$$

then

$$\int_{\{u - u_j \geq d\}} dd^c v_k \wedge S \geq \frac{a_p A_p}{4} := A_{p+1} \quad \text{for } j > k > k_2,$$

which finishes the proof of the inductive step and that of Lemma 2.4.  $\square$

We will now finish the proof of Theorem 2.2. It is enough to prove condition (2.4) in Lemma 2.3. We argue by contradiction. Suppose that it is not true. Then the assumptions of Lemma 2.4 are valid and its statement for  $p = m$  tells that for a fixed  $k > k_1$

$$\int_{\{u - u_j \geq a_m\}} (dd^c v_k)^m \wedge \beta^{n-m} > A_m \quad \text{when } j > k.$$

Thus

(2.20)

$$V(\{u - u_j \geq a_m\}) \geq \frac{1}{M_k} \int_{\{u - u_j \geq a_m\}} (dd^c v_k)^m \wedge \beta^{n-m} > \frac{A_m}{M_k} \text{ for } j > k,$$

because  $(dd^c v_k)^m \wedge \beta^{n-m} = g_k \beta^n \leq M_k \beta^n$  for some  $M_k > 0$ . But (2.20) contradicts the fact  $u_j \rightarrow u$  in  $L^1_{loc}$ , i.e. every subsequence of  $\{u_j\}$  also converges to  $u$  in  $L^1_{loc}$ . Thus, the theorem is proved under two extra assumptions.

*General case (we remove two extra assumptions).* **1)** Suppose that  $\varphi \in C(\partial\Omega)$  and the measure  $\mu$  has a compact support in  $\Omega$ . We choose a decreasing sequence  $\varphi_k \in C^2(\partial\Omega)$  converging to  $\varphi$ . Then we obtain a sequence of solutions  $u_k$  satisfying

$$\begin{cases} u_k \in SH_m \cap L^\infty(\Omega), \\ (dd^c u_k)^m \wedge \beta^{n-m} = \mu, \\ \lim_{\zeta \rightarrow z} u_k(\zeta) = \varphi_k(z) \text{ for any } z \in \partial\Omega. \end{cases}$$

It follows from the comparison principle and Corollary 1.15-(a) that  $u_k$  is decreasing and  $u_k \geq v_0$  with  $v_0$  a subsolution without modifications. Set  $u = \lim u_k$ . Then  $u \geq v_0$  and  $(dd^c u)^m \wedge \beta^{n-m} = \mu$  by Proposition 1.11. Thus,  $u$  is the required solution.

**2)** Suppose that  $\mu$  is a finite positive Borel measure,  $\varphi \in C(\partial\Omega)$ . Let  $\chi_j$  be a non-decreasing sequence of cut-off functions  $\chi_j \uparrow 1$  on  $\Omega$ . Since  $\chi_j \mu$  have compact supports in  $\Omega$ , one can find solutions corresponding to  $\chi_j \mu$ , the solutions will be bounded from below by the given subsolution  $v_0$  (by the comparison principle) and they will decrease to the solution by the convergence theorem. Thus we have proved Theorem 2.2.  $\square$

## References

1. Bedford E., Taylor B. A., *The Dirichlet problem for a complex Monge–Ampère equation*, Invent. Math., **37** (1976), 1–44.
2. Bedford E., Taylor B. A., *A new capacity for plurisubharmonic functions*, Acta Math., **149** (1982), 1–40.
3. Blocki Z., *Weak solutions to the complex Hessian equation*, Ann. Inst. Fourier (Grenoble), **55**, No. **5** (2005), 1735–1756.
4. Caffarelli L., Nirenberg L., Spruck J., *The Dirichlet problem for nonlinear second order elliptic equations, III: Functions of the eigenvalues of the Hessian*, Acta Math., **155** (1985), 261–301.
5. Cegrell U., *Pluricomplex energy*, Acta Math., **180**, No. **2** (1998), 187–217.
6. Cegrell U., *The general definition of the complex Monge–Ampère operator*, Ann. Inst. Fourier (Grenoble), **54** (2004), 159–179.
7. Demailly J. P., *Potential theory in several complex variable*, Lecture notes, ICPAM, Nice, 1989.
8. Demailly J. P., *Complex analytic and differential geometry*, available at <http://www-fourier.ujf-grenoble.fr/~demailly/books.html>, (September 2009).

9. Dinew S., *An inequality for mixed Monge–Ampère measures*, Math. Zeit., **262** (2009), 1–15.
10. Dinew S., Kołodziej S., *A priori estimates for the complex Hessian equations*, preprint [arXiv:1112.3063](https://arxiv.org/abs/1112.3063).
11. Gårding L., *An inequality for hyperbolic polynomials*, J. Math. Mech., **8** (1959), 957–965.
12. Hörmander L., *Notions of convexity*, Birkhauser, Boston, 1994.
13. Kołodziej S., *The range of the complex Monge–Ampère operator II*, Indiana U. Math. J., **44**, No. **3** (1995), 765–782.
14. Kołodziej S., *The complex Monge–Ampère equation and pluripotential theory*, Memoirs Amer. Math. Soc., **178** (2005).
15. Li S.-Y., *On the Dirichlet problems for symmetric function equations of the eigenvalues of the complex Hessian*, Asian J. Math., **8** (2004), 87–106.
16. Trudinger N. S., Wang X.-J., *Hessian measures II*, Ann. of Math., **150** (1999), 579–604.
17. Wang X.-J., *The  $k$ -Hessian equation*, Lect. Not. Math., **1977** (2009).

*Received August 7, 2012*

Institute of Mathematics  
Jagiellonian University  
Łojasiewicza 6  
30-348 Kraków, Poland  
*e-mail:* [Nguyen.Ngoc.Cuong@im.uj.edu.pl](mailto:Nguyen.Ngoc.Cuong@im.uj.edu.pl)