

*Plant Pathol. J.* 33(1) : 21-33 (2016)  
<https://doi.org/10.5423/PPJ.OA.2016.01.001>  
 pISSN 1598-2254 eISSN 2093-5263

Research Article Open Access

The Plant Pathology Journal

©The Korean Society of Plant Pathology

## Characterization of *Pseudomonas syringae* pv. *syringae*, Causal Agent of Citrus Blast of Mandarin in Montenegro

Žarko Ivanović<sup>1\*</sup>, Tatjana Hrnčić<sup>3</sup>, Tatjana Popović<sup>1</sup>, Jovana Blagojević<sup>2</sup>, Nenad Trkulja<sup>1</sup>, and Snježana Hrnčić<sup>3</sup>

<sup>1</sup>Institute for Plant Protection and Environment, Belgrade 11040, Serbia

<sup>2</sup>Scholar of Ministry of Education, Science and Technological Development of the Republic of Serbia, Department of Plant Diseases, Institute for Plant Protection and Environment, Belgrade 11040, Serbia

<sup>3</sup>Biotechnical Faculty, University of Montenegro, Podgorica, Podgorica 81000, Montenegro

(Received on August 2, 2016; Revised on September 22, 2016; Accepted on November 1, 2016)

Citrus blast caused by bacterium *Pseudomonas syringae* is a very important disease of citrus occurring in many areas of the world, but with few data about genetic structure of the pathogen involved. Considering the above fact, this study reports genetic characterization of 43 *P. syringae* isolates obtained from plant tissue displaying citrus blast symptoms on mandarin (*Citrus reticulata*) in Montenegro, using multilocus sequence analysis of *gyrB*, *rpoD*, and *gapI* gene sequences. Gene sequences from a collection of 54 reference pathotype strains of *P. syringae* from the Plant Associated and Environmental Microbes Database (PAMDB) was used to establish a genetic relationship with our isolates obtained from mandarin. Phylogenetic analyses of *gyrB*, *rpoD*, and *gapI* gene sequences showed that *P. syringae* pv. *syringae* causes citrus blast in mandarin in Montenegro, and belongs to genomospecies 1. Genetic homogeneity of isolates suggested that the Montenegrin population might be clonal which indicates a possible common source of infection. These findings may assist in further epidemiological studies of this pathogen and for determining mandarin breeding strategies for *P. syringae* control.

**Keywords :** bacteria, *Citrus reticulata*, multilocus se-

quence analysis, pathogen, phylogeny

In Mediterranean countries, phytopathogenic bacteria do not generally cause serious damage in citrus groves, excluding the incidence of the blast and black pit, caused by *Pseudomonas syringae* pv. *syringae* van Hall 1902. This bacterium infects more than 180 different plant species (Bradbury, 1986), among them *Citrus* spp. where it causes bacterial blast of orange (*Citrus sinensis*) and mandarin (*Citrus reticulata*) (Smith and Fawcett, 1930; Whiteside et al., 1988) and black pit of orange fruits (Mirik et al., 2005; Whiteside et al., 1988). *P. syringae* pv. *syringae* causes damage especially under cool and humid conditions in spring when the development and spread of the bacterial blast happens quicker and more easily (Gorlenko, 1965), and when the shoots or fruits are damaged by wind, hail or thorns. Characteristic disease symptoms appearing as water-soaked lesions extended to the mid-vein and to the twigs surrounding the base of the petiole. Ultimately, the leaves dry and curl, while still firmly attached, and eventually fall without petioles. The necrotic areas on twigs further enlarge and the twigs are eventually killed within three to four weeks.

*P. syringae* pv. *syringae* is an organism which has been researched on non-woody hosts extensively compared to wood plant like citrus. Only a few studies have been done that take this topic with citrus into account (Mirik et al., 2005; Shigeta and Nakata, 1995; Smith and Fawcett, 1930; Thomidis et al., 2005; Timmer et al., 2000; Vanarelli et al., 2010). The lack of genetic data that considers this citrus pathogen is evident in literature. Scortichini et al. (2003) ascertained genetic variability of *P. syringae* pv.

\*Corresponding author.

Phone) +381-112660049, FAX) +381-112669860

E-mail) zarko.ivanovic@yahoo.com

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Articles can be freely viewed online at [www.ppjonline.org](http://www.ppjonline.org).

*syringae* strains isolated from apple, pear, apricot, *Citrus* spp. and cereals and determined that no clear relation was observed between the host plant and the bacterial genomic fingerprint (BOX-PCR). Despite the evidence of the occurrence of *P. syringae* pv. *syringae* in citrus growing areas in different parts of the world, the lack of molecular studies of pathogen variation is also a factor that limits the better understanding of the pathogen. In Montenegro, the symptoms of citrus blast have been observed for the first time in 1987, when severe outbreaks were found in many citrus nurseries, with disease incidence approaching 10% to 30% (Vučinić, 1987). In a survey for bacterial blast in Montenegro during the spring seasons of 2013 and 2014, the incidence of disease was at least 10%. Given the ubiquity of *P. syringae* pv. *syringae*, a better understanding of the epidemiology and pathogen population structure could be used in predicting disease and developing predictive models of outbreaks. For this reason, a collection of *P. syringae* pv. *syringae* isolates obtained from diseased Montenegrin mandarin (*Citrus reticulata* cv. Owari) were studied using *syxB* analysis which determines the toxin involved in virulence of *P. syringae* pv. *syringae* (Sorensen, 1998), partial 16S rDNA sequencing (Gironde and Manceau, 2012; Hilario et al., 2004) and multilocus sequence analysis (MLSA) using three different housekeeping genes (Bull et al., 2011). Furthermore, pathogenicity testing was performed in order to conclude Koch's postulates. This paper also shows the general similarity and certain differences to other *P. syringae* reference strains obtained from different hosts deposited in the Plant Associated and Environmental Microbes Database (PAMDB) data base ([www.pamdb.org](http://www.pamdb.org)) and to bring together certain information regarding the aforementioned.

## Materials and Methods

**Bacterial isolates.** For the pathogen isolation, 20 samples of necrotic buds from diseased twigs were taken. Small pieces of tissue were taken from the margins of necrotic lesions and were first disinfected with 70% ethanol, then cut and crushed using a mortar and pestle in sterile phosphate-buffered saline (0.27% Na<sub>2</sub>HPO<sub>4</sub>, 0.04% NaH<sub>2</sub>PO<sub>4</sub>, 0.8% NaCl). After 20 min, the suspensions were plated on King's medium B (King et al., 1954). After 2 days of incubation at 26°C, a total of 43 fluorescent colonies were selected and purified on Sucrose Nutrient Agar (SNA) (Difco nutrient agar, supplemented with 0.5% sucrose; Difco Laboratories, Detroit, MI, USA) and maintained at -80°C in Luria Bertani medium supplemented with 20% glycerol (Bertani, 1951). The strains CFBP 11 (*P. syringae* pv. *syringae*), CFBP 1582 (*P. syringae* pv. *syringae*), and CFBP 2119 (*P. syringae* pv. *morsprunorum*), were

used as reference strains (Collection Francaise de Bacteries Phytopathogenes [CFBP], Angers, France).

**Pathogenicity.** Pathogenicity testing of the obtained isolates were checked by artificial inoculation of unripe green lemon (*Citrus limon*) fruits, bean pods (*Phaseolus vulgaris*) and leaves of mandarin and lilac (*Syringa vulgaris*) using the methods described by Lelliott and Stead (1987) and Klement (1990). Each experiment was repeated twice for all tested isolates. The bacterial cells were grown for 2 days on SNA at 25°C, and bacterial suspensions were prepared in sterile distilled water (SDW) and photometrically adjusted to the final concentration of 10<sup>8</sup> cfu/ml. Control fruits and leaves were also included in tests, but treated with reference strains and SDW as positive and negative controls, respectively.

The lemons were surface sterilized by being dipped into 2% NaClO solution for 2 min and then shortly rinsed with SDW. Four punctures were made on the fruits (1 mm wide, 3 mm deep), by using a sterile hypodermic needle and per 20 µl of the bacterial suspension (10<sup>6</sup> cfu/ml) was placed on each rupture. Bean pods were inoculated by injection of a bacterial suspension (10<sup>7</sup> cfu/ml) into young bean pods with a hypodermic syringe. Inoculated fruits and pods were kept in plastic bags and incubated at 25°C for 7 days under conditions of high humidity (70–80%). For each isolate, 12 sites involving three fruits/pods were inoculated. Pathogenic reactions were assessed seven days after inoculation (DAI).

Lilac (*Syringa vulgaris* cv. Sensation) and mandarin (cv. Owari) were ruptured with a sterile hypodermic needle by pricking the leaf midrib and into areas on the wing of the petiole of fully expanded leaves, and inoculated with a 20 µl drop of bacterial suspension (10<sup>6</sup> cfu/ml) deposited on a newly made rupture. Inoculated leaves and the petioles were placed in plastic bags and kept under room temperature conditions and high humidity (80%). For each tested isolate and controls, five different leaves and petioles were inoculated. Pathogenic reactions were assessed 14 DAI.

In addition, twigs of mandarin trees were longitudinally ruptured (1 cm) using a sterile scalpel, and per 100 µl of the bacterial suspension (10<sup>6</sup> cfu/ml) was placed onto the cuts. Twigs were kept in clear polyethylene bags for 24 h under room temperature conditions and high humidity (70%–80%). For each tested isolate and controls, three stems were inoculated. Symptoms were assessed 28 DAI.

**Biochemical characteristics.** Bacterial isolates and reference strains were tested for LOPAT (levan production, oxidase reaction, potato soft rot, arginine dihydrolase activity, and tobacco hypersensitivity) tests (Lelliott and

Stead, 1987). Levan production was estimated by growing isolates on SNA medium. For oxidase reaction test, bacteria were grown on SNA medium for 24 h at 25°C until colonies were obtained. Colonies were picked with a loop and rubbed on filter-paper impregnated with 1% (w/v) aqueous tetramethyl-p-phenylene diamine dihydrochloride solution. Production of a purple colour within 10 s was recorded as positive, its development in 10 to 60 s as delayed positive, and the absence of coloration or its still later development as negative (Kovacs, 1956). Pectinase activity was tested by adding bacterial suspension ( $10^6$  cfu/ml) onto the potato slices and incubated for 24 h at 25°C. Each isolate was tested for arginine dihydrolase production by the method of Thornley (Thornley, 1960), by growing in medium with L-arginine. They were then covered with a 5-mm layer of sterile liquid paraffin, tightly capped, and incubated at 25°C for 7 days. A positive test for the presence of arginine dihydrolase was indicated by a deep pink to red color, whereas a negative test was demonstrated by an absence of color change in the medium. Tobacco hypersensitivity, was tested on leaves of the tobacco. Inoculum were prepared in SDW as described for the pathogenicity test. Reaction to Gram stain, oxidative/fermentative metabolism of glucose (O/F) and fluorescent pigment production on the King's medium B, were also tested (Lelliott and Stead, 1987; Schaad et al., 2001). In addition, GATT tests (Gelatin and Aesculine hydrolysis, formation of Tyrosinase activity and Tartrate metabolism) for the differentiation within the species *P. syringae* were carried out (Kaluzna et al., 2012). Gelatin liquefaction was done by growing bacteria in tubes containing 12% (w/v) gelatin, which were incubated up to 15 days at 25°C and then held at 4°C for 15 min before determining liquefaction.  $\beta$ -Glucosidase activity of tested isolates was estimated by growing isolates on 0.1% (w/v) aesculin agar medium and incubating four days at 25°C. Positive reaction is dark brown color of the medium after incubation. Tyrosinase activity of isolates were estimated on agar medium containing 0.1% (w/v) L-tyrosine by incubating 4 days at 25°C. A positive test is red dark brown color of the medium after incubation. Utilization of tartarate as a sole carbon source on Simmons basal medium (Holding and Collee, 1971) containing 0.2% (w/v) sodium tartarate. After 21 days of incubation at 25°C if reaction is positive medium turn color from green to blue.

**Genomic DNA extraction.** Total genomic DNA was extracted using a modification of the procedure of Ausubel et al. (1992). All tested 43 Montenegrin mandarin isolates and *P. syringae* reference strains were grown on SNA solid medium for 48 h at 25°C. Bacterial cells were washed in 1.5 ml of SDW and centrifuged at 4,000g for

10 min at 4°C. The resulting pellet was washed twice in 500  $\mu$ l 0.85% NaCl, recentrifuged and washed once in 0.1 M NaPO<sub>4</sub> buffer, pH 6.8. Cells were treated with 10% sodium dodecyl sulfate (SDS) and with proteinase K at 37°C for 1 h. DNA was purified by adding 100  $\mu$ l 5 M NaCl and 80  $\mu$ l solution of 10% hexadecyltrimethyl ammonium bromide (CTAB) in 1 M NaCl at 65°C for 10 min extracted with an equal volume of chloroform and centrifuged for 10 min at 8,000g. The aqueous phase was transferred to a new tube and precipitated by adding 0.6 vol of isopropanol. The DNA was then washed with 70% ethanol and re-dissolved in 100  $\mu$ l Tris-EDTA (TE; 10 mM Tris, 1 mM EDTA, pH 8.0), and quantified spectrophotometrically at 260 nm.

**Detection of the *syxB* gene.** In order to detect the possible presence of the *syxB* gene in all tested isolates, the primer pair B1 (5'-CTTTCCGTGGTCTTGATGAGG-3') and B2 (5'-TCGATTTGCCGTGATGAGTC-3') was used. PCR amplification procedure was described by Sorensen (1998). For negative control, template was replaced with the same volume of SDW. Additionally, reference strains of *P. syringae* pv. *syringae* (CFBP 11 and CFBP 1582) and *P. syringae* pv. *morsprunorum* (CFBP 2119) were tested as templates for PCR amplification. The PCR was programmed for 35 cycles, denaturation at 94°C for 1.5 min, primer annealing at 60°C for 1.5 min, DNA extension for 3.0 min at 72°C and additional extension at 72°C for 10 min. PCR amplifications were performed in a Mastercycler personal model (Eppendorf, Hamburg, Germany).

**Repetitive PCR genomic fingerprinting.** An enterobacterial repetitive intergenic consensus (ERIC)-PCR with primer pair ERIC1R (5'-ATGTAAGCTCCTGGGGATTCAC-3') and ERIC2 (5'-AAGTA AGTGAAGTGGGGT-GAGCG-3') have been used for differentiation among the isolates (Lupski and Weinstock, 1992). ERIC-PCR conditions were as previously described by de Bruijn (1992). Amplification of PCR was performed with a Mastercycler personal model (Eppendorf) by using the following cycles: 1 initial cycle at 95°C for 7 min; 30 cycles of denaturation at 94°C for 1 min, annealing at 52°C for 1 min, and extension at 65°C for 8 min with a single final extension cycle at 65°C for 15 min and a final soak at 4°C. The PCR amplifications were performed in triplicate. Amplified PCR products were separated by gel electrophoresis on 1% agarose gels in 0.5 $\times$  TAE buffer for 1 h at 5 V/cm, stained with 0.05  $\mu$ l/ml ethidium bromide and visualized under UV illumination. Fingerprints generated from different strains were compared visually.



**16S rDNA sequence analysis.** Amplicons for the partial 16S rDNA sequences were generated using universal primers (5'-GAGAGTTTGATCCTGGCTCAG-3') and P6 (5'-CTACGGCTACCTTGTACGA-3') (Grifoni et al., 1995) and reaction conditions described by Scortichini et al. (2005). PCR products were purified with the QIAquick PCR purification kit (Qiagen, Valencia, CA, USA) following the manufacturer's instructions. The sequencing was performed by Macrogen Inc. (Seoul, Korea) and the sequences were deposited in the GenBank database. The MEGA6 software package (Tamura et al., 2013) was used to align sequences from forward and reverse strands and generate consensus sequences. A basic local alignment search tool BLAST (Altschul et al., 1997) was used to compare unidentified sequences with sequences in public databases. Neighbor joining (NJ) analyses were performed using 16S rDNA sequences which were assembled and edited using FINCHTV v.1.4.0 (<http://www.geospiza.com>). Multiple alignments and comparisons with reference strains were performed using CLUSTALW integrated into MEGA6 software (Tamura et al., 2013).

**Multilocus sequence analysis.** MLSA was performed in order to reveal the relationship between the *P. syringae* pv. *syringae* isolates obtained from mandarin and the *P. syringae* reference strains. Gene fragments of three housekeeping genes such as, *gyrB*, *rpoD*, and *gap1* coding for DNA gyrase B, sigma factor 70 and glyceraldehyde-3-phosphate dehydrogenase, respectively, were amplified from genomic DNA and sequenced with primers described by Sarkar and Guttman (2004). PCR products were purified with the QIAquick PCR purification kit (Qiagen) following the manufacturer's instructions. Sequencing was performed by Macrogen Inc. and the sequences were deposited in the GenBank database. Sequences were edited, aligned, and compared with other *Pseudomonas* sp. sequences as is described in the study given by Sarkar and Guttman (2004).

**Data analysis.** In order to estimate the evolutionary relation between *P. syringae* pv. *syringae* isolates obtained from mandarin and the pathotype reference strains obtained from PAMDB database, phylogenetic trees were built using NJ analyses. Gene sequences were assembled and edited using FINCHTV v.1.4.0 (<http://www.geospiza.com>). Multiple alignments and comparisons with reference strains for each of the genes were performed using CLUSTAL W integrated into MEGA6 software (Tamura et al., 2013). *Pseudomonas fluorescens* was used as out-group.

Minimum spanning tree (MST) was generated, where isolates are the nodes of a tree using SplitsTree software

(Huson and Bryant, 2006). The method estimates a probability of ancestry for each individual isolate from each of the groups. The individual isolates are assigned to one cluster or jointly to two or more clusters if their genotypes indicate that they were admixed.

## Results

**Bacterial isolation and characterization.** During the spring of 2013 and 2014, severe outbreaks of citrus blast were observed in mandarin (cv. Owari) in the regions of Bar and Ulcinj in Montenegro (Fig. 1). Isolations of bacteria from necrotic buds yielded almost pure cultures of whitish, translucent bacterial colonies that were fluorescent on King's medium B after 48 h of incubation. A total of 43 separate bacterial colonies were selected and purified on SNA for further analysis (Table 1). On SNA, isolates formed large (2–3 mm in diameter), convex (levan positive), cream-whitish, mucoid colonies 2–3 DAI.

*P. syringae* strains isolated from diseased mandarin tissues induced typical symptoms of *P. syringae* pv. *syringae* on an inoculated unripe lemon, appearing as deep, black necrotic pits. On dwarf bean pods, isolates produced water-soaked lesions with reddish margins 3 DAI. In syringe-inoculated lilac leaves, necrotic areas were seen 2–4 DAI. Inoculated petioles showed water-soaked or black areas, which expanded along the mid vein of the leaves, 7–10 DAI. Symptoms on mandarin leaves and petioles were observed 2 DAI in the form of a brown necrotic area at the point of inoculation. Lesions enlarged slightly on leaves until 10 DAI and brown-black necroses were recorded. On inoculated mandarin twigs there was typical and extensive tissue necrosis, while wilting and dieback were observed 4 weeks after inoculation. No symptoms



**Fig. 1.** Symptoms of citrus blast disease caused by *Pseudomonas syringae* pv. *syringae* on mandarin natural infection.

**Table 1.** Original source and main characteristics of *Pseudomonas syringae* pv. *syringae* isolates

Isolates	Species	Area of origine	Detection of <i>syxB</i>	LOPAT tests results				LOPAT group	Glucose (O/F) metabolism	Gelatin liquefaction	Aesculin hydrolysis	Tyrosinase activity	Tartrate utilisation
				L	O	P	A						
IZB 200	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	+	+	-	-	-	+	O	+	-	-	
IZB 201	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	+	+	-	-	-	+	O	+	-	-	
IZB 202	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	+	+	-	-	-	+	O	+	-	-	
IZB 203	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	+	+	-	-	-	+	O	+	-	-	
IZB 204	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	+	+	-	-	-	+	O	+	-	-	
IZB 205	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	+	+	-	-	-	+	O	+	-	-	
IZB 206	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	+	+	-	-	-	+	O	+	-	-	
IZB 207	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	+	+	-	-	-	+	O	+	-	-	
IZB 208	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	+	+	-	-	-	+	O	+	-	-	
IZB 209	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	+	+	-	-	-	+	O	+	-	-	
IZB 210	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	+	+	-	-	-	+	O	+	-	-	
IZB 211	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	+	+	-	-	-	+	O	+	-	-	
IZB 212	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	+	+	-	-	-	+	O	+	-	-	
IZB 213	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	+	+	-	-	-	+	O	+	-	-	
IZB 214	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	+	+	-	-	-	+	O	+	-	-	
IZB 215	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	+	+	-	-	-	+	O	+	-	-	
IZB 216	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	+	+	-	-	-	+	O	+	-	-	
IZB 217	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	+	+	-	-	-	+	O	+	-	-	
IZB 218	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Iliino)	+	+	-	-	-	+	O	+	-	-	
IZB 219	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Iliino)	+	+	-	-	-	+	O	+	-	-	
IZB 220	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Iliino)	+	+	-	-	-	+	O	+	-	-	
IZB 221	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Iliino)	+	+	-	-	-	+	O	+	-	-	
IZB 222	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Iliino)	+	+	-	-	-	+	O	+	-	-	
IZB 223	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Iliino)	+	+	-	-	-	+	O	+	-	-	
IZB 224	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	O	+	-	-	
IZB 225	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	O	+	-	-	
IZB 226	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	O	+	-	-	
IZB 227	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	O	+	-	-	
IZB 228	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	O	+	-	-	
IZB 229	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	O	+	-	-	
IZB 230	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	O	+	-	-	
IZB 231	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	O	+	-	-	
IZB 232	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	O	+	-	-	
IZB 233	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	O	+	-	-	
IZB 234	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	O	+	-	-	

Table 1. Continued

Isolates	Species	Area of origine	Detection of <i>syrB</i>	LOPAT tests results							Glucose (O/F) metabolism	Gelatin liquefaction	Aesculin hydrolysis	Tyrosinase activity	Tartrate utilisation
				L	O	P	A	T	LOPAT group						
IZB 235	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	Ia	O	+	+	-	-	
IZB 236	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	Ia	O	+	+	-	-	
IZB 237	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	Ia	O	+	+	-	-	
IZB 238	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	Ia	O	+	+	-	-	
IZB 239	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	Ia	O	+	+	-	-	
IZB 240	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	Ia	O	+	+	-	-	
IZB 241	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	Ia	O	+	+	-	-	
IZB 242	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	+	+	-	-	-	+	Ia	O	+	+	-	-	

L, levan production; O, oxidase reaction; P, potato soft rot; A, arginine dihydrolase activity; T, tobacco hypersensitivity; O/F, oxidative/fermentative metabolism of glucose.

were observed on SDW inoculated plants.

All isolates showed similar biochemical and physiological characteristics that were typical of *P. syringae*. They were Gram-negative, fluorescent on King's medium B, showed strictly aerobic metabolism of glucose, produced levan, and none produced oxidase, arginine dehydrolase and pectinase. All bacterial isolates caused a hypersensitive reaction on the tobacco leaves. These reactions correspond to LOPAT Ia group which includes *P. syringae* pathovars.

All isolates were positive for gelatin and aesculin hydrolysis, but negative for tyrosinase activity and metabolism of tartrate, demonstrating typical characteristics for bacteria *P. syringae* pv. *syringae*.

**Genetic characterization.** PCR amplification with primers B1 and B2 amplified the specific 752 bp band for the *syrB* gene in all tested 43 isolates and two reference *P. syringae* pv. *syringae* strains, CFBP 11 and CFBP 1582. The strain of *P. syringae* pv. *morsprunorum* CFBP 2119 failed to show amplification and hence did not possess the *syrB* gene.

To determine genetic diversity of the 43 *P. syringae* pv. *syringae* isolates, ERIC-PCR genomic fingerprints were generated. DNA banding patterns were identical among tested isolates and *P. syringae* pv. *syringae* reference strains, and did not show any genetic diversity among them. Considering the fact that there was no difference among the isolates, a further genetic study was performed with a reduced number of isolates (12 in total) on the basis of the geographic origin of the isolates.

Molecular identification of isolates from mandarin was performed using 16SrDNA gene sequencing and MLSA. The sequence analysis of the conserved 16SrDNA gene deposited in NCBI GenBank database under accession numbers KP099969 to KP099980 (Table 2), confirmed that isolates originated from mandarin belong to *P. syringae* pv. *syringae* showing 100% homology with *P. syringae* pv. *syringae* strain ICMP 3023 from GenBank (accession number NR\_117820).

The sequences generated with three housekeeping genes (*gyrB*, *rpoD*, and *gapI*), were deposited in the GenBank database under accession numbers KP099981 to KP100028 (Table 2). A NJ tree constructed for the concatenated data set showed that the mandarin isolates were grouped together with the following reference strains from the database; *P. syringae* pv. *aceris* LMG 2106, *P. syringae* pv. *atofaciens* LMG 5095, *P. syringae* pv. *coryli* NCBI 4273, *P. syringae* pv. *japonica* MAFF 301072, *P. syringae* pv. *lapsa* LMG 2206, *P. syringae* pv. *pisi* NCP-PB 2585, *P. syringae* pv. *solidagae* ICMP 16925, and *P. syringae* pv. *syringae* LMG 1247 (Fig. 2).

**Table 2.** Bacterial isolates and strains used in this study and their geographic origin and GenBank accession numbers

Isolates and reference strains	Species and pathovars	Area of origin	GenBank accession numbers			
			16srDNA	<i>gyrB</i>	<i>rpoD</i>	<i>GapI</i>
IZB 200 (50)*	<i>Pseudomonas syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	KP099969	KP100017	KP099981	KP100005
IZB 201 (50)	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	KP099970	KP100018	KP099982	KP100006
IZB 202	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	-	-	-	-
IZB 203	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	-	-	-	-
IZB 204 (50)	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	KP099971	KP100019	KP099983	KP100007
IZB 205	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	-	-	-	-
IZB 206	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	-	-	-	-
IZB 207	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Dobra voda)	-	-	-	-
IZB 208 (50)	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	KP099972	KP100020	KP099984	KP100008
IZB 209 (50)	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	KP099973	KP100021	KP099985	KP100009
IZB 210	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	-	-	-	-
IZB 211	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	-	-	-	-
IZB 212 (50)	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	KP099974	KP100022	KP099986	KP100010
IZB 213	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	-	-	-	-
IZB 214 (50)	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	KP099975	KP100023	KP099987	KP100011
IZB 215	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	-	-	-	-
IZB 216	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	-	-	-	-
IZB 217	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Bjeliši)	-	-	-	-
IZB 218 (50)	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Ilino)	KP099976	KP100024	KP099988	KP100012
IZB 219 (50)	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Ilino)	KP099977	KP100025	KP099989	KP100013
IZB 220	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Ilino)	-	-	-	-
IZB 221 (50)	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Ilino)	KP099978	KP100026	KP099990	KP100014
IZB 222	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Ilino)	-	-	-	-
IZB 223	<i>P. syringae</i> pv. <i>syringae</i>	Bar (Ilino)	-	-	-	-
IZB 224	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 225	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 226	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 227	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 228 (50)	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	KP099979	KP100027	KP099991	KP100015
IZB 229	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 230	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 231 (50)	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	KP099980	KP100028	KP099992	KP100016
IZB 232	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 233	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 234	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 235	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 236	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 237	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 238	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 239	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 240	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 241	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
IZB 242	<i>P. syringae</i> pv. <i>syringae</i>	Ulcinj (Štoj)	-	-	-	-
LMG 2106 (45)	<i>P. syringae</i> pv. <i>aceris</i>	United Kingdom	-	-	-	-
ICMP 3023	<i>P. syringae</i> pv. <i>syringae</i>	United Kingdom	-	-	-	-
MAFF301072 (47)	<i>P. syringae</i> pv. <i>japonica</i>	Japan	-	-	-	-

Table 2. Continued

Isolates and reference strains	Species and pathovars	Area of origin	GenBank accession numbers			
			16srDNA	<i>gyrB</i>	<i>rpoD</i>	<i>GapI</i>
ATCC19322 (13)	<i>P. syringae</i> pv. <i>morsprunorum</i>	USA	-	-	-	-
Cit7 (42)	<i>P. syringae</i>	-	-	-	-	-
CFBP 2103 (4)	<i>P. syringae</i> pv. <i>apii</i>	USA	-	-	-	-
CFBP 1573 (1)	<i>P. syringae</i> pv. <i>persicae</i>	France	-	-	-	-
CFBP 1727 (5)	<i>P. syringae</i> pv. <i>berberidis</i>	New Zealand	-	-	-	-
CFBP 2341 (16)	<i>Pseudomonas cannabina</i> pv. <i>cannabina</i>	Hungary	-	-	-	-
CFBP 6866 (15)	<i>P. cannabina</i> pv. <i>alisalensis</i>	USA	-	-	-	-
NCPPB 600 (56)	<i>P. syringae</i> pv. <i>coronafaciens</i>	United Kingdom	-	-	-	-
LMG 5095 (48)	<i>P. syringae</i> pv. <i>atrofaciens</i>	New Zealand	-	-	-	-
ICMP 12471 (14)	<i>P. syringae</i> pv. <i>coriandricola</i>	Germany	-	-	-	-
LMG 1247 (51)	<i>P. syringae</i> pv. <i>syringae</i>	United Kingdom	-	-	-	-
LMG 2184 (35)	<i>P. syringae</i> pv. <i>eriobotryae</i>	USA	-	-	-	-
LMG 2206 (49)	<i>P. syringae</i> pv. <i>lapsa</i>	-	-	-	-	-
LMG 2245 (26)	<i>P. syringae</i> pv. <i>phaseolicola</i>	Canada	-	-	-	-
LMG 2276 (19)	<i>P. syringae</i> pv. <i>ribicola</i>	-	-	-	-	-
LMG 2289 (30)	<i>P. syringae</i> pv. <i>sesami</i>	Yugoslavia	-	-	-	-
LMG 2349 (34)	<i>P. syringae</i> pv. <i>ulmi</i>	Yugoslavia	-	-	-	-
LMG 2351 (8)	<i>P. syringae</i> pv. <i>vibruni</i>	USA	-	-	-	-
LMG 5062 (41)	<i>P. syringae</i> pv. <i>dysoxli</i>	New Zealand	-	-	-	-
LMG 5064 (54)	<i>P. syringae</i> pv. <i>garcae</i>	Brazil	-	-	-	-
LMG 5066 (25)	<i>P. syringae</i> pv. <i>glycinea</i>	New Zealand	-	-	-	-
LMG 5067 (37)	<i>P. syringae</i> pv. <i>helianthi</i>	Mexico	-	-	-	-
LMG 5072 (29)	<i>P. syringae</i> pv. <i>mellea</i>	Japan	-	-	-	-
LMG 5074 (28)	<i>P. syringae</i> pv. <i>mori</i>	Hungary	-	-	-	-
LMG 5076 (43)	<i>P. syringae</i> pv. <i>papulans</i>	Canada	-	-	-	-
NCPPB 2585 (52)	<i>P. syringae</i> pv. <i>pisi</i>	New Zealand	-	-	-	-
LMG 5090 (38)	<i>P. syringae</i> pv. <i>tagetis</i>	Zimbabwe	-	-	-	-
LMG 5092 (11)	<i>P. syringae</i> pv. <i>theae</i>	Japan	-	-	-	-
LMG 5185 (9)	<i>P. syringae</i> pv. <i>passiflorae</i>	New Zealand	-	-	-	-
LMG 5381 (7)	<i>P. syringae</i> pv. <i>delphini</i>	New Zealand	-	-	-	-
LMG 5541 (40)	<i>P. syringae</i> pv. <i>ciccaronei</i>	Italy	-	-	-	-
LMG 5668 (40)	<i>P. syringae</i> pv. <i>myricae</i>	Japan	-	-	-	-
LMG 10912 (55)	<i>P. syringae</i> pv. <i>oryzae</i>	Japan	-	-	-	-
CFBP 2212 (6)	<i>P. syringae</i> pv. <i>tomato</i>	United Kingdom	-	-	-	-
CFBP 1657 (3)	<i>P. syringae</i> pv. <i>maculicola</i>	United Kingdom	-	-	-	-
CFBP 6463 (21)	<i>P. syringae</i> pv. <i>lachrymans</i>	Hungary	-	-	-	-
NCPPB 3257 (17)	<i>P. syringae</i> pv. <i>philadelphii</i>	United Kingdom	-	-	-	-
NCPPB 3688 (24)	<i>P. syringae</i> pv. <i>photiniae</i>	Japan	-	-	-	-
NCPPB 1427 (32)	<i>P. syringae</i> pv. <i>tabaci</i>	Hungary	-	-	-	-
NCPPB 3618 (39)	<i>P. syringae</i> pv. <i>rhapiolepidis</i>	Japan	-	-	-	-
CFBP 6109 (23)	<i>P. syringae</i> pv. <i>cerasicola</i>	Japan	-	-	-	-
ICMP 13650 (27)	<i>P. syringae</i> pv. <i>broussonetiae</i>	Japan	-	-	-	-
NCPPB 3681 (33)	<i>P. syringae</i> pv. <i>aesculi</i>	India	-	-	-	-
ICMP 16925 (44)	<i>P. syringae</i> pv. <i>solidagae</i>	Japan	-	-	-	-
ICMP 16929 (2)	<i>P. syringae</i> pv. <i>spinaceae</i>	Japan	-	-	-	-
ICMP 11894 (36)	<i>P. syringae</i> pv. <i>cunninghamiae</i>	China	-	-	-	-



**Table 2.** Continued

Isolates and reference strains	Species and pathovars	Area of origin	GenBank accession numbers			
			16srDNA	<i>gyrB</i>	<i>rpoD</i>	<i>GapI</i>
NCPPB 3682 (31)	<i>P. syringae</i> pv. <i>hibisci</i>	USA	-	-	-	-
NCPPB 3690 (53)	<i>P. syringae</i> pv. <i>zizaniae</i>	USA	-	-	-	-
LMG 5030 (57)	<i>P. syringae</i> pv. <i>atropurpurea</i>	Japan	-	-	-	-
NCPPB 4273 (46)	<i>P. syringae</i> pv. <i>coryli</i>	Italy	-	-	-	-
NCPPB 3487 (12)	<i>P. syringae</i> pv. <i>avellanae</i>	Greece	-	-	-	-
LMG 2209 (22)	<i>P. syringae</i> pv. <i>savastanoi</i>	Yugoslavia	-	-	-	-
NCPPB 3617 (20)	<i>P. syringae</i> pv. <i>daphniphylli</i>	Japan	-	-	-	-
LMG 2252 (18)	<i>P. syringae</i> pv. <i>primulae</i>	United Kingdom	-	-	-	-
NCPPB 3739 (10)	<i>P. syringae</i> pv. <i>actinidiae</i>	Japan	-	-	-	-
Pf-5	<i>Pseudomonas fluorescens</i>	-	-	-	-	-

-, no data.

\*Number of the strain in minimum spanning tree (MST) (Fig. 3).

An MST using the MLSA data using statistical parsimony with split network revealed no ambiguous connections between pathovars and confirmed clustering of testing isolates and reference strains associated within the same genomospecies 1 (Fig. 3). The first cluster (colored in black) included all of our 43 mandarin isolates and members of genomospecies 1. The second cluster included members of genomospecies 2 (colored in blue). The other six clusters present the strains which belong in genomospecies 3, 4 and 6–9 (represented in red, green, light blue, orange, white-green, and white-red, respectively).

## Discussion

The occurrence of citrus blast disease, caused by *P. syringae* pv. *syringae* in Montenegro was reported for the first time by Vučinić (1987), but a detailed analysis of the pathogen population structure was not performed until now. Based on the observations during epidemics on mandarin in Montenegro and on earlier findings in literature related to environmental influences of citrus blast disease, it is postulated that *P. syringae* pv. *syringae* probably easily colonizes buds when spring is cool and moist and enters through wounds when the tissue is previously damaged by frost (Cao et al., 1999; Crosse, 1966; Gross et al., 1983; Klement et al., 1984; Nejad et al., 2004; Sule and Seemuller, 1987).


The relationship of frost damage and *P. syringae* pv. *syringae* diseases of woody plants has been demonstrated previously in pear blossom blight (Panagopoulos and Crosse, 1964), leaf spot of sour cherry (Sule and Seemuller, 1987), and bacterial canker of peach (Cao et al., 1999), apricot (Klement et al., 1984), prune, almond (Cao et al., 1999), and willow (Nejad et al., 2004). It has

been suggested that winter damage is a contributing factor to outbreaks on many host plants (Buttner and Amy, 1989; Cambours et al., 2005; Gross et al., 1988; Kennelly et al., 2007; Lindow et al., 1982; Sule and Seemuller, 1987). In cases where frost damage incites a greater degree of blast and black pit citrus diseases, cultural management strategies against frost damage may be a viable method of minimizing losses. If fruits and leaves were kept damp using sprinklers, it would allow thin layers of ice to form on plant surfaces rather than inside them where the tissue can get damaged. As ice crystals form, the latent heat of water is released and prevents frost damage (Cambours et al., 2005; Gross et al., 1988; Kennelly et al., 2007; Lindow et al., 1982).


In our study 43 isolates originated from diseased necrotic mandarin buds were identified and characterized using pathogenicity, biochemical tests and for the first time molecular analysis of pathogen. Based on obtained results, they could be assigned to the phytopathogenic bacterium *P. syringae* pv. *syringae*. This bacterium has been previously reported as the causal agent of citrus blast of mandarin in Italy, Japan, Iran and Turkey (De Cicco et al., 1978; Mirik et al., 2005; Salerno and Cutuli, 1985; Shams-Bakhsh and Rahimian, 1997).

Using biochemical and GATT differentiation tests our isolates could be discriminated as *P. syringae* pv. *syringae*, showing reactions typical for *P. syringae* pv. *syringae* as it is described by Lelliott and Stead (1987).

All isolates produced black pit lesions on lemons by needle pricks. The characteristics that positive lesions produced suggest that lemons might be readily inoculated with all strains of this bacteria obtained from any host (Bryan, 1928; Lelliott and Stead, 1987; Smith and Fawcett, 1930). Typical black pit lesions were also formed on



**Fig. 2.** Neighbor joining phylogenetic tree of *Pseudomonas syringae* pv. *syringae* isolates from mandarin and reference strains derived from a concatenation of three housekeeping genes (*gyrB*, *rpoD*, and *gapI*). Bar—estimated nucleotide substitutions per site is 0.02.



**Fig. 3.** Mini spanning tree splits genealogical networks for 12 *Pseudomonas syringae* pv. *syringae* isolates originating from mandarin colored in white-black (50) and representative pathotype *P. syringae* strains. Colored in black genomospecies 1, colored in blue genomospecies 2, colored in red genomospecies 3, colored in green genomospecies 4, colored in light blue to genomospecies 6, colored in orange genomospecies 7, colored in white-green genomospecies 8, colored in white-red genomospecies 9.

lilac and mandarin leaves and leaf petioles and on mandarin twigs as is described by Bryan (1928).

*P. syringae* pv. *syringae* is a known bacterium which produces different phytotoxins depending on the host origin. Isolates from the stone fruits or peach produced syringomycin, isolates from citrus trees yielded syringotoxin while isolates from lilac blight produced syringostatin (Fukuchi et al., 1990; Gross and Cody, 1985; Isogai et al., 1989; Mitchell, 1981; Mo and Gross, 1991). Syringomycin was detected in our tested isolates originating from mandarin.

It should be noted that the lack of molecular data in literature sources concerning the genetic identity and structure of *P. syringae* pv. *syringae* populations is evident. The present study based on ERIC-PCR and 16S rDNA gene sequencing showed a genetically homogeneous population of *P. syringae* pv. *syringae* causing citrus blast disease in Montenegro. A phylogenetic tree obtained from MLSA, showed that our isolates belong to the previously described genomospecies 1 of *P. syringae* (Berge et al., 2014; Gardan et al., 1999; Sarkar and Guttman, 2004).

Results of MST showed that *P. syringae* pv. *japonica* is defined as a central node of the cluster and point of diversification for other strains. *P. syringae* pv. *syringae* isolates from mandarin were closely related to other *P. syringae* pvs. such as *atrofaciens*, *lapsa*, *syringae*, *japonica*, and *pisi*. We consider that hybridization events could have occurred between them or their ancestors. MST shows, from the clustering of the *P. syringae* pv. *syringae* isolates

originated from mandarin, that these isolates are genetically diversified from *P. syringae* pv. *syringae* pathotype strain LMG 1247. *P. syringae* pv. *syringae* isolates from mandarin represent a clearly clonal complex and further investigation is needed to understand the origin of this pathogen. In conclusion, our study revealed that *P. syringae* pv. *syringae* isolates originated from mandarin which caused citrus blast in Montenegro and that they belong to genomospecies 1. The homogeneity found in all used tests suggest that tested isolates may have originated from a single inoculum source such as planting material.

Further detailed biological studies, using more strains of *P. syringae* pv. *syringae* which cause citrus blast in other parts of the world should be performed to elucidate their origin and to further resolve their taxonomic status. This might be very important for efficient management of outbreaks and the prevention of further spreading of this pathogen.

## Acknowledgments

This work was supported by the Ministry of Education and Science, Republic of Serbia (Grants No. TR31018) and the Ministry of Science, Montenegro (Grant “Invasive species”).

## References

- Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* 25:3389-3402.
- Ausubel, F. M., Brent, R., Kingston, R. E., Moore, D. D., Seidman, J. G., Smith, J. A. and Struhl, K. 1992. Current protocols in molecular biology. Vol. I. Greene Publishing Associates and Wiley-Interscience, New York, NY, USA.
- Berge, O., Monteil, C. L., Bartoli, C., Chandeysson, C., Guilbaud, C., Sands, D. C. and Morris, C. E. 2014. A user's guide to a data base of the diversity of *Pseudomonas syringae* and its application to classifying strains in this phylogenetic complex. *PLoS One* 9:e105547.
- Bertani, G. 1951. Studies on lysogenesis. I. The mode of phage liberation by lysogenic *Escherichia coli*. *J. Bacteriol.* 62: 293-300.
- Bradbury, J. F. 1986. Guide to plant pathogenic bacteria. CAB International, Farnham Royal, Great Britain.
- Bryan, M. K. 1928. Lilac blight in the United States. *J. Agric. Res.* 36:225-235.
- Bull, C. T., Clarke, C. R., Cai, R., Vinatzer, B. A., Jardini, T. M. and Koike, S. T. 2011. Multilocus sequence typing of *Pseudomonas syringae* sensu lato confirms previously described genomospecies and permits rapid identification of *P. syringae* pv. *coriandricola* and *P. syringae* pv. *apii* causing bacterial leaf spot on parsley. *Phytopathology* 101:847-858.
- Buttner, M. P. and Amy, P. S. 1989. Survival of ice nucleation-active and genetically engineered non-ice-nucleating *Pseudomonas syringae* strains after freezing. *Appl. Environ. Microbiol.* 55:1690-1694.
- Cambours, M. A., Nejad, P., Granhall, U. and Ramstedt, M. 2005. Frost-related dieback of willows. Comparison of epiphytically and endophytically isolated bacteria from different *Salix* clones, with emphasis on ice nucleation activity, pathogenic properties and seasonal variation. *Biomass Bioenerg.* 28:15-27.
- Cao, T., Sayler, R. J., Dejong, T. M., Kirkpatrick, B. C., Bostock, R. M. and Shackel, K. A. 1999. Influence of stem diameter, water content, and freezing-thawing on bacterial canker development in excised stems of dormant stone fruit. *Phytopathology* 89:962-966.
- Crosse, J. E. 1966. Epidemiological relations of the pseudomonad pathogens of deciduous fruit trees. *Annu. Rev. Phytopathol.* 4:291-310.
- de Bruijn, F. J. 1992. Use of repetitive (repetitive extragenic palindromic and enterobacterial repetitive intergeneric consensus) sequences and the polymerase chain reaction to fingerprint the genomes of *Rhizobium meliloti* isolates and other soil bacteria. *Appl. Environ. Microbiol.* 58:2180-2187.
- De Cicco, V., Luisi, N. and Salerno, M. 1978. Epidemiology and control of citrus blast. In: *Proceedings of third meeting of the International Society of Citriculture*, pp. 204-208. August 15-23, 1978, International Society of Citriculture, Sydney, Australia.
- Fukuchi, N., Isogai, A., Nakayama, J. and Suzuki, A. 1990. Structure of syringotoxin B, a phytotoxin produced by citrus isolates of *Pseudomonas syringae* pv. *syringae*. *Agric. Biol. Chem.* 54:3377-3379.
- Gardan, L., Shafik, H., Belouin, S., Broch, R., Grimont, F. and Grimont, P. A. 1999. DNA relatedness among the pathovars of *Pseudomonas syringae* and description of *Pseudomonas tremae* sp. nov. and *Pseudomonas cannabina* sp. nov. (ex Suttic and Dowson 1959). *Int. J. Syst. Evol. Bacteriol.* 49:469-478.
- Gironde, S. and Manceau, C. 2012. Housekeeping gene sequencing and multilocus variable-number tandem-repeat analysis to identify subpopulations within *Pseudomonas syringae* pv. *maculicola* and *Pseudomonas syringae* pv. *tomato* that correlate with host specificity. *Appl. Environ. Microbiol.* 78:3266-3279.
- Gorlenko, M. V. 1965. Bacterial diseases of plants. 2nd ed. Israel Program for Scientific Translations, Jerusalem, Israel.
- Grifoni, A., Bazzicalupo, M., Di Serio, C., Fancelli, S. and Fani, R. 1995. Identification of *Azospirillum* strains by restriction fragment length polymorphism of the 16S rDNA and of the histidine operon. *FEMS Microbiol. Lett.* 127:85-91.
- Gross, D. C. and Cody, Y. S. 1985. Mechanisms of plant pathogenesis by *Pseudomonas* species. *Can. J. Microbiol.* 31:403-410.
- Gross, D. C., Cody, Y. S., Proebstring, E. L., Radamaker, G. K. and Spotts, R. A. 1983. Distribution, population dynamics, and characteristics of ice nucleation-active bacteria in deciduous fruit tree orchards. *Appl. Environ. Microbiol.* 46:1370-

- 1379.
- Gross, D. C., Proebstring, E. L. and MacCrimble-Zimmerman, H. 1988. Development, distribution, and characteristics of intrinsic, nonbacterial ice nuclei in *Prunus* wood. *Plant Physiol.* 88:915-922.
- Hilario, E., Buckley, T. R. and Young, J. M. 2004. Improved resolution on the phylogenetic relationships among *Pseudomonas* by the combined analysis of *atpD*, *carA*, *recA* and 16S rDNA. *Antonie van Leeuwenhoek* 86:51-64.
- Holding, A. J. and Collee, J. G. 1971. Routine biochemical tests. In: *Methods in microbiology*, Vol. 6, eds. by J. R. Norris and D. W. Ribbons, pp. 2-30. Academic Press, New York, NY, USA.
- Huson, D. H. and Bryant, D. 2006. Application of phylogenetic networks in evolutionary studies. *Mol. Biol. Evol.* 23:254-267.
- Isogai, A., Fukuchi, N., Yamashita, S., Suyama, K. and Suzuki, A. 1989. Syringostatins, novel phytotoxins produced by *Pseudomonas syringae* pv. *syringae*. *Agric. Biol. Chem.* 53: 3117-3119.
- Kaluzna, M., Janse, J. D. and Young, J. M. 2012. Detection and identification methods and new tests as developed and used in the framework of COST 873 for bacteria pathogenic to stone fruits and nuts. *J. Plant Pathol.* 94(1 Suppl):S1.117-S1.126.
- Kennelly, M. M., Cazorla, F. M., de Vicente, A., Ramos, C. and Sundin, G. W. 2007. *Pseudomonas syringae* diseases of fruit trees: progress toward understanding and control. *Plant Dis.* 91:4-17.
- King, E. O., Ward, M. K. and Raney, D. E. 1954. Two simple media for the demonstration of pyocyanin and fluorescin. *J. Lab. Clin. Med.* 44:301-307.
- Klement, Z. 1990. Inoculation of plant tissues. Cancer and dieback disease. In: *Methods in phytobacteriology*, eds. by Z. Klement, K. Rudolph and D. C. Sands, pp. 105-106. Akademiai Kiado, Budapest, Hungary.
- Klement, Z., Rozsnyay, D. S., Báló, E., Pánczél, M. and Prilesky, G. 1984. The effect of cold on development of bacterial canker in apricot trees infected with *Pseudomonas syringae* pv. *syringae*. *Physiol. Plant Pathol.* 24:237-246.
- Kovacs, N. 1956. Identification of *Pseudomonas pyocyanea* by the oxidase reaction. *Nature* 178:703.
- Lelliott, R. A. and Stead, D. E. 1987. Methods for the diagnosis of bacterial diseases of plants. Blackwell Scientific Publications, Oxford, UK.
- Lindow, S. E., Arny, D. C. and Upper, C. D. 1982. Bacterial ice nucleation: a factor in frost injury to plants. *Plant Physiol.* 70:1084-1089.
- Lupski, J. R. and Weinstock, G. M. 1992. Short, interspersed repetitive DNA sequences in prokaryotic genomes. *J. Bacteriol.* 174:4525-4529.
- Mirik, M., Baloglu, S., Aysan, Y., Cetinkaya-Yildiz, R., Kusek, M. and Sahin, F. 2005. First outbreak and occurrence of citrus blast disease, caused by *Pseudomonas syringae* pv. *syringae*, on orange and mandarin trees in Turkey. *Plant Pathol.* 54:238.
- Mitchell, R. E. 1981. Structure: bacterial. In: *Toxins in plant diseases*, ed. by R. D. Durbin, pp. 259-291. Academic Press, New York, NY, USA.
- Mo, Y. Y. and Gross, D. C. 1991. Plant signal molecules activate the *syrB* gene, which is required for syringomycin production by *Pseudomonas syringae* pv. *syringae*. *J. Bacteriol.* 173:5784-5792.
- Nejad, P., Ramstedt, M. and Granhall, U. 2004. Pathogenic ice-nucleation active bacteria in willows for short rotation forestry. *For. Pathol.* 34:369-381.
- Panagopoulos, C. G. and Grosse, J. E. 1964. Frost injury as a predisposing factor in blossom blight of pear caused by *Pseudomonas syringae* van Hall. *Nature* 202:1352.
- Perrot, P. 1998. A to Z of thermodynamics. Oxford University Press, Oxford, UK.
- Salerno, M. and Cutuli, G. 1985. Le malattie degli agrumi: batteriosi (blast and black pit). *Inf. Fitopatol.* 35:27-28.
- Sarkar, S. F. and Guttman, D. S. 2004. Evolution of the core genome of *Pseudomonas syringae*, a highly clonal, endemic plant pathogen. *Appl. Environ. Microbiol.* 70:1999-2012.
- Schaad, N. W., Jones, J. B. and Chun, W. 2001. Laboratory guide for identification of plant pathogenic bacteria. 3rd ed. APS Press, St. Paul, MN, USA. 373 pp.
- Scortichini, M., Marchesi, U., Dettori, M. T. and Rossi, M. P. 2003. Genetic diversity, presence of the *syrB* gene, host preference and virulence of *Pseudomonas syringae* pv. *syringae* strains from woody and herbaceous host plants. *Plant Pathol.* 52:277-286.
- Scortichini, M., Rossi, M. P., Loreti, S., Bosco, A., Fiori, M., Jackson, W., Stead, D. E., Aspin, A., Marchesi, U., Zini, M. and Janse, J. D. 2005. *Pseudomonas syringae* pv. *coryli*, the causal agent of bacterial twig dieback of *Corylus avellana*. *Phytopathology* 95:1316-1324.
- Shams-Bakhsh, M. and Rahimian, H. 1997. Comparative study on agents of citrus blast and bacterial canker of stone fruits in Mazandaran. *Iran. J. Plant Pathol.* 33:132-143.
- Shigeta, S. and Nakata, E. 1995. Bacterial brown spot of citrus caused by *Pseudomonas syringae* pv. *syringae* van Hall 1902. *J. Ann. Phytopathol. Soc. Jpn.* 61:150-157.
- Smith, C. O. and Fawcett, H. S. 1930. A comparative study of the citrus blast bacterium and some other allied organisms. *J. Agric. Res.* 41:233-246.
- Sorensen, K. N., Kim, K. H. and Takemoto, J. Y. 1998. PCR detection of cyclic lipodepsinonapeptide-producing *Pseudomonas syringae* pv. *syringae* and similarity of strains. *Appl. Environ. Microbiol.* 61:226-230.
- Sule, S. and Seemuller, E. 1987. The role of ice formation in the infection of sour cherry leaves by *Pseudomonas syringae* pv. *syringae*. *Phytopathology* 77:173-177.
- Tamura, K., Stecher, G., Peterson, D., Filipowski, A. and Kumar, S. 2013. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol. Biol. Evol.* 30:2725-2729.
- Thomidis, T., Tsiouridis, C., Exadaktylou, E. and Drogoudi, P. 2005. Comparison of three laboratory methods to evaluate the pathogenicity and virulence of ten *Pseudomonas syringae* pv. *syringae* strains on apple, pear, cherry and peach trees. *Phytoparasitica* 33:137-140.
- Thornley, M. J. 1960. The differentiation of *Pseudomonas* from



- other Gram-negative bacteria on the basis of arginine metabolism. *J. Appl. Bacteriol.* 23:37-52.
- Timmer, L. W., Garnsey, S. M. and Graham, J. H. 2000. Compendium of citrus diseases. APS Press, St. Paul, MN, USA.
- Vanarelli, S., Rizzo, D., Stefani, L. and Paoli, M. (2010 onwards). Avvizzimento batterico degli agrumi--Piticchia batterica (Citrus blast/black pit). URL [http://www.cespevi.it/servfito/pdf/Scheda\\_Pseudomonas\\_syringae.pdf](http://www.cespevi.it/servfito/pdf/Scheda_Pseudomonas_syringae.pdf) [25 January 2014].
- Vučinić, Z. 1987. La necrose bactérienne des agrumes sur le littoral monténégrin. In: *50 Years of Agricultural Institute Titograd*, ed. by Ž. Kalezić, pp. 59-67. Agricultural Institute, Titograd, Yugoslavia (in Serbian).
- Whiteside, J. O., Garnsey, S. M. and Timmer, L. W. 1988. Compendium of citrus diseases. APS Press, St. Paul, MN, USA.