

Note Added in Proof. When the reaction of **1** with PhSCl in a 1:3 molar ratio is carried out in the absence of a solvent, the yield of the diazene **2e** is $\geq 90\%$.

Acknowledgment. We thank the NSERC (Canada) for support in the form of operating and infrastructure grants, Mr. Glen Bigam (University of Alberta) for recording the ESR spectra, and Drs. A. Rauk and T. G. Back for helpful discussions. V.C. and M. N.S.R. thank IIT Kanpur and IIT Madras, respectively, for a sabbatical leave.

Registry No. **1**, 24261-90-3; **2a**, 136276-65-8; **2b**, 136276-66-9; **2c**, 136276-67-0; **2d**, 136276-68-1; **2e**, 136276-69-2; **3a**, 136276-70-5; **3b**,

136276-71-6; **3c**, 136276-72-7; **4b**, 136276-73-8; **4c**, 136276-74-9; **5a**, 86602-13-3; **5b**, 136396-53-7; **6a**, 136276-75-0; **6b**, 136276-76-1; **6c**, 136276-77-2; PhSCl, 931-59-9; Cl₃CSCl, 594-42-3; 2,4-(NO₂)₂C₆H₃SCl, 528-76-7; Cl₃SeMe, 37826-07-6; TMS-N=C[N(TMS)₂]C₆H₄-*p*-Me, 117357-77-4; 2,4-(NO₂)₂C₆H₃SN=C(Ph)NH₂, 136276-78-3; TMS-N=C[N(TMS)₂]C₆H₄-*p*-CF₃, 117357-81-0; 2,4-(NO₂)₂C₆H₃SN=C(Ph)N=C(Ph)=NS-2,4-(NO₂)₂C₆H₃, 136276-79-4; PhSeCl, 5707-04-0; PhSeCl₃, 42572-42-9.

Supplementary Material Available: For **4b**, listings of non-hydrogen and hydrogen coordinates, anisotropic thermal parameters, bond distances, bond angles, torsion angles, and experimental details for the X-ray analyses (7 pages); listings of structure factor amplitudes (23 pages). Ordering information is given on any current masthead page.

Contribution from the Institut für Anorganische Chemie, Universität des Saarlandes, WD-6600 Saarbrücken, Germany

Halide-Complexing Characteristics of Germa- and Stannamine Cages


Michael Veith,* Joachim Fischer, Timothy R. Prout, Martin Nötzel, Peter Hobein, and Volker Huch

Received April 30, 1991

The cations $M_3(N-t-Bu)_4H_3^+$ ($M = Ge, Sn$) coordinate very efficiently via hydrogen bridges to chloride, bromide, and iodide anions. These molecular complexes have the general formula $M_3(N-t-Bu)_4H_3X$ ($M = Ge, X = Cl$ (**1**), Br (**2**), I (**3**); $M = Sn, X = Cl$ (**4**), Br (**5**), I (**6**)) and are obtained by three different methods. The newly synthesized compounds have been thoroughly characterized by IR, NMR, and mass spectroscopy. X-ray crystal structure determinations of **1**, **3**, **4**, and **5** reveal the molecules to be built of a M_3N_4 -norcubane-type skeleton, which is attached by nitrogen-hydrogen bonds to a H_3X trigonal pyramid. The overall closed cage " $M_3N_4H_3X$ " has an approximate $3m$ (C_{3v}) symmetry with one nitrogen and the halogen atom situated on the 3-fold axis. The compounds **3** and **4** crystallize isotypically with 0.5 benzene/molecule in the space group $R\bar{3}$ with $a = b = 11.990$ (7) Å, $c = 34.40$ (1) Å, $\alpha = \beta = 90^\circ$, $\gamma = 120^\circ$, $Z = 6$, and $V = 4283$ (2) Å³ and $a = b = 11.720$ (8) Å, $c = 35.109$ (15) Å, $\alpha = \beta = 90^\circ$, $\gamma = 120^\circ$, $Z = 6$, and $V = 4176$ (2) Å³, respectively, with the final R factors being 0.057 (1369 reflections) and 0.032 (1152 reflections). Compound **1** crystallizes in the monoclinic space group $P2_1/n$ with $a = 10.188$ (6) Å, $b = 19.923$ (8) Å, $c = 12.099$ (7) Å, $\beta = 91.54$ (5)°, $Z = 4$, and $V = 2455$ (1) Å³. A total of 2859 independent data were collected, giving a final R factor of 0.035. Compound **5** is orthorhombic (space group $Pnma$) with $a = 21.471$ (9) Å, $b = 11.753$ (6) Å, $c = 10.476$ (5) Å, $Z = 4$, and $V = 2644$ (1) Å³. Refinement of this structure, using 1921 independent reflections, gave an R factor of 0.033. The compounds can be described as being composed of a spherical lipophilic part ($M_3(N-t-Bu)_4$) and a somewhat protruding lipophobic part (H_3X pyramid). These properties seem to account for the ease with which these cages exchange the halogen anions in metathesis reactions with simple salts like NaBr, NaI, and TlCl.

Introduction

Of the molecular amino and imino cages of Ge(II), Sn(II) and Pb(II),¹ obtained by aminolysis of bis(amino)germylenes, -stannylenes, and -plumbylenes,² the compounds $Ge_3(N-t-Bu)_4H_2$ and $Sn_3(N-t-Bu)_4H_2$ are peculiar in several respects. They form "open" polycycles (i.e. the skeleton of the molecules may be visualized as a cube with a missing corner, the seven remaining corners being alternately occupied by Ge/Sn and N atoms; for other compounds of this type see also ref 3) and show bond fluctuation behavior as established by variable-temperature NMR studies.⁴ One of these movements is a mutual intramolecular displacement of the two hydrogen atoms with respect to the three nitrogen atoms in the "open" corner of the cube. Taking the crystallographic dimensions of $Sn_3(N-t-Bu)_4H_2$ into account,⁵ it seemed clear to us that by introduction of a simple proton this intramolecular displacement should be arrested, forming a cation $Sn_3(N-t-Bu)_4H_3^+$. The whole charged entity should behave as an excellent counterion to simple anions, as the three hydrogen atoms are pointing rigidly in one direction (see formula A). Since macrocyclic counterions or cryptands for simple anions like chloride, bromide, and iodide are less commonly available than for cationic species,⁶⁻¹⁰ it seemed



interesting to us to study the potential tridentate function of $Ge_3(N-t-Bu)_4H_3^+$ or $Sn_3(N-t-Bu)_4H_3^+$ toward these halogen anions. At the same time we intended to study the properties of the neutral complexes from a structural and chemical point of view.

Experimental Section

Apparatus and Materials. All operations were carried out under an inert atmosphere of nitrogen by using a modified Stock vacuum apparatus and Schlenk techniques. Infrared, ¹H NMR, and mass spectra were obtained by using Perkin-Elmer IR 883, Bruker WP 80 (80 MHz), and Finnigan MAT 90 (DCI⁺ using isobutane) spectrometers, respectively. ¹H NMR chemical shifts as δ values were measured relative to internal Me₄Si. Elemental analyses were performed by Beller, Mi-

- (1) (a) Veith, M. *Chem. Rev.* **1990**, *90*, 3. (b) Veith, M. *Angew. Chem.* **1987**, *99*, 1; *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 1.
- (2) Veith, M. *Comments Inorg. Chem.* **1985**, *4*, 179.
- (3) Holmes, R. R. *Acc. Chem. Res.* **1989**, *22*, 190.
- (4) Veith, M. *Phosphorus, Sulfur and Silicon* **1989**, *41*, 195.
- (5) Veith, M. *Z. Naturforsch.* **1980**, *35B*, 20.
- (6) (a) Lehn, J. M. *Angew. Chem.* **1990**, *102*, 1347. (b) Hosseini, M. W.; Blacker, A. J.; Lehn, J. M. *J. Am. Chem. Soc.* **1990**, *112*, 3896. (c) Schmidchen, F. P. *Nachr. Chem. Tech. Lab.* **1988**, *36*, 8. (d) Hosseini, M. W.; Lehn, J. M. *Helv. Chim. Acta* **1988**, *71*, 749. (e) Lehn, J. M.; Sonveaux, E.; Willard, A. K. *J. Am. Chem. Soc.* **1987**, *109*, 4914. (f) Pascal, R. A., Jr.; Spergel, J.; Van Engen, D. *Tetrahedron Lett.* **1986**, *27*, 4099. (g) Pierre, J. L.; Barat, P. *Bull. Soc. Chim. Fr.* **1983**, II-367. (h) Lehn, J. M. *Acc. Chem. Res.* **1978**, *11*, 49.

- (7) Dietrich, B.; Guilhem, J.; Lehn, J. M.; Pascard, C.; Sonveaux, E. *Helv. Chim. Acta* **1984**, *67*, 91.
- (8) Graf, E.; Lehn, J. M. *J. Am. Chem. Soc.* **1976**, *98*, 6403.
- (9) Metz, B.; Rosalky, J. M.; Weiss, R. *J. Chem. Soc., Chem. Commun.* **1976**, 533.
- (10) (a) Park, C. H.; Simmons, H. E. *J. Am. Chem. Soc.* **1968**, *90*, 2429. (b) Bell, R. A.; Christoph, G. G.; Fronzek, F. R.; Marsh, R. E. *Science* **1975**, *190*, 151.

Table I. Crystallographic Data for 1, 3-0.5C₆H₆, 4-0.5C₆H₆, and 5

	1	3-0.5C ₆ H ₆	4-0.5C ₆ H ₆	5
mol formula	C ₁₆ H ₃₉ ClGe ₃ N ₄	C ₁₉ H ₄₂ Ge ₃ IN ₄	C ₁₉ H ₄₂ ClIn ₄ Sn ₃	C ₁₆ H ₃₉ BrN ₄ Sn ₃
fw	540.79	671.31	718.10	723.49
cryst dimens, mm	0.35 × 0.4 × 0.4	0.4 × 0.4 × 0.4	0.2 × 0.2 × 0.4	0.3 × 0.4 × 0.25
cryst color	colorless	colorless	colorless	light yellow
cryst syst	monoclinic	rhombohedral	rhombohedral	orthorhombic
space group	P2 ₁ /n	R $\bar{3}$	R $\bar{3}$	Pnma
a, Å	10.188 (6)	11.990 (7)	11.720 (8)	21.471 (9)
b, Å	19.923 (8)	11.990 (7)	11.720 (8)	11.753 (6)
c, Å	12.099 (7)	34.40 (1)	35.109 (15)	10.476 (5)
β or γ, deg	91.54 (5)	120.0	120.0	90.0
V, Å ³	2455 (1)	4283 (2)	4176 (2)	2644 (1)
Z	4	6	6	4
δ, g cm ⁻³	1.463	1.56	1.71	1.818
no. of indep data	2859	1369	1152	1921
no. of data with F _o > nσ	2666 (n = 3)	965 (n = 2)	966 (n = 2)	1723 (n = 3)
no. of variables	260	73	105	158
μ, cm ⁻¹	36.92	40.69	25.71	40.93
R(F _o)	0.035	0.057	0.032	0.0334
R _w (F _o)	0.037	0.057	0.031	0.0374

kroanalytisches Laboratorium, Theaterstr. 23, Göttingen, Germany. The compounds Me₂Si(*t*-BuN)₂Ge,¹¹ Me₂Si(*t*-BuN)₂Sn,¹² Me₂Si(*t*-BuN)(*t*-BuNH)GeX (X = Cl, Br),¹³ Me₂Si(*t*-BuN)(*t*-BuNH)SnX (X = Cl, Br, I),¹⁴ Ge₃(*t*-BuNH)₂Cl₂,¹⁵ Sn₂(*t*-BuNH)₂X₂ (X = Cl, Br),¹⁴ Ge₃(*N*-*t*-Bu)₄H₂,¹¹ and Sn₃(*N*-*t*-Bu)₄H₂¹⁵ were prepared as previously described.

tert-Butylamine (Merck) was dried over CaH₂ and distilled. HCl(g) and HBr(g) (Messer, Griesheim) were dried by using previously published methods¹⁶ and subsequently bubbled through toluene or benzene. The molarities of these solutions were determined by titration with both NaOH and AgNO₃.¹⁷ The solvents benzene, diethyl ether, and toluene were dried with LiAlH₄ and/or sodium-benzophenone, distilled, and stored over Na wire. NaBr, NaI, and TiCl₄ (Merck) were recrystallized and dried by published methods.¹⁶

The single crystals obtained were transferred to glass capillaries of appropriate sizes (0.1–0.5-mm diameter) and sealed under nitrogen. Crystal quality was examined by using either a Weissenberg camera (Stoe; Darmstadt, Germany) or a Buerger precession camera (Huber; Rimsting, Germany). The capillaries were then transferred to an automatic four-circle diffractometer (Siemens, type AED2) where data were collected by using the Wyckoff- ω -scan technique with Mo K α radiation. Space group determinations and measurements of the cell constants were calculated on the basis of the photographs and then checked and refined again on the diffractometer. The details of data collections and structure determinations are summarized in Table I. All structures were solved by using standard Patterson techniques. Extinction corrections were not applied; however, the intensities were corrected for Lp effects. The absorption effects were taken into account by using reflections from a complete ψ scan (empirical absorption correction). The hydrogen atoms on the methyl groups were treated together with the central carbon as a rigid group with tetrahedral local symmetry and standard bond lengths (C–H = 1.08 Å). Final atomic positional parameters as well as the isotropic *B* values calculated from *U*_{ij} values are given in Table II. The most significant bond lengths and angles are collected in Table III. Anisotropic thermal parameters (Table SI), a full table of bond lengths and angles (Table SII), and a listing of observed and calculated structure factors are available as supplementary material. For computation a VAX-II computer was used with the programs SHELX,¹⁸ ORTEP,¹⁹ and SCHKAL.²⁰

Formation of M₃(*N*-*t*-Bu)₄H₂X. (A) M₃(*N*-*t*-Bu)₄H₂ with HX/*Benzene*. To a stirred, dilute benzene solution (10 mL) of M₃(*N*-*t*-Bu)₄H₂ (M = Ge, Sn) (1 mmol) was added an HX/benzene solution (X

= Cl, Br), e.g. 9.1 mL of 0.11 M benzene/HCl, dropwise over 2 h. As the solution was added, 1, 2, 4, and 5 precipitated as white solids. After 2 h of stirring, the volatiles were removed in vacuo resulting in 98–99% analytically pure 0.5 benzene adducts of 1, 2, 4, and 5. The physical and spectroscopic data of 1–6 are collected in Table IV (see below for formation of 3 and 6).

(B) Me₂Si(*N*-*t*-Bu)₂M and Me₂Si(*t*-BuNH)(*N*-*t*-Bu)MX with *t*-BuNH₂. *tert*-Butylamine (1.26 mL, 12 mmol) was added dropwise to a stirred mixture of Me₂Si(*N*-*t*-Bu)₂Ge or Me₂Si(*N*-*t*-Bu)₂Sn (3 mmol) and Me₂Si(*t*-BuNH)(*N*-*t*-Bu)GeCl or Me₂Si(*t*-BuNH)(*N*-*t*-Bu)SnX (X = Cl, Br, I) (6 mmol) in benzene (5 mL). The yellow or orange solution changed to colorless or slightly yellow shortly after addition of the amine as a crystalline solid precipitated. The solution was refluxed for 2 h in order to ensure completion of the reaction. The volatiles were removed under vacuum resulting in colorless crystalline solids. Recrystallization from benzene resulted in pure products: 1-0.5C₆H₆ (90%), 4-0.5C₆H₆ (87%), 5-0.5C₆H₆ (95%), and 6-0.5C₆H₆ (94%). When 1 and 5 were recrystallized from toluene, solvent-free crystals were obtained (91%).

(C) Me₂Si(*N*-*t*-Bu)₂M and [(*t*-BuNH)MX]₂ with *t*-BuNH₂. *tert*-Butylamine (0.95 mL, 9 mmol) was added dropwise to a mixture of Me₂Si(*N*-*t*-Bu)₂M (6.0 mmol) and [(*t*-BuNH)GeCl]₂ or [(*t*-BuNH)SnX]₂ (X = Cl, Br, I) (1.5 mmol) in benzene (3 mL). Again the orange color of the solution changed to a light yellow with addition of the amine and a colorless crystalline solid precipitated. The solution was refluxed for 2 h. All volatiles were removed in vacuo and the solids recrystallized from benzene (5 mL): 1-0.5C₆H₆ (88%), 4-0.5C₆H₆ (92%), 5-0.5C₆H₆ (93%), 6-0.5C₆H₆ (93%).

For physical and spectroscopic data of 1, 2, 4, 5, and 6, see Table IV. Anal. Calc for C₁₆H₃₉ClGe₃N₄, *M*_r = 540.79 (1): C, 35.54; H, 7.27; Cl, 6.56; N, 10.36. Found: C, 35.51; H, 7.22; Cl, 6.62; N, 10.23. Calc for C₁₉H₄₂ClGe₃N₄, *M*_r = 579.80 (1-0.5C₆H₆): C, 39.36; H, 7.30; Cl, 6.11; N, 9.66. Found: C, 40.23; H, 7.41; Cl, 6.01; N, 9.76. Calc for C₁₆H₃₉BrGe₃N₄, *M*_r = 585.25 (2): C, 32.84; H, 6.72; N, 9.57. Found: C, 33.54; H, 6.78; N, 9.42. Calc for C₁₉H₄₂ClIn₄Sn₃, *M*_r = 718.10 (4-0.5C₆H₆): C, 31.78; H, 5.90; Cl, 4.94; N, 7.80. Found: C, 32.07; H, 5.45; Cl, 4.86; N, 7.66. Calc for C₁₉H₄₂BrIn₄Sn₃, *M*_r = 762.55 (5-0.5C₆H₆): C, 29.93; H, 5.50; Br, 10.48; N, 7.35. Found: C, 30.13; H, 5.55; Br, 10.00; N, 7.41. Calc for C₁₉H₄₂IN₄Sn₃, *M*_r = 809.54 (6-0.5C₆H₆): C, 28.19; H, 5.23; I, 15.68; N, 6.92. Found: C, 28.46; H, 5.20; I, 15.46; N, 6.97.

Formation of [(*t*-BuNH)SnI]₂ (7). Me₂Si(*t*-BuNH)(*N*-*t*-Bu)SnI with *t*-BuNH₂. To a stirred solution of Me₂Si(*t*-BuNH)(*N*-*t*-Bu)SnI (1.430 g, 3.2 mmol) was added *tert*-butylamine (0.34 mL, 3.2 mmol) dropwise. After 2 days of stirring at room temperature all volatiles were removed in vacuo producing a slightly yellow microcrystalline solid of [(*t*-BuNH)SnI]₂ (7): yield 1.23 g, 89%; mp 124 °C. Anal. Calc for C₈H₂₀I₂N₂Sn₂, *M*_r = 635.48: C, 15.12; H, 3.17; I, 39.94; N, 4.41. Found: C, 14.89; H, 3.38; I, 39.45; N, 4.47. ¹H NMR (C₆D₆): δ 0.91 [s, 18 H, C(CH₃)₃]. Molecular mass: calc, 635.48 g/mol; found (by cryoscopy in benzene), 653.8 g/mol.

Anion-Exchange Reactions. (A) (*N*-*t*-Bu)(*t*-BuNH)₃M₃Cl with NaX (X = Br, I). In a typical experiment, a 0.02 M solution of 1 or 4 in benzene or toluene was added to an equimolar amount of the crystalline salt. The suspension was vigorously stirred and periodically monitored by ¹H NMR spectroscopy. Upon completion of the reaction the crystalline solid was separated by filtration and was identified as NaCl by X-ray powder diffraction. The filtrate was concentrated in vacuo and

- (11) Veith, M.; Grosser, M. *Z. Naturforsch.* **1982**, *37B*, 1375.
- (12) Veith, M. *Angew. Chem.* **1975**, *87*, 287; *Angew. Chem., Int. Ed. Engl.* **1975**, *14*, 263.
- (13) Veith, M.; Hobein, P.; Rösler, R. *Z. Naturforsch.* **1989**, *44B*, 1067.
- (14) Veith, M.; Jarczyk, M.; Huch, V. *Chem. Ber.* **1988**, *121*, 347.
- (15) Veith, M.; Sommer, M.-L.; Jäger, D. *Chem. Ber.* **1979**, *112*, 2581.
- (16) Perrin, D. D.; Armarego, W. L. F. *Purification of Laboratory Chemicals*, 3rd ed.; Pergamon Press: Oxford, England, 1988.
- (17) Jander, G.; Blasius, E. *Einführung in das anorganisch-chemische Praktikum*; S. Hirzel Verlag: Stuttgart, Germany, 1980.
- (18) Sheldrick, G. M. *SHELX-76/86, Program for Crystal Structure Determination*; University of Cambridge: Cambridge, U.K., 1976.
- (19) Johnson, C. K. *ORTEP-II*; Report ORNL-3794; Oak Ridge National Laboratory: Oak Ridge, TN, 1971.
- (20) Keller, E. *SCHKAL-88, Programm zur Darstellung von Kristallstrukturen*; Universität Freiburg: Freiburg, Germany, 1988.

Table II. Atomic Positional Parameters and Isotropic *B* Values (Å²) (Calculated from *U*_{ij}) of **1**, 3-0.5C₆H₆, 4-0.5C₆H₆, and **5**

atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>B</i>
1				
Ge(1)	0.83746 (5)	0.09487 (3)	0.25669 (4)	3.46 (3)
Ge(2)	0.59245 (5)	0.07038 (2)	0.11631 (4)	2.99 (3)
Ge(3)	0.58944 (5)	0.04807 (3)	0.36332 (4)	3.32 (3)
Cl	0.5630 (2)	0.25813 (6)	0.2879 (1)	4.82 (7)
N(1)	0.6990 (4)	0.0237 (2)	0.2351 (3)	2.9 (2)
N(2)	0.7243 (4)	0.1450 (2)	0.1447 (3)	3.3 (2)
H(1)	0.691 (5)	0.173 (2)	0.172 (3)	2 (1)
N(3)	0.4872 (3)	0.1015 (2)	0.2467 (3)	3.0 (2)
H(2)	0.5056 (0)	0.1471 (0)	0.2648 (0)	2.5 (9)
N(4)	0.7199 (4)	0.1245 (2)	0.3802 (3)	3.4 (2)
H(3)	0.6712 (0)	0.1696 (0)	0.3621 (0)	5 (1)
C(1)	0.7399 (5)	-0.0475 (3)	0.2206 (4)	3.8 (2)
C(11)	0.8303 (7)	-0.0531 (3)	0.1227 (5)	5.4 (3)
C(12)	0.6193 (6)	-0.0913 (3)	0.1992 (5)	4.9 (3)
C(13)	0.8137 (7)	-0.0715 (3)	0.3244 (5)	5.8 (3)
C(2)	0.7904 (5)	0.1748 (3)	0.0466 (4)	4.0 (2)
C(21)	0.6851 (6)	0.2052 (3)	-0.0308 (5)	5.9 (3)
C(22)	0.8832 (6)	0.2301 (3)	0.0844 (5)	5.7 (3)
C(23)	0.8638 (6)	0.1200 (3)	-0.0133 (5)	5.5 (3)
C(3)	0.3398 (5)	0.0937 (3)	0.2404 (4)	3.9 (2)
C(31)	0.3054 (6)	0.0213 (3)	0.2121 (5)	5.1 (3)
C(32)	0.2870 (6)	0.1123 (4)	0.3535 (5)	5.7 (3)
C(33)	0.2873 (6)	0.1412 (3)	0.1521 (5)	5.5 (3)
C(4)	0.7840 (5)	0.1370 (3)	0.4926 (4)	4.2 (3)
C(41)	0.6755 (6)	0.1554 (4)	0.5723 (5)	6.0 (3)
C(42)	0.8803 (7)	0.1941 (4)	0.4827 (5)	6.4 (4)
C(43)	0.8514 (7)	0.0740 (3)	0.5312 (5)	5.9 (3)
3-0.5C₆H₆				
I	0.0000 (0)	0.0000 (0)	0.16486 (3)	7.16 (9)
Ge	0.0841 (1)	0.1683 (1)	0.29224 (3)	6.5 (1)
N(1)	0.0000 (0)	0.0000 (0)	0.3209 (3)	6.5 (8)
N(2)	0.1573 (8)	0.0785 (9)	0.2601 (2)	6.0 (8)
C(1)	0.0000 (0)	0.0000 (0)	0.3643 (4)	7 (1)
C(2)	-0.070 (1)	-0.139 (1)	0.3796 (5)	9.0 (3)
C(3)	0.302 (1)	0.150 (1)	0.2525 (3)	7 (1)
C(4)	0.336 (1)	0.062 (1)	0.2289 (4)	8.6 (3)
C(5)	0.335 (1)	0.272 (1)	0.2289 (4)	8.6 (3)
C(6)	0.371 (1)	0.185 (1)	0.2918 (4)	8.9 (4)
C(7)	0.013 (5)	0.123 (3)	0.5011 (8)	15.3 (8)
C(8)	0.207 (4)	0.103 (4)	0.503 (1)	2.3 (8)
4-0.5C₆H₆				
Sn	-0.09523 (3)	-0.19054 (3)	0.21101 (1)	3.01 (4)
Cl	0.0000 (0)	0.0000 (0)	0.32469 (8)	4.6 (1)
N(1)	0.0886 (4)	-0.0879 (4)	0.2452 (1)	3.0 (3)
N(2)	0.0000 (0)	0.0000 (0)	0.1814 (2)	2.8 (3)
C(1)	0.1606 (6)	-0.1596 (6)	0.2548 (2)	4.1 (5)
C(11)	0.0694 (7)	-0.2818 (6)	0.2776 (2)	5.2 (5)
C(12)	0.2812 (7)	-0.0688 (7)	0.2777 (2)	5.7 (6)
C(13)	0.2006 (8)	-0.1988 (8)	0.2177 (2)	5.8 (6)
C(2)	0.0000 (0)	0.0000 (0)	0.1388 (3)	2.9 (4)
C(21)	0.0709 (6)	-0.0689 (6)	0.1236 (2)	4.3 (5)
C(3)	-0.599 (1)	-0.1997 (8)	-0.3337 (3)	7.0 (7)
H(1)	0.068 (5)	-0.059 (5)	0.269 (2)	3 (1)
H(2)	-0.224 (8)	-0.105 (8)	0.003 (3)	9 (2)
5				
Sn(1)	0.43441 (2)	0.10678 (4)	0.90635 (4)	3.73 (3)
Sn(2)	0.40438 (3)	0.2500 (0)	0.63787 (6)	4.01 (3)
Br	0.23506 (5)	0.2500 (0)	0.9154 (1)	4.69 (5)
N(1)	0.3600 (2)	0.1199 (4)	0.7610 (5)	3.3 (2)
H(1)	0.325 (3)	0.156 (5)	0.800 (5)	3 (1)
N(2)	0.3902 (3)	0.2500 (0)	1.0111 (7)	3.2 (3)
H(2)	0.349 (4)	0.2500 (0)	0.986 (8)	2 (2)
N(3)	0.4713 (3)	0.2500 (0)	0.7969 (7)	3.7 (4)
C(1)	0.3381 (3)	0.0133 (6)	0.6980 (7)	4.5 (3)
C(11)	0.3110 (4)	-0.0663 (7)	0.7962 (9)	6.5 (5)
C(12)	0.2877 (5)	0.0419 (8)	0.6007 (8)	6.5 (5)
C(13)	0.3911 (4)	-0.0447 (8)	0.631 (1)	7.9 (6)
C(2)	0.3948 (5)	0.2500 (0)	1.154 (1)	5.2 (6)
C(21)	0.4653 (6)	0.2500 (0)	1.193 (1)	6.3 (7)
C(22)	0.3632 (5)	0.3563 (9)	1.2074 (9)	8.4 (6)
C(3)	0.5404 (5)	0.2500 (0)	0.763 (1)	5.4 (6)
C(31)	0.5806 (5)	0.2500 (0)	0.885 (1)	7.0 (8)
C(32)	0.5562 (4)	0.352 (1)	0.689 (1)	8.0 (6)


the crystalline material spectroscopically (¹H NMR, IR, MS) characterized. In the case of **2** (98%) and **3** (98%) elemental analyses were performed. Further data for **2** and **3** are found in Table IV.

(B) *t*-BuN(*t*-BuNH)₃M₃I with TICI. These reactions were carried out similarly to method A above. In these reactions I⁻ exchanged with Cl⁻ forming TII and the compounds **1** and **4**, respectively (yield 98%, characterized by ¹H NMR and IR). Anal. Calc for C₁₉H₄₂IGe₃N₄, *M_r* = 671.31 (3-0.5C₆H₆): C, 33.99; H, 6.31; N, 8.35. Found: C, 33.31; H, 6.34; N, 8.44.








Reaction of Sn₃(*N-t*-Bu)₄H₂ (**10**) with Excess HCl. Sn₃(*N-t*-Bu)₄H₂ (2 mmol) in benzene (10 mL) was treated with an HCl/benzene solution in molar portions. The first molar equivalent produced **4**. As excess HCl was added (Sn₃(*N-t*-Bu)₄H₂:HCl = 1:6), SnCl₂·H₂N-*t*-Bu was obtained: yield 1.18 g, 75%. The elemental analysis and a singlet in the ¹H NMR spectrum at δ 0.73 are consistent with those previously reported.¹⁴ After further addition of HCl (Sn₃(*N-t*-Bu)₄H₂:HCl = 1:9), *t*-BuNH₃⁺SnCl₃⁻ was obtained: yield 1.6 g, 89%. This product was characterized by ¹H NMR spectroscopy [δ 1.33 (CD₃CN)] and its melting point (135 °C).¹⁴

Results and Discussion

Syntheses. As discussed in the Introduction the iminogermylene and -stannylenes (*N-t*-Bu)₂(*t*-BuNH)₂Ge₃¹¹ and (*N-t*-Bu)₂(*t*-BuNH)₂Sn₃¹⁵ react quantitatively with dilute HX solutions in benzene forming *t*-BuN(*t*-BuNH)₃M₃X (**1**, **2**, **4**, **5**) (eq 1). Other




synthetic routes to these compounds are depicted in eqs 2 and 3.



These alternative methods require more easily obtainable starting materials, such as Me₂Si(*N-t*-Bu)₂M and Me₂Si(*N-t*-Bu)(*t*-BuNH)MX, and avoid the tedious preparation of anhydrous HI.

In eq 2 the cyclic bis(amino)germylene or -stannylene is treated with *tert*-butylamine in the presence of Me₂Si(*t*-BuNH)(*N-t*-Bu)MX (M = Ge, X = Cl; M = Sn, X = Cl, Br, I). In all cases complete aminolysis occurs with the metallacycle Me₂Si(*t*-BuNH)(*N-t*-Bu)MX serving as the HX carrier. The only side product is the volatile silazane Me₂Si(*t*-BuNH)₂, which is easily removed from the mixture in vacuo. Equation 3 is very similar to eq 2, the only exception being that the HX carrier now is [(*t*-BuNH)MX]₂. The fact that the product [(*t*-BuNH)SnI]₂ (**7**) is formed from aminolysis of Me₂Si(*t*-BuNH)(*N-t*-Bu)MX with *t*-BuNH₂, as depicted in eq 4 (in analogy to former procedures),^{13,14} is a strong indication that [(*t*-BuNH)MX] is an intermediate in the formation of **1**, **4**, **5**, and **6** in eq 2.



The products **1**, **2**, **4**, **5**, and **6** are initially identified by their ¹H NMR spectra as two resonances in a ratio of 3:1. This spectral pattern is consistent with the molecules possessing C_{3v} (3*m*) symmetry in solution. The three equivalent *tert*-butylamino groups occur at low field (δ 1.43 for **1–3** and 1.41 for **4–6**) as expected,^{12,15} while the resonance for the remaining *tert*-butylamino group (labeled R' in formula A) is found at higher field (δ range 1.16–1.06; for specific values, see Table IV). This high-field shift indicates there is an inductive effect as the size of the anion

Table III. Selected Bond Lengths (Å) and Angles (deg) (Some Averaged^a) of Compounds 1, 3-0.5C₆H₆, 4-0.5C₆H₆, 5, and Sn₃(N-*t*-Bu)₄H₂⁵ for Comparison

	1 ^a	3-0.5C ₆ H ₆	4-0.5C ₆ H ₆	5 ^a	Sn ₃ (N- <i>t</i> -Bu) ₄ H ₂
N'-M ^b	2.005 (6)	2.007 (6)	2.195 (4)	2.191 (7)	2.207
N-M ^c	2.027 (4)	2.025 (8)	2.221 (4)	2.217 (5)	2.208
N-H	0.91 (9)		0.96 (6)	0.94 (3)	
N-C	1.504 (8)	1.51 (2)	1.50 (1)	1.51 (1)	1.50
N...X	3.281 (20)	3.661 (8)	3.316 (5)	3.484 (3)	
H...X	2.379 (13)		2.36 (5)	2.55 (3)	
M...M	3.020 (2)	3.027 (4)	3.350 (1)	3.350 (8)	3.344
N...N	2.875 (5)	2.829 (9)	3.103 (6)	3.087 (9)	2.961
N...N'	2.668 (5)	2.654 (6)	2.868 (6)	2.855 (6)	2.869
cg...N ^d	2.089 (7)	2.092 (7)	2.240 (6)	2.230 (7)	2.304
cg...X	2.830 (5)	3.276 (5)	2.791 (5)	2.993 (5)	
cg...N	1.660 (1)	1.633 (2)	1.791 (2)	1.782 (1)	1.71
N'-M-N	82.85 (6)	82.4 (3)	80.9 (2)	80.8 (4)	81.1
N-M-N	90.3 (2)	88.8 (1)	88.7 (1)	88.3 (6)	84.2
M-N'-M	97.7 (1)	97.9 (4)	99.4 (2)	99.7 (7)	98.5
M-N-M	96.3 (1)	96.9 (5)	98.1 (3)	98.1 (5)	98.4
C-N-M	118.0 (2)	117.8 (7)	118.9 (3)	117.7 (6)	120.0
C-N-H	106.0 (9)		108 (3)	110 (2)	
N...H...X	171.07 (9)		175 (4)	178 (1)	
N...X...N	51.97 (6)	45.5 (3)	55.8 (3)	52.6 (3)	
H...X...H	55.5 (9)		57.0 (9)	53.1 (9)	

^a Most of the distances and angles have been calculated from several chemically identical bonds. The mean deviation in parentheses is given here instead of the standard deviation, which may be smaller. ^b N' designates the N atom bonded to three metal atoms, which is situated on the (quasi) 3-fold axis (compare formula A). ^c N designates the N atoms to which H atoms are bonded (compare formula A). ^d Cg = center of gravity of the three hydrogen-linked N atoms.

Table IV. Physical and Spectroscopic Data for 1-6

	mp, °C	¹ H NMR: δ	mass spectrum	IR: ν(NH), cm ⁻¹
1	150 dec ^a 280 dec ^b	1.16 (9 H) 1.43 (27 H)	<i>m/z</i> = 505 C ₁₆ H ₃₉ N ₄ ⁷² Ge ₂ ⁷⁴ Ge	3067 3080
2	190 dec ^a 280 dec ^b	1.13 (9 H) 1.43 (27 H)	<i>m/z</i> = 505	3098
3	160 dec ^a 270 dec ^b	1.11 (9 H) 1.43 (27 H)	<i>m/z</i> = 505	3066
4	289 dec ^b	1.11 (9 H) 1.41 (27 H)	<i>m/z</i> = 643 C ₁₆ H ₃₉ N ₄ ¹¹⁸ Sn ₂ ¹²⁰ Sn	3087
5	290 dec ^b	1.08 (9 H) 1.41 (27 H)	<i>m/z</i> = 643	3116
6	195 dec ^a 270 dec ^b	1.06 (9 H) 1.41 (27 H)	<i>m/z</i> = 643	

^a With 0.5 C₆H₆/molecule. ^b Without solvent in the crystal.

increases from Cl⁻ to I⁻. The compounds are only slightly soluble in noncoordinating solvents such as cyclohexane, benzene, or toluene, which may be taken as an indication of their polar bonds (see below). The compounds 1, 2, 4, 5, and 6 as well as 3 (for formation, see below) are furthermore characterized by mass and IR spectra and elemental analysis in addition to ¹H NMR spectra. Some of these data are collected for comparison in Table IV. The desorption chemical ionization (DCI⁺) mass spectra display parent ion envelopes centered at *m/z* = 505 for 1-3 (M = Ge) and 642 for 4-6 (M = Sn) with characteristic isotope patterns for the calculated cations [Ge₃(N-*t*-Bu)₄H₃]⁺ and [Sn₃(N-*t*-Bu)₄H₃]⁺, respectively. The repeated appearance of these ions and their stability within the mass spectrometer may be taken as a further hint at the cation/anion nature of these complexes. Another interesting feature is found in the ν(N-H) wavenumbers of the compounds. There is an increase in these numbers, when Cl⁻ is substituted by Br⁻ or I⁻, respectively, for both the Ge series (1-3) and the Sn series (4-6). This fact seems to reflect the longer H...X distances, which result from the bigger halogens and which therefore should lead to less effective hydrogen bridging. In the structural part of this paper (see below), it can be shown that these wavenumbers are linearly correlated to the ionic radii of Cl, Br, and I or to the intramolecular distances obtained from the X-ray structure determinations.


It seemed at this point evident that the molecules 1, 2, 4, 5, and 6 are composed of a spherical lipophilic part (M₃(N-*t*-Bu)₄) dominated by a quasitrahedral arrangement of the *tert*-butyl

groups and a lipophobic part (H₃X) containing most of the polarity. Such species have directional behavior at the phase boundaries, with the "organic part" pointing into a noncoordinating organic solvent and the polar H₃X arrangement pointing in the opposite direction. To test this assumption, we treated benzene solutions of 1 and 4 with the salts NaBr and NaI in heterogeneous reactions. As depicted in eqs 5 and 6, 1 and 4 are transformed to 2, 3, 5, and 6, respectively, NaCl being the byproduct in all cases.



The iodine compound 3 is obtained here for the first time. Its data are compatible with those for the other homologous compounds (see Table IV). The yields for these reactions are almost quantitative although reaction time varies considerably. Sodium chloride can be identified by X-ray powder diffraction. This reaction is comparable to the transformation of NaBr to NaCl by interaction of gaseous HCl with solid NaBr.²¹ The transformation is ascribed to the higher polar nature and crystal lattice energy of NaCl compared to NaBr.²²

If the crystal energy is a driving force in these reactions, we thought it should be possible to retransform 3 to 1 using a metal halide whose iodide is more stable than the chloride. As depicted in eq 7, thallium chloride quantitatively transforms 3 back to 1,



with formation of TlI. Here, as before, 1 is identified by ¹H NMR

(21) Harrison, L. G.; Siddiqui, R. A. *Trans. Faraday Soc.* 1962, 58, 982.

(22) Huheey, J. E. *Anorganische Chemie*; W. de Gruyter: Berlin, 1988; p 72.





Figure 1. Graphic representation of $\text{Ge}_3(\text{N-}t\text{-Bu})_4\text{H}_3\text{Cl}$ (**1**) based on the single-crystal X-ray structure determination.

spectra and TII by its X-ray powder diffraction pattern.

The ratio of the reactants in eq 1 plays an important role. When an excess of HX was used, the cages react, forming products similar to those found in the HCl addition to $\text{Sn}_4(\text{N-}t\text{-Bu})_4$.¹⁴ In eq 8 the products formed by subsequent addition of HCl to



$\text{Sn}_3(\text{N-}t\text{-Bu})_4\text{H}_2$ are summarized. The HCl adduct of $\text{Sn}_3(\text{N-}t\text{-Bu})_4\text{H}_2$ (**4**) is formed first. Further addition of HCl ($\text{Sn}_3(\text{N-}t\text{-Bu})_4\text{H}_2:\text{HCl} = 1:6$) destroys the cage **4** to form $t\text{-Bu-NH}_2 \cdot \text{SnCl}_2$ and finally $t\text{-BuNH}_3^+ \text{SnCl}_3^-$ ($\text{Sn}_3(\text{N-}t\text{-Bu})_4\text{H}_2:\text{HCl} = 1:9$). The four-membered ring $(t\text{-BuN(H)SnCl})_2$ has not been found to be stable under these conditions, in contrast to the reaction of $\text{Sn}_4(\text{N-}t\text{-Bu})_4$ with HCl.¹⁴

Structural Details. X-ray structural determinations have been carried out on single crystals of **1**, $3 \cdot 0.5\text{C}_6\text{H}_6$, $4 \cdot 0.5\text{C}_6\text{H}_6$, and **5**. While $3 \cdot 0.5\text{C}_6\text{H}_6$ and $4 \cdot 0.5\text{C}_6\text{H}_6$ are isotypical, crystallizing in the rhombohedral space group $R\bar{3}$, **1** crystallizes in a monoclinic ($P2_1/n$) and **5** in an orthorhombic ($Pnma$) space group. The crystals of **1** and **5** contain no interstitial benzene and form closed-packed (**1**) or mostly closed-packed (**5**) van der Waals lattices.²³ **1** has no crystallographic symmetry, whereas **5** is situated on a mirror plane of the space group. A closer look at the bond lengths and angles, however, reveals there is only a minor deviation from the higher C_{3v} ($3m$) symmetry. The molecules thus lose the higher symmetry, which exists in solution, as they are packed in the crystal lattice following the principles formulated by Kitaigorodskii.²³

In Figure 1 the compound $\text{Ge}_3(\text{N-}t\text{-Bu})_4\text{H}_3\text{Cl}$ (**1**) is shown to illustrate the general overall structure of the molecules. Figure 2 gives a view down the 3-fold axis of $\text{Sn}_3(\text{N-}t\text{-Bu})_4\text{H}_3\text{Cl}$ (**4**), showing the high symmetry of the molecules. As can be seen from these representations, the central cage of the molecules $\text{M}_3(\text{N-}t\text{-Bu})_4\text{H}_3\text{X}$ can be described as a $\text{M}_3\text{N}_4\text{X}$ "cube". Here the corner X is displaced from the center but held by three hydrogen atoms bridged to the three nitrogen atoms. The same cage may also be described as an N_4 tetrahedron with three faces pyramidally capped by the metallic elements and one face capped by a H_3X trigonal pyramid.

In the rhombohedral system the molecules **3** and **4** adopt the relatively high point symmetry C_3 (**3**), losing only the mirror plane. Figure 3 shows a section of the packing of $4 \cdot 0.5\text{C}_6\text{H}_6$. When viewed down the 3-fold axis of the lattice, "pairs of molecules" are separated by benzene molecules, which are situated around 3 centers. These benzene molecules are in between two *tert*-butyl




Figure 2. View down the 3-fold axis of $\text{Sn}_3(\text{N-}t\text{-Bu})_4\text{H}_3\text{Cl}$ (**4**) with Cl as the nearest atom.




Figure 3. Section of the packing of **4** to show the interstitial benzene molecules (left and right). The *tert*-butyl amino groups in the middle of the picture belong to six different molecules, which are omitted for clarity.

groups of different molecules. Interestingly, these "pairs" have opposing halogens (**3**, $\text{I} \cdots \text{I} = 11.34 \text{ \AA}$; **4**, $\text{Cl} \cdots \text{Cl} = 12.31 \text{ \AA}$) at very long distances. The long separation of the halogen atoms of the molecule pair is caused by *tert*-butyl groups of other molecules (Figure 3).

It is interesting to note that **1**, as well as **5** and **6**, when crystallized from benzene, form crystals isotypical to $3 \cdot 0.5\text{C}_6\text{H}_6$ and $4 \cdot 0.5\text{C}_6\text{H}_6$ (rhombohedral lattices: $1 \cdot 0.5\text{C}_6\text{H}_6$, $a = b = 11.495$ (7) \AA , $c = 34.54$ (2) \AA , $V = 3953 \text{ \AA}^3$; $5 \cdot 0.5\text{C}_6\text{H}_6$, $a = b = 11.807$ (6) \AA , $c = 35.27$ (2) \AA , $V = 4258 \text{ \AA}^3$; $6 \cdot 0.5\text{C}_6\text{H}_6$, $a = b = 11.996$ (5) \AA , $c = 35.53$ (2) \AA , $V = 4428 \text{ \AA}^3$).

At a first view, the bond lengths of **4** and **5** are only slightly different from those in $\text{Sn}_3(\text{N-}t\text{-Bu})_4\text{H}_2$ and, generally speaking, the $\text{M}_3\text{N}_4\text{H}_3\text{X}$ cages, when germanium is a component, are smaller than those where $\text{M} = \text{Sn}$. A closer look reveals nevertheless that important changes are taking place, if a molecule of HX is added to the neutral $\text{Sn}_3(\text{N-}t\text{-Bu})_4\text{H}_2$ moiety: One can distinguish clearly between two kinds of metal–nitrogen bonds, the bonds between the nitrogen on the 3-fold axis and the metal atoms being shorter than these between the hydrogen-bearing nitrogen and metal atoms. While the nonbonding distances $\text{N} \cdots \text{N}'$ (for labeling compare formula A) in **4** and **5** differ only slightly compared to those in the neutral molecule $\text{Sn}_3(\text{N-}t\text{-Bu})_4\text{H}_2$, the $\text{N} \cdots \text{N}$ distances are much longer. The incorporation of HX in the polycycle seems to occur at the expense of the $\text{M}-\text{N}$ bonding and leads to an "opening" of the hole in the M_3N_4 cage (compare also $\text{N}-\text{M}-\text{N}$ angles with $\text{N}-\text{M}-\text{N}'$). This place is needed to accommodate the third hydrogen atom, which, of course, should have a repulsive force toward the other two hydrogen atoms already present. In **1**, $4 \cdot 0.5\text{C}_6\text{H}_6$, and **5** the positions of the hydrogen atoms bonded to nitrogen can be located. They have in the three structures similar intramolecular distances ($\text{H} \cdots \text{H} = 2.23$ (**1**), 2.23 ($4 \cdot 0.5\text{C}_6\text{H}_6$), 2.27 \AA (**5**)). Despite the different compositions of the compounds the interatomic $\text{H} \cdots \text{H}$ distances in **1** are equal to those found in $4 \cdot 0.5\text{C}_6\text{H}_6$!

The halides in these compounds are clearly pyramidally coordinated by three hydrogen atoms. As far as determined, the $\text{H} \cdots \text{X} \cdots \text{H}$ angles are quite acute and are somewhat greater than the $\text{N} \cdots \text{X} \cdots \text{N}$ angles. This implies that the hydrogen bridges are situated somewhat outside the N_3X pyramid although the $\text{N} \cdots \text{H} \cdots \text{X}$ angles are very close to 180° (compare Table III). The

(23) Kitaigorodskii, A. I. *Molecular Crystals and Molecules*; Academic Press: New York, 1973.




Figure 4. N...X distances as a function of the $\nu(\text{N-H})$ wavenumbers.

nearly linear N...H...X arrangements suggest relatively strong bridges. This becomes evident, if the N...X distances found here are compared to the CsCl-type structures (disorder of the NH_4^+ ions) of ammonium chloride ($\text{N}\cdots\text{Cl} = 3.356 \text{ \AA}$), ammonium bromide ($\text{N}\cdots\text{Br} = 3.507 \text{ \AA}$), and ammonium iodide ($\text{N}\cdots\text{I} = 3.737 \text{ \AA}$).^{24,25} Contrarily the N...Cl distances in $\text{CH}_3\text{NH}_3\text{Cl}$ (CN at Cl = 4) are 3.18 \AA , those in $(\text{CH}_3)_2\text{NH}_2\text{Cl}$ (CN at Cl = 2) are 3.11 \AA , and those in $(\text{CH}_3)_3\text{NHCl}$ (CN at Cl = 1) are 3.005 \AA , respectively, and therefore considerably shorter than those in 1 and 4.²⁶ Compared to the complexes formed by Cl^- and Br^- with cryptands of a different charge,^{7,9,10b} the N...X distances found in 1, 4, and 5 are again too long. A typical value for four-coordinate chloride N...Cl is $3.09 (2) \text{ \AA}$, and for six-coordinate chloride and bromide N...Cl/Br values are around $3.30 (9)$ and $3.39 (4) \text{ \AA}$, respectively. This discrepancy cannot be explained by the different numbers of hydrogen atoms, which coordinate the chloride (3 versus 6, 4, 2, or 1). The compounds 1 and 4 constitute a special case (for explanation see below).

The N-H bond lengths (compare Table III) are routinely determined too short in X-ray analyses and should be near 1.02 \AA (cf. the literature²⁵). They nevertheless indicate that the N...H...X bridge is highly unsymmetrical, the hydrogen atom being principally bonded to nitrogen. In summary, it seems to be clear

from the geometric arrangements of H and X atoms that an attractive force must be assumed between hydrogen and the halogen atoms, which is presumably of the ionic type with a strong directional character. The N...X distances correlate well with the $\nu(\text{N-H})$ wavenumbers, as may be seen from Figure 4.

At this stage a comparison of the molecules 1 and 4 may improve the understanding of the bonding. As can be seen in Table III, the N...Cl distances in 1 are shorter than those in 4, but the distance from the center of gravity (cg) of the three hydrogen-bonded nitrogen atoms to Cl is longer in 1 than in 4. Clearly, the Sn_3N_4 polycycle is better suited to accommodate a chloride or even a bromide than the Ge_3N_4 polycycle. This is also reflected in the N...X...N angles, which increase in the order $3 < 1 < 5 < 4$, the germanium compound 1 interestingly displaying a smaller angle than the tin compounds 4 and even 5. To use another picture, the halogen atoms are drawn further into the tin cage than into the germanium cage (see also distances $\text{cg}\cdots\text{X}$ in Table III), as the hole in the tin cage is deeper ($\text{cg}\cdots\text{X}$ distances) and larger (N...N or N...X distances) than that of the germanium analogue. The astonishingly long N...X distances found in 1 and 4, compared to the methylammonium chlorides or to cryptates (see before), must be attributed to the fact that the amino-hydrogen atoms are close together (see above). Besides attractive forces between Cl and H repulsive forces between Cl and the three nitrogen atoms, to which hydrogen is bonded, must be considered.

Conclusion

The physical, spectral (δ (^1H NMR) ν (IR)), and crystallographic data of 1-6 are in agreement with a description as molecular complexes with a high degree of "zwitter ionic" character between the $\text{M}_3(\text{N-}t\text{-Bu})_4\text{H}_3^+$ and the X^- part of the cage molecule. The exchange of the halogen atoms in biphasic reactions can be attributed to the special geometry and the polar structures of the compounds as well as to the bonding. The possible use of 1-6 as phase-transfer catalysts seems evident. Further experiments concerning the reactivity of these molecules are under investigation.

Acknowledgment. Financial support from the Fonds der Chemischen Industrie and the DFG is gratefully acknowledged. T.R.P. thanks the University of the Saarland for financial support. We thank Dr. L. Stahl for reading the manuscript.

Supplementary Material Available: Tables giving crystallographic details, thermal parameters, and bond distances and angles (10 pages); listings of structure factors (37 pages). Ordering information is given on any current masthead page.

(24) Jones, K. The Chemistry of Nitrogen. In *Comprehensive Inorganic Chemistry*; Pergamon Press: Oxford, England, 1975; Chapter 19, p 245.

(25) Parsonage, N. G.; Staveley, L. A. *Disorder in Crystals*; Clarendon Press: Oxford, England, 1978.

(26) Lindgren, J.; Olovson, I. *Acta Crystallogr.* 1968, B24, 554.