

Chromosome numbers of 36 cultivated taxa of *Prunus* subg. *Cerasus* in Japan

著者	Iwatsubo Yoshikane, Seng Yoo, Naruhash Naohiro
journal or publication title	植物地理・分類研究 = The journal of phytogeography and toxonomy
volume	52
number	1
page range	73-76
year	2004-07-25
URL	http://hdl.handle.net/2297/48661

Yoshikane Iwatsubo¹, Yoo Sengi² and Naohiro Naruhashi¹: Chromosome numbers of 36 cultivated taxa of *Prunus* subg. *Cerasus* in Japan

¹Department of Biology, Faculty of Science, Toyama University, Gofuku 3190, Toyama 930-8555, Japan;

²Ishikawa Prefecture Forestry Products Research Institute, Sannomiya, Tsurugi-machi, Ishikawa 920-2114, Japan

The latest guide to the flowering cherries of Japan (Nihon Hana no Kai 2002) carries 350 taxa including both wild and cultivated taxa. The chromosome number of flowering cherry in Japan had been reported by Ishikawa (1916), Okabe (1927, 1928), Sax (1931), Tanaka and Oginuma (1975), Oginuma and Tanaka (1976), Nishikawa (1985), Oginuma (1987 a,b, 1988, 1993) and Somego (2000), which showed that the flowering cherry in Japan had diploids, triploids and tetraploids. The latest and preceding papers (Iwatsubo et al. 2002, 2003) of this series about the chromosome numbers of flowering cherries in Japan showed that, on the basis of the chromosome numbers of 217 taxa, the cherries in Japan had diploid, triploid, tetraploid and hexaploid plants, many of them were diploid plants followed by triploid, tetraploid and hexaploid plants. The authors found the chromosome numbers of more than one hundred taxa listed in the latest guide (Nihon Hana no Kai 2002) are unknown.

This is the third paper of a series of chromosome numbers of *Prunus* subg. *Cerasus* (Iwatsubo et al. 2002, 2003), which reports on the chromosome numbers of 36 taxa conserved in the Arboretum of Ishikawa Forest Experiment Station, Ishikawa Pref., central Japan.

Materials and methods

Meristems of leaf buds of 36 taxa in *Prunus* subg. *Cerasus* were collected from the plants conserved in the Arboretum of Ishikawa Forest Experiment Station. One plant from each taxon was used for the study. The leaf buds of cherry trees cut into halves with a razor were pre-

treated in a 2 mM 8-hydroxyquinoline solution for one hour at room temperature and kept at ca. 5°C for about 15 h. They were fixed and kept in Newcomer's fluid at room temperature until used. After being immersed in 1 N HCl for a few hours, the leaf buds were macerated in 1 N HCl at 60°C for 10 minutes, and then washed in tap water. They were stained in a drop of 1.5% lacto-propionic orcein on the slide glass and ordinary squash technique was applied in preparation. Herbarium specimens made from the plants conserved in the Arboretum of Ishikawa Forest Experiment Station will be deposited in the Toyama Science Museum (TOYA).

Results and discussion

The studied taxa and their chromosome counts determined in the present study as well as counts reported previously by other authors are listed in Table 1. As seen in the Table, the chromosome counts registered so far showed an euploid series, based on $x=8$, of diploid ($2n=16$), triploid ($2n=24$) and tetraploid ($2n=32$). In the 36 taxa studied, 31 taxa are diploids, four taxa are triploids, and one taxon is a tetraploid. In these taxa, chromosome counts are reported here for the first time for 23 taxa: 20 are diploids and three are triploids. In the taxa studied, *P. ×introrsa* 'Introrsa' is reported as $2n=16$ by Somego (2000), however, the preceding two papers of this series (Iwatsubo et al. 2002, 2003) as well as the present result all show this taxon has $2n=24$ chromosomes. In 'Introrsa' the triploid ($2n=24$) form may be more widely cultivated than the diploid ($2n=16$) form. Okabe (1928) reported that *P. pendula* 'Pendula' had two forms

Table 1. List of studied taxa of *Prunus* subg. *Cerasus*, Japanese name, and present and previous chromosome counts

Taxon	Japanese name	Present count(2n)	Previous report	Reference
<i>P. incisa</i> Thunb.	'Akane-yae' (茜八重)	16		
	'Micantha'	16		
<i>P. ×introrsa</i> Yagi ex Ohwi	'Introrsa'	24	16	Somego 2000
			24	Iwatsubo et al. 2002, 2003
<i>P. jamaesakura</i> Siebold ex Koidz.	'Geba-zakura'	16		
	'Octopes'	16		
	'Sendaiya'	16	16	Iwatsubo et al. 2003
	'Zenshoji-kikuzakura'	16	16	Iwatsubo et al. 2002
cv.		16		
cv.		16		
var. <i>chikusiensis</i> (Koidz.) Ohwi		16		
f. <i>pubescens</i> (Makino) Ohwi		16		
<i>P. lannesiana</i> (Carrère) E.H.Wilson	'Amayadori'	16		
	'Beni-tamanishiki'	16	16	Iwatsubo et al. 2003
	'Fasciculata'	16		
	'Horinji'	16		
	'Kusimana'	16		
	'Moutan'	16		
	'Polycarpa'	24		
	'Purpurea-plena'	16		
	'Shibayama'	24		
var. <i>spectosa</i> (Koidz.) Makino	'Akami-obshima'	16	16	Iwatsubo et al. 2003
	'Cataracta'	16		
	'Gozanomaioi'	16		
	'Kanzaki-obshima'	16		
f. <i>plena</i> Y. Kimura		16		
<i>P. pendula</i> Maxim.	'Pendula'	16		
		16		Oginuma 1987 a
				Okabe 1928 (as <i>P. itosakura</i> Siebold); Oginuma 1987 a;
				Iwatsubo et al. 2002
<i>P. pseudocerasus</i> Lindl.		16	24	Okabe 1928 (as <i>P. itosakura</i> Siebold)
	'Pendula-rosea'	16	16	Iwatsubo et al. 2002
		32	24	Oginuma 1987 a,b
<i>P. sargentii</i> Rehd.		16	16	Okabe 1928; Oginuma 1987 b, 1988; Iwatsubo et al. 2002
				Okabe 1927, 1928 (as <i>P. sachalinensis</i> Miyoshi); Oginuma and Tanaka 1976; Oginuma 1987 a; Iwatsubo et al. 2003
<i>P. serrulata</i> Lindl. f.	'Longipes Miyoshi'	16	16	Iwatsubo et al. 2002, 2003
<i>P. ×subhirtella</i> Miq.	'Autumnalis'	16	24	Oginuma and Tanaka 1976; Oginuma 1987 a
			16	Okabe 1928 (as <i>P. subhirtella</i>); Sax 1931 (8 II, as <i>P. subhirtella</i>); Iwatsubo et al. 2002, 2003
	'Subhirtella'	16	24	Oginuma and Tanaka 1976; Oginuma 1987 a
<i>P. ×yedoensis</i> Matsum.	'Candida'	16		
	'Hayazaki-oshima'	16		
	'Kurama-zakura'	16		
<i>Prunus</i> sp.	'Izutaga-shiro' (伊豆多賀白)	16	16	Iwatsubo et al. 2002

of cytotypes: $2n=16$ and 24 chromosomes. However, recent (Oginuma 1987 a; Iwatsubo et al. 2002) and present studies for this taxon all showed $2n=16$ chromosomes. The diploid ($2n=16$) form appears to be more widely cultivated than the triploid ($2n=24$) form. The plant of *P. × subhirtella* ‘Subhirtella’ used in the study was a diploid plant with $2n=16$ chromosomes, but this taxon is known to have two cytological forms; one diploid with $2n=16$ chromosomes (Okabe 1928; Sax 1931; Iwatsubo et al. 2002, 2003) and the other triploid with $2n=24$ chromosomes (Oginuma and Tanaka 1976; Oginuma 1987 a). ‘Subhirtella’ is considered as a hybrid between *P. incisa* and *P. pendula* f. *ascendens* (Kawasaki 1994), both of which are known to be diploid taxa (Okabe, 1927, 1928; Sax 1931; Tanaka and Oginuma 1975; Oginuma and Tanaka 1976; Oginuma 1987 a; Iwatsubo et al. 2003). Thus, in the two cytological forms of ‘Subhirtella’, the diploid form appears to be a hybrid plant having one chromosome set from each of the two parent species.

References

- Ishikawa, M. 1916. A list of the number of chromosomes. Bot. Mag. Tokyo **30**: 404–448.
- Iwatsubo, Y., Kawasaki, T. and Naruhashi, N. 2002. Chromosome numbers of 193 cultivated taxa of *Prunus* subg. *Cerasus* in Japan. J. Phytogeogr. Taxon. **50**: 21–34.
- Iwatsubo, Y., Kawasaki, T. and Naruhashi, N. 2003. Chromosome numbers of 41 cultivated taxa of *Prunus* subg. *Cerasus* in Japan. J. Phytogeogr. Taxon. **51**: 165–168.
- Kawasaki, T. 1994. Nihon no Sakura (Flowering cherries of Japan). 383 pp. Yama-kei Publishers, Tokyo. (in Japanese)
- Nihon Hana no Kai. 2002. The guide on the flowering cherries (Sakura zukan) (CD-ROM). Nihon Hana no Kai (The Flower Association of Japan), Tokyo. (in Japanese)
- Nishikawa, T. 1985. Chromosome counts of flowering plants of Hokkaido (8). J. Hokkaido Univ. Educ., Sect. 2 B, **35**: 31–42. (in Japanese)
- Oginuma, K. 1987 a. Karyomorphological studies on *Prunus* in Japan. J. Sci. Hiroshima Univ., Ser. B, Div. 2, Bot. **21**: 1–66.
- Oginuma, K. 1987 b. Karyomorphological studies on some species of Japanese *Prunus*. Pl. Chromos. Res. 1987: 131–134.
- Oginuma, K. 1988. A cytological study of the tetraploid *Prunus pseudo-cerasus* ($2n=32$). La Kromosomo II–51–52: 1710–1714.
- Oginuma, K. 1993. Chromosomes of *Prunus zippeiana* (Rosaceae: Laurocerasus). Bull. Coll. Child Develop., Kochi Women’s Univ. **17**: 107–109.
- Oginuma, K. and Tanaka, R. 1976. Karyomorphological studies on some cherry trees in Japan. J. Jpn. Bot. **51**: 104–109.
- Okabe, S. 1927. Cytological studies on *Prunus*. Bot. Mag. Tokyo **41**: 398–404. (in Japanese with English summary)
- Okabe, S. 1928. Zur Cytologie der Gattung *Prunus*. Sci. Repts. Tohoku Univ., Ser. IV, Biol. **3**: 733–743.
- Sax, K. 1931. The origin and relationships of the Pomoideae. J. Arn. Arb. **12**: 3–22.
- Somego, M. 2000. Cytogenetic studies of the *Prunus yedoensis* Matsum. (Somei-yoshino). Sakura Science **7**: 14–22. (in Japanese with English summary)
- Tanaka, R. and Oginuma, K. 1975. Karyological study in *Prunus × yedoensis*. Chrom. Inf. Serv. **19**: 13–15.

(Received March 8, 2004; accepted April 26, 2004)

岩坪美兼¹・千木 容²・鳴橋直弘¹: サクラ属 *Cerasus* 亜属 36 分類群の染色体数

サクラ属 *Cerasus* 亜属 36 分類群 (野生種を含む) について、染色体数の観察を行った。31 分類群は二倍体、4 分類群は三倍体、そして 1 分類群は四倍体であった。観察の結果は以下のとおりである。これらのうち 23 分類群の染色体数は初めての報告である (*印は初報告の分類群)。

二倍体 ($2n=16$) のサクラ:

アカネヤエ*, アカミオオシマ, アマヤドリ*, イズタガシロ*, イトククリ*, ウスゲオオシマ*, ウスゲヤマザクラ*, オオヤマザクラ, オクミヤコ*, カンザキオオシマ*, クシマザクラ*, クラマザクラ, ゲバザクラ*, ゴザノマニオイ*, コヒガン, コマメザクラ*, シダレザクラ, ジュウガツザクラ, スジガクヤマザクラ*, センショウジキクザクラ, センダイヤ, タキニオイ*, ツクシザクラ*, ハヤザキオオシマ*, ベニシダレ, ベニタマニシキ, ホウリンジ*, ミドリヤマザクラ*, ヤエノオオシマザクラ,

ヤエムラサキザクラ*, ヤツブサザクラ*。

三倍体 ($2n=24$) のサクラ:

シバヤマ*, ツバキカンザクラ, フクザクラ*, ボタン*。

四倍体 ($2n=32$) のサクラ:

シナミザクラ。

以上の 36 分類群のなかで, ツバキカンザクラには二倍体の報告があり, コヒガン, シダレザクラ, シナミザクラ, ジュウガツザクラには三倍体の報告

がある。今回の石川県林業試験場に保存されているサクラの観察ではコヒガン, シダレザクラ, ジュウガツザクラは二倍体, ツバキカンザクラは三倍体, そしてシナミザクラは四倍体であり, いずれもサクラの染色体数に関するこのシリーズの以前の報告 (Iwatsubo et al. 2002, 2003) と一致した。

(¹〒930-8555 富山市五福 3190 富山大学理学部生物学科; ²〒920-2114 石川県石川郡鶴来町三宮 石川県林業試験場)