

# Activation of Bombesin receptor subtype-3 influences activity of orexin neurons by both direct and indirect pathways

著者	Furutani Naoki, Hondo Mari, Tsujino Natsuko, Sakurai Takeshi
journal or publication title	Journal of Molecular Neuroscience
volume	42
number	1
page range	106-111
year	2010-09-01
URL	<a href="http://hdl.handle.net/2297/25023">http://hdl.handle.net/2297/25023</a>

doi: 10.1007/s12031-010-9382-5

1  
2  
3 **Activation of Bombesin Receptor Subtype-3 Influences Activity of Orexin Neurons**  
4  
5  
6  
7 **by Both Direct and Indirect Pathways**  
8  
9

10 Naoki Furutani, Mari Hondo, Natsuko Tsujino, Takeshi Sakurai  
11  
12

13 Department of Molecular Neuroscience and Integrative Physiology, Graduate School of  
14  
15  
16  
17 Medical Science, Kanazawa University  
18  
19  
20

21 *This manuscript is related to the Yakushima 2009 symposium.*  
22  
23

24 a) Running title:  
25  
26

27  
28 BRS3 and orexin neurons  
29  
30

31 b) Takeshi Sakurai, M.D., Ph.D.  
32  
33

34  
35 Department of Molecular Neuroscience and Integrative Physiology, Graduate School of  
36  
37  
38  
39 Medical Science, Kanazawa University, 13-1 Takaramachi, Kanazawa-shi, Ishikawa  
40  
41  
42 920-8640, Japan  
43  
44

45  
46 e-mail: [tsakurai@med.kanazawa-u.ac.jp](mailto:tsakurai@med.kanazawa-u.ac.jp)  
47  
48

49  
50 Tel: +81-76-265-2173  
51

52  
53 Fax: +81-76-234-4224  
54  
55

56 **Number of Figures: 3**  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Number of Pages:** 20

**Keywords:** orexin, BRS3, patch clamp, calcium imaging, food intake, sleep

**Acknowledgements**

This study was supported by Grants-in-aid for Scientific Research (B), and for Young Scientists (B). Authors thank Dr. Wendy Gray for critical reading.

1  
2  
3 **Abstract**  
4  
5  
6

7 The neuropeptides orexin A and orexin B (also known as hypocretin 1 and hypocretin 2),  
8  
9  
10 produced in lateral hypothalamic neurons, are critical regulators of feeding behavior, the  
11  
12 reward system and sleep/wake states. Orexin-producing neurons (orexin neurons) are  
13  
14 regulated by various factors involved in regulation of energy homeostasis and  
15  
16 sleep/wakefulness states. Bombesin receptor subtype 3 (BRS3) is an orphan receptor that  
17  
18 might be implicated in energy homeostasis, and is highly expressed in the hypothalamus.  
19  
20  
21 However, the neural pathway by which BRS3 regulates energy homeostasis is largely  
22  
23  
24 unknown. We examined whether BRS3 is involved in the regulation of orexin neurons.  
25  
26  
27 Using a calcium imaging method, we found that a selective BRS3 agonist  
28  
29  
30 (Ac-Phe-Trp-Ala-His-( $\tau$ Bzl)-Nip-Gly-Arg-NH<sub>2</sub>) increased the intracellular calcium  
31  
32  
33 concentration of orexin neurons. However, intracellular recordings from slice  
34  
35  
36 preparations revealed that the BRS3 agonist hyperpolarized orexin neurons. The BRS3  
37  
38  
39 agonist depolarized orexin neuron in the presence of tetrodotoxin. Moreover, in the  
40  
41  
42 presence of GABA receptor blockers, picrotoxin and CGP55845, the BRS3 agonist  
43  
44  
45 induced depolarization and increased firing frequency. Additionally, double-label in situ  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 hybridization study revealed that *Brs3* mRNA was expressed in almost all orexin neurons  
4  
5  
6  
7 and many cells around these neurons. These findings suggest that the BRS3 agonist  
8  
9  
10 indirectly inhibited orexin neurons through GABAergic input and directly activated  
11  
12  
13 orexin neurons. Inhibition of activity of orexin neurons through BRS3 might be an  
14  
15  
16 important pathway for regulation of feeding and sleep/wake states. This pathway might  
17  
18  
19  
20  
21 serve as a novel target for the treatment of obesity.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 **Introduction**  
4  
5  
6

7 The neuropeptides orexin A and orexin B were identified as endogenous ligands for two  
8  
9  
10 orphan G-protein-coupled receptors (GPCRs) (Sakurai, 1998). Orexin-producing  
11  
12  
13 neurons (orexin neurons) are localized exclusively in the lateral hypothalamic area  
14  
15  
16 (LHA), which is known as the feeding center. Various evidence has suggested that this  
17  
18  
19  
20  
21 neuropeptide is involved in the regulation of feeding behavior (Sakurai, 1998; Edwards,  
22  
23  
24 1999; Haynes, 2000; Haynes, 2002). Subsequently, the finding that orexin deficiency  
25  
26  
27  
28 causes narcolepsy in humans and animals suggested that these hypothalamic  
29  
30  
31  
32 neuropeptides also play a critical role in regulating sleep and wakefulness (Chemelli,  
33  
34  
35 1999; Lin, 1999; Peyron, 2000; Thannickal, 2000; Hara, 2001).  
36  
37

38  
39         Recent studies of the efferent and afferent systems of orexin-producing neurons,  
40  
41  
42 as well as phenotypic characterization of mice with genetic alterations in the orexin  
43  
44  
45 system, have suggested further roles of orexin in the coordination of emotion, energy  
46  
47  
48 homeostasis, reward, drug addiction, and arousal. Orexin neurons receive abundant input  
49  
50  
51  
52 from the limbic system, preoptic sleep-promoting neurons, and hypothalamic neurons  
53  
54  
55  
56 implicated in energy homeostasis. Orexin neurons are also regulated by peripheral  
57  
58  
59  
60  
61

1  
2  
3 metabolic cues, including ghrelin, leptin and glucose, suggesting that orexin neurons  
4  
5  
6  
7 might provide a link between energy homeostasis and sleep/wake states (Yamanaka,  
8  
9  
10 2003b).

11  
12  
13  
14 Bombesin receptor subtype 3 (BRS3) is a member of the bombesin receptor  
15  
16  
17 subfamily of GPCRs, which also includes the neuromedin B- and gastrin-releasing  
18  
19  
20 peptide receptors (NMBR and GRPR) (Gorbulev, 1992). BRS3 couples to endogenous  
21  
22  
23 Gq family proteins and increases intracellular calcium concentration. BRS3 has a unique  
24  
25  
26 pharmacological profile and does not bind with high affinity with any known  
27  
28  
29 bombesin-related peptides. Although a number of molecules that can activate human  
30  
31  
32 BRS3 have been developed (Mantey, 1997; Liu, 2002; Weber, 2003; Boyle, 2005;  
33  
34  
35 Gonzalez, 2008; Zhang, 2009), its natural ligands are still unknown (Jensen, 2008).  
36  
37  
38  
39  
40  
41

42 BRS3 is primarily expressed in the CNS, and is present in the highest amounts  
43  
44  
45 in the hypothalamic area where orexin neurons are localized (Ohki-Hamazaki, 1997a;  
46  
47  
48 Yamada, 1999; Liu, 2002; Jennings, 2003; Sano, 2004). Although the normal  
49  
50  
51 physiological function of BRS3 is largely unknown, mice lacking the BRS3 gene develop  
52  
53  
54 late onset obesity, accompanied by a reduced metabolic rate and increased food intake  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 (Ohki-Hamazaki, 1997b). In good agreement with the *Brs3*<sup>-/-</sup> mouse study, a BRS3  
4  
5  
6  
7 agonist, Bag-1, reduced food intake and increased metabolic rate (Guan, 2010). These  
8  
9  
10 studies suggested that BRS3 might be an important regulator of feeding behavior and  
11  
12  
13 energy homeostasis. A recent study showed that the anorectic effects of the BRS3 agonist  
14  
15  
16 remained intact in *neuropeptide Y*<sup>-/-</sup>, *agouti-related peptide*<sup>-/-</sup>, *melanocortin 4 receptor*<sup>-/-</sup>,  
17  
18  
19 *cannabinoid receptor 1*<sup>-/-</sup> and *leptin receptor*<sup>db/db</sup> mice (Guan, 2010). These results  
20  
21  
22 suggest that the effect of the BRS3 agonist might be mediated through other feeding  
23  
24  
25 mechanisms. However, the neural pathways of feeding behavior mediated by BRS3 are  
26  
27  
28 largely undefined. The orexin pathway is one of the possible pathways by which the  
29  
30  
31 BRS3 agonist evokes its effects, because orexin neurons are regulated by several factors  
32  
33  
34 involved in energy metabolism, and expression of BRS3 were found in the LHA. To  
35  
36  
37 investigate this possibility, we evaluated the effect of a BRS3 agonist on orexin neurons.  
38  
39  
40  
41  
42 In this study, we found that a BRS3 agonist activated orexin neurons directly, while it  
43  
44  
45  
46  
47  
48  
49 also inhibited these neurons through activation of local GABAergic interneurons.  
50  
51  
52  
53  
54  
55

## 56 **Materials and Methods**

57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3 **Drugs**  
4  
5  
6

7 The BRS3 agonist, neuromedin B (Peptide Institute, Osaka, Japan), tetrodotoxin (TTX)  
8  
9  
10 (Wako, Osaka, Japan), picrotoxin (Sigma, St. Louis, MO), and CGP55845 (Wako) were  
11  
12  
13 dissolved in extracellular solution. We also used a synthesized a selective BRS3 agonist  
14  
15  
16  
17 (Ac-Phe-Trp-Ala-His ( $\tau$ Bzl)-Nip-Gly-Arg-NH<sub>2</sub>) (Boyle, 2005).  
18  
19  
20

21 **Animals**  
22  
23

24 All experimental procedures involving animals were approved by the Kanazawa  
25  
26  
27  
28 University Animal Care and Use Committee and were in accordance with NIH guidelines.  
29  
30

31 All efforts were made to minimize animal suffering and discomfort and to reduce the  
32  
33  
34  
35 number of animals used. Genetically-modified mice used in this study were  
36  
37  
38  
39 *Orexin/YC2.1* mice (Tsuji, 2005) and *Orexin/EGFP* mice (Yamanaka, 2003a).  
40  
41

42 ***In situ hybridization***  
43  
44

45 Double in situ hybridization was performed according to procedures previously described  
46  
47  
48  
49 (Mieda, 2006). Digoxigenin (DIG)-labeled riboprobes for *mBrs3* were synthesized from  
50  
51  
52  
53 a 1196 bp fragment of murine *Brs3* cDNA (nucleotides -172 - +1367 from the initiation  
54  
55  
56  
57 codon) containing 1196 bp of the whole coding region cloned into the pCRII vector  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 (Invitrogen). Fluorescein (FITC)-labeled riboprobes for *prepro-Orexin* were synthesized  
4  
5  
6  
7 by in vitro transcription. pBluescript II SK (+) containing a 0.5 kb fragment encoding  
8  
9  
10 Gln33-Val130 of *prepro-Orexin* was used as a template for in vitro transcription (Sakurai,  
11  
12  
13 1998). The DIG and FITC-labeled probes were detected by means of anti-DIG (1/2000)  
14  
15  
16 and anti-FITC (1/1000) antibodies conjugated with alkaline phosphatase (Roche  
17  
18 Diagnostics, Basel, Switzerland). Alkaline phosphatase activity was detected with  
19  
20  
21 NBT/BCIP and INT/BCIP (Roche Diagnostics).  
22  
23  
24  
25  
26

### 27 ***Brain slice preparation***

28  
29  
30 *Orexin/YC2.1* mice (3-8 weeks old) and *Orexin/EGFP* mice (3-6 weeks old) were  
31  
32 anesthetized with forane (Abbott, Osaka, Japan). The mice were decapitated under deep  
33  
34  
35 anesthesia. Brains were isolated in ice-cold cutting solution consisting of (mM): 280  
36  
37  
38 sucrose, 2 KCl, 10 HEPES, 0.5 CaCl<sub>2</sub>, 10 MgCl<sub>2</sub>, 10 glucose, pH 7.4, bubbled with 100%  
39  
40  
41 O<sub>2</sub>. Brains were cut coronally into 300- $\mu$ m slices with a vibratome (VTA-1000S, Leica,  
42  
43  
44 Germany). Slices containing the LHA were transferred for 1 hr to an incubation chamber  
45  
46  
47  
48  
49  
50  
51  
52  
53 at room temperature filled with physiological solution containing (mM): 125 NaCl, 2.5  
54  
55  
56 KCl, 1.25 NaH<sub>2</sub>PO<sub>4</sub>, 2.0 CaCl<sub>2</sub>, 1.0 MgSO<sub>4</sub>, 26 NaHCO<sub>3</sub>, 11 glucose, pH 7.4, bubbled  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 with 95% O<sub>2</sub>/5% CO<sub>2</sub>.  
4  
5

### 6 7 ***Calcium imaging of orexin neurons*** 8 9

10 Optical recordings were performed on a fluorescence microscope (BX51WI, Olympus,  
11 Tokyo, Japan) equipped with a cooled charge-coupled device (CCD) camera (CoolSNAP  
12 HQ2, Roper Scientific, Tucson, AZ) controlled by MetaFluor 5.0.7 software (Universal  
13 Imaging, West Chester, PA). YC2.1 was excited through a 440DF20 filter, and its  
14 fluorescent image was subjected to dual emission ratio imaging through two emission  
15 filters (480DF30 for ECFP, 535DF26 for EYFP) controlled by a filter changer (ProscanII,  
16 Prior Scientific Instruments, Cambridge, UK).  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

### 34 35 ***Electrophysiological recording*** 36 37

38 *Orexin/EGFP* mice were used for whole cell recordings. The slices were transferred to a  
39 recording chamber (RC-27L, Warner Instrument Corp., CT, USA) at room temperature  
40 on a fluorescence microscope stage (BX51WI, Olympus, Tokyo, Japan). Neurons that  
41 showed GFP fluorescence were used for patch-clamp recordings. The fluorescence  
42 microscope was equipped with an infrared camera (C-3077, Hamamatsu Photonics,  
43 Hamamatsu, Japan) for infrared differential interference contrast (IR-DIC) imaging and a  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 CCD camera (JK-TU53H, Olympus) for fluorescent imaging. Each image was displayed  
5  
6  
7 separately on a monitor. Recordings were carried out with an Axopatch 700B amplifier  
8  
9  
10 (Axon Instruments, Foster City, CA) using a borosilicate pipette (GC150-10, Harvard  
11  
12  
13 Apparatus, Holliston, MA) prepared by a micropipette puller (P-97, Sutter Instruments,  
14  
15  
16 Pangbourne, UK) and filled with intracellular solution (4-10 M $\Omega$ ), consisting of (mM):  
17  
18 125 K-gluconate, 5 KCl, 1 MgCl<sub>2</sub>, 10 HEPES, 1.1 EGTA-Na<sub>3</sub>, 5 MgATP, 0.5 Na<sub>2</sub>GTP,  
19  
20  
21 pH7.3 with KOH. Osmolarity of the solution was checked by a vapor pressure osmometer  
22  
23  
24 (model 5520, Wescor, Logan, UT). The osmolarity of the internal and external solutions  
25  
26  
27 was 280-290 and 320-330 mOsm/l, respectively. The liquid junction potential of the  
28  
29  
30 patch pipette and perfused extracellular solution was estimated to be -16.2 mV and was  
31  
32  
33 applied to the data. The recording pipette was under positive pressure while it was  
34  
35  
36 advanced toward an individual cell in the slice. A tight seal of 0.5-1.0 G $\Omega$  was made by  
37  
38  
39 applying negative pressure. The membrane patch was then ruptured by suction. The  
40  
41  
42 series resistance during recording was 10-25 M $\Omega$  and was compensated. The reference  
43  
44  
45 electrode was an Ag-AgCl pellet immersed in the bath solution. During recordings, cells  
46  
47  
48 were superfused with extracellular solution at a rate of 1.0-2.0 ml/min using a peristaltic  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 pump (K.T. Lab, Japan) at RT.  
4  
5  
6  
7  
8  
9

## 10 **RESULTS**

### 11 ***Selective BRS3 agonist influences activity of orexin neurons by direct and indirect*** 12 13 14 15 16 17 **pathways**

18  
19  
20  
21 Recent studies have suggested that the activity of orexin neurons is influenced by various  
22  
23  
24 factors involved in sleep/wake states or feeding behavior. It is also possible that BRS3  
25  
26  
27 may regulate the activity of orexin neurons, because *Brs3*<sup>-/-</sup> mice developed obesity and  
28  
29  
30 impairment of glucose metabolism (Ohki-Hamazaki, 1997b). To examine the possibility  
31  
32  
33 that BRS3 is involved in regulation of the activity of orexin neurons, we applied a  
34  
35  
36 selective BRS3 agonist (Ac-Phe-Trp-Ala-His (τBzl)-Nip-Gly-Arg-NH<sub>2</sub>) to orexin  
37  
38  
39 neurons using brain slice preparations of *Orexin/YC2.1* transgenic mice, in which orexin  
40  
41  
42 neurons specifically express the calcium-indicator protein, yellow cameleon (YC) 2.1  
43  
44  
45 (Tsuji, 2005), and observed the change in intracellular calcium concentration ([Ca<sup>2+</sup>]<sub>i</sub>)  
46  
47  
48  
49 of orexin neurons. In the presence of tetrodotoxin (TTX), the BRS3 agonist increased the  
50  
51  
52  
53  
54  
55  
56 YFP/CFP ratio, showing that [Ca<sup>2+</sup>]<sub>i</sub> was increased (Fig. 1A). The BRS3 agonist-induced  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 increase in  $[Ca^{2+}]_i$  was concentration-dependent (Fig. 1B) (n=3-6). Of YFP-positive  
4  
5  
6  
7 neurons tested, 91.8% were activated by the BRS3 agonist (n=49), but they showed no  
8  
9  
10 response to neuromedin B, an NMBR agonist (n=11).  
11

12  
13  
14 We also studied the effect of the BRS3 agonist on orexin neurons in detail by  
15  
16  
17 slice patch clamp recording. Unexpectedly, BRS3 agonist (1 $\mu$ M) bath application  
18  
19  
20  
21 potently hyperpolarized and decreased the firing frequency of orexin neurons under the  
22  
23  
24 whole cell current clamp mode (n=7, Fig. 2A). On the other hand, the BRS3 agonist  
25  
26  
27 (1 $\mu$ M) induced depolarization of orexin neurons in the presence of TTX (Fig. 2B).  
28  
29  
30  
31 Furthermore, the BRS3 agonist (1 $\mu$ M) induced depolarization and increased the firing  
32  
33  
34 frequency of orexin neurons in the presence of both picrotoxin and CGP55845 (GABA<sub>A</sub>  
35  
36  
37 and GABA<sub>B</sub> receptor blockers, respectively) (Fig. 2C). These observations suggest that  
38  
39  
40  
41  
42 the BRS3 agonist induced depolarization of orexin neurons directly, but more potently  
43  
44  
45 hyperpolarized these neurons through activation of GABAergic interneurons.  
46  
47

### 48 ***Brs3* and *Orexin* are colocalized in lateral hypothalamic neurons**

49  
50  
51 To examine the tissue localization of BRS3, we performed double-label in situ  
52  
53  
54 hybridization. We found that BRS3-expressing neurons were widely distributed in the  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 CNS. Relatively large numbers of cells were found in the cortex, hippocampus, amygdala  
4  
5  
6  
7 and LHA (Fig. 3A and B). We found that almost all neurons expressing *Orexin* mRNA  
8  
9  
10 expressed *Brs3* mRNA (Fig. 3C and D).  
11

## 12 13 14 **Discussion** 15

16  
17 Although BRS3 still remains an orphan receptor, it is thought to be a possible drug target  
18  
19  
20  
21 for treating obesity and the metabolic syndrome, because BRS3-deficient mice show late  
22  
23  
24  
25 onset obesity and impaired glucose metabolism.  
26

27  
28 In the present study, we found that a selective BRS3 agonist hyperpolarized all  
29  
30  
31 orexin neurons under current clamp recording (Fig. 2A) but depolarized them in the  
32  
33  
34 presence of GABA receptor antagonists, picrotoxin and CGP55845 (Fig. 2C). This result  
35  
36  
37  
38 suggests that the inhibitory effect on orexin neurons was mediated by activation of  
39  
40  
41  
42 GABAergic input. Consistently, under inhibition of neural input by TTX, the effect of the  
43  
44  
45 agonist on orexin neurons was excitatory, and intracellular  $Ca^{2+}$  of orexin neurons was  
46  
47  
48 increased presumably through activation of the Gq-subfamily of G- proteins (Fig. 2B). In  
49  
50  
51  
52 situ hybridization study revealed that *Brs3* mRNA was expressed widely in hypothalamic  
53  
54  
55  
56 areas (Fig. 3A and B), in agreement with previous studies (Ohki-Hamazaki, 1997a;  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 Yamada, 1999; Guan, 2010; Jennings, 2003; Sano, 2004). Double-label in situ  
4  
5  
6  
7 hybridization study showed that almost all orexin neurons also expressed *Brs3* mRNA  
8  
9  
10 (Fig. 3C and D). Both electrophysiological and histochemical analyses suggested that the  
11  
12  
13  
14 BRS3 agonist activated orexin neurons directly through activation of BRS3 on orexin  
15  
16  
17 neurons. However, the direct depolarizing effect of the BRS3 agonist appeared only when  
18  
19  
20  
21 neural input was suppressed by TTX or a GABA blocker. This suggests that the indirect  
22  
23  
24  
25 inhibitory effect predominated over the direct excitatory effect. Consistently, we  
26  
27  
28 observed many BRS3-positive neurons in the LHA by in situ hybridization, which could  
29  
30  
31  
32 possibly be GABAergic interneurons, although further studies are required to reveal the  
33  
34  
35 origin of GABAergic input to orexin neurons.  
36

37  
38  
39 From a physiological viewpoint, unknown natural ligands of BRS3 also might  
40  
41  
42 inhibit the activity of orexin neurons and suppress feeding behavior. Although a recent  
43  
44  
45 report suggested that a BRS3 agonist increased inhibitory synaptic transmission of the  
46  
47  
48 arcuate nucleus (Guan, 2010), other pathways regulating feeding behavior have not yet  
49  
50  
51  
52 been clarified. This study revealed a new pathway of a BRS3 agonist. Our results also  
53  
54  
55  
56 suggest the new possibility that BRS3 might be an important factor in sleep/wake  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

regulation, because the orexin system is considered to be a critical regulator of  
sleep/wake regulation.

1  
2  
3 **Figure legends**  
4  
5  
6

7 Fig. 1  
8  
9

10 FRET calcium imaging study showing that the BRS3 agonist influenced orexin neuron  
11 activity, using a hypothalamic slice from *Orexin/YC2.1* mice. A, In the presence of  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

FRET calcium imaging study showing that the BRS3 agonist influenced orexin neuron activity, using a hypothalamic slice from *Orexin/YC2.1* mice. A, In the presence of tetrodotoxin (TTX), bath application of the BRS3 agonist (1 $\mu$ M) increased the YFP/CFP ratio, suggesting that the BRS3 agonist (1 $\mu$ M) increased [Ca<sup>2+</sup>]<sub>i</sub> in orexin neurons. B, The BRS3 agonist induced a concentration-dependent increase in [Ca<sup>2+</sup>]<sub>i</sub> (EC<sub>50</sub>: 45.7  $\pm$  2.4 nM).

Fig. 2

A, Under whole cell current clamp mode, the BRS3 agonist (1 $\mu$ M) induced hyperpolarization of orexin neurons and decreased firing frequency. B, In the presence of TTX, orexin neurons were depolarized by the BRS3 agonist (1 $\mu$ M). C, In the presence of both picrotoxin (GABA<sub>A</sub> blocker) and CGP55845 (GABA<sub>B</sub> blocker), the BRS3 agonist (1 $\mu$ M) depolarized orexin neurons and increased their firing frequency.

1  
2  
3 Fig. 3  
4  
5  
6

7 Double-label in situ hybridization histochemistry showing distributions of *Orexin*  
8 (brown) and *Brs3* (blue) mRNA. A, *Brs3* mRNA was distributed widely in the CNS,  
9  
10 especially in the cortex, hippocampus, amygdala and LHA. B, Schematic representation  
11 of distribution of *Brs3* and *Orexin* mRNA (bregma -1.34 mm). C. High power view of  
12  
13 lateral hypothalamic area. D, High power view of region in rectangle in C. *Orexin* mRNA  
14  
15 colocalized with *Brs3* mRNA. Yellow arrowheads show colocalization.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3 **References**


- 4 Boyle RG, Humphries J, Mitchell T et al. (2005) The design of a new potent and selective  
5 ligand for the orphan bombesin receptor subtype 3 (BRS3). *J Pept Sci.* 11,  
6 136-141.  
7  
8  
9 Chemelli RM, Willie JT, Sinton CM et al. (1999) Narcolepsy in orexin knockout mice:  
10 molecular genetics of sleep regulation. *Cell.* 98, 437-451.  
11  
12 Edwards CM, Abusnana S, Sunter D, Murphy KG, Ghatei MA, Bloom SR (1999) The  
13 effect of the orexins on food intake: comparison with neuropeptide Y,  
14 melanin-concentrating hormone and galanin. *J Endocrinol.* 160, R7-12.  
15  
16 Gonzalez N, Hocart SJ, Portal-Nunez S et al. (2008) Molecular basis for agonist  
17 selectivity and activation of the orphan bombesin receptor subtype 3 receptor. *J*  
18 *Pharmacol Exp Ther.* 324, 463-474.  
19  
20 Gorbulev V, Akhundova A, Buchner H, Fahrenholz F (1992) Molecular cloning of a new  
21 bombesin receptor subtype expressed in uterus during pregnancy. *Eur J Biochem.*  
22 208, 405-410.  
23  
24 Guan XM, Chen H, Dobbelaar PH et al. (2010) Regulation of energy homeostasis by  
25 bombesin receptor subtype-3: selective receptor agonists for the treatment of  
26 obesity. *Cell Metab.* 11, 101-112.  
27  
28 Hara J, Beuckmann CT, Nambu T et al. (2001) Genetic ablation of orexin neurons in mice  
29 results in narcolepsy, hypophagia, and obesity. *Neuron.* 30, 345-354.  
30  
31 Haynes AC, Jackson B, Chapman H et al. (2000) A selective orexin-1 receptor antagonist  
32 reduces food consumption in male and female rats. *Regul Pept.* 96, 45-51.  
33  
34 Haynes AC, Chapman H, Taylor C et al. (2002) Anorectic, thermogenic and anti-obesity  
35 activity of a selective orexin-1 receptor antagonist in ob/ob mice. *Regul Pept.* 104,  
36 153-159.  
37  
38 Jennings CA, Harrison DC, Maycox PR, Crook B, Smart D, Hervieu GJ (2003) The  
39 distribution of the orphan bombesin receptor subtype-3 in the rat CNS.  
40 *Neuroscience.* 120, 309-324.  
41  
42 Jensen RT, Battey JF, Spindel ER, Benya RV (2008) International Union of  
43 Pharmacology. LXVIII. Mammalian bombesin receptors: nomenclature,  
44 distribution, pharmacology, signaling, and functions in normal and disease states.  
45 *Pharmacol Rev.* 60, 1-42.  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3 Lin L, Faraco J, Li R et al. (1999) The sleep disorder canine narcolepsy is caused by a  
4 mutation in the hypocretin (orexin) receptor 2 gene. *Cell*. 98, 365-376.  
5  
6 Liu J, Lao ZJ, Zhang J et al. (2002) Molecular basis of the pharmacological difference  
7 between rat and human bombesin receptor subtype-3 (BRS-3). *Biochemistry*. 41,  
8 8954-8960.  
9  
10 Mantey SA, Weber HC, Sainz E et al. (1997) Discovery of a high affinity radioligand for  
11 the human orphan receptor, bombesin receptor subtype 3, which demonstrates  
12 that it has a unique pharmacology compared with other mammalian bombesin  
13 receptors. *J Biol Chem*. 272, 26062-26071.  
14  
15 Mieda M, Williams SC, Richardson JA, Tanaka K, Yanagisawa M (2006) The  
16 dorsomedial hypothalamic nucleus as a putative food-entrainable circadian  
17 pacemaker. *Proc Natl Acad Sci U S A*. 103, 12150-12155.  
18  
19 Ohki-Hamazaki H, Wada E, Matsui K, Wada K (1997a) Cloning and expression of the  
20 neuromedin B receptor and the third subtype of bombesin receptor genes in the  
21 mouse. *Brain Res*. 762, 165-172.  
22  
23 Ohki-Hamazaki H, Watase K, Yamamoto K et al. (1997b) Mice lacking bombesin  
24 receptor subtype-3 develop metabolic defects and obesity. *Nature*. 390, 165-169.  
25  
26 Peyron C, Faraco J, Rogers W et al. (2000) A mutation in a case of early onset narcolepsy  
27 and a generalized absence of hypocretin peptides in human narcoleptic brains. *Nat*  
28 *Med*. 6, 991-997.  
29  
30 Sakurai T, Amemiya A, Ishii M et al. (1998) Orexins and orexin receptors: a family of  
31 hypothalamic neuropeptides and G protein-coupled receptors that regulate  
32 feeding behavior. *Cell*. 92, 573-585.  
33  
34 Sano H, Feighner SD, Hreniuk DL et al. (2004) Characterization of the bombesin-like  
35 peptide receptor family in primates. *Genomics*. 84, 139-146.  
36  
37 Thannickal TC, Moore RY, Nienhuis R et al. (2000) Reduced number of hypocretin  
38 neurons in human narcolepsy. *Neuron*. 27, 469-474.  
39  
40 Tsujino N, Yamanaka A, Ichiki K et al. (2005) Cholecystokinin activates  
41 orexin/hypocretin neurons through the cholecystokinin A receptor. *J Neurosci*. 25,  
42 7459-7469.  
43  
44 Weber D, Berger C, Eickelmann P, Antel J, Kessler H (2003) Design of selective  
45 peptidomimetic agonists for the human orphan receptor BRS-3. *J Med Chem*. 46,  
46 1918-1930.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65


- 1  
2  
3 Yamada K, Wada E, Imaki J, Ohki-Hamazaki H, Wada K (1999) Hyperresponsiveness to  
4 palatable and aversive taste stimuli in genetically obese (bombesin receptor  
5 subtype-3-deficient) mice. *Physiol Behav.* 66, 863-867.  
6  
7  
8 Yamanaka A, Muraki Y, Tsujino N, Goto K, Sakurai T (2003a) Regulation of orexin  
9 neurons by the monoaminergic and cholinergic systems. *Biochem Biophys Res*  
10 *Commun.* 303, 120-129.  
11  
12  
13 Yamanaka A, Beuckmann CT, Willie JT et al. (2003b) Hypothalamic orexin neurons  
14 regulate arousal according to energy balance in mice. *Neuron.* 38, 701-713.  
15  
16  
17 Zhang L, Nothacker HP, Wang Z, Bohn LM, Civelli O (2009) Pharmacological  
18 characterization of a selective agonist for bombesin receptor subtype-3. *Biochem*  
19 *Biophys Res Commun.* 387, 283-288.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Figure 1

[Click here to download high resolution image](#)




Furutani et al. Figure 1



Furutani et al. Figure 2





Furutani et al. Figure 3