

# Maxillary stability after Le Fort I osteotomy using three different plate systems

著者	Ueki Koichiro, Okabe Katsuhiko, Moroi Akinori, Marukawa Kohei, Sotobori Megumi, Ishihara Yuri, Nakagawa Kiyomasa
journal or publication title	International Journal of Oral and Maxillofacial Surgery
volume	41
number	8
page range	942-948
year	2012-08-01
URL	<a href="http://hdl.handle.net/2297/31962">http://hdl.handle.net/2297/31962</a>

doi: 10.1016/j.ijom.2012.02.023

## **Maxillary stability after Le Fort I osteotomy using 3 different plate systems**

KOICHIRO UEKI<sup>1</sup>, KATSUHIKO OKABE<sup>1</sup>, AKINORI MOROI<sup>1</sup>, KOHEI MARUKAWA<sup>1</sup>,  
MEGUMI SOTOBORI<sup>1</sup>, YURI ISHIHARA<sup>1</sup>, KIYOMASA NAKAGAWA<sup>1</sup>

### **Key words:**

Le Fort I osteotomy, Unsintered hydroxyapatite (u-HA) / poly-L-lactic acid (PLLA) plate,  
Poly-L-lactic acid (PLLA) plate, Titanium plate, Stability

Address correspondence to: Koichiro Ueki, DDS, PhD.

<sup>1</sup>Department of Oral and Maxillofacial Surgery, Graduate School of Medicine, Kanazawa  
University, 13-1 Takaramachi, Kanazawa 920-8641, Japan.

Tel: +81-76-265-2444; Fax: +81-76-234-4268

E-mail: [kueki@med.kanazawa-u.ac.jp](mailto:kueki@med.kanazawa-u.ac.jp)

## **Abstract**

The purpose of this study was to compare postoperative changes in maxillary stability after Le Fort I osteotomy with an unsintered hydroxyapatite (u-HA) / poly-L-lactic acid (PLLA) plate, a PLLA plate versus a titanium plate.

Subjects comprised 60 Japanese patients diagnosed with mandibular prognathism. All patients underwent Le Fort I osteotomy and bilateral sagittal split ramus osteotomy (SSRO). All patients were randomized in groups of 20 to a u-HA/PLLA group, a PLLA plate group and a titanium plate group. Changes in postoperative time intervals between the plate groups were compared using lateral and posteroanterior (PA) cephalography.

The uHA/PLLA group had significantly larger value than the PLLA group regarding change of mx1-S perpendicular to SN between 3 and 12 months (T3) ( $P=0.0269$ ). The uHA/PLLA group had a significantly larger value than PLLA group regarding change of S-A perpendicular to SN between baseline and 1 month (T1) ( $P=0.0257$ ). However there was no significant difference in the other measurements.

This study suggested that maxillary stability with satisfactory results could be obtained in the u-HA/PLLA group, PLLA plate group and titanium plate group, although there was a slight difference between the u-HA/PLLA plate system and PLLA plate system in Le Fort I osteotomy.

Comparison of the stability of Le Fort I advancement using biodegradable poly-p-dioxanon thread with titanium miniplates demonstrated good stability of both fixation devices in the anterior-posterior plane, but a tendency to relapse in the vertical dimension.<sup>17</sup>

Poly-L-lactic acid (PLLA) is one of various absorbable materials that has been used for fixation after Le Fort I osteotomy and sagittal split ramus osteotomy (SSRO). PLLA miniplates promote osteosynthesis of the oral and maxillofacial skeleton, and PLLA screws have been used in patients undergoing orthognathic surgery<sup>9,26,29</sup>. In our previous study, it was found that PLLA plates and screws (Fixorb<sup>®</sup>-MX, Takiron Co., Osaka, Japan) were useful in Le Fort I osteotomy with SSRO and intraoral vertical ramus osteotomy (IVRO), as well as the conventional titanium plate system.<sup>27</sup> Furthermore, fixation plate system (Super-FIXSORB<sup>®</sup>-MX, Takiron Co. Ltd, Osaka) has been newly developed for use in orthopedic or cranio-facial, oral and maxillofacial or plastic and reconstructive surgeries.<sup>20-22</sup> These devices are made from composites of uncalcined and unsintered hydroxyapatite (u-HA) particles and PLLA, and they are produced by a forging process, which is a unique compression molding, and machining treatment. They have a modulus of elasticity close to that of natural cortical bone, and can retain a high strength during the period required for bone healing. They can also show optimal degradation and resorption behavior, osteoconductivity, and bone bonding capability.

Material character and strength of plate were different in each product so that, it is difficult to compare with other studies that used different plate systems. Therefore, it is important to compare different types of plate systems in a study. In the previous study on stability after sagittal split ramus osteotomy, there were no significant differences in postoperative time-course changes between the u-HA/PLLA plate system, PLLA plate system and conventional titanium plate system.<sup>25</sup> However, there is no report of a study that examined the stability after Le Fort I osteotomy using the u-HA/PLLA plate system. The present study compared time-course changes in maxillary stability after Le Fort I osteotomy with the u-HA/PLLA plate system, PLLA plate system and conventional titanium plate system.

## **Patients and Methods**

### Patients:

Subjects comprised 60 Japanese adults (16 men, 44 women) presenting with jaw deformities diagnosed as mandibular prognathism with maxillary retrognathism. At the time of orthognathic surgery, mean patient age ranged from 16 to 48 years), with a mean age and standard deviation of  $23.9 \pm 6.9$  years. Time frame of the operation was from 2001 August to 2010 April. The cases whose pre and post-operative cephalograms could be actually obtained and other procedures were not performed were selected in this study. Adequate total sample number calculated by the power analysis software (G\*Power Version 3.12; program written by Franz Faul, University Kiel, Germany)<sup>7</sup> was more than 54, when the effect size was 0.25. The number of sample size 60 in the repeated measure ANOVA in this study was considered to be valid as a prospective study. Informed consent was obtained from the patients and the study was approved by Kanazawa University Hospital. This study was performed according to the guideline of Helsinki declaration.

### Surgery:

All 60 patients underwent Le Fort I osteotomy and bilateral SSRO (by the Obwegeser method) to advance the maxilla with and without impaction and set back the mandible. The distribution in the direction and amount of maxillary advancement in three groups was not significantly different. All patients received orthodontic treatment before and after surgery by two orthodontists. All patients received the operation by two surgeons (K.U. and K.N).

In 20 (men: 9, women:11) of the 60 patients, 2 uHA/PLLA L-type mini-plates (10×22×1.4 mm with 4 screws (2×8 mm), Super-Fixorb<sup>®</sup>-MX; Takiron Co., Osaka, Japan) and 2 straight uHA/PLLA plates (28×4.5×1.4 mm with 4 screws (2×8 mm),

Super-Fixorb<sup>®</sup>-MX; Takiron Co.) were used to fix the advanced maxilla and 2 uHA/PLLA mini-plates (28×4.5×1.5 mm with 4 screws (2×8 mm), Super-Fixorb<sup>®</sup>-MX; Takiron Co.) were used for bilateral internal fixation of the mandible (u-HA/PLLA Group). The patients in the u-HA/PLLA group ranged in age from 16 to 48 years, with a mean age and standard deviation of 26.4±8.6 years.

In 20 patients (men: 4, women:16), 2 PLLA L-type mini-plates (10×22×1.5 mm with 4 screws (2×8 mm), Fixorb<sup>®</sup>-MX; Takiron Co., Osaka, Japan) and 2 straight PLLA plates (28×4.5×1.5 mm with 4 screws (2×8 mm), Fixorb<sup>®</sup>-MX; Takiron Co.) were used to fix the advanced maxilla and 2 PLLA mini-plates (28×4.5×1.5 mm with 4 screws (2×8 mm), Fixorb<sup>®</sup>-MX; Takiron Co.) were used for bilateral internal fixation of the mandible (PLLA Group). The patients in the PLLA group ranged in age from 16 to 34 years, with a mean age and standard deviation of 23.8±6.4 years.

In the remaining 20 patients(men: 4, women:16), 2 L-type titanium mini-plates and 2 straight titanium mini-plates (4 holes / thickness 0.55 mm with 4 screws (2×5 mm), Würzburg titanium miniplate system; Leibinger Co., Freiburg, Germany) were used to fix the advanced maxilla and 2 titanium mini-plates (4 holes / thickness 0.55 mm with 4 screws (2×7 mm), Würzburg titanium miniplate system, Leibinger Co.) were used for bilateral internal-fixation of mandible (titanium group). The patients in the titanium group ranged in age from 16 to 32 years, with a mean age and standard deviation of 21.6±4.4 years.

After a few days of inter maxillary fixation (IMF), elastic was placed to maintain an ideal occlusion in the same manner in all the groups.

#### Cephalographic assessment:

All patients underwent lateral and posteroanterior (PA) cephalography to assess skeletal changes at 1, 3, and 12 months postoperative (Fig. 1). To assess maxillary stability, arbitrary points for the anterior nasal spine (ANS), and posterior nasal spine (PNS), point A and incisor edge were defined and measured as follows: from the preoperative images,

and subsequently transferred to all remaining radiographs. One skilled observer performed all digitization to minimize errors in the cephalometric method and was acceptable for the purposes of this study. Error analysis by digitization and remeasurement of 10 randomly selected cases generated an average error of less than 0.4 mm for the linear measurements and 0.5 degree for the angular measurements.

#### Lateral cephalometric analysis

S-A parallel to SN: distance between point A and sella parallel to SN plane

S-A perpendicular to SN: distance between point A and sella perpendicular to SN plane

S-PNS parallel to SN: distance between the arbitrary PNS and sella parallel to SN plane

S-PNS perpendicular to SN: distance between the arbitrary PNS and sella perpendicular to SN plane

mx1-S parallel to SN: distance between the incisor edge and sella parallel to SN plane

mx1-S perpendicular to SN: distance between the incisor edge and sella perpendicular to SN plane

S-ANS parallel to SN: distance between the arbitrary ANS and sella parallel to SN plane

S-ANS perpendicular to SN: distance between the arbitrary ANS and sella perpendicular to SN plane

#### PA cephalometric analysis

Mx-Md Midline: angle between the ANS-Menton line and the line perpendicular to the bilateral zygomatic frontal suture line.

right mx6 to Zy-Zy: distance between the most buccal point at the right molar crown and the line connecting the most lateral points of the bilateral zygomatic arches (Zy-Zy).

left mx6 to Zy-Zy: distance between the most buccal point at the left molar crown and Zy-Zy.

Occlusal cant: angle between Zy-Zy and the line from the most buccal point at the right

first molar crown to the most buccal point at the left molar crown.

#### Statistical analysis:

Data were statistically analyzed with StatView software, version 4.5 (ABACUS Concepts, Inc., Berkeley, CA, USA).

Each serial period was defined, and the differences between measurements were calculated as shown below.

T1: (baseline to 1 month)

T2: (1 month to 3 months)

T3: (3 months to 1 year)

The statistic calculation with repeated measure analysis of variance (ANOVA) was performed using class category (uHA/PLLA group, PLLA group and titanium group) and time-course (T1, T2 and T3). Then, comparisons among three groups in each time period (T1, T2 and T3) were performed using Scheffe's method. Significant differences were considered significant at  $p < 0.05$ .

#### **Results**

After surgery, no patient experienced complications such as wound infection or dehiscence, bone instability, or long-term malocclusion. There was no significant difference among three groups in the distribution in men and women with chi-square test. There was no significant difference among three groups in age with chi-square test (Table. 1). Mean setback was  $6.5 \pm 3.5$  mm on the right and  $6.1 \pm 3.7$  mm on the left in the uHA/PLLA group,  $5.9 \pm 3.1$  mm on the right and  $6.1 \pm 2.5$  mm on the left in the PLLA group, and  $6.3 \pm 3.5$  mm on the right and  $6.0 \pm 2.6$  mm on the left in the titanium group. There was no significant



difference between the three groups with student's t-test.

From the results of repeated measure ANOVA, significant differences were identified among the three groups in S-A perpendicular to SN (between subjects;  $F=11.310$ ;  $df=2$ ;  $P<0.0001$ ), mx1-S perpendicular to SN (between subjects;  $F=5.712$ ;  $df=2$ ;  $P=0.0055$ ) and S-ANS perpendicular to SN (between subjects;  $F=4.867$ ;  $df=2$ ;  $P=0.0112$ ) (Fig.1, Tables 2 and 3).

From the multiple comparison in each time period, the uHA/PLLA group had significantly larger value than the PLLA group regarding change of mx1-S perpendicular to SN between 3 and 12 months (T3) ( $P=0.0269$ ). The uHA/PLLA group had a significantly larger value than PLLA group regarding change of S-A perpendicular to SN between baseline and 1 month (T1) ( $P=0.0257$ ).

No significant differences were identified among the three groups in the other measurements on lateral cephalometric analysis and in all measurements in the PA cephalometric analysis (Fig. 1, Tables 2 and 4).

## **Discussion**

Most studies on stability in Le Fort I osteotomy with SSRO have suggested that no appreciable difference exists in wire versus plate and screw fixation for single-piece maxillary impactions and/or advancements.<sup>6,18,23,30</sup> However, inferior repositioning of the maxilla has been shown to be an unstable move regardless of the fixation method used.<sup>1,17,18</sup> Many studies have also illustrated that mandibular surgery in combination with maxillary surgery does not affect maxillary stability.<sup>2,10,11,19,30</sup>

Norholt et al<sup>15</sup> found significant differences in vertical positioning of the maxilla in a lateral cephalometric analysis after 6 weeks as the position became more superior compared with the postoperative situation in a study using Lactosorb® (Lorenz Surgical, Jacksonville, Fla, USA). In the study by Cheung et al.<sup>3</sup>, maxilla with bioresorbable plate fixation (2.0 compact plating system, Inion Ltd., Tampere, Finland) were confirmed to have minimal

relapse compared to titanium plate fixation starting from the 6<sup>th</sup> postoperative week, but vertical instability occurred in the early postoperative period. Costa et al.<sup>4</sup> found that superior displacement of the maxilla occurred mainly within the first 8 postoperative weeks in the study with Lactorsb®. On the other hand, in our previous study using the PLLA plate (Fixsorb®-MX), we found that the maxilla was stable in the horizontal plane but tended to displace superiorly following Le Fort I osteotomy with SSRO or IVRO.<sup>27</sup> The plate systems used in each study were different so that it was difficult to compare the results. However, it seemed that superior displacement at the anterior part of maxilla could occur after Le Fort I osteotomy with an absorbable plate.

The most commonly used polymers are homopolymers of PLLA and polyglycolides (PGA) and copolymers of polyglycolide-poly lactide (PLGA) or copolymers of L- and D-lactide, poly-L/D-lactide (PLDLA)<sup>24</sup>. There are methods to obtain strong malleable devices from PLLA or PLDLA, such as self-reinforced (SR) composite (Biosorb FX, Bionximplants Ltd, Tampere, Finland), as-polymerized PLLA and drawn PLLA (Fixorb®-MX)<sup>13, 14</sup>.

Fixorb®-MX, bioabsorbable ultra-high-strength PLLA developed for internal fixation of fractures, was fabricated by a drawing technique developed by Matsusue et al<sup>13, 14</sup>. The bending strength and anti-pull-out strength of Fixorb are higher than those of human cortex and lower than those of titanium plates. In vitro, Fixorb plates can maintain 80% of the early bending strength until 12 weeks postoperative. Fixorb requires a longer period to disappear than PGA/PLA copolymers. However, it has a higher strength than PGA/PLA copolymers such that it can be used for loading regions<sup>13, 14</sup>. To produce the PLLA for use in the miniplate system, after dissolution and molding of the PLLA as a biomechanical polymer (molecular weight, about 400 kDa), it was mechanical processed into rods made through uniaxial extension into various forms. The bending strength of PLLA used was 240 MPa, and the bending modulus was 13 GPa<sup>13</sup>. This strength is considerably greater than that of PLLA used in other system<sup>5</sup>. Maurer et al estimated that the stress of the material was postulated to have reached threshold values for stability. Maximum chewing force as determined by the finite element method analysis was 132N for Fixsorb®, 115N for

Biosorb® (Bionximplants Ltd., Tampere, Finland), and 46.4N for Lactosorb®<sup>15</sup>. Theoretically, it was considered that Fixorb also had sufficient strength to fix the maxilla-mandibular bone in orthognathic surgery.

The PLLA plates could be easily bent with a forceps at room temperature and stayed in the desired position without use of a heating device. PLLA plates are therefore easy to use for fixation of bone segments after osteotomy. Even after the PLLA plate was bent, it was strong enough to fix bone segments after mandibular osteotomy. However, the stability and strength of bent PLLA plates remain unclear. The fact that the PLLA plate bent easily at room temperature suggests that after bending, slight distortion may occur in vivo, although data supporting this assumption are lacking. On the other hand, the u-HA/PLLA could be bent with a forceps in 60 °C hot water. The stability of bent u-HA/PLLA seemed to be higher than bent PLLA, although this study could not detect the difference between the materials per se

However, there remain some problems, such as those given below, which still need to be solved in the search for better resorbable devices. The rigidity should be increased, although bending strength is sufficiently high in some devices. The degradation rate of high strength PLLA devices should be enhanced and period of time until the complete resorption should be shortened. Bioactivity such as bone conduction and bone bonding capability should be made available by using other bioactive materials. For the reasons, u-HA/PLLA (Super-Fixorb®-MX) was developed to overcome these problems.<sup>21</sup>

The u-HA denotes an inorganic compound which is neither calcined at 800-900 °C nor sintered at 1000-1400 °C, and is a raw material of a HA ceramic, it has almost the same composition as natural bone. Sintered HA is surface bioactive, but not bioresorbable. On the other hand, u-HA is bioabsorbable so that the u-HA/PLLA plate and screw could be absorbed in vivo. u-HA/PLLA plate system which have completed clinical tests in orthopedic, oral and maxillofacial surgeries exhibit total resorbability and osteological bioactivity such as the ability to directly bond to bone and osteoconductivity<sup>12,22,31</sup> as well as good biocompatibility and high stiffness retainable for a long period of time to achieve bone union.<sup>8</sup> The screw and plate of the u-HA/PLLA system contain 30 and 40 weight % of

u-HA in each. The u-HA/PLLA plate was higher than the PLLA plate (Fixorb<sup>®</sup>-MX) and human cortical bone in the bending strength (u-HA/PLLA: 200-270 MPa, PLLA: 200-250 MPa, Cortical bone: 100-200MPa) and the shearing strength (u-HA/PLLA: 120-145 MPa, PLLA: 90-95 MPa, Cortical bone: 100 MPa). Shikinami et al. documented the complete process of bioresorption and bone replacement of rods made of forged composites of unsintered hydroxyapatite particles/poly L-lactide (F-u-HA/PLLA) implanted in the femoral medullary cavities of rabbits.<sup>20</sup> From the results, it was found that morphological changes during biodegradation and bone replacement in the proximal medullary cavity took up to 4.5 years, while molecular weight and bending strength had decreased to 50 KDa and 200 MPa, respectively, after 6 months. Therefore, if the strength of the absorbable plate decreases and the bony healing between segments is not complete by at least 6 months after osteotomy, the skeletal stability can not sustain for a long time. However, they have a modulus of elasticity close to that of natural cortical bone, and they can retain a high strength during the period required for bone healing. They can also show optimal degradation and resorption behavior, osteoconductivity, and bone bonding capability, because of the HA content. Furthermore, the u-HA/PLLA plate and screw could be recognized in the computed tomography image, although the PLLA plate was completely radiolucent. Therefore, it is easy to judge whether the u-HA/PLLA plate or screw breaks or becomes displaced. However, it was thought that breakage of the screw head by the driver of the system device in the u-HA/PLLA occurred more frequently than with the PLLA system. Perhaps the u-HA content might reduce elasticity of the screw.

In this study, predictor variables were three types of plate (uHA/PLLA, PLLA, titanium plate) and cephalometric measurements in each time interval. Age, gender, race etc. were considered as the confounding variables. However, the significant difference was not found in the distribution of age, gender, race and setback amount so that the effects by confounding variables could be excluded. The difference in the preoperative skeletal pattern and advancement amount in maxilla did not identified among three groups as shown in Tables 2 and 4. Therefore, it could be considered that three groups with similar skeletal pattern were randomly selected in this study.

In the lateral cephalometric analysis, there were significant differences in the time-course changes in S-A perpendicular to SN, mx1-S perpendicular to SN and S-ANS perpendicular to SN among the three groups. For these reasons, it could be understood that postoperative superior displacement at the anterior part in maxilla tended to occur in the PLLA group. The uHA/PLLA group had a significantly larger value than PLLA group regarding change of S-A perpendicular to SN between baseline and 1 month. This suggested that the maxillary movement the PLLA group contained maxillary advancement with impaction. However, in the other time interval, except for mx1-S perpendicular to SN in T3, there was no significant difference among the groups, and differences in the changes were not clinically appreciable.

Analysis of the PA cephalography can be used to measure dentofacial asymmetry and determine occlusal cant, although advance movement was dominant in all patients in this study. A line was drawn connecting the most buccal points of the left and right maxillary first molars in this study. Standard PA cephalometric analyses do not include evaluation of the relationships of the occlusal plane to the horizontal. This represents an important deficiency, as leveling the occlusal plane, when necessary, should be a goal of surgical and orthodontic therapy. However, occlusal cant alone is insufficient to evaluate asymmetry, so Mx-Md Midline was also added for a more accurate evaluation. In the time-course of change in the result of the PA cephalogram, no significant differences were identified among the three groups. This suggests that uHA/PLLA and PLLA plates are as strong as titanium plate to fix the maxillary segment from the frontal view.

Bone strength of the maxilla should be considered when stability after Le Fort I osteotomy is examined. Our previous study showed that there were no significant differences in area of bone defect between the segmental gap after Le Fort I osteotomy among the plate types using 3-dimensional computed tomography.<sup>28</sup> Various factors such as preoperative bone thickness, occlusion, inter-maxillary traction, moving direction and amount, age and gender etc. were considered to be associated with healing at the anterior and lateral walls of the maxilla. The change in stress distribution at the region of the space between segments might also affect the change in bone area. At least, use of bone graft or

an alternative material may not be decided solely on the basis of the amount of bony gap by movement of the maxillary segment. It will also be necessary to examine the relationship between new bone formation and type of plate.

In conclusion, this study suggested that there was no significant difference time-course changes in most measurements among three groups, although there was significant difference in mx1-S perpendicular to SN in T3 between the u-HA/PLLA plate system and PLLA plate system in Le Fort I osteotomy. Clinically, maxillary skeletal and occlusal stability with satisfactory results could be obtained in the u-HA/PLLA group, PLLA plate group and titanium plate group without complication.

## References

- 1) Baker DL, Stoelinga PJW, Blijdorp PA, Brous JJI. Long-term stability after inferior maxillary repositioning by miniplate fixation. *Int J Oral Maxillofac Surg* 1992; 21: 320-6.
- 2) Bothur S, Blomqvist JE, Isaksson S. Stability of Le Fort I osteotomy with advanced: A comparison of single maxillary surgery and two-jaw procedure. *J Oral Maxillofac Surg* 1998; 56:1029-33.
- 3) Cheung LK, Yip IH, Chow RL. Stability and mobility of Le Fort I osteotomy with bioabsorbable fixation: a randomized controlled trial. *Int J Oral Surg* 2008; 37: 232-41.
- 4) Costa F, Robiony M, Zorzan E, Zerman N, Politi M. Stability of skeletal Class III malocclusion after combined maxillary and mandibular procedures: Titanium versus resorbable plates and screws for maxillary fixation. *J Oral Maxillofac Surg* 2006; 64: 642-51.
- 5) Daniel AU, Chang MKO, Andriano KP: Mechanical properties of biodegradable polymers and composites proposed for internal fixation of bone. *J Appl Biomater* 1: 57-78, 1990.
- 6) Egbert M, Hepworth B, Myall R, West R. Stability of Le Fort I osteotomy with maxillary advancement: A comparison of combined wire fixation and rigid fixation. *J Oral Maxillofac Surg* 1995; 53: 243-8.
- 7) Faul F, Erdfelder E, Buchner A, Lang AG. G\*power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods* 2007; 39: 175-191.
- 8) Furukawa T, Matsusue Y, Yasunaga T, Nakagawa Y, Okada Y, Shikinami Y, Okuno M, Nakamura T. Histomorphometric study on high-strength hydroxyapatite/poly (L-lactide) composite rods for internal fixation of bone fractures. *J Biomed Mater Res* 2000; 50: 410-9.
- 9) Harada K, Enomoto S. Stability after surgical correction of mandibular prognathism

- using the sagittal split ramus osteotomy and fixation with poly-L-lactic acid (PLLA) screws. *J Oral Maxillofac Surg* 1997; 55: 464-8.
- 10) Hennes JA, Wallen TR, Bloomquist DS, Crouch DL. Stability of simultaneous mobilization of the maxilla and mandible utilizing internal rigid fixation. *Int J Adult Orthod Orthognath Surg* 1988; 3: 127-41.
  - 11) Law JH, Rotskoff KS, Smith RJ. Stability following combined maxillary and mandibular osteotomies treated with rigid internal fixation. *J Oral Maxillofac Surg* 1989; 47: 128-36.
  - 12) Matsumoto M, Chosa E, Nabeshima K, Shikinami Y, Tajima N: Influence of bioresorbable, unsintered hydroxyapatite/poly-L-lactide composite films on spinal corc, nerve roots, and epidural space. *J Biomed Mater Res* 2002; 60: 101-9.
  - 13) Matsusue Y, Yamamuro T, Oka M, Shikinami Y, Hyon SH, Ikada Y. In vitro and in vivo studies on bioabsorbable ultra-high-strength poly (L-lactide) rods. *J Biomed Mater Res* 1992; 26: 1553-67.
  - 14) Matsusue Y, Yamamuro T, Yoshii S, Oka M, Ikada Y, Hyon S, Shikinami Y. Biodegradable screw fixation of rabbit tibia proximal osteotomies. *J Appl Biomater* 1991; 2: 1-12.
  - 15) Maurer P, Holweg S, Knoll WD, Schubert J. Study by finite element method of the mechanical stress of selected biodegradable osteosynthesis screws in sagittal ramus osteotomy. *Br J Oral Maxillofac Surg* 2002; 40: 76-83.
  - 16) Norholt SE, Pedersen TK. Le Fort I miniplate osteosynthesis: a randomized, prospective study comparing resorbable PLLA/PGA with titanium. *Int J Oral Maxillofac Surg* 2004; 33: 245-52.
  - 17) Obwegeser JA. Osteosynthesis using biodegradable Poly-p-dioxanon (PDS II) in Le Fort I osteotomy without postoperative intermaxillary fixation. *J Craniomaxillofac Surg* 1994; 22: 129-37.
  - 18) Proffit WR, Phillips C, Prewitt JW, Turvey TA. Stability after surgical-orthodontic correction of skeletal class III malocclusion. II. Maxillary advancement. *Int J Adult Orthod Orthognath Surg* 1991; 6: 71-80.




- 19) Proffit WR, Turvey TA, Phillips C. Orthognathic surgery: A hierarchy of stability. *Int J Adult Orthod Orthognath Surg* 1996; 11: 191-204.
- 20) Shikinami Y, Matsusue Nakamura T. The complete process of bioresorption and bone replacement using devices made of forged composite of raw hydroxyapatite particles/poly L-lactide (F-u-HA/PLLA). *Biomaterials* 2005; 26: 5542-51.
- 21) Shikinami Y, Okuno M. Bioresorbable devices made of forged composites of hydroxyapatite (HA) particles and poly-l-lactide (PLLA): Part I. Basic characteristics. *Biomaterials* 1999; 20: 859-77.
- 22) Shikinami Y, Okuno M. Bioresorbable devices made of forged composites of hydroxyapatite (HA) particles and poly-l-lactide (PLLA): Part II. Practical properties of miniscrews and miniplates. *Biomaterials* 2001; 22: 3197-211.
- 23) Skoczlas LJ, Ellis E, Fonseca RJ, Gallo WJ. Stability of simultaneous maxillary intrusion and mandibular advancement: A comparison of rigid and nonrigid fixation techniques. *J Oral Maxillofac Surg* 1988; 46: 1056-64.
- 24) Törmälä P. Biodegradable self-reinforced composite materials: manufacturing structure and mechanical properties. *Clin Mater* 1992; 10: 29-34.
- 25) Ueki K, Okabe K, Miyazaki M, Mukozawa A, Moroi A, Marukawa K, Nakagawa K, Yamamoto E: Skeletal stability after mandibular setback surgery: Comparisons among unsintered hydroxyapatite (u-HA) / poly-L-lactic acid (PLLA) plate, PLLA plate and titanium plate. *J Oral Maxillifac Surg* 2011 Jan 6 [Epub ahead of print].
- 26) Ueki K, Hashiba Y, Marukawa K, Okabe K, Nakagawa K, Alam S, Yamamoto E. Evaluation of bone formation after sagittal split ramus osteotomy with bent plate fixation using computed tomography. *J Oral Maxillofac Surg* 2009; 67: 1062-8.
- 27) Ueki K, Marukawa K, Shimada M, Nakagawa K, Alam S, Yamamoto E. Maxillary stability following Le Fort I osteotomy in combination with sagittal split ramus osteotomy and intraoral vertical ramus osteotomy: a comparative study between titanium miniplate and poly-L-lactic acid plate. *J Oral Maxillofac Surg* 2006; 64: 74-78.
- 28) Ueki K, Miyazaki M, Okabe K, Mukozawa K, Moroi A, Nakagawa K, Yamamoto E.

Assessment of bone healing after Le Fort I osteotomy with 3-dimensional computed tomography. *J Cranio-Maxillofac Surg* 2010 Aug 10 [Epub ahead of print].

- 29) Ueki K, Nakagawa K, Marukawa K, Takazakura D, Shimada M, Takatsuka S, Yamamoto E. Changes in condylar long axis and skeletal stability after bilateral sagittal split ramus osteotomy with poly-L-lactic acid or titanium plate fixation. *Int J Oral Maxillofac Surg* 2005; 34: 627-34.
- 30) Van Sickels JE, Richardson DA. Stability of orthognathic surgery: A review of rigid fixation. *Br J Oral Maxillofac Surg* 1996; 34: 279-85.
- 31) Yasunaga T, Matsusue Y, Furukawa T, Shikinami Y, Okuno M, Nakamura T. Bonding behavior of ultrahigh strength unsinistered hydroxyapatite particles/poly (L-lactide) composites to surface of tibia cortex in rabbits. *J Biomed Mater Res* 1999; 47: 412-9.

## Legends

Fig.1 Measurement points on PA and lateral cephalograms



**Table 1.** Patient demographic data. There was no significant difference among three groups in the distribution in men and women. There was no significant difference among three groups in age

	Number			Age(years)			
	Total	Men	Women	Mean	SD	Minimum	Maximum
uHA/PLLA group	20	9	11	26.4	8.6	16	48
PLLA group	20	4	16	23.8	6.4	16	34
Titanium group	20	4	16	21.6	4.4	16	32

**Table 2.** Cephalometric measurements. SD indicates standard deviation.

			Baseline		1 month		3 months		1 year	
			Mean	SD	Mean	SD	Mean	SD	Mean	SD
uHA/PLLA group	SNA	(dg)	81.6	2.8	82.9	3.6	83.2	2.7	83.3	3.3
	S-A parallel to SN	(mm)	59.9	4.0	62.1	4.8	62.3	4.3	62.8	4.9
	S-A perpend to SN	(mm)	64.8	4.6	65.5	5.0	64.4	4.2	64.9	4.2
	S-PNS parallel to SN	(mm)	16.2	2.5	17.0	2.7	17.6	2.9	18.0	3.8
	S-PNS perpend to SN	(mm)	49.7	3.8	48.1	4.6	47.9	3.6	48.6	4.8
	mx1-S parallel to SN	(mm)	62.5	6.2	64.5	5.6	65.5	5.7	65.5	6.5
	mx1-S perpend to SN	(mm)	69.6	2.9	69.9	2.6	69.7	2.7	70.3	2.9
	S-ANS parallel to SN	(mm)	65.4	4.3	68.2	4.9	68.0	4.5	68.7	5.2
	S-ANS perpend to SN	(mm)	57.8	3.5	57.9	3.6	57.1	3.1	57.8	3.2
	Mx-Md Midline	(dg)	-2.1	5.6	-1.0	1.8	-1.3	2.1	-0.9	1.9
	right mx6 to Zy-Zy	(mm)	49.5	5.9	46.8	5.3	47.4	4.2	48.5	5.5
	left mx6 to Zy-Zy	(mm)	49.9	5.8	47.5	4.7	48.7	5.2	49.8	6.1
	Occlusal cant	(dg)	1.4	2.6	1.5	1.7	1.9	2.2	1.9	2.1
PLLA group	SNA	(dg)	80.2	5.2	81.2	2.7	82.4	2.9	82.7	3.8
	S-A parallel to SN	(mm)	54.5	5.7	57.0	4.9	57.9	5.1	56.9	5.0
	S-A perpend SN	(mm)	61.2	4.6	58.6	5.3	58.6	5.1	57.2	5.0
	S-PNS parallel to SN	(mm)	12.9	4.2	15.2	4.3	15.0	3.8	15.0	3.2
	S-PNS perpend to SN	(mm)	47.7	3.5	45.1	4.0	45.5	4.0	45.5	3.7
	mx1-S parallel to SN	(mm)	57.6	7.0	60.3	6.1	61.4	6.9	59.8	6.3
	mx1-S perpend to SN	(mm)	65.6	3.7	65.5	4.1	65.1	3.7	64.2	3.8
	S-ANS parallel to SN	(mm)	57.6	5.1	59.6	4.4	60.5	4.8	59.9	5.2
	S-ANS perpend to SN	(mm)	54.5	3.9	53.3	5.2	52.5	4.6	51.8	4.5
	Mx-Md Midline	(dg)	1.5	4.5	0.1	2.0	0.2	2.0	0.2	2.3
	right mx6 to Zy-Zy	(mm)	47.3	5.1	45.1	5.7	44.3	5.6	43.6	5.9
	left mx6 to Zy-Zy	(mm)	48.1	4.6	46.5	5.7	45.9	4.6	44.6	6.7
	Occlusal cant	(dg)	-0.1	2.4	2.4	2.8	2.1	3.0	2.4	1.8
Titanium group	SNA	(dg)	80.7	3.6	82.9	4.0	83.2	4.1	83.5	3.7
	S-A parallel to SN	(mm)	58.1	6.6	60.8	6.6	61.2	6.4	61.3	6.0
	S-A perpend to SN	(mm)	62.7	4.0	61.8	4.7	62.6	5.0	63.0	4.5
	S-PNS parallel to SN	(mm)	14.1	3.7	15.9	3.6	15.7	3.4	14.9	2.6
	S-PNS perpend to SN	(mm)	49.3	4.0	48.0	4.6	48.2	4.2	48.6	4.4
	mx1-S parallel to SN	(mm)	61.8	6.6	64.6	7.1	65.4	7.6	65.0	7.3

mx1-S perpend to SN	(mm)	61.4	7.1	60.8	7.3	60.7	7.9	61.0	7.3
S-ANS parallel to SN	(mm)	61.1	7.5	63.4	7.1	63.6	6.8	63.9	6.5
S-ANS perpend to SN	(mm)	56.9	3.9	56.1	4.2	56.7	4.5	57.0	4.3
Mx-Md Midline	(dg)	2.6	7.5	1.2	2.0	0.7	2.0	1.5	1.7
right mx6 to Zy-Zy	(mm)	45.8	5.8	44.4	6.9	46.2	5.4	45.1	5.9
left mx6 to Zy-Zy	(mm)	47.4	7.2	45.9	6.8	46.7	5.0	46.4	5.1
Occlusal cant	(dg)	-0.2	3.5	1.5	1.9	1.2	2.0	0.8	1.6

**Table 3.** The results of repeated measure ANOVA. \*:Significant differences (between subjects) were identified among the three groups.

S-A perpendicular to SN	Degree of freedom	Sum of squares	Mean sum of squares	F-value	P-value
Class	2	79.950	39.975	11.310	<.0001*
Groups residual	57	201.474	3.535		
Category for time course	2	24.161	12.080	1.076	0.344
Category for time course*Class	4	102.999	25.750	2.294	0.064
Category for time course*Groups residual	114	1279.000	11.224		

mx1-S perpendicular to SN	Degree of freedom	Sum of squares	Mean sum of squares	F-value	P-value
Class	2	15.311	7.655	5.712	0.0055*
Groups	57	76.394	1.340		
Category for time course	2	2.473	1.239	0.311	0.733
Category for time course*Class	4	22.060	5.515	1.384	0.244
Category for time course*Groups residual	114	454.192	3.984		

ANS-S perpendicular to SN	Degree of freedom	Sum of squares	Mean sum of squares	F-value	P-value
Class	2	34.846	17.423	4.867	0.0112*
Groups	57	204.053	3.580		
Category for time course	2	15.921	7.961	1.019	0.364
Category for time course*Class	4	30.563	7.641	0.978	0.423
Category for time course*Groups residual	114	890.612	7.812		

**Table 4.** Time interval value of cephalometric data. SD indicates standard deviation. T1: (baseline to 1 month), T2: (1 month to 3 months), T3: (3 months to 1 year). \* shows significant difference between groups at P<0.05.

			T1		T2		T3	
			Mean	SD	Mean	SD	Mean	SD
uHA/PLLA group	SNA	(dg)	1.3	2.6	0.3	1.9	0.0	2.0
	S-A parallel to SN	(mm)	2.2	1.6	0.2	2.0	0.5	2.7
	S-A perpend to SN	(mm)	0.6*	3.2	-1.1	2.1	0.4	1.9
	S-PNS parallel to SN	(mm)	0.8	2.3	0.6	2.9	0.4	4.2
	S-PNS perpend to SN	(mm)	-1.6	1.9	-0.1	1.9	0.6	2.1
	mx1-S parallel to SN	(mm)	2.0	1.9	1.1	2.0	0.0	3.1
	mx1-S perpend to SN	(mm)	0.3	1.2	-0.3	1.5	0.6 *	1.7
	S-ANS parallel to SN	(mm)	2.8	1.8	-0.2	2.5	0.6	2.6
	S-ANS perpend to SN	(mm)	0.1	2.1	-0.8	1.7	0.7	1.7
	Mx-Md Midline	(dg)	1.1	5.2	-0.3	2.0	0.4	1.9
	right mx6 to Zy-Zy	(mm)	-2.7	4.9	0.6	3.2	1.1	3.1
	left mx6 to Zy-Zy	(mm)	-2.4	5.2	1.2	3.9	1.1	4.2
	Occlusal cant	(dg)	0.2	3.0	0.3	2.5	0.0	2.2
PLLA group	SNA	(dg)	1.1	4.5	1.2	2.5	0.3	2.3
	S-A parallel to SN	(mm)	2.5	2.6	0.8	3.2	-1.0	1.3
	S-A perpend to SN	(mm)	-2.6*	4.7	0.0	2.1	-1.4	1.5
	S-PNS parallel to SN	(mm)	2.3	3.5	-0.2	2.7	0.0	2.3
	S-PNS perpend to SN	(mm)	-2.7	2.9	0.4	1.2	0.1	1.1
	mx1-S parallel to SN	(mm)	2.8	3.3	1.0	3.0	-1.5	2.2
	mx1-S perpend to SN	(mm)	-0.1	2.9	-0.4	1.1	-0.9*	1.6
	S-ANS parallel to SN	(mm)	2.0	2.2	0.9	3.0	-0.7	2.1
	S-ANS perpend to SN	(mm)	-1.3	4.1	-0.8	2.5	-0.8	1.6
	Mx-Md Midline	(dg)	-1.4	3.9	0.1	1.6	-0.1	1.9
	right mx6 to Zy-Zy	(mm)	-2.2	4.5	-0.8	2.9	-0.7	6.3
	left mx6 to Zy-Zy	(mm)	-1.6	5.1	-0.6	3.7	-1.3	6.6
	Occlusal cant	(dg)	2.5	3.8	-0.3	3.1	0.2	2.5
Titanium group	SNA	(dg)	2.2	2.9	0.3	3.5	0.3	2.7
	S-A parallel to SN	(mm)	2.7	2.6	0.4	3.5	0.2	3.1
	S-A perpend to SN	(mm)	-0.8	2.9	0.7	3.2	0.4	3.6
	S-PNS parallel to SN	(mm)	1.8	4.0	-0.2	2.8	-0.8	2.2
	S-PNS perpend to SN	(mm)	-1.3	2.1	0.2	1.3	0.4	2.1



mx1-S parallel to SN	(mm)	2.9	2.6	0.7	3.9	-0.4	2.8
mx1-S perpend to SN	(mm)	-0.6	1.8	-0.1	1.3	0.3	2.1
S-ANS parallel to SN	(mm)	2.2	2.6	0.2	3.4	0.4	2.6
S-ANS perpend to SN	(mm)	-0.8	2.4	0.6	3.2	0.3	2.6
Mx-Md Midline	(dg)	-1.4	6.7	-0.4	2.1	0.8	1.9
right mx6 to Zy-Zy	(mm)	-1.4	7.1	1.7	4.3	-1.0	4.1
left mx6 to Zy-Zy	(mm)	-1.5	8.6	0.7	4.7	-0.3	3.8
Occlusal cant	(dg)	1.6	3.5	-0.3	2.3	-0.4	2.0