

# One-pot oxidative carbon-carbon bond formation of 3-benzylic and 3-allylic indoles with carbon nucleophiles

著者	Matsuo Jun-ichi, Tanaki Yumi, Ishibashi Hiroyuki
journal or publication title	Tetrahedron
volume	64
number	22
page range	5262-5267
year	2005-05-01
URL	<a href="http://hdl.handle.net/2297/9912">http://hdl.handle.net/2297/9912</a>

doi: 10.1016/j.tet.2008.03.033


## Graphical Abstract

To create your abstract, type over the instructions in the template box below.  
Fonts or abstract dimensions should not be changed or altered.

### One-pot oxidative carbon-carbon bond formation of 3-benzylic and 3-allylic indoles with carbon nucleophiles

Jun-ichi Matsuo,\* Yumi Tanaki and Hiroyuki Ishibashi\*

*Division of Pharmaceutical Sciences, Graduate School of Natural Science and Technology, Kanazawa University, Kakuma-machi, Kanazawa 920-1192, Japan.*




Leave this area blank for abstract info.



for derivatizing them, and, to the best of our knowledge, such one-pot bond formation has not been reported to date.

We would like to report here a new method for generating free indolenine **2** by dehydrogenation of 3-alkylindoles **5** with **4** at  $-78$  °C, and unexpectedly high reactivity of indolenine **2** toward carbon nucleophiles such as ethyl cyanoacetate. The reaction profile of organometallic agents toward **2** is also described.



Scheme 1. Generation of indolenine **2** by oxidation of **5** with **4** and one-pot carbon-carbon bond formation.

## 2. Results and discussion

### 2.1. One-pot C-C bond formation of 3-benzylic and 3-allylic indoles with active methylene compounds

First, suitable reaction conditions for oxidation of 3-benzylindole (**5a**) with **4** were investigated (Table 1). Lithiation of **5a** with *n*-BuLi followed by reaction with **4** at  $-78$  °C proceeded immediately. In order to trap the indolenine intermediate **2a**, ethyl cyanoacetate was added. Although gramine derivatives were reported to react with active methylene compounds above at room temperature,<sup>3,4</sup> the present C-C bond formation proceeded at surprisingly low temperature (at  $-78$  °C) to afford adduct **3a** in 81% yield (entry 1). Moreover, the intermediate **2a** reacted even at  $-100$  °C (entry 2). Elevation of reaction temperature from  $-78$  °C to room temperature did not improve the yield of **3a**. The use of lithium hexamethyldisilazide (LHMDS) instead of *n*-BuLi also gave **3a** in 81% yield (entry 3). It was noted that lithium ion was superior to sodium or potassium ion as a counter cation of metallated indole **6**, since sodium hexamethyldisilazide (NHMDS) and potassium hexamethyldisilazide (KHMDS) gave **3a** in 61 and 33% yields, respectively (entries 4 and 5). The effect of metal ion might be explained by regioselectivity (C-3 vs. N) in the reaction of **6** and **4**. The use of diisopropylethylamine as a base gave only a trace amount of the desired product **3a** (entry 6).

Table 1. Generation of free indolenine **2a** by oxidation of **5a** with **4** and successive addition of ethyl cyanoacetate.<sup>a</sup>


entry	base	oxidation temperature	addition conditions	yield (%) <sup>b</sup>
1	<i>n</i> -BuLi	$-78$ °C	$-78$ °C, 30 min	81
2	<i>n</i> -BuLi	$-100$ °C	$-100$ °C, 30 min	63
3	LHMDS	$-78$ °C	$-78$ °C, 30 min	81
4	NHMDS	$-78$ °C	$-78$ °C, 30 min	61
5	KHMDS	$-78$ °C	$-78$ °C, 30 min	33
6	<i>i</i> -Pr <sub>2</sub> NEt	$-78$ °C to rt	$-78$ °C to rt	trace

<sup>a</sup> **5a** (1.0 equiv), base (1.3 equiv), **4** (1.5 equiv) and ethyl cyanoacetate (1.5 equiv) were employed.

<sup>b</sup> Isolated yield. A mixture of diastereomers (D.r.(diastereomer ratio) = 50:50) was obtained in each case.

We then investigated the substrate generality of **4**-mediated oxidative alkylation of 3-alkylindoles (Table 2). In addition to ethyl cyanoacetate, ethyl allylcyanoacetate and ethyl benzylcyanoacetate reacted to afford the corresponding alkylated products in 83% and 78% yields, respectively (entries 1 and 2). Malonic acid esters (entries 3-6), ketoesters (entry 7), and 1,3-diketones (entries 8 and 9) also reacted to give the products in moderate to good yields, whereas the reaction with nitromethane proceeded very sluggishly (12% yield). Next, several 3-benzylic and 3-allylic indoles were subjected to this oxidative alkylation. 3-(4-Chlorobenzyl)indole (**5b**) and 3-(4-methoxybenzyl)indole (**5c**) reacted smoothly to afford the corresponding adducts **3k** (78%) and **3l** (72%), respectively. 3-Cinnamylindole (**5d**) also reacted with ethyl cyanoacetate regioselectively to afford **3m** in 84% yield, while 3-allylindole (**5e**) gave **3n** in 36% yield along with its regioisomer **3n'** (11%). On the other hand, the present oxidative C-C bond formation of 3-methylindole and 3-ethylindole with ethyl cyanoacetate gave only a trace amount of products. The unprecedented high reactivity of **2** toward active methylene compounds may be ascribed to facilitated attack of carbanion to protonated indolenine **8** under neutral conditions (see Scheme 1).

Table 2. Oxidative carbon-carbon bond formation of 3-benzylic and 3-allylic indoles with various carbon nucleophiles bearing active methylene or methine protons.




5a: R = Ph; 5b: R = 4-ClC<sub>6</sub>H<sub>4</sub>; 5c: R = 4-MeOC<sub>6</sub>H<sub>4</sub>; 5d: R = CH=CHPh; 5e: R = CH=CH<sub>2</sub>

entry	indol	NuH	product	yield (%) <sup>a</sup>
1	5a			83 <sup>b</sup>
2	5a			78 <sup>c</sup>
3	5a			66 (78) <sup>d</sup>
4	5a			75
5	5a			55
6	5a			71
7	5a			54 <sup>d,e</sup>
8	5a			56
9	5a			63
10	5b			78 <sup>f</sup>
11	5c			72 <sup>f</sup>
12	5d			84 <sup>g</sup>
13	5e			36 <sup>h</sup>

<sup>a</sup> Isolated yield. <sup>b</sup> D.r. = 64:36. <sup>c</sup> D.r. = 67:33. <sup>d</sup> Determined by <sup>1</sup>H NMR analysis using an internal standard. <sup>e</sup> D.r. = 56:44. <sup>f</sup> D.r. = 50:50.


<sup>g</sup> D.r. = 60:40. <sup>h</sup> D.r. = 60:40. Regioisomer 3n' was also obtained in 11% yield



## 2.2. One-pot C-C bond formation of 3-benzylindole (5a) with organometallic agents

Organometallic agents were next examined as a nucleophile in the present one-pot C-C bond formation of 5a. After screening some organometallic reagents, alkyllithium (MeLi, 25% yield) and Grignard reagents (MeMgBr, 32% yield; PhMgBr, 35% yield) were found to be not suitable for the present carbon-carbon bond formation, while alkylzinc (Me<sub>2</sub>Zn, 50% yield; Et<sub>2</sub>Zn, 70% yield) gave the alkylated products in moderate yields. It was found that higher-order cyanocuprates gave the desired alkylated and arylated products in good yields (Table 3). Methyl, *n*-butyl, *t*-butyl, and phenyl groups were introduced smoothly at the benzylic position of 5a.

Table 3. Oxidative carbon-carbon bond formation of 5a with various higher-order cyanocuprates.<sup>a</sup>



entry	R	product	yield (%) <sup>b</sup>
1	Me	9a	54
2	<i>n</i> -Bu	9b	77
3	<i>t</i> -Bu	9c	54
4	Ph	9d	82

<sup>a</sup> About reaction conditions, see Table 1.

<sup>b</sup> Determined by <sup>1</sup>H NMR analysis.

## 3. Conclusion

3-Benzyl and 3-allyl indoles were activated by oxidation with 4 to form free indolenines 2 at -78 °C, and various carbon nucleophiles such as active methylene compounds and organocuprates reacted with them at -78 °C in a one-pot manner. The present oxidative activation of indoles with 4 will be applicable to various types of nucleophiles and to stereoselective bond formation because of high reactivity observed on 2. The present method can be regarded as a new method for generating free indolenine. Recently, Ir<sup>16</sup>- or Pd<sup>17</sup>-catalyzed direct alkylation of indoles with benzylic or allylic alcohols to the corresponding 3-alkylindoles has been reported.<sup>18</sup> Various 3-alkylindoles are now readily prepared, and the present method will be useful for rapid and further derivatization of 3-alkylindoles. Therefore, a diverse chemical library of 3-substituted indoles can be constructed and it would provide an opportunity to discover a more potent drug candidate.

## 4. Experimental

### 4.1. General

All melting points were determined on a Yanagimoto micro melting point apparatus and are uncorrected. Infrared (IR) spectra were recorded on a Shimadzu FTIR-8100. <sup>1</sup>H

NMR spectra were recorded on a JEOL JNM EX270 (270 MHz) or a JEOL JNM GSX500 (500 MHz) spectrometer; chemical shifts ( $\delta$ ) are reported in parts per million relative to tetramethylsilane. Splitting patterns are designated as s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet.  $^{13}\text{C}$  NMR spectra were recorded on a JEOL JNM GSX500 (500 MHz) spectrometer with complete proton decoupling. Chemical shifts are reported in parts per million relative to tetramethylsilane with the solvent resonance as the internal standard  $\text{CDCl}_3$  and  $\text{DMSO-d}_6$ . High resolution mass spectra (HRMS) were recorded on a JEOL JMS-SX-102A mass spectrometer. Elemental analyses were carried out on a Yanaco CHN Corder MT-5. Analytical TLC was performed on Merck precoated TLC plates (silica gel 60 GF254, 0.25 mm). Silica-gel column chromatography was carried out on silica gel 60N (Kanto Kagaku Co., Ltd., spherical, neutral, 63–210  $\mu\text{m}$ ). Preparative thin-layer chromatography (PTLC) was carried out on silica gel Wakogel B-5F. THF was distilled under argon from sodium/benzophenone ketyl. All oxidative carbon-carbon bond formations were carried out under argon in dried glassware with magnetic stirring.

*N*-*tert*-Butylbenzenesufinimidoyl chloride (**4**) was prepared according to the modified literature procedure<sup>19</sup> of employing 1.3 equivalents of *N,N*-dichloro-*tert*-butylamine, and **1** was stored in a refrigerator. 3-Alkylindoles (**5a**,<sup>20</sup> **5b**,<sup>20</sup> **5d**,<sup>17</sup> and **5e**<sup>17</sup>) were prepared by the reported methods. Ethyl allylcyanoacetate and ethyl benzylcyanoacetate were prepared by alkylation of ethyl cyanoacetate (*t*-BuOK and alkylating agent). Other active methylene compounds and methine compounds were purchased and used after distillation.

#### 4.2. Typical procedure for the oxidative C-C bond formation of 3-alkylindole with active methylene compounds (Table 1, entry 1)

To a stirred solution of **5a** (50 mg, 0.241 mmol) in THF (3 mL) was added dropwise a solution of *n*-BuLi (1.65 *N* in hexane, 0.19 mL, 0.31 mmol) at  $-78^\circ\text{C}$ . After the mixture was stirred for 15 min at the same temperature, a solution of **4** (77 mg, 0.36 mmol) in THF (1 mL) was added at  $-78^\circ\text{C}$ . After the mixture was stirred for 30 min at the same temperature, ethyl cyanoacetate (0.04 mL, 0.38 mmol) was added at  $-78^\circ\text{C}$ , and the reaction mixture was stirred for 30 min at the same temperature. The reaction mixture was then treated with saturated aqueous  $\text{NaHCO}_3$ , and the resulting mixture was extracted with ethyl acetate (three times). The combined organic extracts were dried over anhydrous  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated in vacuo. The crude product was purified by thin layer chromatography on silica gel (hexane/ethyl acetate = 5/1) to afford **3a** (62 mg, 0.195 mmol, 81%) as a white solid.

##### 4.2.1. 2-Cyano-3-(1*H*-indol-3-yl)-3-phenylpropionic Acid Ethyl Ester (**3a**)<sup>8b</sup>

A mixture of diastereomers (63:27); white solid: mp 79.5–80.5  $^\circ\text{C}$  (hexane-ethyl acetate) [lit.<sup>8b</sup> 91–93  $^\circ\text{C}$  (from aqueous AcOH)];  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ , major

diastereomer)  $\delta$  1.10 (t,  $J = 7.5$  Hz, 3H), 4.07–4.20 (m, 2H), 4.32 (d,  $J = 6.7$  Hz, 1H), 5.04 (d,  $J = 6.7$  Hz, 1H), 7.01–7.08 (m, 1H), 7.23–7.39 (m, 6H), 7.46 (d,  $J = 7.3$  Hz, 2H), 8.15 (s, 1H);  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ , minor diastereomer)  $\delta$  1.08 (t,  $J = 7.5$  Hz, 3H), 4.07–4.20 (m, 2H), 4.19 (d,  $J = 6.7$  Hz, 1H), 5.10 (d,  $J = 6.7$  Hz, 1H), 7.01–7.08 (m, 1H), 7.15–7.20 (m, 1H), 7.23–7.39 (m, 6H), 7.50 (d,  $J = 2.4$  Hz, 1H), 8.19 (s, 1H);  $^{13}\text{C}$  NMR (67.8 MHz,  $\text{CDCl}_3$ , major diastereomer)  $\delta$  13.6, 43.1, 44.2, 62.9, 111.3, 113.1, 116.1, 118.6, 119.6, 122.1, 122.5, 126.5, 127.6, 127.9, 128.6, 136.0, 139.6, 165.2;  $^{13}\text{C}$  NMR (67.8 MHz,  $\text{CDCl}_3$ , minor diastereomer)  $\delta$  13.7, 43.1, 45.0, 62.8, 111.3, 114.5, 116.3, 118.8, 119.7, 122.2, 122.4, 126.0, 127.8, 128.3, 128.6, 136.2, 138.6, 165.3.

##### 4.2.2. 2-Allyl-2-Cyano-3-(1*H*-indol-3-yl)-3-phenylpropionic Acid Ethyl Ester (**3b**)

Diastereomers were separated by preparative TLC (hexane/benzene/ethyl acetate = 8/8/1). The less polar diastereomer; white powder: mp 107.5–108.5  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.99 (t,  $J = 7.0$  Hz, 3H), 2.67 (dd,  $J = 6.7$  Hz, 14 Hz, 2H), 2.79 (dd,  $J = 7.9$  Hz, 14 Hz, 1H), 4.01 (q,  $J = 7.0$  Hz, 2H), 4.78 (s, 1H), 5.11–5.15 (m, 2H), 5.72–5.80 (m, 1H), 7.13 (t,  $J = 7.6$  Hz, 1H), 7.20 (t,  $J = 7.3$  Hz, 2H), 7.24–7.28 (m, 2H), 7.36 (d,  $J = 7.9$  Hz, 1H), 7.58 (d,  $J = 7.3$  Hz, 2H), 7.64 (d,  $J = 7.9$  Hz, 1H), 7.71 (d,  $J = 2.4$  Hz, 1H), 8.34 (s, 1H);  $^{13}\text{C}$  NMR (125.7 MHz,  $\text{CDCl}_3$ )  $\delta$  13.7, 41.7, 48.4, 56.7, 62.5, 111.3, 113.1, 118.1, 119.5, 120.0, 120.1, 122.5, 122.5, 127.5, 127.8, 128.5, 128.6, 130.8, 135.3, 139.5, 168.1; IR ( $\text{CHCl}_3$ ,  $\text{cm}^{-1}$ ) 3021, 2359, 1736, 1211; Anal. Calcd for  $\text{C}_{23}\text{H}_{22}\text{N}_2\text{O}_2$ : C, 77.07; H, 6.19; N, 7.82. Found: C, 76.74; H, 6.16; N, 7.68.

The more polar diastereomer; white powder: mp 150.5–151  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.99 (t,  $J = 7.0$  Hz, 3H), 2.40 (dd,  $J = 6.7$  Hz, 13.7 Hz, 1H), 2.72 (dd,  $J = 7.9$  Hz, 13.7 Hz, 1H), 4.01 (q,  $J = 7.0$  Hz, 2H), 4.78 (s, 1H), 5.16 (d,  $J = 4.9$  Hz, 1H), 5.18 (s, 1H), 5.76–5.82 (m, 1H), 7.00 (t,  $J = 7.6$  Hz, 1H), 7.12 (t,  $J = 7.0$  Hz, 1H), 7.25–7.36 (m, 5H), 7.55 (d,  $J = 6.7$  Hz, 2H), 7.80 (d,  $J = 2.4$  Hz, 1H), 8.24 (s, 1H);  $^{13}\text{C}$  NMR (125.7 MHz,  $\text{CDCl}_3$ )  $\delta$  13.7, 42.1, 48.6, 55.3, 62.6, 111.0, 114.2, 118.6, 119.0, 119.6, 120.7, 121.9, 122.4, 126.8, 127.9, 128.6, 129.6, 130.6, 135.5, 138.1, 168.2; IR ( $\text{CHCl}_3$ ,  $\text{cm}^{-1}$ ) 3478, 2359, 1740, 1227; Anal. Calcd for  $\text{C}_{23}\text{H}_{22}\text{N}_2\text{O}_2$ : C, 77.07; H, 6.19; N, 7.82. Found: C, 76.83; H, 6.18; N, 7.76.

##### 4.2.3. 2-Benzyl-2-Cyano-3-(1*H*-indol-3-yl)-3-phenylpropionic Acid Ethyl Ester (**3c**)

Diastereomers were separated by preparative TLC (benzene/ethyl acetate = 100/1). The less polar diastereomer; white amorphous:  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.81 (t,  $J = 7.2$  Hz, 3H), 3.19 (d,  $J = 13.6$  Hz, 1H), 3.36 (d,  $J = 13.6$  Hz, 1H), 3.79–3.89 (m, 2H), 4.94 (s, 1H), 7.14–7.26 (m, 10H), 7.38 (d,  $J = 8.1$  Hz, 1H), 7.61 (d,  $J = 8.1$  Hz, 2H), 7.72 (d,  $J = 8.1$  Hz, 1H), 7.82 (d,  $J = 2.4$  Hz, 1H), 8.38 (s, 1H);  $^{13}\text{C}$  NMR (125.7 MHz,  $\text{CDCl}_3$ )  $\delta$  13.5, 43.2, 49.0, 58.6, 62.4, 111.4, 113.2, 118.2, 119.5, 120.0, 122.6, 122.7, 127.5, 127.6, 127.9, 128.4, 128.5, 128.7,

129.9, 134.5, 135.4, 139.5, 168.0; IR (CHCl<sub>3</sub>, cm<sup>-1</sup>) 3476, 1736; HRMS (EI) Calculated for C<sub>27</sub>H<sub>24</sub>N<sub>2</sub>O<sub>2</sub>: 408.18378. Found: 408.18390.

The more polar diastereomer; white amorphous: <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.80 (t, *J* = 7.1 Hz, 3H), 2.88 (d, *J* = 13.4 Hz, 1H), 3.28 (d, *J* = 13.4 Hz, 1H), 3.80-3.89 (m, 2H), 4.94 (s, 1H), 7.02 (t, *J* = 7.4 Hz, 1H), 7.13 (t, *J* = 7.6 Hz, 1H), 7.20-7.40 (m, 10H), 7.65 (d, *J* = 7.3 Hz, 2H), 7.77 (d, *J* = 2.4 Hz, 1H), 8.14 (s, 1H); <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>) δ 13.5, 43.7, 49.3, 56.9, 62.5, 111.0, 114.3, 118.7, 119.2, 119.7, 121.9, 122.4, 126.9, 127.7, 127.9, 128.4, 128.7, 129.8, 129.8, 134.2, 135.5, 138.2, 168.1; IR (CHCl<sub>3</sub>, cm<sup>-1</sup>) 3478, 2245, 1736, 1242; HRMS (EI) Calculated for C<sub>27</sub>H<sub>24</sub>N<sub>2</sub>O<sub>2</sub>: 408.18378. Found: 408.18410.

#### 4.2.4. Ethyl 2-[1*H*-indol-3-yl(phenyl)methyl]malonate (3d)<sup>8d</sup>

White silky filer: mp 183.5-184.0 °C (hexane-ethyl acetate) [ref.<sup>8d</sup> 165-167 °C (from EtOH)]; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.97 (t, *J* = 7.1 Hz, 3H), 1.00 (t, *J* = 7.1 Hz, 3H), 3.95-4.30 (m, 4H), 4.29 (d, *J* = 11.8 Hz, 1H), 5.08 (d, *J* = 11.8 Hz, 1H), 7.02 (t, *J* = 7.7 Hz, 1H), 7.10-7.14 (m, 3H), 7.20-7.26 (m, 3H), 7.36 (d, *J* = 7.6 Hz, 2H), 7.54 (d, *J* = 8.1 Hz, 1H), 8.09 (s, 1H); <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>) δ 13.7, 13.7, 42.6, 58.4, 61.4, 61.4, 111.0, 116.9, 119.3, 119.4, 120.9, 122.2, 126.7, 126.7, 128.2, 128.3, 136.2, 141.4, 167.8, 168.0.

#### 4.2.5. Ethyl 2-[1*H*-indol-3-yl(phenyl)methyl]-2-methylmalonate (3e)

Pale yellow oil: <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.93 (t, *J* = 7.0 Hz, 3H), 1.10 (t, *J* = 7.0 Hz, 3H), 1.66 (s, 3H), 3.91-3.99 (m, 2H), 4.08 (q, *J* = 7.0 Hz, 2H), 5.36 (s, 1H), 6.99 (t, *J* = 7.6 Hz, 1H), 7.09-7.15 (m, 2H), 7.18-7.21 (m, 2H), 7.27 (d, *J* = 7.9 Hz, 1H), 7.40 (d, *J* = 2.4 Hz, 1H), 7.44 (dd, *J* = 7.0 Hz, 1.5 Hz, 3H), 8.11 (s, 1H); <sup>13</sup>C NMR (67.8 MHz, CDCl<sub>3</sub>) δ 13.5, 13.7, 19.1, 46.7, 59.2, 61.3, 61.3, 110.8, 115.2, 119.1, 119.2, 121.9, 122.4, 126.5, 127.7, 127.9, 130.3, 135.3, 140.2, 171.2, 171.5; IR (CHCl<sub>3</sub>, cm<sup>-1</sup>) 3480, 2350, 1728; HRMS (EI) Calculated for C<sub>23</sub>H<sub>25</sub>NO<sub>4</sub>: 379.17836. Found 379.17920.

#### 4.2.6. Ethyl 2-benzyl-2-[1*H*-indol-3-yl(phenyl)methyl]malonate (3f)

Colorless needles: mp 135.5-136.0 °C; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.85 (t, *J* = 7.3 Hz, 3H), 1.01 (t, *J* = 7.3 Hz, 3H), 3.43 (d, *J* = 14.0 Hz, 1H), 3.60 (d, *J* = 14.0 Hz, 1H), 3.84-4.03 (m, 4H), 5.16 (s, 1H), 6.98-7.01 (m, 3H), 7.10-7.19 (m, 7H), 7.31 (d, *J* = 8.5 Hz, 1H), 7.40 (d, *J* = 7.9 Hz, 3H), 7.72 (d, *J* = 2.4 Hz, 1H), 8.11 (s, 1H); <sup>13</sup>C NMR (67.8 MHz, CDCl<sub>3</sub>) δ 13.4, 13.6, 42.0, 48.3, 60.8, 60.9, 64.4, 110.8, 115.3, 118.8, 119.3, 121.8, 122.9, 126.5, 126.6, 127.5, 127.8, 127.9, 130.2, 130.3, 135.4, 136.8, 140.4, 170.1, 171.1; IR (CHCl<sub>3</sub>, cm<sup>-1</sup>) 3480, 1717; Anal. Calcd for C<sub>27</sub>H<sub>29</sub>NO<sub>4</sub>: C, 76.46; H, 6.42; N, 3.07. Found: C, 76.63; H, 6.51; N, 3.09.

#### 4.2.7. Benzyl 2-[1*H*-indol-3-yl(phenyl)methyl]malonate (3g)

White powder: mp 135.8-136.2 °C; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 4.42 (d, *J* = 12.0 Hz, 1H), 4.90-4.97 (m, 4H),

5.12 (d, *J* = 12.0 Hz, 1H), 6.94 (d, *J* = 7.3 Hz, 2H), 7.02-7.05 (m, 4H), 7.16-7.28 (m, 11H), 7.32 (d, *J* = 7.3 Hz, 2H), 7.51 (d, *J* = 7.9 Hz, 1H), 7.87 (s, 1H); <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>) δ 42.9, 58.3, 67.1, 67.1, 111.0, 116.4, 119.2, 119.5, 121.1, 122.2, 126.5, 126.7, 127.9, 128.0, 128.0, 128.1, 128.1, 128.3, 128.4, 135.0, 135.1, 136.2, 141.1, 167.6, 167.7; IR (CHCl<sub>3</sub>, cm<sup>-1</sup>) 3478, 2386, 1755, 1732; Anal. Calcd for C<sub>32</sub>H<sub>27</sub>NO<sub>4</sub>: C, 78.51; H, 5.56; N, 2.86. Found: C, 78.45; H, 5.60; N, 2.92.

#### 4.2.8. Ethyl 2-[1*H*-indol-3-yl(phenyl)methyl]-3-oxobutanoate (3h)<sup>8c</sup>

A mixture of diastereomers (83:17); white powder: mp 162-162.5 °C (from hexane/benzene = 1:1) [ref.<sup>8c</sup> 162-163 °C (EtOH)]; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>, major diastereomer) δ 0.97 (t, *J* = 7.3 Hz, 3H), 2.04 (s, 3H), 3.94-4.00 (m, 2H), 4.50 (d, *J* = 11.6 Hz, 1H), 5.09 (d, *J* = 11.6 Hz, 1H), 7.01-7.35 (m, 9H), 7.54 (d, *J* = 7.9 Hz, 1H), 8.03 (s, 1H); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>, minor diastereomer) δ 0.99 (t, *J* = 7.3 Hz, minor diastereomer 3H), 2.14 (s, 3H), 3.94-4.00 (m, 2H), 4.38 (d, *J* = 12.3 Hz, 1H), 5.06 (d, *J* = 12.2 Hz, 1H), 7.01-7.35 (m, 9H), 7.54 (d, *J* = 7.9 Hz, 1H), 8.06 (s, 1H); <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>, major diastereomer) δ 13.8, 30.3, 42.6, 61.4, 65.9, 111.0, 117.3, 119.4, 119.5, 120.7, 122.3, 126.8, 128.1, 128.1, 128.4, 128.6, 136.2, 141.3, 168.0.

#### 4.2.9. 3-[1*H*-Indol-3-yl(phenyl)methyl]pentane-2,4-dione (3i)<sup>8d</sup>

Colorless plates: mp 153.5-154.0 °C (from hexane-ethyl acetate = 5:2) [lit.<sup>8d</sup> mp 150-152 °C (from EtOH)]; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 1.93 (s, 3H), 2.05 (s, 3H), 4.64 (d, *J* = 12.2 Hz, 1H), 5.09 (d, *J* = 12.2 Hz, 1H), 7.04 (t, *J* = 7.0 Hz, 1H), 7.11-7.16 (m, 3H), 7.23 (t, *J* = 7.6 Hz, 2H), 7.30 (t, *J* = 7.9 Hz, 3H), 7.53 (d, *J* = 7.9 Hz, 1H), 8.09 (brs, 1H); <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>) δ 27.6, 31.3, 43.2, 75.3, 111.1, 116.6, 119.0, 119.8, 121.2, 122.6, 126.3, 126.8, 128.1, 128.6, 136.2, 141.3, 203.5, 204.1; IR (CHCl<sub>3</sub>, cm<sup>-1</sup>) 3478, 1721, 1696, 1356.

#### 4.2.10. 2-[1*H*-Indol-3-yl(phenyl)methyl]-1,3-diphenylpropane-1,3-dione (3j)

White powder: mp 171-172 °C; <sup>1</sup>H NMR (500 MHz, DMSO) δ 5.34 (d, *J* = 11.5 Hz, 1H), 6.89-7.02 (m, 4H), 7.07 (t, *J* = 7.6 Hz, 2H), 7.20 (d, *J* = 7.8 Hz, 1H), 7.37-7.41 (m, 4H), 7.50-7.57 (m, 5H), 7.61 (s, 1H), 8.03 (d, *J* = 7.0 Hz, 2H), 8.08 (d, *J* = 7.0 Hz, 2H), 10.8 (brs, 1H); <sup>13</sup>C NMR (125.7 MHz, DMSO) δ 43.7, 60.0, 111.1, 116.7, 118.3, 118.5, 121.0, 122.1, 125.9, 126.5, 127.7, 128.5, 128.5, 128.6, 128.6, 128.7, 133.4, 135.8, 136.3, 136.4, 142.6, 193.4, 194.4; IR (CHCl<sub>3</sub>, cm<sup>-1</sup>) 3480, 1728; HRMS (EI) Calculated for C<sub>30</sub>H<sub>23</sub>O<sub>2</sub>N: 429.17288. Found: 429.17405.

#### 4.2.11. 2-Cyano-3-(1*H*-indol-3-yl)-3-(4-chlorophenyl)propionic Acid Ethyl Ester (3k)

A mixture of diastereomers (50:50); colorless oil: <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 1.09 (t, *J* = 7.3 Hz, 3H), 1.13 (t, *J* = 7.3 Hz, 3H), 4.07-4.16 (m, 5H), 4.31 (d, *J* = 6.1 Hz, 1H), 5.02 (d, *J* = 6.7 Hz, 1H), 5.08 (d, *J* = 6.1 Hz, 1H), 7.01-7.10 (m, 2H), 7.17-7.38 (m, 15H), 7.46 (d, *J* = 2.4 Hz, 1H), 8.23 (s, 1H), 8.28 (s, 1H); <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>) δ 13.7,

13.8, 42.4, 42.5, 44.0, 44.8, 63.0, 111.3, 111.4, 112.8, 114.3, 115.9, 116.0, 118.6, 118.8, 119.9, 120.0, 122.0, 122.1, 122.7, 122.8, 125.9, 126.4, 128.9, 129.0, 129.4, 129.8, 133.6, 133.8, 136.1, 136.2, 137.1, 138.2, 165.0, 165.0; IR (CHCl<sub>3</sub>, cm<sup>-1</sup>) 3476, 1745, 1223; HRMS (EI) Calculated for C<sub>20</sub>H<sub>17</sub>N<sub>2</sub>O<sub>2</sub>Cl: 352.09786. Found 352.09807.

#### 4.2.12. 2-Cyano-3-(1*H*-indol-3-yl)-3-(4-methoxyphenyl)propionic Acid Ethyl Ester (3l)

A mixture of diastereomers (50:50); white powder: mp 128-128.5 °C; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 1.08 (t, *J* = 7.0 Hz, 3H), 1.12 (t, *J* = 7.0 Hz, 3H), 3.74 (s, 3H), 3.75 (s, 3H), 4.07-4.16 (m, 5H), 4.30 (d, *J* = 6.1 Hz, 1H), 4.99 (d, *J* = 6.1 Hz, 1H), 5.05 (d, *J* = 6.7 Hz, 1H), 6.80-6.84 (m, 4H), 7.00 (t, *J* = 7.3 Hz, 1H), 7.04 (t, *J* = 7.3 Hz, 1H), 7.15-7.36 (m, 11H), 7.45 (d, *J* = 2.4 Hz, 1H), 8.21 (brs, 1H), 8.25 (brs, 1H); <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>) δ 13.7, 13.8, 42.4, 42.5, 44.5, 45.2, 55.2, 55.2, 62.8, 62.8, 111.2, 111.3, 113.6, 114.0, 114.1, 115.1, 116.2, 116.4, 118.8, 119.0, 119.8, 121.9, 122.0, 122.4, 122.5, 126.1, 126.6, 128.5, 128.6, 128.9, 129.0, 129.1, 129.5, 130.7, 131.8, 136.1, 136.3, 159.0, 159.1, 165.3, 165.3; IR (CHCl<sub>3</sub>, cm<sup>-1</sup>) 3478, 2253, 1744, 1512, 1252; HRMS (EI) Calculated for C<sub>21</sub>H<sub>20</sub>O<sub>3</sub>N<sub>2</sub>: 348.14740. Found: 348.14812.

#### 4.2.13. 2-Cyano-3-(1*H*-indol-3-yl)-5-phenylpenta-4-enoic Acid Ethyl Ester (3m)

A mixture of diastereomers (60:40); colorless oil: <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>, major diastereomer) δ 1.20 (t, *J* = 7.0 Hz, 3H), 4.05-4.08 (m, 1H), 4.21 (q, *J* = 7.0 Hz, 2H), 4.60-4.63 (m, 1H), 6.51-6.70 (m, 2H), 7.11-7.39 (m, 9H), 7.64 (d, *J* = 7.9 Hz, 1H), 8.23 (brs, 1H); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>, minor diastereomer) δ 1.06 (t, *J* = 7.0 Hz, 3H), 4.05-4.08 (m, 3H), 4.60-4.63 (m, 1H), 6.51-6.70 (m, 2H), 7.11-7.39 (m, 9H), 7.64 (d, *J* = 7.9 Hz, 1H), 8.26 (s, 1H); <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>, mixture of diastereomers) δ 13.7, 14.0, 41.2, 41.3, 44.6, 44.7, 62.8, 62.8, 111.4, 111.6, 112.6, 113.7, 115.8, 116.1, 118.3, 118.8, 119.9, 120.0, 122.2, 122.4, 122.5, 122.7, 125.5, 125.6, 126.2, 126.5, 126.6, 127.5, 127.9, 127.9, 128.6, 132.6, 133.8, 136.1, 136.2, 136.3, 136.4, 165.2; IR (CHCl<sub>3</sub>, cm<sup>-1</sup>) 3478, 2359, 1745; HRMS (EI) Calculated for C<sub>22</sub>H<sub>20</sub>N<sub>2</sub>O<sub>2</sub>: 344.15248. Found 344.15233.

#### 4.2.14. 2-Cyano-3-(1*H*-indol-3-yl) penta-4-enoic Acid Ethyl Ester (3n)

A mixture of diastereomers (60:40); colorless oil: <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>, major diastereomer) δ 1.24 (t, *J* = 7.0 Hz, 3H), 3.99 (d, *J* = 6.1 Hz, 1H), 4.22 (q, *J* = 7.0 Hz, 2H), 4.44-4.47 (m, 1H), 5.25-5.37 (m, 2H), 6.13-6.28 (m, 1H), 7.11-7.24 (m, 3H), 7.38 (d, *J* = 7.9 Hz, 1H), 7.59 (t, *J* = 8.2 Hz, 1H), 8.20 (brs, 1H); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>, minor diastereomer) δ 1.08 (t, *J* = 7.0 Hz, 3H), 3.99 (d, *J* = 6.1 Hz, 1H), 4.07 (q, *J* = 7.0 Hz, 2H), 4.44-4.47 (m, 1H), 5.25-5.37 (m, 2H), 6.13-6.28 (m, 1H), 7.11-7.24 (m, 2H), 7.31 (d, *J* = 2.4 Hz, 1H), 7.36 (d, *J* = 7.9 Hz, 1H), 7.59 (t, *J* = 8.2 Hz, 1H), 8.23 (brs, 1H); <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>, mixture of diastereomers) δ 13.7, 13.9, 41.6, 44.2, 44.3, 111.4, 111.5, 112.2, 113.4, 115.6, 116.0, 117.5, 118.2, 118.7, 118.8, 119.7, 119.9, 122.2, 122.4, 122.5, 122.6, 125.6, 126.2, 134.3, 136.0, 136.2, 136.3, 165.2, 165.3; IR (CHCl<sub>3</sub>,

cm<sup>-1</sup>) 3478, 2361, 1743; HRMS (EI) Calculated for C<sub>16</sub>H<sub>16</sub>N<sub>2</sub>O<sub>2</sub>: 268.12118. Found 268.12117.

### 4.3. Typical experimental procedure for oxidative C-C bond formation of 3-benzylindole 5a with higher-order cyanocuprates (Table 3, entry 4)

A solution of Ph<sub>2</sub>Cu(CN)Li<sub>2</sub> was prepared as follows; to a stirred solution of copper(I) cyanide (61.2 mg, 0.68 mmol) in Et<sub>2</sub>O (4 mL) was added dropwise a solution of phenyllithium (1.14 *N* in cyclohexane-ether, 1.06 ml, 1.21 mmol) at -20 °C for 1 h.

To a stirred solution of 5a (50 mg, 0.24 mmol) in THF (3 mL) was added dropwise a solution of *n*-BuLi (1.61 *N* in hexane, 0.20 mL, 0.32 mmol) at -78 °C. After the mixture was stirred at the same temperature for 15 min, a solution of 4 (83 mg, 0.38 mmol) in THF (1 mL) was added at -78 °C. After the mixture was stirred at the same temperature for 30 min, a solution of Ph<sub>2</sub>Cu(CN)Li<sub>2</sub> was added at -78 °C with a cannula, and the reaction mixture was stirred at the same temperature for 30 min. the product 9d was obtained as usual manner.

#### 4.3.1 3-(1-Phenylethyl)indole (9a)<sup>5a</sup>

White solid; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 1.70 (t, *J* = 7.3 Hz, 3H), 4.37 (q, *J* = 7.3 Hz, 1H), 6.98-7.37 (m, 10H), 7.96 (s, 1H); <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>) δ 22.4, 36.9, 111.0, 119.2, 119.7, 121.0, 121.9, 122.3, 125.9, 127.4, 128.3, 128.7, 136.6, 146.8.

#### 4.3.2. 3-(1-Phenylpentyl)indole (9b)

Colorless plates: mp 74.5-75.5 °C; <sup>1</sup>H NMR (270 MHz, CDCl<sub>3</sub>) δ 0.86 (t, *J* = 7.0 Hz, 3H), 1.27-1.35 (m, 4H), 1.96-2.03 (m, 1H), 2.14-2.21 (m, 1H), 4.14 (t, *J* = 7.6 Hz, 1H), 6.97-7.02 (m, 2H), 7.11 (d, *J* = 8.1 Hz, 1H), 7.15 (d, *J* = 7.8 Hz, 1H), 7.23-7.30 (m, 5H), 7.44 (d, *J* = 8.1 Hz, 1H), 7.83 (s, 1H); <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>) δ 14.0, 22.8, 30.3, 35.9, 42.9, 111.0, 119.2, 119.5, 120.7, 120.9, 121.9, 125.8, 127.1, 127.9, 128.2, 136.5, 145.6; IR (CHCl<sub>3</sub>, cm<sup>-1</sup>) 3481, 1456; HRMS (EI) Calculated for C<sub>21</sub>H<sub>19</sub>N: 263.16740. Found 263.16743.

#### 4.3.3. 3-(2,2-Dimethyl-1-phenylpropyl)indole (9c)

White powder: mp 87.5-88.5 °C; <sup>1</sup>H NMR (270 MHz, CDCl<sub>3</sub>) δ 1.04 (s, 9H), 4.15 (s, 1H), 7.02-7.40 (m, 9H), 7.58 (d, *J* = 7.7 Hz, 1H), 7.98 (s, 1H); <sup>13</sup>C NMR (67.8 MHz, CDCl<sub>3</sub>) δ 29.0, 35.2, 53.6, 110.7, 117.8, 119.0, 119.1, 121.0, 121.8, 125.7, 127.5, 128.8, 130.0, 135.1, 143.3; IR (CHCl<sub>3</sub>, cm<sup>-1</sup>) 3481, 1456; HRMS (EI) Calculated for C<sub>21</sub>H<sub>19</sub>N: 263.16740. Found 263.16682.

#### 4.3.4. 3-Diphenylmethylindole (9d)<sup>21</sup>

White needles: mp 124.5-124.8 °C (lit.<sup>21</sup> 121 °C); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 5.66 (s, 1H), 6.54 (d, *J* = 2.2 Hz, 1H), 6.96-6.99 (m, 1H), 7.14-7.28 (m, 12H), 7.33 (d, *J* = 8.1 Hz, 1H), 7.88 (s, 1H); <sup>13</sup>C NMR (125.7 MHz, CDCl<sub>3</sub>) δ 48.8, 111.0, 119.4, 119.9, 122.1, 124.0, 126.2, 127.0, 128.3, 129.0, 136.7, 143.9.



### Acknowledgments

The authors are grateful for the financial support from Takeda Science Foundation, the Research Foundation for Pharmaceutical Sciences and the Sasakawa Scientific Research Grant from The Japan Science Society, and a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

### References

- (a) Sundberg, R. *Indoles*, Academic Press: San Diego, 1996; (b) Robinson, B. *The Fischer Indole Synthesis*, Wiley-Interscience: New York, 1982; (c) Sundberg, R. *The Chemistry of Indoles*, Academic Press: New York, 1970.
- For reviews on the synthesis of indoles: (a) Humphrey, R. R.; Kuethe, J. T. *Chem. Rev.* **2006**, *106*, 2875-2911; (b) Gilchrist, T. L. *J. Chem. Soc., Perkin Trans. 1* **2001**, 2491-2515; (c) Gribble, G. W. *J. Chem. Soc., Perkin Trans. 1* **2000**, 1045-1075; (d) Pindur, U.; Adam, R. *J. Heterocyclic Chem.* **1988**, *25*, 1-8.
- (a) Snyder, H.; Smith, C.; Stewart, J. *J. Am. Chem. Soc.* **1944**, *66*, 200-204; (b) Albright, J. D.; Snyder, H. R. *J. Am. Chem. Soc.* **1959**, *81*, 2239-2245; (c) Howe, E. E.; Zambito, A. J.; Snyder, H. R.; Tishler, M. *J. Am. Chem. Soc.* **1945**, *67*, 38-39; (d) Albertson, N. F.; Archer, S.; Suter, C. M. *J. Am. Chem. Soc.* **1945**, *67*, 36-37; (e) Somei, M.; Karasawa, Y.; Kaneko, C. *Heterocycles* **1981**, *16*, 941-949.
- Review: Semenov, B. B.; Granik, V. G. *Pharm. Chem. J.*, **2004**, *38*, 287-310.
- (a) Ballini, R.; Palmieri, A.; Petrini, M.; Torregiani, E. *Org. Lett.* **2006**, *8*, 4093-4096; (b) Palmieri, A.; Petrini, M. *J. Org. Chem.* **2007**, *72*, 1863-1866.
- (a) Runti, C. *Gazz. Chim. Ital.* **1951**, *81*, 613-620; (b) Runti, C.; Orlando, G. *Ann. Chim. (Rome)* **1953**, *43*, 308-314.
- Poppelsdorf, F.; Holt, S. *J. Chem. Soc.* **1954**, 4094-4101.
- (a) Semenov, B. B.; Novikov, K. A.; Lysenko, K. A.; Kachala, V. V. *Tetrahedron Lett.* **2006**, *47*, 3479-3483; (b) Semenov, B. B.; Smushkevich, V. V. Y. I.; Levina, I. I.; Kurkovskaya, L. N.; Lysenko, K. A.; Kachala, V. V. *Chem. Heterocycl. Compounds* **2005**, *41*, 730-738; (c) Semenov, B. B.; Smushkevich, Y. I.; Grintselev-Knyazev, G. V.; Antipin, M. Y. *Russ. Chem. Bull. Int. Ed.* **2001**, *50*, 570-571; (d) Semenov, B. B.; Smushkevich, Y. I. *Russ. Chem. Bull. Int. Ed.* **2001**, *50*, 543-545.
- Recent examples: (a) Gibbs, T. J. K.; Tomkinson, N. C. O. *Org. Biomol. Chem.* **2005**, *3*, 4043-4045; (b) Shirakawa, S.; Kobayashi, S. *Org. Lett.* **2006**, *8*, 4939-4942; (c) Ke, B.; Qin, Y.; He, Q.; Huang, Z.; Wang, F. *Tetrahedron Lett.* **2005**, *46*, 1751-1753.
- (a) Schellenberg, K. A. *J. Biol. Chem.* **1965**, *240*, 1165-1169; (b) Schellenberg, K. A. *J. Biol. Chem.* **1967**, *242*, 1815-1820; (c) Schellenberg, K. A.; McLean, G. W. *J. Am. Chem. Soc.* **1966**, *88*, 1077-1079; (d) Schellenberg, K. A.; McLean, G. W.; Lipton, H. L.; Lietman, P. S. *J. Am. Chem. Soc.* **1967**, *89*, 1948-1950; (e) Huffman, R. W.; Bruce, T. C. *J. Am. Chem. Soc.* **1967**, *89*, 6243-6251.
- Burr, G. O.; Gortner, R. A. *J. Am. Chem. Soc.* **1924**, *46*, 1224-1246.
- Recent reports: (a) Catino, A. J.; Nicholas, J. M.; Nettles, B. J.; Doyle, M. P. *J. Am. Chem. Soc.* **2006**, *128*, 5648-5649; (b) Murahashi, S.-I.; Komiya, N.; Terai, H. *Angew. Chem., Int. Ed.* **2005**, *44*, 6931-6933; (c) Li, Z.; Li, C.-J. *J. Am. Chem. Soc.* **2005**, *127*, 3672-3673 and references are cited therein.
- (a) Matsuo, J. *J. Synth. Org. Chem. Jpn.* **2004**, *62*, 574-583; (b) Mukaiyama, T. *Angew. Chem., Int. Ed.* **2004**, *43*, 5590-5614.
- (a) Matsuo, J.; Tanaki, Y.; Ishibashi, H. *Org. Lett.* **2006**, *8*, 4371-4374; (b) Matsuo, J.; Tanaki, Y.; Ishibashi, H. *Tetrahedron Lett.* **2007**, *48*, 3233-3236; (c) Matsuo, J.; Kawai, H.; Ishibashi, H. *Tetrahedron Lett.* **2007**, *48*, 3155-3158.
- For application to other one-pot C-C bond formations, see: (a) Matsuo, J.; Aizawa, Y. *Chem. Commun.* **2005**, 2399-2401; (b) Matsuo, J.; Shibata, T.; Kitagawa, H.; Mukaiyama, T. *Arkivoc* **2001**, 58-65; (c) Kerr, W. J.; Pearson, C. M.; Thurston, G. J. *Org. Biomol. Chem.* **2006**, *4*, 47-50; (d) Crawford, J. J.; Henderson, K. W.; Kerr, W. J. *Org. Lett.* **2006**, *8*, 5073-5076.
- Whitney, S.; Grigg, R.; Derrick, A.; Keep, A. *Org. Lett.* **2007**, *9*, 3299-3302.
- Kimura, M.; Futamata, M.; Mukai, R.; Tamaru, Y. *J. Am. Chem. Soc.* **2005**, *127*, 4592-4593.
- (a) Nickel-catalyzed reaction: Pratt, E. T.; Botimer, L. W. *J. Am. Chem. Soc.* **1957**, *79*, 5248-5250; (b) Wenkert, E.; Angell, E. C.; Ferreira, V. F.; Michelotti, E. L.; Piettre, S. R.; Sheu, J.-H.; Swindell, C. S. *J. Org. Chem.* **1986**, *51*, 2343-2351.
- Matsuo, J.; Iida, D.; Tatani, K.; Mukaiyama, T. *Bull. Chem. Soc. Jpn.* **2002**, *75*, 223-234.
- Ernest, F. P.; Laurence, W. B. *J. Am. Chem. Soc.* **1957**, *79*, 5248-5250.
- Yasuda, M.; Somyo, T.; Baba, A. *Angew. Chem., Int. Ed.* **2006**, *45*, 793-796.