

# Total syntheses of ( $\pm$ )-agroclavine-?, ( $\pm$ )-6-nor-chanoclavine-?, and ( $\pm$ )-chanoclavin-?

|                              |   |
|------------------------------|---|
| 著者                           | Somei Masanori, Yamada Fumio, Makita Yoshihiko                                |
| journal or publication title | Heterocycles  |
| volume                       | 26  |
| number                       | 4   |
| page range                   | 895-898   |
| year                         | 1987-01-01  |
| URL                          | <a href="http://hdl.handle.net/2297/4309">http://hdl.handle.net/2297/4309</a> |

TOTAL SYNTHESSES OF (±)-AGROCLAVINE-I, (±)-6-NOR-  
 CHANOCCLAVINE-II, AND (±)-CHANOCCLAVINE-II<sup>1</sup>

Masanori Somei,\* Fumio Yamada, and Yoshihiko Makita  
 Faculty of Pharmaceutical Sciences, Kanazawa University  
 13-1 Takara-machi, Kanazawa 920, Japan

**Abstract** ——— The first total syntheses of (±)-6-nor-  
 chanoclavine-II and (±)-chanoclavine-II are achieved. Total  
 synthesis of (±)-agroclavine-I is also reported.

In the previous papers,<sup>2</sup> we have established a practical and common synthetic method for ergot alkaloids. The community of the synthetic method is now successfully extended to total syntheses of (±)-agroclavine-I (8),<sup>3</sup> (±)-6-nor-chano-clavine-II (11),<sup>4</sup> and (±)-chano-clavine-II (13).<sup>5</sup> In this communication, we describe these results in detail.


First, 4,5-trans-4-nitro-5-(2-methyl-1-propen-1-yl)-1,3,4,5-tetrahydrobenz[cd]indole (2) was prepared in four steps in 42% overall yield starting from 3-indolecarbaldehyde (1) as reported previously.<sup>2</sup> Subsequent treatment of 2 with sodium methoxide in methanol and protonation with 10% aqueous acetic acid readily inverted the stereochemistry of the carbon-4 bound to the nitro group to afford 4,5-cis-4-nitro-5-(2-methyl-1-propen-1-yl)-1,3,4,5-tetrahydrobenz[cd]indole (3)<sup>6a</sup> in 85% yield together with the recovery of 2 in 8% yield.

With the compound (3) in hand, the syntheses of 8, 11, and 13 were carried out by employing our general synthetic operations.<sup>2</sup> Thus, reduction of 3 with zinc amalgam in aqueous hydrochloric acid (HCl) gave the corresponding amine (4)<sup>6b</sup> in 98% yield. Treatment of 4 with methyl chloroformate (ClCOOMe) in methylene chloride (CH<sub>2</sub>Cl<sub>2</sub>) in the presence of triethylamine (NEt<sub>3</sub>) afforded 66% yield of the desired carbamate (5).<sup>6c</sup> Reduction of 5 with lithium aluminum hydride (LiAlH<sub>4</sub>) in absolute tetrahydrofuran (THF) gave (±)-5-epi-6,7-secoagroclavine (6)<sup>6d</sup> in 91% yield. Subsequent regioselective oxidation of Z-methyl<sup>7</sup> in the isobutenyl group at the 5-position was attained by the treatment of 6 with selenium dioxide (SeO<sub>2</sub>) in dioxane and water to

produce *Z*-4,5-*cis*-5-(2-hydroxymethyl-1-propen-1-yl)-4-methylamino-1,3,4,5-tetrahydrobenz[cd]indole [7, (±)-isochanoclavine-II]<sup>6e</sup> in 34% yield. Cyclization of **7** with phosphorus oxychloride in the presence of potassium carbonate<sup>8</sup> produced (±)-agroclavine-I (**8**)<sup>3,6f</sup> in 87% yield. Proton nuclear magnetic resonance (<sup>1</sup>H-NMR) spectrum of **8** was identical with that of the authentic (±)-agroclavine-I, which was synthesized by Ninomiya and co-workers.<sup>3b</sup>

On the other hand, oxidation of **3** with SeO<sub>2</sub> in dioxane and water gave *E*-4,5-*cis*-5-(2-hydroxymethyl-1-propen-1-yl)-4-nitro-1,3,4,5-tetrahydrobenz[cd]indole (**9**)<sup>6g</sup> and *E*-4,5-*cis*-3-(4-nitro-1,3,4,5-tetrahydrobenz[cd]indol-5-yl)-2-methylacrylic aldehyde (**10**)<sup>6h</sup> in 31% and 4% yields,<sup>7</sup> respectively, together with the recovery of **3** in 62% yield. Reduction of **9** with zinc amalgam in aqueous HCl afforded (±)-6-nor-chanoclavine-II (**11**)<sup>4,6i</sup> in 94% yield. Treatment of **11** with ClCOOMe in CH<sub>2</sub>Cl<sub>2</sub> in the presence of NEt<sub>3</sub> afforded 92% yield of the corresponding carbamate (**12**).<sup>6j</sup> Finally, reduction of **13** with LiAlH<sub>4</sub> in absolute THF afforded (±)-chanoclavine-II (**13**)<sup>6k</sup> in 69% yield. <sup>1</sup>H-NMR spectrum of **13** was identical with that of the natural alkaloid, reported in the literature.<sup>5</sup>

Thus, the first and simple total syntheses of **11** and **13** were accomplished.



## ACKNOWLEDGEMENT

The authors express their cordial gratitude to Prof. I. Ninomiya, Kobe Women's College of Pharmacy, for giving us the  $^1\text{H-NMR}$  spectrum of ( $\pm$ )-agroclavine-I.

## REFERENCES AND NOTES

1. This report is part XXXVII of a series entitled "The Chemistry of Indoles". Part XXXVI: M. Somei, E. Iwasa, and F. Yamada, Heterocycles, 24, 3065 (1986).
2. M. Somei, Y. Makita, and F. Yamada, Chem. Pharm. Bull., 34, 948 (1986); M. Somei and F. Yamada, ibid., 32, 504 (1984).
3. As far as we know, two groups succeeded in the total synthesis of ( $\pm$ )-agroclavine-I: a) A.P. Kozikowski and P.D. Stein, J. Am. Chem. Soc., 107, 2569 (1985); b) T. Kiguchi, C. Hashimoto, and I. Ninomiya, Heterocycles, 23, 2891 (1985). Formal total synthesis: W.J. Wheeler, Tetrahedron Lett., 27, 3469 (1986). Isolation: V.G. Sakharovsky and A.G. Kozlovsky, Tetrahedron Lett., 25, 109 (1984).
4. J.M. Cassady, C.I. Abou-Chaar, and H.G. Floss, Lloydia, 36, 390 (1973).
5. D. Stauffacher and H. Tschertter, Helv. Chim. Acta, 47, 2186 (1964).
6. a) mp 147.0-148.0°C. IR (KBr): 3381, 1620, 1606, 1526, 1444, 1378  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) $\delta$ : 1.67 (3H, d,  $J=1.2$  Hz), 1.83 (3H, d,  $J=1.5$  Hz), 3.33 (1H, dd,  $J=15.4$  and 5.3 Hz), 3.59 (1H, ddd,  $J=15.4$ , 10.0, and 1.5 Hz), 4.73 (1H, dd,  $J=10.3$  and 4.4 Hz), 4.96 (1H, ddd,  $J=10.0$ , 5.3, and 4.4 Hz), 5.13 (1H, br d,  $J=10.3$  Hz), 6.75-6.98 (1H, m), 6.92 (1H, br s), 7.00-7.25 (2H, m), 7.98 (1H, br s). b) mp 113.0-114.0°C. IR (KBr): 2920, 1598, 1572, 756  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) $\delta$ : 1.69 (2H, br s,  $\text{NH}_2$ ), 1.81 (3H, d,  $J=1.2$  Hz), 1.85 (3H, d,  $J=1.2$  Hz), 2.81 (1H, ddd,  $J=15.2$ , 7.3, and 1.0 Hz), 3.05 (1H, ddd,  $J=15.2$ , 4.4, and 1.0 Hz), 3.44 (1H, ddd,  $J=7.3$ , 4.4, and 3.5 Hz), 3.98 (1H, dd,  $J=10.0$  and 3.5 Hz), 5.27 (1H, br d,  $J=10.0$  Hz), 6.70-6.91 (1H, m), 6.83 (1H, br s), 6.97-7.19 (2H, m), 8.03 (1H, br s). c) mp 139.0-141.0°C. IR (KBr): 3320, 1683, 1669, 1533  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) $\delta$ : 1.79 (1H, d,  $J=2.0$  Hz), 1.81 (1H, d,  $J=1.5$  Hz), 2.87 (1H, dd,  $J=15.5$  and 7.0 Hz), 3.11 (1H, dd,  $J=15.5$  and 4.2 Hz), 3.61 (3H, s), 4.05 (1H, dd,  $J=9.8$  and 3.7 Hz), 4.30 (1H, dddd,  $J=9.5$ , 7.0, 4.2, and 3.7 Hz), 4.75 (1H, br d,  $J=9.5$  Hz), 5.21 (1H, br d,  $J=9.8$  Hz), 6.68-6.92 (1H, m), 6.85 (1H, br s), 6.96-7.25 (2H, m), 7.93 (1H, br s). d) M. Somei, F. Yamada, Y. Karasawa, and C. Kaneko, Chemistry Lett., 1981, 615. mp 177.0-178.0°C. IR (KBr): 3140, 3090, 1661, 1616, 1437  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) $\delta$ : 1.57 (1H, s, NH), 1.73 (3H, d,  $J=1.2$  Hz), 1.89 (3H, d,  $J=1.2$  Hz), 2.51 (3H, s), 2.77 (1H, ddd,  $J=15.4$ , 10.3, and 1.5 Hz), 2.93-3.22 (2H, m), 4.14 (1H, dd,  $J=10.3$  and 3.7 Hz), 5.21 (1H, br d,  $J=10.3$  Hz), 6.69-6.87 (1H, m), 6.81 (1H, br s), 6.91-7.15 (2H, m), 8.01 (1H, br s, NH). e) oil. IR (film): 3250, 1617, 1604, 1445  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  (pyridine- $d_5$ ) $\delta$ : 1.92 (3H, d,  $J=1.2$  Hz), 2.49 (3H, s), 2.89 (1H, dd,  $J=15.4$  and 11.2 Hz), 3.06-3.35 (2H, m), 4.46 (1H, d,  $J=12.1$  Hz), 4.54 (1H, dd,  $J=10.3$  and 3.7 Hz), 4.78 (1H, d,  $J=12.1$  Hz), 5.54 (1H, d,  $J=10.3$  Hz), 6.98 (1H, d,  $J=6.7$  Hz), 7.05-7.27 (2H, m), 7.34 (1H, d,  $J=7.8$  Hz), 11.50 (1H, br s, NH). f) mp 157.0-158.0°C. IR (KBr): 3400, 3100, 2860, 1618, 1607, 1444  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) $\delta$ : 1.63 (3H, br s), 2.56 (3H, s), 2.78 (1H, ddd,  $J=15.0$ , 10.0, and 1.5 Hz), 2.98 (1H, dd,  $J=15.0$  and 4.5 Hz),

3.07 (2H, br s), 3.24-3.49 (1H, m), 3.78-4.08 (1H, m), 5.48 (1H, br s), 6.74-6.98 (1H, m), 6.79 (1H, br s), 6.98-7.18 (2H, m). g) mp 132.0-134.0°C. IR (KBr); 3496, 3246, 1607, 1535, 1370  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) $\delta$ : 1.86 (3H, d,  $J=1.5$  Hz), 3.37 (1H, dd,  $J=16.0$  and 5.5 Hz), 3.62 (1H, ddd,  $J=16.0$ , 10.3, and 1.7 Hz), 3.94 (2H, br s), 4.81 (1H, dd,  $J=10.3$  and 4.5 Hz), 5.02 (1H, ddd,  $J=10.3$ , 5.5, and 4.5 Hz), 5.46 (1H, dq,  $J=10.3$  and 1.5 Hz), 6.84 (1H, dd,  $J=5.8$  and 2.2 Hz), 6.96 (1H, br s), 7.04-7.28 (2H, m), 8.03 (1H, br s). h) oil. IR (film): 1676 (br), 1636, 1544  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) $\delta$ : 1.96 (3H, d,  $J=1.5$  Hz), 3.32-3.84 (2H, m), 4.89-5.21 (2H, m), 6.41 (1H, br d,  $J=10.2$  Hz), 6.83 (1H, dd,  $J=6.2$  and 1.5 Hz), 7.01 (1H, br s), 6.92-7.34 (2H, m), 8.15 (1H, br s, NH), 9.23 (1H, s). i) mp 208.0-210.0°C (dec.). IR (KBr): 3150, 1614, 1597, 1572, 1420  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  (pyridine- $d_5$ ) $\delta$ : 2.06 (3H, d,  $J=1.0$  Hz), 2.98 (1H, dd,  $J=15.3$  and 7.3 Hz), 3.19 (1H, dd,  $J=15.3$  and 3.9 Hz), 3.50-3.77 (1H, m), 4.28 (1H, dd,  $J=9.8$  and 3.9 Hz), 4.34 (2H, s), 6.05 (1H, br d,  $J=9.8$  Hz), 7.01 (1H, d,  $J=6.8$  Hz), 7.09-7.46 (3H, m), 11.55 (1H, br s). j) oil. IR (film): 3350, 1695  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) $\delta$ : 1.86 (3H, d,  $J=1.2$  Hz), 2.91 (1H, dd,  $J=15.5$  and 6.3 Hz), 3.14 (1H, dd,  $J=15.5$  and 4.5 Hz), 3.60 (3H, s), 4.07 (2H, s), 4.14 (1H, dd,  $J=9.8$  and 3.7 Hz), 4.20-4.57 (1H, m), 4.89 (1H, br d,  $J=9.5$  Hz), 5.56 (1H, br d,  $J=9.8$  Hz), 6.64-6.96 (2H, m), 7.00-7.31 (2H, m), 8.04 (1H, br s). k) mp 86.0-88.0°C (recrystallized from  $\text{CHCl}_3$ ). IR (KBr): 3190, 1620, 1605, 1440  $\text{cm}^{-1}$ .  $^1\text{H-NMR}$  (pyridine- $d_5$ ) $\delta$ : 2.12 (3H, d,  $J=1.0$  Hz), 2.47 (3H, s), 2.95 (1H, dd,  $J=15.5$  and 10.0 Hz), 3.10-3.38 (2H, m), 4.28 (2H, s), 4.43 (1H, dd,  $J=10.2$  and 3.8 Hz), 6.03 (1H, d,  $J=10.2$  Hz), 6.98-7.44 (4H, m), 11.54 (1H, br s).

7. The differences in the direction of oxidation with  $\text{SeO}_2$  would be explained by the formation of a complex with the methylamino group in the compound (6), where the  $\text{SeO}_2$  lie in the vicinity of Z-methyl group. See also reference 2.
8. Combination of these reagents was first applied for the synthesis of benzo[c]-phenanthridine alkaloids by H. Ishii and co-workers: H. Ishii, I.-S. Chen, S. Ueki, T. Masuda, K. Morita, and T. Ishikawa, Symposium Papers, The 28th Symposium on the Chemistry of Natural Products, Sendai, October 1986, p. 457.

Received, 10th December, 1986