

Requirement of Retinoic Acid Receptor for Genipin Derivative-Induced Optic Nerve Regeneration in Adult Rat Retina

著者	Koriyama Yoshiki, Takagi Yusuke, Chiba Kenzo, Yamazaki Matsumi, Sugitani Kayo, Arai Kunizo, Suzuki Hirokazu, Kato Satoru
journal or publication title	PLoS ONE
volume	8
number	8
page range	e71252
year	2013-08-06
URL	http://hdl.handle.net/2297/35629

doi: 10.1371/journal.pone.0071252

Requirement of Retinoic Acid Receptor β for Genipin Derivative-Induced Optic Nerve Regeneration in Adult Rat Retina

Yoshiki Koriyama^{1*}, Yusuke Takagi², Kenzo Chiba³, Matsumi Yamazaki³, Kayo Sugitani⁴, Kunizo Arai², Hirokazu Suzuki⁵, Satoru Kato¹

1 Department of Molecular Neurobiology, Graduate School of Medicine, Kanazawa University, Kanazawa, Japan, **2** Department of Clinical Drug Informatics, Faculty of Pharmacy, Kanazawa University, Kanazawa, Japan, **3** Department of Biochemistry Faculty of Pharmaceutical Sciences, Hokuriku University, Kanazawa, Japan, **4** Division of Health Sciences, Graduate School of Medicine, Kanazawa University, Kanazawa, Japan, **5** Department of Medicinal Chemistry, Faculty of Pharmaceutical Sciences, Hokuriku University, Kanazawa, Japan

Abstract

Like other CNS neurons, mature retinal ganglion cells (RGCs) are unable to regenerate their axons after nerve injury due to a diminished intrinsic regenerative capacity. One of the reasons why they lose the capacity for axon regeneration seems to be associated with a dramatic shift in RGCs' program of gene expression by epigenetic modulation. We recently reported that (1R)-isoPropyloxygenipin (IPRG001), a genipin derivative, has both neuroprotective and neurite outgrowth activities in murine RGC-5 retinal precursor cells. These effects were both mediated by nitric oxide (NO)/S-nitrosylation signaling. Neurotogenic activity was mediated by S-nitrosylation of histone deacetylase-2 (HDAC2), which subsequently induced retinoic acid receptor β (RAR β) expression via chromatin remodeling *in vitro*. RAR β plays important roles of neural growth and differentiation in development. However, the role of RAR β expression during adult rat optic nerve regeneration is not clear. In the present study, we extended this hypothesis to examine optic nerve regeneration by IPRG001 in adult rat RGCs *in vivo*. We found a correlation between RAR β expression and neurite outgrowth with age in the developing rat retina. Moreover, we found that IPRG001 significantly induced RAR β expression in adult rat RGCs through the S-nitrosylation of HDAC2 processing mechanism. Concomitant with RAR β expression, adult rat RGCs displayed a regenerative capacity for optic axons *in vivo* by IPRG001 treatment. These neurotogenic effects of IPRG001 were specifically suppressed by siRNA for RAR β . Thus, the dual neuroprotective and neurotogenic actions of genipin via S-nitrosylation might offer a powerful therapeutic tool for the treatment of RGC degenerative disorders.

Citation: Koriyama Y, Takagi Y, Chiba K, Yamazaki M, Sugitani K, et al. (2013) Requirement of Retinoic Acid Receptor β for Genipin Derivative-Induced Optic Nerve Regeneration in Adult Rat Retina. PLoS ONE 8(8): e71252. doi:10.1371/journal.pone.0071252

Editor: Steven Barnes, Dalhousie University, Canada

Received: February 28, 2013; **Accepted:** June 27, 2013; **Published:** August 6, 2013

Copyright: © 2013 Koriyama et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was partly supported by research grants from the Ministry of Education, Culture, Sports, Science and Technology of Japan (Nos 25462753 to YK, 23650163 to SK and 23618006 to KS). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: koriyama@med.kanazawa-u.ac.jp

Introduction

In contrast to the PNS, neurons of the adult CNS do not normally display axon regeneration after nerve injury. The neonatal nervous system of mammals retains some capacity for CNS regeneration but lose this ability after birth [1]. In newborn stages, RGCs can regrow injured axon within 1–2 weeks after birth [2], [3]. This switch of capacity for axon regeneration is associated with a dramatic shift in RGCs' program of gene expression [4]. The normal development induces restrictive expression of growth associated genes [5] and regeneration associated genes [6]. One of cell-intrinsic change that may contribute to RGCs' loss of regenerative potential is a decrease of histone acetylation. Histone acetylation declines in rat RGCs within 2–3 weeks after birth and decreases even further after nerve injury [7]. Thus, we hypothesize that re-expression of regeneration associated genes which show highly expression in neonatal but not in adulthood in RGCs are able to regenerate optic nerve after injury.

Genipin, an herbal iridoid, has been shown to have both neuroprotective and neurotogenic activity in PC12h cells and Neuro2a cells [8–12]. Recently, we extended this activity to RGC-5, retinal precursor cell line using (1R)-isoPropyloxygenipin (IPRG001), a long-acting genipin derivative. IPRG001 showed significant neuroprotective activity in RGC-5 cells against oxidative stress, such as hydrogen peroxide exposure [13]. Furthermore, IPRG001 sufficiently promoted staurosporine-induced neurite outgrowth from RGC-5 cells in a dose-dependent manner [14]. However, whether IPRG001 can be used to enhance axon regeneration after injury *in vivo* has not been shown. The molecular mechanism of genipin-induced neuroprotective and neurite outgrowth activity was initially believed to be neural nitric oxide synthase (nNOS)/nitric oxide (NO) -dependent because of its structural similarity to tetrahydrobiopterin, which is a cofactor for NOS enzymatic activity [15]. Indeed, both the neuroprotective and neurotogenic effects of IPRG001 in RGC-5 cells were all nNOS/NO-dependent [13,14]. As protein S-nitrosylation is a

sequential event following NO generation by nNOS activation, we focused on this modification in RGC-5 cells because of the ineffectiveness of NO/cGMP signaling [14]. However, the target proteins of S-nitrosylation for neuroprotection and neurite outgrowth were different for each activity. The neuroprotective target is Kelch-like ECH-associated protein-1 (Keap1)/NF-E2 related factor2 (Nrf2), leading to the activation of antioxidative protein expression, whereas the neurotogenic target on the RGC-5 cells is histone deacetylase 2 (HDAC-2), leading to the induction of histone acetylation and retinoic acid receptor β (RAR β) expression [13], [14]. Retinoic acid signaling plays essential roles in neural development, growth, and cellular differentiation [16] via members of the nuclear receptor family including RARs. Especially, the expression levels of RAR β are restricted in adult retina [17]. However, the role of RAR β upon adult rat optic nerve regeneration is unknown. Therefore, in the present study, we examined the neurotogenic properties of genipin on optic nerve regeneration in adult rat retinal ganglion cells (RGCs) *in vivo* after nerve injury. IPRG001 induced RAR β expression in adult rat RGCs through the NO/S-nitrosylation pathway. Concomitant with RAR β expression, IPRG001 successfully regenerated optic axons from matured rat RGCs *in vivo*. These findings may be useful for development of novel therapeutic strategies for CNS regeneration.

Materials and Methods

Ethics Statement

All animal care and handling procedures were approved by the Animal Care and Use Committee of Kanazawa University (No.111848). All surgery was performed under sodium pentobarbital anesthesia, and all efforts were made to minimize suffering.

Chemicals

Genipin was purchased from Wako Pure Chemical Industries, Ltd. (Osaka, Japan). (1R)-*iso*Propyloxygenipin (IPRG001) was synthesized from genipin, as previously described [18], and dissolved in dimethylsulfoxide. 2-(4-carboxyphenyl)-4, 4, 5, 5-tetramethylimidazoline-1-oxyl-3-oxide sodium salt (c-PTIO) and S-ethyl-N-[4-(trifluoromethyl) phenyl] isothiurea (ETPI), were obtained from Alexis Co. (San Diego, CA, USA).

Animals and Surgery

Sprague–Dawley male rats (postnatal 1 day to adult (body weight, 250–300 g)) were used throughout this study. Rats were by intraperitoneal injection of sodium pentobarbital (30–40 mg/kg body weight). Intravitreal injections of various reagents were performed with a Hamilton microsyringe (Hamilton Syringe, Whitteier, CA, USA) 30G needle. Injections were performed by the sclera and retina with a needle 1–2 mm superior to the optic nerve head to avoid lens injury which itself leads to considerable regeneration as a consequence of one or more trophic factors associated with inflammatory cells [19]. As the control of each experiment, we injected vehicles to eye ball. The volume of injection was set at 5 μ l of total volume after pre-suction of the same volume of vitreal fluid. After lateral canthotomy of rats, the conjunctiva was incised laterally to the cornea, the retractor bulbi muscle was separated, and the optic nerve exposed under the binocular-operating microscope. Optic nerve was crushed 2 mm behind the eye with angled jeweler's forceps (Dumont # 5) for 10 sec, avoiding injury to the ophthalmic artery as previously described [20]. Rats were housed in clear plastic cages and maintained under the 12 h light/dark cycles at 23°C.

Retinal Explant Culture

Rat retinal explant cultures were performed as previously described [14]. Retinas were cut into small pieces (0.5 mm squares) and cultured in medium containing DMEM (Sigma-Aldrich, St Louis, MO, USA), 10% fetal bovine serum and penicillin–streptomycin (100 μ g/ml–100 U/ml, respectively) in collagen gel (Cellmatrix; Nitta Gelatin, Osaka, Japan) on a 35 mm culture dish. We observed neurite outgrowth in each explant from total 30–40 explants per dish using phase contrast microscopy. Positive neurite outgrowth was defined on the basis of the length (>200 μ m) of the neurites, following the description in our previous study [14]. We showed the percentage of explants with positive neurites. Four independent experiments were repeated.

Immunohistochemistry

Tissue fixation and cryosectioning were performed as previously described [21]. Briefly, the eyes and optic nerves were isolated and fixed in 4% paraformaldehyde containing 0.1 M phosphate buffer (pH 7.4) and 5% sucrose for 2 h at 4°C. Sucrose concentration was gradually increased from 5 to 20%. The eyes were then embedded in optimal cutting temperature compound, Tissue Tek (Sakura Finetek, Tokyo Japan) and cryosectioned at 12 μ m thickness. The frozen sections were mounted and blocked with Blocking One (Nacalai Tesque, Japan) and retinal sections were incubated with primary anti-nNOS (1:500), anti-RAR β (1:500) (Santa Cruz Biotechnology, Santa Cruz, CA, USA), anti-NeuN (1:500, Millipore, Billerica, MA, USA) and anti-acetyl histone H3 (1:500, Cell Signaling Technology, Tokyo, Japan) antibodies. The sections were then incubated with Alexafluoro anti-IgG (Molecular probe, Eugene, OR, USA).

NADPH Diaphorase Staining

Tissue fixation and cryosectioning were performed as previously described [22]. The frozen sections were mounted onto silane-coated glass slides and air-dried. The slides were then brought to 23°C and incubated overnight in 0.1 M Tris-HCl (pH 8.0) containing 0.3% Triton-X 100. Each sample was stained in buffer containing NADPH and 4-nitroblue tetrazolium chloride (Roche Diagnostics Corporation, Indianapolis, IN, USA) for 2–3 h at 37°C.

Subcellular Fractions for the Extraction of RAR β Protein

Retinas were lysed in hypotonic buffer containing 10 mM HEPES-KOH (pH7.9), 10 mM KCl, 1.5 mM MgCl₂, 1 mM DTT, 0.5 mM phenylmethylsulfonyl fluoride, protease inhibitor cocktail and centrifuged at 10,000 g for 15 min at 4°C. The supernatants were used as the cytoplasmic fraction and the pellets were incubated with a nuclear lysis buffer containing 20 mM HEPES-KOH (pH7.9), 400 mM NaCl, 1.5 mM MgCl₂, 0.2 mM EDTA, 1 mM DTT, 5% glycerol, and protease inhibitor cocktail (Sigma-Aldrich) for 30 min on ice. The lysates were centrifuged at 18,000 g for 15 min at 4°C. The supernatants contained the nuclear fraction. Immunoblotting analysis of β -actin and histone H4 was performed to ensure no contamination of cytoplasmic and nuclear fractions.

Western Blot Analysis

Retinal explants or retinas from various conditions were extracted and 30 μ g of protein were subjected to polyacrylamide gel electrophoresis using a 5–20% gradient gel as previously described [22]. The separated proteins were transferred to a nitrocellulose membrane and incubated with primary and

secondary antibodies (Santa Cruz Biotechnology). Protein bands (nNOS (1:500), RAR β (1:500)) were detected using a BCIP/NBT Kit (Funakoshi, Tokyo, Japan). Antibodies against β -actin (1:500, Gene Tex, San Antonio, TX) and histone H4 (1:500, Cell Signaling Technology, Tokyo, Japan) were used as an internal standard. Protein bands isolated from cells cultured under various conditions were analyzed densitometrically using Scion Image Software (Scion Corp. Frederick, MD, USA). All experiments were repeated at least three times.

S-Nitrosylation Analyses of HDAC2

S-Nitrosylation of HDAC2 was assessed by a modified of biotin switch assay [23] using the S-nitrosylated Protein Detection Assay Kit (Cayman Chemical, Ann Arbor, MI). Retinas exposed to agents for 1 day were harvested and lysed at 4°C. Free thiols were blocked by adding S-methyl methanethiosulfonate and biotinylation of nitrosothiols was performed using maleimide-biotin. Biotinylated proteins were further purified by overnight incubation with neutravidin-coupled agarose beads (Pierce-Thermo Scientific, Rockford, IL, USA). After incubation, beads were washed three times with PBS. Isolated proteins were recovered from beads by the addition of Laemmli sample buffer, and heated at 85°C for 10 min. The amount of S-nitrosylated HDAC2 protein in the samples was analyzed by western blot analysis using anti-HDAC2 antibody (1:500, Cell Signaling Technology, Tokyo, Japan).

Application of siRNA for RAR β

Small interfering RNA (siRNA) for the target region of RAR β mRNA were as follows: 5'-GGAGCCGACUGCAAUACAAG-3' (sense); 5'-UGUAUUUGCAGUCGGUCUCAA-3' (antisense) (Sigma-Aldrich, Japan); and a randomly shuffled sequence: 5'-AGUCGUCGUAUACGGUAUAUC -3' (sense); 5'-UAUACCGUAUACGACGACUUC -3' (antisense). In retinal explant culture, transfection of siRNA (100 pmol) into retinal explants was carried out using Lipofectamine 2000 (Invitrogen Corporation, Carlsbad, CA, USA) for 4 h before culture. In rat retina, knockdown of RAR β expression was performed by injecting siRNA (2 μ g) using a 30-gauge syringe to eyes as described [24], [25]. After injection, five square 80 V pulses of 50 ms duration with 950 ms intervals were applied by NEPA21 (Nepagene, Chiba, Japan) with forcep-type electrodes. After 5 days of electroporation, retinal cells were cultured, and optic nerves were crushed.

Quantitation of Optic Nerve Regeneration *in vivo*

Rats were sacrificed at 14 days after optic nerve injury and were perfused with 4% paraformaldehyde. Optic nerves were impregnated with 5% and then 30% sucrose, embedded in Optimal Cutting Temperature compound (Sakura Finetechnical, Tokyo, Japan) and cut into longitudinal sections of 14 μ m thickness. Regenerating optic axons were visualized by staining with mouse anti-GAP43 antibody (1:250, Santa Cruz Biotechnology, Santa Cruz, CA) followed by a fluorescently-labeled secondary antibody and captured by fluorescent microscopy (BZ-9000, Keyence, Osaka, Japan). Axons were counted manually in at least 8 sections per conditions (6 rats of each treatment) at prespecified distances (250 μ m and 500 μ m) away from the injury site. The numbers of regenerating axons were counted as described by Leon et al. [19].

Counting of Surviving RGCs

RGC survival was evaluated in flat-mounted retinas by immunohistochemistry using a mouse antibody against β III-tubulin (TUJ1, R&D Systems, Inc., Minneapolis, USA 1:500),

followed by a fluorescent secondary antibody after 10 days of injury. Images of 8 prespecified retinal areas of 3 mm away from the optic disc (middle region of the retina) were captured by fluorescent microscopy (under x200 magnification; E600, Nikon) and positive cells were counted using ImageJ software (Wayne Rasband, NIH, Bethesda, MD). Cell densities were calculated as mean values of surviving RGCs per 0.14 mm² in the 8 specified areas. Data are presented as means \pm S.E.M on five rats per group.

Statistics

All results were reported as mean \pm SEM for 3–5 experiments. Differences between groups were analyzed using one-way ANOVA, followed by Dunnett's multi-comparison test with PASW Software (SPSS Inc., Chicago, IL, USA). P values <0.05 were considered statistically significant.

Result

Correlation between Axonal Regeneration and RAR β Expression in the Developing Rat Retina

Newborn rat retina at postnatal day 1 (P1) showed spontaneous axonal regeneration in explant cultures forming long neurites more than 200 μ m in length (Fig. 1A). In contrast, rat retina at postnatal day 14 (P14, Fig. 1B), and day 60 (P60, Fig. 1C) lost the capacity for axonal regeneration in explant culture. Figure 1D shows a graphical representation of neurite outgrowth in P1, P14 and P60 rat retinal explant cultures. In P14 rat retina, neurite outgrowth was about 30% of P1 retina. In P60 rat retina, neurite outgrowth was less than 10% of P1 retina. To ascertain the levels of retinoic acid receptor β (RAR β) in rat retina after birth, we performed western blot analysis in the total retina at P1, P5, P14 and P60. Figure 1E shows a graphical representation of RAR β protein expression in developing rat retinas. In P5 rat retina, the levels of RAR β expression were 68% of P1 rat. In P14 rat retina, the level of RAR β expression was about 30% of P1 rat. In P60 rat retina, the expression was less than 30% of P1 rat. Histone H4 protein did not change during this period (Fig. 1E). Similarly, we performed immunohistochemistry for RAR β protein expression in the developing rat retina. At P1 rat retina, strong immunoreactivity for RAR β could be seen in the retinal ganglion cell layer (Fig. 1F). The localization of RAR β in RGCs was further confirmed using anti-NeuN, a potent marker of RGCs [26], [27] (Fig. 1G and Fig. 1H merged). In P14 rat retina, RAR β expression in the RGCs was decreased compared to P1 rat retina (Figs. 1 I, J, K). In adult P60 rat retina, RAR β expression in the RGCs was barely visible (Figs. 1 L, M, N). From these results similarity between neurite outgrowth and RAR β expression in development retina, we next examined the effect of RAR β knockdown with retinal explant culture on neurite outgrowth. A specific siRNA for RAR β suppressed the neurite outgrowth to 63% (RAR β expression to 52%) compared with no treated control of P1 retina (Figs. 1P and Q), whereas scrambled RNA did not suppress both neurite outgrowth and RAR β expression (Fig. 1O and Q).

Increased NOS Activity and nNOS Expression in rat RGCs by IPRG001 Treatment

As we showed that IPRG001 induced neurite outgrowth in RGC-5 cells through nNOS/NO signaling [14], we examined NOS activity and nNOS expression in the rat retina after intraocular injection of IPRG001. To examine NOS activity, we studied histochemical staining of NADPH diaphorase activity, which was associated with NOS activity [22], [28]. Intraocular injection of IPRG001 (100 pmol/eye) significantly increased




Figure 1. Correlation between axon outgrowth and RAR β expression in rat retina after birth. (A) P1 retinal explant, (B) P14 retinal explant, (C) P60 retinal explant, Scale = 100 μ m. (D) Quantification of axon outgrowth from rat retinal explants at various days after birth. Y axis shows % of explants with neurites longer than 200 μ m. * P <0.01 vs P1 retina. (n = 3, 5 rats of each stage). (E) Quantification of RAR β expression in rat retina during development. * P <0.05, ** P <0.01 vs P1 (n = 3). (F–N) RAR β and NeuN expression in the rat retina during development. (F, I, L) RAR β expression at P1 (F), P14 (I) and P60 (L). Scale = 50 μ m. (G, J, M) NeuN-positive RGCs at P1 (G), P14 (J), and P60 (M). (H, K, N) Merged images of each day at P1 (H), P14 (K), and P60 (N). (O) P1 retina treated with scrambled siRNA (P) siRNA for RAR β in P1. Scale = 100 μ m. (Q) siRNA of RAR β suppressed expression of RAR β and neurite outgrowth in P1 retina. * P <0.01 vs vehicle control (Con) and scrambled siRNA(Scr) (RAR β expression), * P <0.01 vs Con and Scr (neurite outgrowth), (n = 3, 5 rats of each treatment). doi:10.1371/journal.pone.0071252.g001

NADPH diaphorase staining in RGCs (Fig. 2B) 1 day after treatment compared to vehicle control retina (Fig. 2A). Next, we studied the effects of IPRG001 on nNOS protein expression in the adult rat retina by western blot. Intraocular IPRG001 significantly increased nNOS protein levels 2.5-fold at 1 day after treatment compared to vehicle control (Fig. 2C). The increase in nNOS expression returned to control levels by 5 days after treatment. Levels of β -actin expression did not change during this period (Fig. 2C). We further compared nNOS expression with or without IPRG001 injection at 1 day following optic nerve injury (Fig. 2D).

Optic nerve injury did not change the levels of nNOS expression at this period compared to control as reported by Lee et al. [29]. IPRG001 increased a similar level of nNOS protein at 1 day after treatment with or without optic nerve injury. To determine nNOS protein expression by IPRG001, we performed double-staining in adult rat retina with anti-nNOS and anti-NeuN antibodies. At 1 day after IPRG001 treatment, a strong immunoreactivity for nNOS could be seen in RGCs (Figs. 2F, H, J) compared to the control retina (Figs. 2E, G, I). The localization of increased




Figure 2. IPRG001 activates NADPHd staining and nNOS protein expression in the rat retina. (A, B) NADPHd staining in the retina increased at 1 day post-treatment with 100 pmol/eye of IPRG001 (B) as compared to vehicle control (A). Scale = 100 μ m. (C) Western blot analysis of nNOS protein after treatment of IPRG001. * $P < 0.01$ vs control ($n = 3$). (D) Levels of nNOS expression by IPRG001 with or without optic nerve injury. * $P < 0.01$ vs control ($n = 3$). (E–J) nNOS and NeuN expression in the retina after treatment of IPRG001. (E and F) Immunoreactivity of nNOS protein was increased in RGCs at 1 day after IPRG001 treatment (F) compared to vehicle control (E). Scale = 50 μ m. (G and H) NeuN staining of E and F. (I and J) Merged images.

doi:10.1371/journal.pone.0071252.g002

expression of nNOS by IPRG001 was also observed in RGCs after nerve injury (data not shown).

Increased RAR β Expression in Adult Rat RGCs with IPRG001 Treatment through nNOS/NO Signaling

In our previous study, we found that IPRG001 induced RAR β expression through nNOS/NO signaling in RGC-5 cells [14]. In this study, we investigated the effects of IPRG001 on RAR β expression in the adult rat retina. The changes of RAR β protein expression in the rat retina were examined for 5 days after treatment (Fig. 3A). RAR β rapidly increased by 3-fold of control levels in the retina at 1 day and gradually declined to lower levels by day 5 post-treatment (Fig. 3A). Histone H4 expression did not change during this period. IPRG001 induced a similar level of RAR β protein at 1 day after treatment with or without optic nerve injury (Fig. 3B). Next, we performed immunohistochemistry for RAR β protein expression in the rat retina after treatment of IPRG001. In vehicle control retina, weak RAR β expression was seen in RGCs with anti-RAR β and anti-NeuN antibodies (Figs. 3C, D, E). Intraocular IPRG001 (100 pmol/eye) increased RAR β expression in RGCs at 1 day post-treatment (Figs. 3F, G, H). The localization of RAR β expression by IPRG001 was seen in RGCs at 1 day after nerve injury (data not shown). The induction

of RAR β expression by IPRG001 in the rat retina was significantly blocked by a NO scavenger, c-PTIO (500 pmol) or a specific nNOS inhibitor, ETPI (200 pmol) (Fig. 3I). Neither c-PTIO nor ETPI had any effects on RAR β expression in vehicle control retina. These data strongly indicate that the IPRG001-induced RAR β expression in the adult rat RGCs was nNOS/NO-dependent.

IPRG001-induced S-nitrosylation of Histone Deacetylase 2 (HDAC2) and Acetylation of Histone H3

In our previous study, we showed that IPRG001-induced nNOS/NO signaling occurred after the S-nitrosylation of HDAC2 leading to increased RAR β expression and neurogenesis in RGC-5 cells [14]. Therefore, we investigated the S-nitrosylation of HDAC2 after IPRG001 treatment. Figure 4A shows a 2.6-fold increase in the S-nitrosylation of HDAC2 at 1 day after IPRG001 (100 pmol/eye) treatment. The S-nitrosylation of HDAC2 was also nNOS/NO-dependent. c-PTIO or ETPI significantly blocked HDAC2 S-nitrosylation (Fig. 4A). Figure 4B illustrates a graphical representation of acetylated histone H3 (AcH3) with IPRG001 (100 pmol/eye) treatment. IPRG001 increased AcH3 2.3-fold in the rat retina at 1 day and returned to control levels by 5 days post treatment. The increase of AcH3 by intraocular IPRG001 was




Figure 3. Upregulation of RAR β expression in RGCs after IPRG001 treatment. (A) Western blot analysis of RAR β protein expression increased at 1 day after treatment of IPRG001. * $P < 0.01$ vs control ($n = 3$). (B) Levels of RAR β expression by IPRG001 with or without optic nerve injury. * $P < 0.01$ vs control ($n = 3$). (C–H) RAR β and NeuN expression in the retina after treatment of IPRG001. (C and F) Immunoreactivity of RAR β protein was increased in RGCs 1 day after treatment with IPRG001 (F) compared to vehicle control (C). Scale = 50 μm . (D and G) NeuN staining of C and F. (E and H) Merged images. (I) nNOS/NO-dependent expression of RAR β by IPRG001. * $P < 0.05$, ** $P < 0.01$ vs control, + $P < 0.01$ vs IPRG001 treatment alone ($n = 3$).

doi:10.1371/journal.pone.0071252.g003

almost blocked by ETPI (Fig. 4C). Immunohistochemical staining revealed that IPRG001 significantly enhanced immunoreactivity of ACh3 in rat RGCs at 1 day after treatment (Figs. 4G, H, I) compared to vehicle control retina (Figs. 4D, E, F). The data indicate that there is a good correlation between the S-

nitrosylation of HDAC2 and ACh3 in the rat retina for 1 day treatment of IPRG001.





Figure 4. S-Nitrosylation of HDAC2 and the acetylation of histone H3 in the rat retina by IPRG001. (A) S-Nitrosylation of HDAC2 1 day after intraocular administration of IPRG001. NO scavenger, c-PTIO or nNOS inhibitor, ETPI was treated 1 h before IPRG001 treatment. Biotinylated proteins were mixed with avidin beads, eluted and analyzed by western blotting with anti-HDAC2 antibody. ** $P < 0.01$ vs control. + $P < 0.01$ vs IPRG001 treatment alone ($n = 3$). (B) Levels of acetylated histone H3 (AcH3) after treatment of IPRG001. * $P < 0.05$, ** $P < 0.01$ vs control ($n = 3$). (C) nNOS dependent histone H3 acetylation by IPRG001. ETPI was treated 1 h before IPRG001. * $P < 0.05$, ** $P < 0.01$ vs control. + $P < 0.01$ vs IPRG001 treatment alone ($n = 3$). (D–I) Immunohistochemistry of AcH3 and NeuN in rat RGCs. Levels of AcH3 were increased 1 day after IPRG001 treatment (G–I) compared to vehicle control (D–F). (D and G) Immunoreactivity of AcH3. (E and H) NeuN staining. (F and I) Merged images. Scale = 20 μm . doi:10.1371/journal.pone.0071252.g004

Intraocular IPRG001 Promotes Optic Nerve Regeneration from Adult Rat RGCs through a RAR β -dependent Mechanism after Nerve Injury

As IPRG001 enhanced neurite outgrowth from retinal explants *in vitro* [14], we further confirmed this activity *in vivo* preparation after nerve injury. Intraocular IPRG001 (100 pmol/eye) induced optic nerve regeneration *in vivo* (Fig. 5C) compared to vehicle control (Fig. 5A), which was revealed by GAP43 staining. siRNA for RAR β significantly suppressed the effects of IPRG001 (Fig. 5E).

c-PTIO also attenuated the IPRG001-induced optic nerve regeneration (Fig. 5G) although c-PTIO alone did not change the optic nerve regeneration of no treatment (data not shown). Figures 5B, D, F, H show enlarged images of the areas enclosed within the white boxes in Figs. 5A, C, E, G, respectively. IPRG001 (Fig. 5D) showed many regenerating fibers compared to control (Fig. 5B), IPRG001 plus siRNA (Fig. 5F), or IPRG001 plus c-PTIO (Fig. 5H). Figure 5I illustrates the quantitative data of optic nerve regeneration *in vivo* at 250 μm and 500 μm away from the

crush site of optic nerve (asterisk). At both sites, siRNA for RAR β significantly cancelled the effects of IPRG001. In contrast, scramble siRNA did not affect the axon elongation effects of IPRG001. siRNA and scramble siRNA alone did not affect the basal level of optic nerve regeneration without IPRG001 treatment (Fig. 5I).

RAR β Induction by IPRG001 did not Promote RGCs Survival after Nerve Injury

IPRG001 sufficiently rescued rat RGCs against optic nerve injury through antioxidative mechanism [13]. Here, we further examined whether or not the protective effect of IPRG001 was mediated through a RAR β expression mechanism. We therefore counted the number of surviving RGCs by staining of anti-TUJ1

antibody after nerve injury. Optic nerve injury drastically reduced the number of surviving RGCs at 10 days after nerve injury (Fig. 6B) as compared to the vehicle control (Fig. 6A). IPRG001 certainly rescued the RGCs cell death after injury (Fig. 6C). However, neither siRNA for RAR β (Fig. 6D) nor scrambled siRNA affects this protection of RGCs cell death by IPRG001 (Fig. 6E). Fig. 6E illustrates a graphical representation of the number of surviving RGCs after optic nerve injury with or without siRNA for RAR β . These results suggest that this RAR β induction by IPRG001 is targeted only to the neurotogenic action, not to survival action.

Discussion

A Correlation between Neurite Outgrowth and RAR β Expression in the Developing Rat Retina

Immature retina has a higher capacity for optic nerve regeneration rather than mature retina *in vitro* [30]. However, the rat retina gradually loses this capacity for regeneration with development, although the reason remains unsolved. Goldberg et al. [3] reported that rat retinal ganglion cells (RGCs) of embryonic day 20 (E20) had the most intensive capacity for extending optic axons and they showed a higher extension of optic axons 7-fold than the RGCs at postnatal day 8 (P8). We confirmed these results with developing rat retinas of postnatal days 1, 14 and 60 (Fig. 1). It is also known that retinoic acid (RA) signaling is involved in neural development and regeneration regulated through nuclear RA receptors (RARs). In the dorsal root ganglia and spinal cord, a particular role of RAR β expression was implicated in nerve regeneration after nerve injury. Grondona et al. [31] reported that RAR β expression in the mouse retina could be seen in RGCs from postnatal days 0 to 21 (P0–P21). Mori et al. [32] also reported RAR β expression in the developing rat retina. RAR β expression in P14 or P60 rat RGCs was less than




Figure 5. IPRG001-induced rat optic nerve regeneration under RAR β -dependent conditions *in vivo*. (A, C, E, G) Longitudinal sections of the adult rat optic nerve showing GAP-43 positive axons extending over the injury site (asterisks) after 2 weeks after optic nerve injury. (B, D, F, H) Enlarged image of the area enclosed within the white box of A, C, E, G. Scale = 50 μ m. (A) Vehicle control, Scale = 250 μ m. (C) IPRG001, (E) IPRG001 plus siRNA for RAR β . (G) IPRG001 plus c-PTIO (500 pmol). (I) Quantification of axonal regrowth at two indicated proximal points from the injury site (250 μ m and 500 μ m). ** $P < 0.01$ vs control, + $P < 0.01$ vs injury plus IPRG001 (n = 8, 6 rats per each group). doi:10.1371/journal.pone.0071252.g005




Figure 6. No effect of RAR β expression on RGCs survival induced by IPRG001. (A-D) Surviving RGCs stained by TUJ1 in whole mount retina. (A) Vehicle control. Scale = 50 μ m. (B) 10 days after injury. (C) Injury plus IPRG001. (D) Injury plus IPRG001 with RAR β -specific siRNA. (E) Quantification of surviving RGCs after nerve injury with or without siRNA for RAR β . * $P < 0.01$ vs control, + $P < 0.01$ vs injury (n = 8). doi:10.1371/journal.pone.0071252.g006

30% of those at P1 (Fig. 1E). The cellular localization of RAR β in RGCs was confirmed by double-staining of NeuN labeling [26]. This time course of RAR β expression in the developing rat retina was very comparable to that of our data (cf. Fig. 1D and Fig. 1E). Furthermore, we indicated that specific siRNA for RAR β significantly suppressed neurite outgrowth in P1 retina. These results suggest that RAR β is one of the important molecules for neuritogenesis in developing retina. The data of developing rat retina further encouraged us to investigate the neuritogenic activity of RAR β in adult rat retina *in vivo*.

RAR β Expression Induced by IPRG001 is Mediated via NO/S-nitrosylation of HDAC2

In the previous study with RGC-5 cells, IPRG001 promoted staurosporine-induced neurite outgrowth in an nNOS/NO-dependent manner [14]. RGC-5 cells were treated with staurosporine to arrest cell growth and induce cell differentiation [33]. RGC-5 cells are known as neuronal precursor cells of retinal origin. Recently, Van Bergen et al. [34] reported the re-characterization of RGC-5 cells. Differentiated by staurosporine, RGC-5 cells express the RGCs markers, Thy1 and the NMDA receptor [35,36]. Thus, we used staurosporine as a reagent for RGC-5 differentiation as an *in vitro* model of RGCs. However, whether IPRG001 can be used to enhance axon regeneration of RGCs after injury *in vivo* has not been shown. The neurite outgrowth activity of IPRG001 in RGC-5 cells is mediated by the S-nitrosylation of HDAC2. It is well known that the inhibition of HDAC2 induces chromatin remodeling that leads to the transcriptional activation of genes including RAR β [37–39]. Nott et al. [40] further reported that the S-nitrosylation of HDAC2 also caused chromatin remodeling in the neural cells like HDAC2 inhibition. In our examination of RGC-5 cells *in vitro*, the induction of RAR β by IPRG001 was nNOS/NO- and HDAC2 S-nitrosylation-dependent [14]. In the present study, IPRG001 significantly S-nitrosylated HDAC2 in rat RGCs at 1 day post treatment. The HDAC2 S-nitrosylation in the rat retina was also nNOS/NO-dependent. A NO scavenger and nNOS inhibitor clearly suppressed RAR β expression. HDAC inhibitor, which can induce histone H3 acetylation (AcH3) and RAR β expression [38], dramatically promotes neurite outgrowth in retinal explant culture [41]. Moreover, it has reported that HDAC inhibitor, trichostatin A strongly induced histone H3 acetylation, neuritogenesis in RGCs [7], [42] and RAR β expression [43], [44]. Our data of IPRG001 well agree with their results. IPRG001 increased AcH3 in the rat RGC 2.6-fold at 1 day after optic nerve injury by western blot analysis (Fig. 4C). This increase of AcH3 was nNOS-dependent. ETPI, a nNOS inhibitor completely blocked the IPRG001-induced increase of AcH3. These data strongly indicate that IPRG001 leads chromatin remodeling like an HDAC2 inhibitor through an NO/S-nitrosylation mechanism. Recently, Gaub et al. [7] reported that the levels of histone acetyltransferase P300 protein and histone acetylation (AcH3) in the rat RGCs decreased with maturation. This property might be one of the reasons why mammalian optic nerves lost the capacity for optic nerve regeneration after nerve injury [41].

References

- Bregman BS, Goldberger ME (1982) Anatomical plasticity and sparing of function after spinal cord damage in neonatal cats. *Science* 217: 553–555.
- Chen DF, Jhaveri S, Schneider GE (1995) Intrinsic changes in developing retinal neurons result in regenerative failure of their axons. *Proc Natl Acad Sci USA* 92: 7287–7291.

Role of RAR β Expression in Rat RGCs for Optic Nerve Regeneration

In general, successful CNS regeneration after nerve injury is mainly dependent on neuritogenesis. As for neuronal cell survival, the overexpression of Bcl-2, an antiapoptotic molecule in mouse RGCs, sufficiently maintains the cell survival of RGCs, but does not promote axonal regeneration from RGCs [45], [46]. There are many reports that RAR β expression is involved in axonal regeneration in the CNS neurons, including the spinal cord, dorsal root ganglia [47–49] and we could show the important role of RAR β expression during optic nerve regeneration in this study. On the other hand, the lack of regenerative capacity is attributable in part to inhibitory factors in myelin [50]. There are a few inhibitory environments on neonatal axonal regeneration within 2 days after birth [51]. Myelin basic protein, a marker protein of myelin, start to express from few days after birth and inhibitory factors from myelin inhibit axon regeneration [52]. Interestingly, retinoic acid/its receptor, RAR β counteracts myelin-dependent inhibition of axon outgrowth by repression of Lingo-1, one of the coreceptor of inhibitory Nogo receptor [53]. Although we do not know the exact mechanism behind AcH3-induced RAR β expression and its downstream signaling pathway, the cooperation with histone acetyltransferase and RAR is involved in promotion of gene expression and cell differentiation in the spinal motor neuron development [54], [55]. Further studies are needed to elucidate the mechanism of axon outgrowth through AcH3/RAR β expression. On the other hand, there are very few reports of RAR β expression resulting in neuroprotection or survival in neurons [56]. To know the effects of RAR β induction on IPRG001-induced RGCs survival after nerve injury, we used specific siRNA for RAR β . In our experimental conditions, siRNA for RAR β did not affect the neuroprotective effects of IPRG001 in rat RGCs after injury (Fig. 6). These results suggested that cell survival effect by IPRG001 in RGCs after injury was irrelevant to RAR β expression by IPRG001. In our previous study [13], IPRG001 supports RGC survival *in vivo* after optic nerve injury through the S-nitrosylation of another target protein, Keap1, leading to the induction of an antioxidative protein, heme oxygenase-1. Thus RAR β induction by IPRG001 seems to be unconnected with the survival effect against RGCs after injury. The present results in connection with our previous study [14] strongly indicate that IPRG001 activates nNOS/NO/S-nitrosylation signaling, which leads to distinct neuroprotective and neuritogenic actions on damaged adult rat RGCs *in vivo*. The concurrent neuroprotective and neuritogenic actions of IPRG001 could provide a new therapeutic tool for CNS disorders, including brain ischemia, age-related brain disorders, spinal cord injury, and RGC degenerative diseases, such as glaucoma.

Acknowledgments

We thank Ms Sachiko Higashi and Tomoko Kano for their administrative and technical assistance.

Author Contributions

Conceived and designed the experiments: YK. Performed the experiments: YK YT KC MY KS KA SK. Analyzed the data: YK YT MY KS SK. Contributed reagents/materials/analysis tools: KC HS. Wrote the paper: YK.

3. Goldberg JL, Klassen MP, Hua Y, Barres BA (2002) Amacrine-signaled loss of intrinsic axon growth ability by retinal ganglion cells. *Science* 296: 1860–1864.
4. Wang JT, Kunzevitzky NJ, Dugas JC, Cameron M, Barres BA, et al. (2007) Disease gene candidates revealed by expression profiling of retinal ganglion cell development. *J Neurosci* 27: 8593–8603.
5. Benowitz LI, Routtenberg A (1997) GAP-43: an intrinsic determinant of neuronal development and plasticity. *Trends Neurosci* 20: 84–91.
6. Moore DL, Blackmore MG, Hu Y, Kaestner KH, Bixby JL, et al. (2009) KLF family members regulate intrinsic axon regeneration ability. *Science* 326: 298–301.
7. Gaub P, Joshi Y, Wuttke A, Naumann U, Schnichels S, et al. (2011) The histone acetyltransferase p300 promotes intrinsic axonal regeneration. *Brain* 134: 2134–2148.
8. Yamazaki M, Chiba K, Mohri T (1996) Neurotogenic effect of natural iridoid compounds on PC12h cells and its possible relation to signaling protein kinases. *Biol Pharm Bull* 19: 791–795.
9. Yamazaki M, Chiba K, Mohri T, Hatanaka H (2001) Activation of the mitogen-activated protein kinase cascade through nitric oxide synthesis as a mechanism of neurotogenic effect of genipin in PC12h cells. *J Neurochem* 79: 45–54.
10. Yamazaki M, Chiba K, Mohri T, Hatanaka H (2004) Cyclic GMP-dependent neurite outgrowth by genipin and nerve growth factor in PC12h cells. *Eur J Pharmacol* 488: 35–43.
11. Yamazaki M, Chiba K, Satoh K (2008) Neuro2a Cell Death Induced by 6-Hydroxydopamine is Attenuated by Genipin. *J Health Sci* 54: 638–644.
12. Yamazaki M, Chiba K (2005) Neurotrophic Effects of Genipin on Neuro2a Cells. *J of Health Sci* 51: 687–692.
13. Koriyama Y, Chiba K, Yamazaki M, Suzuki H, Muramoto K, et al. (2010) Long-acting genipin derivative protects retinal ganglion cells from oxidative stress models in vitro and in vivo through the Nrf2/antioxidant response element signaling pathway. *J Neurochem* 115: 79–91.
14. Koriyama Y, Takagi Y, Chiba K, Yamazaki M, Arai K, et al. (2011) Neurotogenic activity of a genipin derivative in retinal ganglion cells is mediated by retinoic acid receptor β expression through nitric oxide/S-nitrosylation signaling. *J Neurochem* 119: 1232–1242.
15. Suzuki H, Yamazaki M, Chiba K, Sawanishi H (2007) Characteristic properties of genipin as an activator in neuronal nitric oxide synthase. *J Health Sci* 53: 730–733.
16. Chambon P (1996) A decade of molecular biology of retinoic acid receptors. *FASEB J* 10: 940–954.
17. Perez-Juste G, Aranda A (1999) Differentiation of neuroblastoma cells by phorbol esters and insulin-like growth factor 1 is associated with induction of retinoic acid receptor beta gene expression. *Oncogene* 18: 5393–5402.
18. Suzuki H, Yamazaki M, Chiba K, Uemori Y, Sawanishi H (2010) Neurotogenic activities of 1-alkyloxygenipins. *Chem Pharm Bull* 58: 168–171.
19. Leon S, Yin Y, Nguyen J, Irwin N, Benowitz LI (2000) Lens injury stimulates axon regeneration in the mature rat optic nerve. *J Neurosci* 20: 4615–4626.
20. Koriyama Y, Tani H, Ohno M, Kimura T, Kato S (2008) A novel neuroprotective role of a small peptide from flesh fly, 5-S-GAD in the rat retina in vivo. *Brain Res* 1240: 196–203.
21. Koriyama Y, Ohno M, Kimura T, Kato S (2009b) Neuroprotective effects of 5-S-GAD against oxidative stress-induced apoptosis in RGC-5 cell. *Brain Res* 1296: 187–195.
22. Koriyama Y, Yasuda R, Homma K, Mawatari K, Nagashima M (2009a) Nitric oxide-cGMP signaling regulates axonal elongation during optic nerve regeneration in the goldfish in vitro and in vivo. *J Neurochem* 110: 890–901.
23. Jaffrey SR, Snyder SH (2001) The biotin switch method for the detection of S-nitrosylated proteins. *Sci STKE* 12: 1–9.
24. Matsuda T, Cepko CL (2004) Electroporation and RNA interference in the rodent retina in vivo and in vitro. *Proc Natl Acad Sci USA* 10: 16–22.
25. Matsuda T, Cepko CL (2007) Controlled expression of transgenes introduced by in vivo electroporation. *Proc Natl Acad Sci USA* 104: 1027–1032.
26. Wolf HK, Buslei R, Schmidt-Kastner R, Schmidt-Kastner PK, Pietsch T, et al. (1996) NeuN: a useful neuronal marker for diagnostic histopathology. *J. Histochem. and Cytochem.* 44: 1167–1171.
27. Park HY, Kim JH, Park CK (2012) Activation of autophagy induces retinal ganglion cell death in a chronic hypertensive glaucoma model. *Cell Death and Dis* 3: e290.
28. Dawson TM, Bredt DS, Fotuhi M, Hwang PM, Snyder SH (1991) Nitric oxide synthase and neuronal NADPH diaphorase are identical in brain and peripheral tissues. *Proc Natl Acad Sci USA* 88: 7797–7801.
29. Lee EJ, Kim KY, Gu TH, Moon JI, Kim IB, et al. (2003) Neuronal nitric oxide synthase is expressed in the axotomized ganglion cells of the rat retina. *Brain Res* 986: 174–180.
30. Symonds AC, Rodger J, Tan MM, Dunlop SA, Beazley LD, et al. (2001) Reinnervation of the superior colliculus delays down-regulation of ephrin A2 in neonatal rat. *Exp Neurol* 170: 364–370.
31. Grondona JM, Kastner P, Gansmuller A, Décimo D, Chambon P, et al. (1996) Retinal dysplasia and degeneration in RARbeta2/RARgamma2 compound mutant mice. *Development* 122: 2173–2188.
32. Mori M, Ghyselinck NB, Chambon P, Mark M (2001) Systematic immunolocalization of retinoid receptors in developing and adult mouse eyes. *Invest Ophthalmol Vis Sci* 42: 1312–1318.
33. Lieven CJ, Millet LE, Hoegger MJ, Levin LA (2007) Induction of axon and dendrite formation during early RGC-5 cell differentiation. *Exp Eye Res* 85: 678–683.
34. Van Bergen NJ, Wood JP, Chidlow G, Trounce IA, Casson RJ, et al. (2009) Recharacterization of the RGC-5 retinal ganglion cell line. *Invest Ophthalmol Vis Sci* 50: 4267–4272.
35. Frassetto LJ, Schlieve CR, Lieven CJ, Utter AA, Jones MV, et al. (2006) Kinase-dependent differentiation of a retinal ganglion cell precursor. *Invest Ophthalmol Vis Sci* 47: 427–438.
36. Hironaka K, Inokuchi Y, Fujisawa T, Shimazaki H, Akane M, et al. (2011) Edaravone-loaded liposomes for retinal protection against oxidative stress-induced retinal damage. *Eur J Pharm Biopharm* 79: 119–125.
37. Qian DZ, Ren M, Wei Y, Wang X, van de Geijn F, et al. (2005) In vivo imaging of retinoic acid receptor beta2 transcriptional activation by the histone deacetylase inhibitor MS-275 in retinoid-resistant prostate cancer cells. *Prostate* 64: 20–28.
38. De los Santos M, Zambrano A, Sánchez-Pacheco A, Aranda A (2007) Histone deacetylase inhibitors regulate retinoic acid receptor beta expression in neuroblastoma cells by both transcriptional and posttranscriptional mechanisms. *Mol Endocrinol* 21: 2416–2426.
39. Watson PM, Riccio A (2009) Nitric oxide and histone deacetylases: A new relationship between old molecules. *Commun Integr Biol* 2: 11–13.
40. Nott A, Watson PM, Robinson JD, Crepaldi L, Riccio A (2008) S-Nitrosylation of histone deacetylase 2 induces chromatin remodelling in neurons. *Nature* 455: 411–415.
41. Biermann J, Grieshaber P, Goebel U, Martin G, Thanos S, et al. (2010) Valproic acid-mediated neuroprotection and regeneration in injured retinal ganglion cells. *Invest Ophthalmol Vis Sci* 51: 526–534.
42. Schwechter BR, Millet LE, Levin LA (2007) Histone deacetylase inhibition-mediated differentiation of RGC-5 cells and interaction with survival. *Invest Ophthalmol Vis Sci* 48: 2845–2857.
43. Minucci S, Horn V, Bhattacharyya N, Russanova V, Ogryzko VV, et al. (1997) A histone deacetylase inhibitor potentiates retinoid receptor action in embryonal carcinoma cells. *Proc Natl Acad Sci U S A* 94: 11295–11300.
44. Hayashi K, Yokozaki H, Goodison S, Oue N, Suzuki T, et al. (2001) Inactivation of retinoic acid receptor beta by promoter CpG hypermethylation in gastric cancer. *Differentiation* 68: 13–21.
45. Inoue T, Hosokawa M, Morigiwa K, Ohashi Y, Fukuda Y (2002) Bcl-2 overexpression does not enhance in vivo axonal regeneration of retinal ganglion cells after peripheral nerve transplantation in adult mice. *J Neurosci* 22: 4468–4477.
46. Isenmann S, Kretz A, Cellerino A (2003) Molecular determinants of retinal ganglion cell development, survival, and regeneration. *Prog Retin Eye Res* 22: 483–543.
47. Corcoran J, Maden M (1999) Nerve growth factor acts via retinoic acid synthesis to stimulate neurite outgrowth. *Nat Neurosci* 2: 307–308.
48. Corcoran J, Shroot B, Pizzey J, Maden M (2000) The role of retinoic acid receptors in neurite outgrowth from different populations of embryonic mouse dorsal root ganglia. *J Cell Sci* 113: 2567–2574.
49. Corcoran J, So PL, Barber RD, Vincent KJ, Mazarakis ND, et al. (2002) Retinoic acid receptor beta2 and neurite outgrowth in the adult mouse spinal cord in vitro. *J Cell Sci* 115: 3779–3786.
50. Caroni P, Schwab ME (1988) Antibody against myelin-associated inhibitor of neurite growth neutralizes nonpermissive substrate properties of CNS white matter. *Neuron* 1: 85–96.
51. Shewan D, Berry M, Cohen J (1995) Extensive regeneration in vitro by early embryonic neurons on immature and adult CNS tissue. *J Neurosci* 3: 2057–2062.
52. Colella RJ, Pott U, Schwab ME (1994) The role of oligodendrocytes and myelin on axon maturation in the developing rat retinofugal pathway. *J Neurosci* 1: 2594–2605.
53. Puttagunta R, Schmandke A, Floriddia E, Gaub P, Fomin N, et al. (2011) RAR- β counteracts myelin-dependent inhibition of neurite outgrowth via Lingo-1 repression. *J Cell Biol* 193: 1147–1156.
54. Kawasaki H, Eckner R, Yao TP, Taira K, Chiu R, et al. (1998) Distinct roles of the co-activators p300 and CBP in retinoic-acid-induced F9-cell differentiation. *Nature* 393: 284–289.
55. Lee S, Lee B, Lee JW, Lee SK (2009) Retinoid signaling and neurogenin2 function are coupled for the specification of spinal motor neurons through a chromatin modifier CBP. *Neuron* 62: 641–654.
56. Jarvis CI, Goncalves MB, Clarke E, Dogruel M, Kalindjian SB, et al. (2010) Retinoic acid receptor- α signalling antagonizes both intracellular and extracellular amyloid- β production and prevents neuronal cell death caused by amyloid- β . *Eur. J. Neurosci.* 32: 1246–1255.