

Assessment of Olfactory Nerve by SPECT-MRI Image with Nasal Thallium-201 Administration in Patients with Olfactory Impairments in Comparison to Healthy Volunteers

著者	Shiga Hideaki, Taki Junichi, Washiyama Kohshin, Yamamoto Junpei, Kinase Sakae, Okuda Koichi, Kinuya Seigo, Watanabe Naoto, Tonami Hisao, Koshida Kichiro, Amano Ryohei, Furukawa Mitsuru, Miwa Takaki
journal or publication title	PLoS ONE
volume	8
number	2
page range	e57671
year	2013-02-02
URL	http://hdl.handle.net/2297/34159

doi: 10.1371/journal.pone.0057671

Assessment of Olfactory Nerve by SPECT-MRI Image with Nasal Thallium-201 Administration in Patients with Olfactory Impairments in Comparison to Healthy Volunteers

Hideaki Shiga^{1,2*}, Junichi Taki³, Kohshin Washiyama², Junpei Yamamoto¹, Sakae Kinase⁴, Koichi Okuda³, Seigo Kinuya³, Naoto Watanabe⁵, Hisao Tonami⁵, Kichiro Koshida², Ryohei Amano², Mitsuru Furukawa⁶, Takaki Miwa¹

1 Department of Otorhinolaryngology, Kanazawa Medical University, Ishikawa, Japan, **2** Department of Quantum Medical Technology, Graduate School of Medical Science, Kanazawa University, Ishikawa, Japan, **3** Department of Biotracer Medicine, Graduate School of Medical Science, Kanazawa University, Ishikawa, Japan, **4** Risk Analysis and Applications Research Group, Nuclear Safety Research Center, Japan Atomic Energy Agency, Ibaraki, Japan, **5** Department of Diagnostic and Therapeutic Radiology, Kanazawa Medical University, Ishikawa, Japan, **6** Headquarter, Kanazawa University, Ishikawa, Japan

Abstract

Purpose: The aim of this study was to assess whether migration of thallium-201 (²⁰¹Tl) to the olfactory bulb were reduced in patients with olfactory impairments in comparison to healthy volunteers after nasal administration of ²⁰¹Tl.

Procedures: 10 healthy volunteers and 21 patients enrolled in the study (19 males and 12 females; 26–71 years old). The causes of olfactory dysfunction in the patients were head trauma (n = 7), upper respiratory tract infection (n = 7), and chronic rhinosinusitis (n = 7). ²⁰¹TlCl was administered unilaterally to the olfactory cleft, and SPECT-CT was conducted 24 h later. Separate MRI images were merged with the SPECT images. ²⁰¹Tl olfactory migration was also correlated with the volume of the olfactory bulb determined from MRI images, as well as with odor recognition thresholds measured by using T&T olfactometry.

Results: Nasal ²⁰¹Tl migration to the olfactory bulb was significantly lower in the olfactory-impaired patients than in healthy volunteers. The migration of ²⁰¹Tl to the olfactory bulb was significantly correlated with odor recognition thresholds obtained with T&T olfactometry and correlated with the volume of the olfactory bulb determined from MRI images when all subjects were included.

Conclusions: Assessment of the ²⁰¹Tl migration to the olfactory bulb was the new method for the evaluation of the olfactory nerve connectivity in patients with impaired olfaction.

Citation: Shiga H, Taki J, Washiyama K, Yamamoto J, Kinase S, et al. (2013) Assessment of Olfactory Nerve by SPECT-MRI Image with Nasal Thallium-201 Administration in Patients with Olfactory Impairments in Comparison to Healthy Volunteers. PLoS ONE 8(2): e57671. doi:10.1371/journal.pone.0057671

Editor: Thomas Hummel, Technical University of Dresden Medical School, Germany

Received: November 19, 2012; **Accepted:** January 23, 2013; **Published:** February 28, 2013

Copyright: © 2013 Shiga et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was funded in part by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture of Japan (C21592174 to H.S.). No additional external funding received for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: shigah@kanazawa-med.ac.jp

Introduction

Many physicians find it difficult to detect malingering among olfaction-impaired patients. Current olfactory function tests (the University of Pennsylvania Smell Identification Test [1], the Connecticut Chemosensory Clinical Research Center Test [2], Sniffin sticks [3], and T&T olfactometry [4]) are useful for the analysis of olfactory function in olfaction-impaired patients. However, an abnormality in the olfactory function tests has not been used as an index for the connectivity of olfactory nerve axons passing through the cribriform plate of the ethmoid bone to enter glomeruli within the olfactory bulb in patients because it is difficult to directly view the connectivity of the peripheral olfactory nerve

with current magnetic resonance imaging. On the other hand, it has previously been shown that olfactory bulb volume is correlated with odor threshold determined by using Sniffin sticks in patients with postinfectious olfactory loss [5].

The radioisotope thallium-201 (²⁰¹Tl) migrates to the olfactory bulb after nasal administration in rodents [6], and this migration is significantly decreased by the transection of olfactory nerve fibers in mice [7]. Furthermore, The ability to detect odors and the rate of migration of ²⁰¹Tl in the olfactory nerve are correlated in mice [8].

Nasally administered thallium-201 (²⁰¹Tl) migrates to the olfactory bulb 24 h after ²⁰¹Tl administration in subjects, as has been shown in healthy volunteers using a combination of single

photon emission computed tomography (SPECT), X-ray computed tomography (CT), and magnetic resonance imaging (MRI) [9]. The biological safety of nasal ^{201}Tl administration has also previously been shown [10].

In this study, we applied the technique of nasal administration of ^{201}Tl followed by SPECT-CT and MRI imaging to patients with hyposmia due to head trauma, upper respiratory tract infection, or chronic rhinosinusitis, which are major causes of olfactory dysfunction [11], and to healthy volunteers. ^{201}Tl olfactory migration in patients with impaired olfaction was compared to that in healthy volunteers. ^{201}Tl olfactory migration was also correlated with the volume of the olfactory bulb determined from MRI images, as well as with odor recognition thresholds measured by using T&T olfactometry.

Methods

Subjects

Participants were 10 healthy volunteers (7 males and 3 females, 30–66 years old; Table 1) recruited from medical workers in the Kanazawa Medical University Hospital and Kanazawa University Hospital with normal olfactory thresholds as determined with T&T olfactometry (i.e., the odor recognition threshold score of each nostril was less than 2.0) and 21 patients with olfactory disorders receiving treatment at the smell clinic in the Kanazawa Medical University Hospital (26–71 years old; 12 males and 9 females; odor recognition threshold score: 4.2 ± 1.4 [mean \pm S.D.]; Table 1; i.e., the odor recognition threshold score of each nostril was 2.0 or more than 2.0 with self-reported olfactory disturbance). The mean age was not significantly different in the patients with head trauma, upper respiratory infection, or chronic sinusitis than in the healthy volunteers (Table 1). The numbers of smokers were not significantly different among healthy volunteer group and each patient group (Table 1).

Causes of olfactory dysfunction in the patients were head trauma ($n = 7$), upper respiratory tract infection ($n = 7$), and chronic rhinosinusitis ($n = 7$). For head trauma subjects, types of head trauma were concussion ($n = 2$), cerebral contusion ($n = 2$), and cerebral contusion with intracranial hemorrhage ($n = 3$). One patient with hyposmia due to head trauma showed a shallow olfactory sulcus. Another patients with hyposmia due to head trauma showed no abnormal frontal lobe findings on MRI.

The duration of olfactory deficits at the time of the examination ranged from 4 weeks to 10 years (58 months \pm 53 months [mean \pm S.D.]). Patients with chronic rhinosinusitis were assessed in this trial if they continued to suffer olfactory impairments even though receiving medical and/or surgical treatment. Olfactory clefts were open in all subjects.

All participants were informed of the objectives of the study and possible side effects (allergic reactions to $^{201}\text{TlCl}$, irritation of the

digestive system, blood pressure fluctuation, or asthmatic crisis), and have given written informed consent. Subjects were excluded if they were pregnant or lactating. Exclusion criteria also included a history of kidney disease, liver injury, or other serious illness. Healthy volunteers who might have olfactory dysfunction due to head trauma, upper respiratory tract infection, or chronic rhinosinusitis were excluded in this study. Both groups of healthy volunteers (medical workers) and patients did not include nuclear medicine specialists. Therefore, their knowledge about ^{201}Tl imaging may be same. The Medical Ethics Committees of Kanazawa Medical University and Kanazawa University approved this trial in advance.

T&T Olfactometry

Each odorant was dissolved in mineral oil to form a graded series of concentrations, and then applied liberally to blotting paper and presented to subjects by using T&T olfactometry (Daiichi Yakuhin Sangyo, Tokyo, Japan). In Japan, T&T olfactometry is now a standard means of measuring olfactory thresholds. The normal odor recognition threshold score of each nostril is less than 2.0. The recognition threshold was obtained with T&T olfactometry on the side ^{201}Tl nasally administered in this study.

Merging of ^{201}Tl SPECT Images with MRI Images and Analysis of Olfactory Bulb Volume

All procedures involving the handling of radioisotopic materials were performed in the Division of Radiology at Kanazawa Medical University or the Division of Radioisotopes at Kanazawa University Hospital. A solution of $^{201}\text{TlCl}$ in saline (74 MBq/mL) was obtained from Nihon Medi-Physics (Tokyo, Japan). For each subject, 0.3 mL $^{201}\text{TlCl}$ saline solution (22 MBq) was instilled via syringe into the olfactory cleft in the right or left larger nasal cavity. The unilateral ^{201}Tl nasal administration was performed in this study, because uptake of ^{201}Tl in the right or left distinction may be unclear in the case of bilateral ^{201}Tl nasal administration. Under endoscopic findings, we choose the side of larger olfactory cleft to be assessed for the certain nasal administration to the olfactory cleft. After ^{201}Tl nasal administration, subjects laid on their sides for 30 min. Uptake of ^{201}Tl was assessed from SPECT scans performed 24 h after ^{201}Tl administration. It has been shown that ^{201}Tl migration from the olfactory epithelium to the olfactory bulb is visible 24 h after ^{201}Tl nasal administration in healthy volunteers [9]. Each SPECT scan was 30 min in duration and was followed by an X-ray CT scan to localize the nasal cavity and olfactory bulb. The data acquisition was started using a dual-headed SPECT-CT hybrid system (Symbia T6; Siemens Japan Healthcare, Japan) equipped with low-energy, high-resolution collimators. Data were acquired over 360 degrees, with 90

Table 1. Odor recognition thresholds measured by using T&T olfactometry.

Subject Group	Age, years (Mean \pm S.D.)	Gender (Male/Female)	Non-smokers/Smokers or ex-smokers	Odor recognition threshold (Mean \pm S.D.)
Healthy volunteer (n = 10)	44.7 \pm 11.9	7/3	7/3	0.9 \pm 0.6
Head trauma (n = 7)	43.0 \pm 13.3	3/4	3/4	4.9 \pm 0.9*
Upper respiratory tract infection (n = 7)	50.4 \pm 10.0	4/3	6/1	3.4 \pm 1.5*
Chronic rhinosinusitis (n = 7)	54.7 \pm 15.1	5/2	4/3	4.4 \pm 1.4*

* $P < 0.05$ (unpaired t-tests: comparison with healthy volunteers).

doi:10.1371/journal.pone.0057671.t001

projections (30 s per projection in continuous rotation), 128×128 matrix (with 2.29-times zoom, pixel size 2.1 mm), and 72 KeV photopeak with a 30% main window and 7% subwindow on both sides. Data reconstruction underwent by using ordered subsets expectation maximization (9 subsets and 8 iterations) with 3-dimensional resolution recovery, CT-based attenuation correction, and scatter correction [12,13] (Flash 3D, Siemens Japan Healthcare, Japan).

For each subject, MRI T2 weighted images (Syngo MR B17 3.0T; Siemens Japan Healthcare, Japan, or Signa HDx 3.0T; GE Healthcare-Japan, Japan) were collected separately from the SPECT scan, one day before. Each MRI image was merged with the corresponding SPECT image through fusion with the CT part of the SPECT-CT by using the mutual information-based registration method [14]. Experienced radiologists, (N.W. and H.T.) who were blind to the olfactory test data, separately determined the volume of the olfactory bulb by manual segmentation of MRI coronal slices (2-mm slice thickness) through the olfactory bulb on the side of nasal administration. The change of diameter at the beginning of the olfactory tract was used as the proximal demarcation of the olfactory bulb. The average scores of those were used for the statistical analysis. Olfactory bulb volume was not regarded as criteria for the selection of patients.

Analysis of ^{201}Tl Counts on the SPECT-MRI Image

Two regions of interest for the nasal turbinate area and the anterior skull base (olfactory bulb area) were set on the ^{201}Tl SPECT-MRI fusion image. On 3 sequential fused images in both sagittal and coronal planes, large regions of interest were tentatively set manually on the nasal turbinate area to cover all of the residual ^{201}Tl activity in the nasal cavity. The nasal cavity region of interest was defined as the area bounded by the 50% threshold of the peak ^{201}Tl count and delineated. Then, oval regions of interest were set manually to delineate the olfactory bulb on the side of nasal administration of the tracer by referencing the MRI T2 weighted images. The size of the oval olfactory bulb region of interest was approximately 8 to 9 pixels on the long axis and 3 to 4 pixels on the short axis in the sagittal image and 5 to 6 pixels by 2 to 3 pixels in the coronal image. The regions of interest were set excluding the sphenoidal sinus area and the nasopharyngeal area.

The index of ^{201}Tl migration from the olfactory epithelium to the olfactory bulb was determined as the ratio of the total ^{201}Tl counts in the olfactory bulb region of interest to the total ^{201}Tl counts in the nasal turbinate region of interest, expressed as a mean percentage of the values calculated from both sagittal and coronal images. Experienced nuclear radiologists (J.T. and K.O.) who were blind to the olfactory test data determined separately the two regions of interest on the SPECT-MRI image. The average scores of those were used for the statistical analysis.

Phantom Study

A phantom study was performed with the same data acquisition and reconstruction method used in the clinical study. In a cylindrical phantom (20 cm in diameter) filled with water, a ^{201}Tl (40 μCi) spherical source (4 mm in diameter) was positioned on the center of the rotation. Then the counts of pixels on the line passing through the center of the source and at a right angle to the axis of camera rotation were measured. Full width at half maximum (FWHM) was 7.0 mm. When normalized by the maximum pixel count (7972 counts) of the spherical source, pixel counts at 1 (2.1 mm), 2 (4.2 mm), 3 (6.3 mm), 4 (8.4 mm), and 5 (10.5 mm) pixels from the center of the source were 0.76, 0.34, 0.09, 0.016, 0.001, respectively. Thus, when the olfactory bulb is

located more than 4 pixels (8.4 mm) from the high radioactivity in the nasal cavity, the count contamination to the olfactory bulb will be less than 2% of the activity of the nasal cavity.

Statistical Analysis

Two-tailed Spearman correlations, unpaired t-tests, Bonferroni's multiple comparison test, and Kruskal-Wallis tests were performed using Prism 5 software (GraphPad, San Diego, CA, USA). *P* values less than 0.05 were considered significant. A sample size of 31 subjects gave 99.9% power ($\alpha = .05$, two-tailed) for comparing nasal ^{201}Tl migration to the olfactory bulb in 4 groups (JMP software, Cary, NC, USA).

Results

Reductions in Nasal ^{201}Tl Migration to the Olfactory Bulb in Patients

To determine whether the viability or function of the peripheral olfactory nerve was reduced in the patients with impaired olfaction, we assessed migration of nasally administered ^{201}Tl to the olfactory bulb in the patients and healthy volunteers. Migration of nasal ^{201}Tl to the olfactory bulb was significantly lower in the patients with head trauma, upper respiratory infection, or chronic sinusitis than in the healthy volunteers (Fig. 1; Kruskal-Wallis test for comparing 4 groups, $P = 0.0004$; unpaired t-tests for comparing 2 groups: head trauma, $P = 0.0005$; upper respiratory tract infection, $P = 0.0001$; chronic sinusitis, $P = 0.0003$; Bonferroni's multiple comparison test for comparing 2 groups: head trauma, $P < 0.0001$; upper respiratory tract infection, $P < 0.0001$; chronic sinusitis, $P < 0.0001$). There were no significant differences between each patient group in migration of nasally administered ^{201}Tl to the olfactory bulb.

Representative cases are shown in Fig. 2 and summarized in Table 2. A 60-year-old healthy male volunteer showed good olfactory function and a high level of nasal ^{201}Tl migration to the olfactory bulb (Fig. 2A, Table 2), a 44-year-old female with hyposmia after head trauma showed severe olfactory dysfunction and a low level of nasal ^{201}Tl migration to the olfactory bulb (Fig. 2B, Table 2), a 42-year-old female with hyposmia after upper respiratory tract infection showed moderate olfactory dysfunction and a low level of nasal ^{201}Tl migration to the olfactory bulb (Fig. 2C, Table 2), and a 67-year-old female with hyposmia from chronic rhinosinusitis showed moderate olfactory dysfunction and a low level of nasal ^{201}Tl migration to the olfactory bulb (Fig. 2D, Table 2). MRI images of the representative cases are shown in Fig. 3.

Nasal ^{201}Tl Migration to the Olfactory Bulb and Olfactory Function Assessed by T&T Olfactometry

To assess whether nasal ^{201}Tl migration to the olfactory bulb reflects olfactory function, we examined the relationships between the recognition threshold obtained with T&T olfactometry and the amount of nasal ^{201}Tl that migrated to the olfactory bulb in the subjects.

Nasal ^{201}Tl migration to the olfactory bulb was correlated with the odor recognition threshold obtained by T&T olfactometry (Spearman $r = -0.62$, $P = 0.0002$) in the healthy volunteers and the olfaction-impaired patients evaluated as a single group (Fig. 4). Nasal ^{201}Tl migration to the olfactory bulb was not correlated with duration of olfactory deficits at the time of the examination in the patients with impaired olfaction (Spearman $r = 0.33$, $P = 0.15$).




Figure 1. ^{201}Tl migration in patients with impaired olfaction in comparison to healthy volunteers. Nasal ^{201}Tl migration to the olfactory bulb in healthy volunteers ($n=10$) and patients with impaired olfaction due to head trauma ($n=7$), upper respiratory tract infection (respiratory infection; $n=7$), or chronic rhinosinusitis ($n=7$). P values were obtained with Bonferroni's multiple comparison test and the Kruskal-Wallis test. Bars: Mean \pm S.D.

doi:10.1371/journal.pone.0057671.g001




Figure 2. SPECT-MRI fusion image (coronal view) of nasal ^{201}Tl migration to the olfactory bulb. White arrows indicate the olfactory bulb and olfactory nerve. (A) A 60-year-old healthy male volunteer. (B) A 44-year-old female with hyposmia after head trauma. (C) A 42-year-old female with hyposmia after upper respiratory tract infection. (D) A 67-year-old female with hyposmia due to chronic rhinosinusitis. The index of ^{201}Tl migration from the olfactory epithelium to the olfactory bulb in the selected subjects was shown in Table 2.

doi:10.1371/journal.pone.0057671.g002

Table 2. Functional olfactory measurements in selected subjects.

Selected subjects	Nasal ²⁰¹ Tl migration to the olfactory bulb, %	Odor recognition threshold
60-year-old healthy male volunteer	29.0	1.4
44-year-old female with hyposmia after head trauma	4.2	5.8
42-year-old female with hyposmia after upper respiratory tract infection	4.5	4.8
67-year-old female with hyposmia due to chronic rhinosinusitis	5.0	3.2

doi:10.1371/journal.pone.0057671.t002

Nasal ²⁰¹Tl Migration to the Olfactory Bulb and Olfactory Bulb Volume Determined from MRI Images

To assess whether nasal ²⁰¹Tl migration to the olfactory bulb reflects olfactory bulb size, we examined the relationships between the olfactory bulb volume determined from MRI images and the amount of nasal ²⁰¹Tl that migrated to the olfactory bulb in the subjects. Olfactory bulb volume was significantly lower in the patients with head trauma, upper

respiratory infection, or chronic sinusitis than in the healthy volunteers (Table 3).

Nasal ²⁰¹Tl migration to the olfactory bulb was correlated with the olfactory bulb volume determined from MRI images (Spearman $r=0.73$, $P<0.0001$) in the healthy volunteers and the olfaction-impaired patients evaluated as a single group (Fig. 5).




Figure 3. MRI image (T2 weighted, coronal view). White arrows indicate the olfactory bulb and olfactory nerve. (A) A 60-year-old healthy male volunteer. (B) A 44-year-old female with hyposmia after head trauma. (C) A 42-year-old female with hyposmia after upper respiratory tract infection. (D) A 67-year-old female with hyposmia due to chronic rhinosinusitis.

doi:10.1371/journal.pone.0057671.g003




Figure 4. Correlation between olfactory ^{201}Tl migration and olfactory function as assessed by T&T olfactometry. Nasal ^{201}Tl migration to the olfactory bulb as a function of odor recognition threshold in patients and healthy volunteers ($n=31$). Spearman $r=-0.62$, $P=0.0002$. Closed circles, patients; Open squares, healthy volunteers. doi:10.1371/journal.pone.0057671.g004

Discussion

Nasal ^{201}Tl migration to the olfactory bulb was reduced in the patients with impaired olfaction due to head trauma, upper respiratory tract infection, and chronic rhinosinusitis, which are major causes of olfactory dysfunction [11] relative to the values in healthy volunteers. The assessment of olfactory nerve damage using our imaging method may be useful for scanning the lesion of the olfactory nerve connectivity in patients with impaired olfaction. Patients with impaired olfaction due to major causes of olfactory dysfunction would have damage of olfactory mucosa, because histological changes have been reported in the olfactory mucosa of patients with olfactory deficits due to head trauma [15,16], upper respiratory tract infection [17], or chronic sinusitis [18]. The degree of axon degeneration in human olfactory mucosa correlates with olfactory function [19]. Decreased peripheral olfactory neuron in olfactory mucosa of the patients with impaired olfaction may reduce the nasal ^{201}Tl migration to the olfactory bulb.

The migration of ^{201}Tl to the olfactory bulb was correlated with odor recognition thresholds obtained with T&T olfactometry when all subjects were included. Furthermore, the volume of the olfactory bulb determined by using MR images was positively correlated with the level of ^{201}Tl migration from the nasal turbinate area to the olfactory bulb on SPECT-MRI fusion images in both the healthy volunteers and patients with impaired olfaction. Our results suggest that the connectivity of olfactory neurons between the olfactory bulb and olfactory mucosa reflect

the volume of the olfactory bulb. Increasing olfactory stimulation leads to decreased cell mortality in the olfactory bulb *in vivo* [20]. Reduced connectivity of olfactory neurons between the olfactory bulb and olfactory mucosa may effect on olfactory bulb volume with a decrease of olfactory stimulation in the patients with impaired olfaction.

The nasal ^{201}Tl migration to the olfactory bulb was not significantly correlated with olfactory bulb volume determined by using MR images in the healthy volunteers or in the patients with impaired olfaction evaluated as separate groups (data not shown), likely because of the small number of subjects in each group. It has been shown that olfactory bulb volume is larger in men than in women [21]. It warrants further investigation whether gender would effect on the nasal ^{201}Tl migration to the olfactory bulb.

In this study, the decrease of the migration of ^{201}Tl to the olfactory bulb was not significantly different among the patients with head trauma, upper respiratory tract infection, or chronic sinusitis. The degree of peripheral olfactory nerve degeneration may be similar in the patient groups included in this study, because we observed no significant different olfactory thresholds among the patient groups.

In this study, we observed no adverse effects in the subjects. Unlike intravenous administration of radiopharmaceutical agents, which would deliver only small amounts of radiation to the nasal cavity, nasal administration of ^{201}Tl delivers a high radiation dose to the nasal cavity. Therefore, the estimation of the absorbed dose in the nasal cavity and neighboring organ such as brain and lens are important. We calculated the absorbed dose of ^{201}Tl in the

Table 3. Olfactory bulb volume measurements in subjects.

Subject Group	Olfactory bulb volume, mm^3 (Mean \pm S.D.)
Healthy volunteer (n=10)	83.2 \pm 25.5
Head trauma (n=7)	22.4 \pm 7.5*
Upper respiratory tract infection (n=7)	35.8 \pm 12.3*
Chronic rhinosinusitis (n=7)	33.3 \pm 10.5*

* $P<0.05$ (unpaired t-tests: comparison with healthy volunteers).

doi:10.1371/journal.pone.0057671.t003




Figure 5. Correlation between olfactory ²⁰¹Tl migration and unilateral olfactory bulb volume as determined by using MRI images. Nasal ²⁰¹Tl migration to the olfactory bulb as a volume of olfactory bulb in patients and healthy volunteers (n=31). Spearman $r=0.73$, $P<0.0001$. Closed circles, patients; Open squares, healthy volunteers.

doi:10.1371/journal.pone.0057671.g005

nasal cavity in a previous clinical study, and its dose was too low for acute radiation effects [9]. The brain and eyes are separated from the nasal cavity by the nasal bone, ethmoturbinals, and the sphenoidal bone. A Monte Carlo simulation using the ICRP reference adult male [22] showed that almost all conversion electrons and Auger electrons emitted from ²⁰¹Tl in the nasal cavity were stopped in the nasal cavity and were not absorbed by neighboring organs (data not shown). Therefore, gamma-rays and X-rays from the ²⁰¹Tl in the nasal cavity contributed the doses of ²⁰¹Tl absorbed in the brain and lens. Our preliminary calculation showed that the absorbed doses in the brain and lens after nasal administration of 22 MBq ²⁰¹Tl were 0.067 mGy and 0.59 mGy, respectively. Therefore, acute radiation effects in the brain and lens can be also avoided.

The majority of ²⁰¹Tl administered intranasally in this study migrated to the nasopharyngeal region and then was swallowed. It has been shown that intravenously administered ²⁰¹Tl is hardly absorbed by the central nervous system, including the olfactory bulb, *in vivo* [6]. Therefore, the potential systemic effects of

swallowed ²⁰¹Tl can be ignored. Residual activity in the olfactory epithelium may be independent from mucosal clearance of the administered ²⁰¹Tl, because mature olfactory neurons have immotile cilia [23].

It will be important to study whether an increase in ²⁰¹Tl migration to the olfactory bulb during treatment is correlated with a decrease in odor recognition thresholds in patients with olfactory disorders. The recovery rate of olfactory function in patients with traumatic olfactory dysfunction is less than 30% [24]. In this study, we did not assess follow-up data in the subjects, but ²⁰¹Tl migration can be used to visualize olfactory nerve regeneration *in vivo* [25]. Our imaging method could be especially helpful for assessing the connectivity of the peripheral olfactory nerve in patients with olfactory disorders during treatment and predicting the course of olfactory impairment. Patients with intact olfactory nerve fibers may be well selected by means of a new isotope imaging technique for the long-term treatment of olfactory dysfunction.

Conclusions

Nasal ²⁰¹Tl migration to the olfactory bulb was reduced in patients with impaired olfaction due to head trauma, upper respiratory tract infection, and chronic rhinosinusitis, which are major causes of olfactory dysfunction when compared to the ²⁰¹Tl migration in healthy volunteers. Assessment of the ²⁰¹Tl migration to the olfactory bulb was the new method for the evaluation of the olfactory nerve connectivity in patients with olfactory disorders.

Acknowledgments

We are grateful to the staff in the Division of Radioisotopes, Kanazawa University, and the Division of Radiology, Kanazawa Medical University, for their technical support.

Author Contributions

Conceived and designed the experiments: HS JT TM. Performed the experiments: HS JT JY KW. Analyzed the data: HS JT S. Kinase KO NW HT. Contributed reagents/materials/analysis tools: TM S. Kinuya KK RA MF. Wrote the paper: HS JT KW TM.

References

- Doty RL, Shaman P, Kimmelman CP, Dann MS (1984) University of Pennsylvania Smell Identification Test: a rapid quantitative olfactory function test for the clinic. *Laryngoscope* 94: 176–178.
- Cain WS, Gent JF, Goodspeed RB, Leonard G (1988) Evaluation of olfactory dysfunction in the Connecticut Chemosensory Clinical Research Center (CCCRC). *Laryngoscope* 98: 83–88.
- Hummel T, Sekinger B, Wolf S, Pauli E, Kobal G (1997) ‘Sniffin’ sticks’: olfactory performance assessed by the combined testing of odor identification, odor discrimination and olfactory threshold. *Chem Senses* 22: 39–52.
- Takagi SF (1987) A standardized olfactometer in Japan. A review over ten years. *Ann NY Acad Sci* 510: 113–118.
- Rombaux P, Mouraux A, Bertrand B, Nicolas G, Duprez T, et al. (2006) Olfactory function and olfactory bulb volume in patients with postinfectious olfactory loss. *Laryngoscope* 116: 436–439.
- Kanayama Y, Enomoto S, Irie T, Amano R (2005) Axonal transport of rubidium and thallium in the olfactory nerve of mice. *Nucl Med Biol* 32: 505–512.
- Kinoshita Y, Shiga H, Washiyama K, Ogawa D, Amano R, et al. (2008) Thallium transport and the evaluation of olfactory nerve connectivity between the nasal cavity and olfactory bulb. *Chem Senses* 33: 73–78.
- Shiga H, Kinoshita Y, Washiyama K, Ogawa D, Amano R, et al. (2008) Odor detection ability and thallium-201 transport in the olfactory nerve of traumatic olfactory-impaired mice. *Chem Senses* 33: 633–637.
- Shiga H, Taki J, Yamada M, Washiyama K, Amano R, et al. (2011) Evaluation of the olfactory nerve transport function by SPECT-MRI fusion image with nasal thallium-201 administration. *Mol Imaging Biol* 13: 1262–1266.
- Washiyama K, Shiga H, Hirota K, Tsuchida A, Yamamoto J, et al. (2011) Biological safety of nasal thallium-201 administration: a preclinical study for olfacto-scintigraphy. *J. Rad. Res. (Tokyo)* 52: 450–455.
- Holbrook EH, Leopold DA (2006) An updated review of clinical olfaction. *Curr Opin Otolaryngol Head Neck Surg* 14: 23–28.
- Zeintl J, Vija AH, Yahil A, Hornegger J, Kuwert T (2010) Quantitative accuracy of clinical 99mTc SPECT/CT using ordered-subset expectation maximization with 3-dimensional resolution recovery, attenuation, and scatter correction. *J Nucl Med* 51: 921–928.
- Stansfield EC, Sheehy N, Zurakowski D, Vija AH, Fahey FH, et al. (2010) Pediatric 99mTc-MDP bone SPECT with ordered subset expectation maximization iterative reconstruction with isotropic 3D resolution recovery. *Radiology* 257: 793–801.
- Ken S, Di Gennaro G, Giuletta G, Sebastiano F, De Carli D, et al. (2007) Quantitative evaluation for brain CT/MRI coregistration based on maximization of mutual information in patients with focal epilepsy investigated with subdural electrodes. *Magn Reson Imaging* 25: 883–888.
- Hasegawa S, Yamagishi M, Nakano Y (1986) Microscopic studies of human olfactory epithelia following traumatic anosmia. *Arch Otorhinolaryngol* 243: 112–116.
- Jafek BW, Eller PM, Esses BA, Moran DT (1989) Post-traumatic anosmia. Ultrastructural correlates. *Arch Neurol* 46: 300–304.
- Jafek BW, Hartman D, Eller PM, Johnson EW, Strahan RC, et al. (1990) Postviral olfactory dysfunction. *Am J Rhinol* 4: 91–100.
- Kern RC (2000) Chronic sinusitis and anosmia: pathologic changes in the olfactory mucosa. *Laryngoscope* 110: 1071–1077.
- Holbrook EH, Leopold DA, Schwob JE (2005) Abnormalities of axon growth in human olfactory mucosa. *Laryngoscope* 115: 2144–2154.

20. Rochefort C, Gheusi G, Vincent JD, Lledo PM (2002) Enriched odor exposure increases the number of newborn neurons in the adult olfactory bulb and improves odor memory. *J Neurosci* 22: 2679–2689.
21. Buschhüter M, Smitka S, Puschmann S, Gerber JC, Witt M, et al. (2008) Correlation between olfactory bulb volume and olfactory function. *Neuroimage* 42: 498–502.
22. Menzel HG, Clement C, DeLuca P (2009) ICRP Publication 110. Realistic reference phantoms: an ICRP/ICRU joint effort. A report of adult reference computational phantoms. *Ann. ICRP* 39: 1–164.
23. Mair RG, Gesteland RC, Blank DL (1982) Changes in morphology and physiology of olfactory receptor cilia during development. *Neuroscience* 7: 3091–3103.
24. Fujii M, Fukazawa K, Takayasu S, Sakagami M (2002) Olfactory dysfunction in patients with head trauma. *Auris Nasus Larynx* 29: 35–40.
25. Shiga H, Washiyama K, Hirota K, Amano R, Furukawa M, et al. (2009) Use of thallium transport to visualize functional olfactory nerve regeneration in vivo. *Rhinology* 47: 460–464.