

# Prostaglandin E2 signaling and bacterial infection recruit tumor-promoting macrophages to mouse gastric tumors

著者	Oshima Hiroko, Hioki Kyoji, Popivanova Boryana K., Oguma Keisuke, Rooijen Nico van, Ishikawa Tomoo, Oshima Masanobu
journal or publication title	Gastroenterology
volume	140
number	2
page range	596-607
year	2011-02-01
URL	<a href="http://hdl.handle.net/2297/26532">http://hdl.handle.net/2297/26532</a>

doi: 10.1053/j.gastro.2010.11.007

# Prostaglandin E<sub>2</sub> signaling and bacterial infection recruit tumor-promoting macrophages to mouse gastric tumors

Hiroko Oshima<sup>#</sup>, Kyoji Hioki<sup>\*</sup>, Boryana K. Popivanova<sup>#</sup>, Keisuke Oguma<sup>#,‡</sup>, Nico van Rooijen<sup>§</sup>, Tomo-o Ishikawa<sup>#</sup>, Masanobu Oshima<sup>#</sup>

<sup>#</sup>Division of Genetics, Cancer Research Institute, Kanazawa University, Kanazawa, Japan.

<sup>\*</sup>Gnotobiology Laboratory, Central Institute for Experimental Animals, CIEA, Kawasaki, Japan. <sup>‡</sup>Research Fellow of the Japan Society for the Promotion of Science, Japan.

<sup>§</sup>Department of Molecular Cell Biology, Vrije Universiteit, Amsterdam, The Netherlands

**Short title:** PGE<sub>2</sub> and infection in tumorigenesis

## **Grant Support & Assistance:**

Supported by Grants-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan, and the Ministry of Health, Labour and Welfare of Japan.

The authors thank Manami Watanabe for her excellent technical assistance.

## **Correspondence:**

Masanobu Oshima, D.V.M., Ph.D.,

Division of Genetics, Cancer Research Institute, Kanazawa University,  
Kakuma-machi, Kanazawa, 920-1192 Japan.

Phone: +81-76-264-6760, FAX: +81-76-234-4519

e-mail: [oshimam@kenroku.kanazawa-u.ac.jp](mailto:oshimam@kenroku.kanazawa-u.ac.jp)

**Conflict of Interest:** No conflicts of interest exist.

**Author Contributions:** Hiroko Oshima performed in vivo experiments. Kyoji Hioki generated germfree mouse colonies in the CIEA. Boryana Popivanova performed chemokine expression analysis. Keisuke Oguma performed Wnt analysis. Nico van Rooijen prepared clodronate liposome. Tomo-o Ishikawa performed the primary culture experiments. Masanobu Oshima designed the study and wrote the manuscript.

**Word count:** 5,998

**Abbreviations:**

COX-2, cyclooxygenase-2; DSS, dextran sodium sulfate; GF, germfree; IL-1 $\beta$ , interleukin-1 $\beta$ ; IL-6, interleukin-6; LPS, lipopolysaccharide; mPGES-1, microsomal prostaglandin E synthase-1; PGE<sub>2</sub>, prostaglandin E<sub>2</sub>; RT-PCR, reverse transcription-polymerase chain reaction; SPF, specific pathogen free; TAM, tumor-associated macrophage; TLR, toll-like receptor; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ .

Abstract:

**Background & Aims:** *Helicobacter pylori* infection induces an inflammatory response, which can contribute to gastric tumorigenesis. Induction of cyclooxygenase-2 (COX-2) results in production of prostaglandin E<sub>2</sub> (PGE<sub>2</sub>), which mediates inflammation. We investigated the role of bacterial infection and the PGE<sub>2</sub> pathway in gastric tumorigenesis.

**Methods:** We generated a germ-free (GF) colony of *K19-Wnt1/C2mE* mice (*Gan* mice), which develop gastric cancer, and examined tumor phenotypes, expression of cytokines and chemokines, and recruitment of macrophages. We also investigated PGE<sub>2</sub> signaling through the receptor EP4 in *Gan* mice injected with specific inhibitors.

**Results:** Gastric tumorigenesis was significantly suppressed in GF-*Gan* mice; reconstitution of commensal flora or infection with *Helicobacter felis* induced gastric tumor development. Macrophage infiltration was significantly suppressed in the stomachs of GF-*Gan* mice and EP4 inhibitor-treated *Gan* mice with decreased expressions of cytokines and chemokines. PGE<sub>2</sub> signaling and bacterial infection or stimulation with lipopolysaccharide induced expression of the chemokine CCL2 (which attracts macrophage) in tumor stromal cells or cultured macrophages, respectively. CCL2 inhibition suppressed macrophage infiltration in tumors and depletion of macrophages from the *Gan* mouse tumors led to signs of tumor regression. Wnt signaling was suppressed in the tumors of GF-*Gan* and *Gan* mice given injections of tumor necrosis factor (TNF)- $\alpha$  neutralizing antibody.

**Conclusions:** Bacterial infection and PGE<sub>2</sub> signaling are required for gastric tumorigenesis in mice, by the cooperative upregulation of CCL2, which recruits macrophage recruitment to gastric tumors. Macrophage-derived TNF- $\alpha$  promotes Wnt signaling in epithelial cells, which contributes to gastric tumorigenesis.

**KEY WORDS:** LPS; stomach cancer, tumor progression, *H. pylori*

## Introduction

Gastric cancer is the second most common cause of cancer-related death in the world, and *Helicobacter pylori* infection is closely associated with gastric cancer development<sup>1</sup>. Infections are estimated to be related to 15% of malignant cancer development, and infection-associated inflammation is a critical component of cancer development<sup>2</sup>. For example, an inflammatory response promotes tumor cell proliferation, metastasis and survival, whereas it suppresses anti-tumor immune responses<sup>2-4</sup>. Moreover, the genetic polymorphisms in genes encoding inflammatory cytokines influence gastric tumorigenesis<sup>5</sup>. These results suggest that an inflammatory cytokine network induced by *H. pylori* infection plays a key role in gastric tumorigenesis.

Cyclooxygenase-2 (COX-2) is an inducible enzyme for prostaglandin biosynthesis, which plays an important role in both inflammation and tumorigenesis<sup>6,7</sup>. Mouse model studies have indicated that induction of the COX-2/prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) pathway accelerates intestinal tumorigenesis through the induction of angiogenesis and suppression of apoptosis<sup>8,9</sup>. Among four PGE<sub>2</sub> receptors, EP1-EP4, EP4 receptor signaling has been shown to play an important role in intestinal tumorigenesis through the activation of epidermal growth factor receptor<sup>10</sup>. The expression of COX-2 is also found in more than 70% of gastric cancers<sup>11</sup>, which is suppressed by the eradication of *H. pylori*<sup>12</sup>, suggesting COX-2 induction by infection in the gastric mucosa. Transgenic mice expressing COX-2 and a PGE<sub>2</sub> converting enzyme, mPGES-1, in the stomach develop hyperplasia with macrophage infiltration, indicating the role of PGE<sub>2</sub> in macrophage recruitment<sup>13</sup>. Tumor-associated macrophages (TAMs) play an important role in tumorigenesis through the enhancement of

angiogenesis, migration, and remodeling<sup>14</sup>. Moreover, the simultaneous activation of Wnt and the PGE<sub>2</sub> pathways in the mouse stomach causes dysplastic tumor development<sup>15</sup>. Accordingly, it is possible that PGE<sub>2</sub>-dependent macrophage recruitment is one of the important mechanisms underlying the *H. pylori* infection-associated inflammation in gastric tumorigenesis. However, the relationship between bacterial infection and PGE<sub>2</sub> signaling in gastric tumorigenesis remains unclear.

Commensal bacteria constitutively stimulate the intestinal mucosa, inducing cytokines and chemokines at a basal level, which is important for homeostasis of the intestinal mucosa<sup>16</sup>. The present study shows that indigenous bacteria constitutively stimulate the gastric mucosa, which is required for tumorigenesis in the Wnt-activated and PGE<sub>2</sub>-induced gastric mucosa. Bacterial colonization and PGE<sub>2</sub> signaling through EP4 receptor cooperatively induced the expression of CCL2, which was a major pathway for macrophage recruitment in the gastric mucosa. Furthermore, depletion of macrophages caused regressive signs of tumors. These results indicate that bacterial infection and PGE<sub>2</sub> signaling cooperatively recruit macrophages to the gastric mucosa, which promotes gastric cancer development.

## Materials and Methods

### Animal experiments

The construction of *K19-Wnt1* [*Tg(Krt19-Wnt1)2Maos*], *K19-C2mE* [*Tg(Krt19-Ptgs2,Krt19-Ptges)8Tko*], and *K19-Wnt1/C2mE* (*Gan* for Gastric neoplasia) [*Tg(Krt19-Wnt1)2Maos/Tg(Krt19-Ptgs2,Krt19-Ptges)8Tko*] transgenic mice was described previously (Supplementary Table 1)<sup>13,15</sup>. All mice used in the present study were backcrossed to C57BL/6 mice more than 12 times (>N12). Germfree mouse colonies of all genotypes and wild-type mice were established by Caesarean derivation at the Central Institute for Experimental Animals (CIEA, Japan), and the mice were raised in germfree isolators under germfree conditions. During the experiments, germfree conditions were monitored by cultures of feces, bedding, and swabs in thioglycollate medium or potato dextrose broth every 2-5 weeks (Supplementary Table 2). SPF-raised mice were maintained in the SPF facility at Kanazawa University. The excluded pathogens in the SPF facility are listed in Supplementary Table 3. Mice were euthanized and examined at 30 and 55 weeks of age (n=4-6), and all experimental groups consisted of both female and male mice. Half of the glandular stomach was used for the histological analysis, and the other half was used for RNA and protein sample preparation. Six germfree *Gan* mice were moved to the SPF facility at 7 weeks of age, reconstituted with commensal flora by co-housing with SPF mice and adding dirty bedding from other cages, and examined at 30 weeks of age. *Helicobacter felis* (ATCC 49179) were inoculated at 10<sup>8</sup>/mouse *p.o.* to germfree *Gan* mice at 30 weeks of age (n=3), and gastric phenotypes were examined at 55 weeks of age. The time course and experimental conditions for each group are shown in Supplementary Figure 1. All animal experiments were carried



out according to the protocol approved by the Committee on Animal Experimentation of Kanazawa University.

### **Gastric pH Measurement**

Gastric pH was measured as described<sup>17</sup>. Briefly, mice were fasted overnight before necropsy. Sterile water (1.5 ml) was injected into the stomach, the stomach was massaged gently, and the pH of the gastric contents was measured using a pH meter.

### **Drug treatment experiments**

The EP4 receptor inhibitor, RQ-00015986/CJ-42794<sup>18</sup> was provided from RaQualia Pharma Inc. (Taketoyo, Japan). *Gain* mice were treated with RQ-00015986 at 100 mg/kg/day *p.o.* from 27 or 52 weeks of age for 3 weeks (n=4-5). For inhibition of TNF- $\alpha$  or CCL2, *Gain* mice were injected with a neutralizing antibody against TNF- $\alpha$  (AB-410-NA, R&D Systems, Minneapolis, MN) at 8 mg/kg/day *i.p.* for 6 days (n=3) or an antibody against CCL2 (AF-479-NA, R&D Systems) at 1 mg/kg/day *i.p.* for 3 days (n=3), respectively. Macrophages were depleted *in vivo* by injection of 200  $\mu$ l of clodronate (dichloromethylene bisphosphonate)-loaded liposomes *i.v.* every 3 days for 2 weeks as previously described<sup>19</sup>. Clodronate was a gift from Roche Diagnostics GmbH (Mannheim, Germany), and was encapsulated in liposomes as described previously<sup>20</sup>.

### **Histology and immunohistochemistry**

Stomach tissues were fixed in 4% paraformaldehyde, paraffin-embedded and sectioned at 4  $\mu$ m-thickness. These sections were stained with H&E or processed for immunostaining. Tissues were also embedded in OCT compound, frozen in liquid nitrogen, and sectioned at 10  $\mu$ m-thickness. The frozen sections were used for CCL2 immunostaining. Antibodies against

the proton pump (MBL, Nagoya, Japan), F4/80 (Serotec, Oxford, UK),  $\beta$ -catenin (Sigma, St. Louis, MO), CCL2 (Hycult Biotech, Uden, The Netherlands), EP4 (MBL), CD44 (Millipore, Billerica, MA), EphB3 (Bioworld Technology, St. Louis Park, MN) and Ki-67 (Dako, Carpinteria, CA) were used as the primary antibodies. Staining signals were visualized using the Vectastain Elite Kit (Vector Laboratories, Burlingame, CA). For fluorescence immunostaining, Alexa Fluor 594 or Alexa Fluor 488 antibody (Molecular Probes, Eugene, OR) was used as the secondary antibody. Apoptosis was examined using the ApopTag Apoptosis Detection Kit (Millipore). The mean index for F4/80 (macrophage), proton pump (parietal cell), Ki-67, or apoptosis was calculated by counting labeled cells per microscopic field (200 $\times$ ) in 5 fields.

### **Scoring tumor volume and preneoplastic lesions**

The mucosal thickness (tumor height) of the gastric tumors of *Gan* mice and the normal stomach of wild-type mice was measured from histology sections. The mucosal thickness relative to that of wild-type mice was calculated. The number of preneoplastic lesions in the whole glandular stomach of *K19-Wnt1* mice was counted under a dissection microscope after staining with 0.05% toluidine blue. The histological characteristics of gastric tumors and preneoplastic lesions were described previously<sup>15</sup>. The histology of these lesions was confirmed after scoring.

### **Cell culture experiments**

The RAW264 macrophage cells (RIKEN BioResource Center, Japan) were treated with lipopolysaccharide (LPS) (Sigma) for 24 h at 1, 10, 50, 100, 1,000, or 10,000 pg/ml with or without treatment with a COX-2 inhibitor, celecoxib, or RQ00015986 at 10  $\mu$ M, and the

expression levels of CCL2 and cytokines were examined. Celecoxib was provided by Pfizer (New York, NY). For the primary culture of gastric epithelial cells, the glandular stomach of *K19-Wnt1* mice was treated with 0.1% collagenase followed by trypsin, and epithelial cells were cultured in matrigel (BD Pharmingen, Franklin Lakes, NJ) with EGF (-) primary culture medium<sup>13</sup>. The primary cultured cells were treated with RQ00015986 at 10  $\mu$ M for 6 days, and organoid structures consisting of epithelial cells larger than 0.2 mm in diameter were counted.

### **Immunoblotting analysis**

The tissues were homogenized and sonicated in lysis buffer. Thereafter, the specimens were centrifuged at 20,000  $\times$  g, and 10  $\mu$ g of the supernatant protein sample was separated in a 10% SDS-polyacrylamide gel. An antibody against unphosphorylated active  $\beta$ -catenin (Millipore) was used as the primary antibody. Anti- $\beta$ -actin (Sigma) was used as the internal control. The ECL detection system (GE Healthcare, Buckinghamshire, UK) was used to detect the signals.

### **Real-time reverse transcription-polymerase chain reaction (RT-PCR)**

Tumor stroma and epithelial cell samples were separately collected from frozen sections using Laser Microdissection (Leica, Wetzlar, Germany). Total RNA was extracted from the tissues or microdissected samples using ISOGEN (Nippon Gene, Tokyo, Japan), reverse transcribed with PrimeScript RT reagent Kit (Takara, Tokyo, Japan), and PCR-amplified by Stratagene Mx300P (Agilent Technologies, Santa Clara, CA) using SYBR Premix ExTaqII (Takara). Primers for the real-time RT-PCR were purchased (Takara), and the primer sequences are shown in Supplementary Table 4.

### **Statistical analysis**

The data were analyzed using the unpaired *t*-test and are presented as the means  $\pm$  standard deviation (s.d.). A value of  $P < 0.05$  was considered to be statistically significant (Asterisks,  $P < 0.05$ ).

## Results

### *Constitutive stimulation of gastric mucosa by indigenous bacteria*

To determine whether indigenous bacteria in the stomach stimulate the gastric mucosa, we examined the expression of cytokines and chemokines in the glandular stomach of germfree (GF) wild-type mice and control SPF mice by real-time RT-PCR (Figure 1A). Importantly, the expression levels of TNF- $\alpha$ , IL-1 $\beta$ , KC and MIP-2 decreased significantly in the GF mouse stomach compared to the SPF mice, while IL-6 expression slightly increased. These results indicate that commensal bacteria constitutively stimulate the gastric mucosa to induce inflammatory cytokines and chemokines at a basal level.

*K19-Wnt1/C2mE* gastric tumor model mice (*Gan* mice for Gastric neoplasia) were previously constructed by crossing *K19-Wnt1* and *K19-C2mE* mice (Supplementary Table 1)<sup>13,15</sup>. We examined the number of parietal cells by immunostaining and measured intragastric pH in these models, because bacterial growth is affected by intragastric acidity. The mean parietal cell index was similar in all strains, and the intragastric pH was around 3.5 (Figure 1B-D). However, in the dysplastic tumors of *Gan* mice, the number of parietal cells decreased significantly, and intragastric pH increased at 50-55 weeks of age. These results indicate that gastric acidity is not changed in *Gan* mice at early stages of disease compared to other models until large tumors develop at the later stages.

### *Suppression of gastric tumorigenesis in germfree mice*

*Gan* mice raised in an SPF facility (SPF-*Gan*) developed large gastric tumors by 55 weeks of age (Figure 2A and B), and the mean tumor height increased to about 1.5-fold when compared to that at 30 weeks of age. In contrast, gastric tumorigenesis was significantly

suppressed in the germfree *Gan* (GF-*Gan*) mice, and the mean mucosal thickness of GF-*Gan* mice was less than 40% of the age-matched SPF-*Gan* mice (Figure 2A and B, Supplementary Figure 2). Approximately 40% of the SPF-*Gan* mice showed a moribund phenotype and thus were euthanized by 60 weeks of age, whereas all GF-*Gan* mice survived by 55 weeks of age (Figure 2C). Importantly, reconstitution of commensal bacteria in GF-*Gan* mice resulted in development of gastric tumors that were significantly larger than those of GF-*Gan* mice (Figure 2A, Supplementary Figure 2), and treatment of SPF-*Gan* mice with antibiotics significantly suppressed gastric tumor growth (Supplementary Figure 3). These results indicate that colonization of indigenous bacteria is required for gastric tumor development.

Moreover, infection with *Helicobacter felis*, separate species of *Helicobacter pylori*, in the GF-*Gan* mouse stomach at 30 weeks of age (GF->*H. felis* mice) induced the development of gastric tumors by 55 weeks of age (Figure 2A and B). The infection of *H. felis* in gastric glands was confirmed by microscopy of histology sections (Supplementary Figure 4). These results also indicate the role of infection in gastric tumorigenesis.

Apoptotic cells were found only on the mucosal surface of gastric tumors (Figure 2D), and the mean apoptosis index was 42.4% and 40.0% on the mucosal surface in SPF-*Gan* and GF-*Gan* mice, respectively. On the other hand, the number of Ki-67-labeled proliferating cells was significantly lower in the GF-*Gan* mouse stomach (Figure 2E and F), suggesting that bacterial colonization contributes to the tumor cell proliferation.

#### ***Suppression of gastric tumorigenesis by inhibition of the PGE<sub>2</sub> receptor, EP4***

There are four PGE<sub>2</sub> receptors, EP1-EP4, and the expression level of EP4 was increased significantly in *Gan* mouse tumors<sup>21</sup>. SPF-*Gan* mice were thus treated with a specific EP4

inhibitor, RQ-00015986, for 3 weeks from 52 weeks of age (Supplementary Figure 1).

Notably, gastric tumors regressed significantly following inhibition of EP4, and the mean tumor size decreased to 23% of that of the age-matched SPF-*Gan* mice (Figure 2A and B).

Moreover, treatment of SPF-*Gan* mice with an EP4 inhibitor during the early stage of tumorigenesis from 27 weeks of age also suppressed tumor development significantly. These results indicate that EP4 signaling plays an important role in gastric tumorigenesis.

### ***Bacterial infection and EP4 signaling for inflammatory responses***

Submucosal lymphocyte infiltration was found in the SPF-*Gan* and GF->*H. felis* mouse tumors but not in GF-*Gan* and EP4 inhibitor-treated *Gan* (EP4i-*Gan*) mice (Figure 3A), suggesting that both bacterial infection or colonization and EP4 signaling are required for inflammatory responses. Moreover, the expression level of proinflammatory cytokines, TNF- $\alpha$ , IL-1 $\beta$ , and IL-6, and chemokines, KC and MIP-2, were significantly elevated in the SPF-*Gan* and GF->*H. felis* gastric tumors (Figure 3B), indicating inflammatory responses in the stomachs of these mice. In contrast, the expression of these cytokines and chemokines was not induced in the GF-*Gan* and EP4i-*Gan* mouse stomachs. The transgenic expression of COX-2 (*Ptgs2*) in the stomach was confirmed in all mouse groups except the wild-type mice, indicating that the PGE<sub>2</sub> level was elevated in all groups of *Gan* mice. These results indicate that both infection and EP4 signaling are required for inflammatory responses in the stomach.

The expression of proinflammatory cytokines was predominantly detected in stromal cells when compared with tumor epithelial cells in SPF-*Gan* tumors (Figure 3C), suggesting that stromal macrophages were the major source of these cytokines. We thus examined macrophage infiltration by immunostaining. As expected, numerous macrophages were

found in the SPF-*Gan* and GF->*H. felis Gan* mouse tumors, while macrophage infiltration was suppressed in the GF-*Gan* and EP4i-*Gan* mice (Figure 3D). The number of macrophages in the GF-*Gan* and EP4i-*Gan* mouse stomachs decreased significantly to 11% and 33% of the SPF-*Gan* mouse level, respectively (Figure 3E). These results suggest that cooperation of bacterial infection or colonization with EP4 signaling contributes to macrophage recruitment to the gastric mucosa.

To examine the role of macrophages in gastric tumorigenesis, SPF-*Gan* mice were treated with clodronate liposomes to deplete macrophages *in vivo*. Macrophage-depleted areas were found by immunostaining in the clodronate liposome-injected *Gan* mouse tumors (Figure 3F). In the macrophage-depleted area, tumors showed regressive signs with atrophic changes of tumor cells or peel-off of tumor epithelial cells from the mucosal surface, suggesting that macrophages are important for the maintenance or survival of tumor epithelial cells.

#### ***Induction of CCL2 by bacterial infection and EP4 signaling***

The expression level of macrophage-tropic chemokines was examined in the SPF-*K19-C2mE* and GF-*K19-C2mE* mouse stomach. The PGE<sub>2</sub> pathway but not Wnt signaling is activated in the glandular stomach of *K19-C2mE* mice (Supplementary Table 1). Importantly, expression of CCL2 and CCL8 increased significantly in the SPF-*K19-C2mE* mouse stomach compared to that in the wild-type mice (Figure 4A). In contrast, expression of CCL2 and CCL8 was not induced in the GF-*K19-C2mE* mouse stomach. Induction of CCL2 and CCL8 expression was also found in gastric tumors of SPF-*Gan* mice but not in GF-*Gan* mice (Figure 4B). These results suggest that PGE<sub>2</sub> signaling and bacterial colonization



cooperatively induce expression of CCL2 and CCL8 in the gastric mucosa.

Because CCL2 is an important chemokine for macrophage infiltration in colon tumors<sup>22</sup>, we further examined CCL2 expression. CCL2-expressing cells were detected by immunostaining in the stroma of SPF-*Gan* mouse tumors where macrophages were accumulated (Figure 4C), suggesting that macrophages express CCL2. The stimulation of RAW264 macrophage cells with LPS induced the expression of CCL2 in a dose-dependent manner (Figure 4D). A low concentration of LPS was used for further experiments to examine the effect of low counts of indigenous bacterial colonization in the stomach. Importantly, treatment of RAW264 macrophages with an EP4 inhibitor significantly suppressed LPS-induced CCL2 expression (Figure 4E), suggesting that both EP4 signaling and LPS stimulation are required for CCL2 induction in macrophages.

To examine the role of CCL2 in macrophage recruitment in gastric tumors, SPF-*Gan* mice were treated with an anti-CCL2 neutralizing antibody. Notably, inhibition of CCL2 suppressed macrophage infiltration in tumors, although a few macrophages were still detected (Figure 4F), suggesting that CCL2 is a major chemokine that recruits macrophages to gastric tumors. In the macrophage-depleted tumor areas caused by CCL2 inhibition, tumors showed regressive signs, which was similar to the observations in the clodronate liposome-treated *Gan* mice (Figure 3F).

### ***M2 type polarization of macrophages in gastric tumors***

TAMs play a pivotal role in tumor development<sup>14</sup>. Macrophage activation is classified as “classically” activated (M1) or “alternatively” activated (M2) type, and TAMs generally express characteristics of M2-polarized macrophages<sup>23</sup>. Interestingly, the expression of M2

macrophage markers, Ym1, Ym2, Arg1, and TGF- $\beta$ , increased significantly in both the SPF-*K19-C2mE* stomach and SPF-*Gan* tumors (Supplementary Figure 5A). Consistently, expression of an M2 marker, mannose receptor, was detected by immunohistochemistry in the SPF-*Gan* mouse tumors (Supplementary Figure 5B). It has been reported that CD4<sup>+</sup> T cells regulate the M2 properties of macrophages<sup>24</sup>. CD4<sup>+</sup> T cells infiltrated into the SPF-*Gan* mouse tumors (Supplementary Figure 5C). However, M2 macrophages were also found in SPF-*Rag2*<sup>-/-</sup> *K19-C2mE* mouse stomachs (Supplementary Figure 5D), suggesting that macrophages in the PGE<sub>2</sub>-induced inflammation can be polarized to the M2 type in the absence of T cells.

#### ***EP4 signaling on macrophages and epithelial cells***

Expression of the EP4 receptor was detected by immunostaining in both tumor epithelial cells and stromal cells of SPF-*Gan* mice, whereas it was rarely detected in the GF-*Gan* or wild-type mouse stomach (Figure 5A). Induction of EP4 in SPF-*Gan* tumors was confirmed by real-time RT-PCR (Figure 5B). When RAW264 macrophages were stimulated with LPS, expression of COX-2 and mPGES-1, as well as proinflammatory cytokines, was elevated (Figure 5C). Importantly, treatment of LPS-stimulated macrophages with an EP4 inhibitor or celecoxib suppressed the induction of COX-2, IL-1 $\beta$  and IL-6. Moreover, inhibition of EP4 suppressed proliferation of the primary cultured gastric epithelial cells in matrigel (Figure 5D). These results suggest that EP4 signaling is also important for macrophage activation and epithelial cell proliferation.

#### ***Wnt promotion by bacterial infection and TNF- $\alpha$ stimulation***

Expression of Wnt-target genes, CD44 and EphB3, was significantly downregulated in

the GF-*Gan* and EP4i-*Gan* mouse stomachs (Figure 6A). In the wild-type mouse stomach, expression of CD44 was found only in the neck of the gastric gland, whereas EphB3 was not detected (Figure 6B). Notably, expression of CD44 and EphB3 was significantly induced in tumor epithelial cells of SPF-*Gan* mice, which was suppressed in GF-*Gan* mice. Consistently, the active  $\beta$ -catenin level was decreased in the GF-*K19-Wnt1* stomach and GF-*Gan* mouse gastric tumors compared to SPF mice (Figure 6C), indicating that Wnt signaling activity is suppressed under GF conditions. We previously showed that macrophage-derived TNF- $\alpha$  promotes Wnt signaling in gastric cancer cells<sup>25</sup>. It is therefore possible that the decreased level of TNF- $\alpha$  in the GF-*Gan* and EP4i-*Gan* mouse stomachs resulted in suppression of Wnt signaling.

*K19-Wnt1* mice develop preneoplastic lesions caused by promotion of Wnt signaling by macrophage-derived TNF- $\alpha$ <sup>25</sup> (Supplementary Table 1). Importantly, the number of preneoplastic lesions decreased significantly in the GF-*K19-Wnt1* mice compared to the SPF-*K19-Wnt1* mice (Figure 6D), suggesting that bacterial colonization is important for macrophage recruitment, which triggers TNF- $\alpha$ -induced Wnt promotion.

We next investigated whether TNF- $\alpha$  promotes Wnt signaling in the gastric tumor tissues. Treatment of the SPF-*Gan* mice with an anti-TNF- $\alpha$  neutralizing antibody resulted in downregulation of TNF- $\alpha$  in tumors, possibly caused by suppression of macrophage activation (Figure 6E). Importantly, expression of EphB3, a Wnt-target gene, was also downregulated. Consistently, the active  $\beta$ -catenin level decreased in gastric tumors by TNF- $\alpha$  inhibition (Figure 6F). These results indicate that macrophage-derived TNF- $\alpha$  enhances Wnt activity in gastric tumors, which may promote gastric tumorigenesis.

## Discussion

Accumulating evidence has indicated that infection-associated inflammation plays an important role in cancer development<sup>2</sup>. Bacterial infection stimulates toll-like receptors (TLRs), which induces activation of the NF- $\kappa$ B pathway. The activation of NF- $\kappa$ B causes tumor promotion through induction of growth factors and suppression of apoptosis<sup>26</sup>. NF- $\kappa$ B activation also induces COX-2 expression, which is followed by induction of the PGE<sub>2</sub> pathway. The COX-2/PGE<sub>2</sub> pathway plays a key role in intestinal tumorigenesis<sup>8,9</sup>. These results indicate that infection plays a key role in the activation of the NF- $\kappa$ B and PGE<sub>2</sub> pathways, which promotes tumorigenesis. In the present study, we infected *H. felis* at 30 weeks of age to examine the effect of infection in tumorigenesis, because we found a significant suppression of tumorigenesis in GF-*Gan* stomach at 30 weeks of age. Importantly, *H. felis* infection to GF-*Gan* mice induced gastric tumor development by 55 weeks. Accordingly, the present study indicates that bacterial infection is still required for tumorigenesis even after the induction of the PGE<sub>2</sub> pathway.

It has been shown that PGE<sub>2</sub> signaling through EP4 receptor is important for intestinal tumorigenesis through the activation of epidermal growth factor receptor<sup>10</sup>. The current results also showed EP4 to play an important role in tumorigenesis. Bacterial colonization and EP4 signaling cooperatively induce expression of macrophage-tropic chemokine, CCL2. It has been shown that CCL2 signaling is important for macrophage infiltration in colon cancers in the intestinal tumorigenesis<sup>22</sup>. Accordingly, it is possible that expression of CCL2 induced by bacterial colonization and EP4 signaling is important for macrophage recruitment in gastric tumorigenesis.

Intestinal commensal bacteria stimulate the TLRs in the mucosa, which is important for the proliferation of undifferentiated epithelial cells<sup>16</sup>. Moreover, macrophages are an important niche component for the proliferation of intestinal progenitor cells in the tissue repair process<sup>27</sup>. Accordingly, it is possible that the innate immune response to commensal bacteria is important for the proliferation of tumor epithelial cells through macrophage recruitment. On the other hand, acquired immunity by T cells is essential for *H. felis*-associated gastric pathology<sup>28</sup>. It is thus possible that T cells play a role in gastric tumorigenesis in *H. felis*-infected GF-*Gan* mice. However, hyperplasia still developed, and the macrophages were polarized to M2, in the SPF-*Rag2*<sup>-/-</sup> *K19-C2mE* mouse stomach<sup>29</sup> (Supplementary Figure 5D), suggesting that increased PGE<sub>2</sub> levels and commensal flora can trigger these gastric phenotypes without T cell response.

We previously showed that macrophage-derived TNF- $\alpha$  promotes Wnt signaling activity in gastric cancer cells, which contributes to gastric tumorigenesis<sup>25</sup>. Moreover, Wnt activation levels correlate with the incidence of intestinal tumorigenesis in *Apc* knockout mice<sup>30</sup>, and promotion of Wnt signaling activity may play an important role in malignant progression<sup>31</sup>. In the present study, inhibition of TNF- $\alpha$  resulted in a decrease in Wnt signaling activity in gastric tumors, confirming that TNF- $\alpha$  functions as a Wnt promoting factor *in vivo*. Notably, Wnt activity in the gastric tumors of GF-*Gan* mice was lower than that of SPF-*Gan* mice, which may have been caused by a decreased level of macrophage-derived TNF- $\alpha$ . Accordingly, it is possible that TNF- $\alpha$ -dependent Wnt promotion is one of the important mechanisms by which macrophages induce tumorigenesis, which is triggered by bacterial colonization and EP4 signaling.

The COX-2/PGE<sub>2</sub> pathway has been shown to suppress the Th1 immune response in the *H. pylori*-infected stomach<sup>32</sup>. Notably, NSAID treatment suppresses gastric carcinogenesis in the INS-GAS gastric tumor model mice. However, NSAID treatment enhances gastritis in *Helicobacter*-infected INS-GAS mice, which may promote gastric tumorigenesis<sup>17</sup>. These results suggest that the PGE<sub>2</sub> pathway suppresses infection-associated carcinogenesis, which appears to be inconsistent with the present results. However, the present results indicate that commensal flora with low bacterial counts can elicit gastritis when the mucosal PGE<sub>2</sub> level is increased, and such commensal flora and PGE<sub>2</sub>-dependent inflammation is important for gastric tumorigenesis. It is therefore possible that the role of PGE<sub>2</sub> for immune responses and tumorigenesis varies according to the level of infection status, such as exogenous aggressive infection by *Helicobacter* or commensal colonization, although this remains to be investigated.

In conclusion, bacterial infection or colonization, in cooperation with PGE<sub>2</sub> signaling through the EP4 receptor, induces expression of CCL2, resulting in macrophage recruitment to gastric mucosa. TNF- $\alpha$  produced by macrophages promotes Wnt signaling in the tumor cells, which may promote gastric tumorigenesis. Accordingly, the eradication and inhibition of the PGE<sub>2</sub> pathway may be an effective strategy for preventing gastric cancer development.

## REFERENCES

1. Correa P, Camargo MC, Piazuelo MB. Overview and pathology of gastric cancer. In: Wang TC, Fox JG, Giraud AS Eds. *The Biology of Gastric Cancer*. New York: Springer, 2008, 1-24.
2. Coussens LM, Werb Z. Inflammation and cancer. *Nature* 2002;420,860-867.
3. Balkwill F, Mantovani A. Inflammation and cancer: back to Virchow. *Lancet* 2001;357,539-545.
4. Mantovani A, Allavena P, Sica A, et al. Cancer-related inflammation. *Nature* 2008;454,436-444.
5. El-Omar EM, Carrington M, Chow WH, et al. Interleukin-1 polymorphisms associated with increased risk of gastric cancer. *Nature* 2000;404,398-402.
6. Oshima M, Dinchuk JE, Kargman SL, et al. Suppression of intestinal polyposis in *Apc*<sup>Δ716</sup> knockout mice by inhibition of cyclooxygenase 2 (COX-2). *Cell* 1996;87:803-809.
7. Gupta RA, DuBois RN. Colorectal cancer prevention and treatment by inhibition of cyclooxygenase-2. *Nat Rev Cancer* 2001;1:11:21.
8. Sonoshita M, Takaku K, Sasaki N, et al. Acceleration of intestinal polyposis through prostaglandin receptor EP2 in *Apc*<sup>Δ716</sup> knockout mice. *Nat Med* 2001;7:1048-1051.
9. Wang D, Wang H, Shi Q, et al. Prostaglandin E<sub>2</sub> promotes colorectal adenoma growth via transactivation of the nuclear peroxisome proliferators-activated receptor δ. *Cancer Cell* 2004;6,285-295.
10. Buchanan F, Gorden DL, Matta P, et al. Role of β-arrestin 1 in the metastatic progression of colorectal cancer. *Proc Natl Acad Sci USA* 2006;103,1492-1497.

11. Saukkonen K, Rintahaka J, Sivula A et al. Cyclooxygenase-2 and gastric carcinogenesis. *APMIS* 2003;111,915-925.
12. Sun WH, Yu Q, Shen H et al. Roles of *Helicobacter pylori* infection and cyclooxygenase-2 expression in gastric carcinogenesis. *World J Gastroenterology* 2004;10,2809-2813.
13. Oshima H, Oshima M, Inaba T et al. Hyperplastic gastric tumors induced by activated macrophages in COX-2/mPGES-1 transgenic mice. *EMBO J* 2004;23,1669-1678.
14. Qian BZ, Pollard JW. Macrophage diversity enhances tumor progression and metastasis. *Cell* 2010;141,39-51.
15. Oshima H, Matsunaga A, Fujimura T, et al. Carcinogenesis in mouse stomach by simultaneous activation of the Wnt signaling and prostaglandin E<sub>2</sub> pathway. *Gastroenterology* 2006;131,1086-1095.
16. Rakoff-Nahoum S, Paglino J, Eslami-Varzaneh F. et al. Recognition of commensal microflora by Toll-like receptors is required for intestinal homeostasis. *Cell* 2004;118,229-241.
17. Lee CW, Rickman B, Rogers AB, et al. Combination of sulindac and antimicrobial eradication of *Helicobacter pylori* prevents progression of gastric cancer in hypergastrinemic INS-GAS mice. *Cancer Res* 2009;69,8166-8174.
18. Takeuchi K, Tanaka A, Kato S. et al. Effect of (S)-4-(1-(5-Chloro-2-(4-fluorophenoxy)benzamido)ethyl) Benzoic Acid (CJ-42794), a selective antagonist of prostaglandin E receptor subtype 4, on ulcerogenic and healing responses in rat gastrointestinal mucosa. *J Pharmacol Exp Ther* 2007;322,903-912.
19. Kaparakis M, Walduck AK, Price JD, et al. Macrophages are mediators of gastritis in acute



- Helicobacter pylori* infection in C57BL/6 mice. *Infect Immun* 2008;76,2235-2239.
20. van Rooijen N, Sanders A. Liposome mediated depletion of macrophages: Mechanism of action, preparation of liposomes and applications. *J Immunol Meth* 1994;174,83-93.
  21. Oshima H, Itadani H, Kotani H, et al. Induction of prostaglandin E<sub>2</sub> pathway promotes gastric hamartoma development with suppression of bone morphogenetic protein signaling. *Cancer Res* 2009;69,2729-2733.
  22. Popivanova BK, Kostadinova FI, Furuichi K, et al. Blockade of a chemokine, CCL2, reduces chronic colitis-associated carcinogenesis in mice. *Cancer Res* 2009;69,7884-7892.
  23. Mantovani A, Sozzani S, Locati M, et al. Macrophage polarization: tumor-associated macrophages as a paradigm for polarized M2 mononuclear phagocytes. *TRENDS Immunol* 2002;23,549-555.
  24. DeNardo DG, Barreto JB, Andreu P, et al. CD4<sup>+</sup> T cell regulates pulmonary metastasis of mammary carcinomas by enhancing protumor properties of macrophages. *Cancer Cell* 2009;16,91-102.
  25. Oguma K, Oshima H, Aoki M, et al. Activated macrophages promote Wnt signaling through tumour necrosis factor- $\alpha$  in gastric tumour cells. *EMBO J* 2008;27,1671-1681.
  26. Greten FR, Eckmann, L, Greten TF, et al. IKK $\beta$  links inflammation and tumorigenesis in a mouse model of colitis-associated cancer. *Cell* 2004;118,285-296.
  27. Pull SL, Doherty JM, Mills JC, et al. Activated macrophages are an adaptive element of the colonic epithelial progenitor niche necessary for regenerative responses to injury. *Proc Natl Acad Sci USA* 2005;102,99-104.
  28. Roth KA, Kapadia SB, Martin SM, et al. Cellular immune responses are essential for the

development of *Helicobacter felis*-associated gastric pathology. J

Immunol 1999;163,1490-1497.

29. Oshima M, Oshima H, Matsunaga A, et al. Hyperplastic gastric tumors with spasmodic polypeptide-expressing metaplasia caused by tumor necrosis factor- $\alpha$ -dependent inflammation in cyclooxygenase-2/microsomal prostaglandin E synthase-1 transgenic mice. Cancer Res 2005;65,9147-9151.
30. Li Q, Ishikawa TO, Oshima M, et al. The threshold level of adenomatous polyposis coli protein for mouse intestinal tumorigenesis. Cancer Res 2005;65,8622-8627.
31. Fodde R, Brabletz T. Wnt/ $\beta$ -catenin signaling in cancer stemness and malignant behavior. Curr Opin Cell Biol 2007;19,150-158.
32. Meyer F, Ramanujam KS, Gobert AP, et al. Cutting Edge: Cyclooxygenase-2 activation suppresses Th1 polarization in response to *Helicobacter pylori*. J Immunol 2003;171,3913-3917.

## Figure Legends

**Figure 1.** Stimulation of normal gastric mucosa by indigenous bacteria. (A) mRNA levels of cytokines and chemokines in the gastric mucosa of the germfree (GF)-wild-type mice (mean Log<sub>2</sub> ratio to the SPF mouse level). Asterisks,  $P < 0.05$ . (B) Immunostaining for proton pumps (parietal cells) in the gastric mucosa of the indicated strains. Scale bars, 100  $\mu\text{m}$ . (C) The parietal cell index relative to the wild-type mouse level (mean  $\pm$  s.d.). Asterisks,  $P < 0.05$  versus the wild-type level (*WT*). *Gan*-NT: *Gan* non-tumor, *Gan*-DT: *Gan* dysplastic tumor. (D) Intra-gastric pH of the indicated mouse strains (mean  $\pm$  s.d.). Asterisks,  $P < 0.05$  versus the wild-type level (*WT*).

**Figure 2.** Suppression of gastric tumorigenesis in germfree (*GF*) *Gan* mice. (A) The gastric mucosal thickness (tumor height) of SPF-*Gan* (*SPF*), *GF*-*Gan* (*GF*), Commensal flora-reconstituted *GF*-*Gan* (*GF*->*SPF*), *H. felis*-infected *GF*-*Gan* (*GF*->*H. felis*), and EP4 inhibitor-treated-*Gan* (*EP4i*) mice at 30 and 55 weeks of age relative to wild-type mice (*WT*). (B) Representative macroscopic photographs of the stomach of the indicated group of *Gan* mice at 55 weeks of age. Closed arrowheads indicate tumors, whereas the white arrowheads indicate suppressed tumorous lesions in the *GF*-*Gan* mouse. (C) The survival rate of the *SPF*-*Gan* and *GF*-*Gan* mice. All *GF*-*Gan* mice were used for experiments at 55 weeks of age. (D) Apoptosis analyses of *SPF*-*Gan* and *GF*-*Gan* mouse gastric tumors. Arrows indicate apoptotic cells on the mucosal surface of tumors. (E) Ki-67 immunostaining in the gastric tumors of *SPF*-*Gan* and *GF*-*Gan* mice. Scale bars in (D) and (E), 100  $\mu\text{m}$ . (F) Relative mean Ki-67 labeling index (mean  $\pm$  s.d.). Asterisks,  $P < 0.05$  versus *SPF* level.

**Figure 3.** Macrophage recruitment in the *Gan* mouse tumors. (A) H&E staining of the SPF-*Gan*, GF-*Gan*, *H. felis*-infected GF-*Gan* (GF->*H. felis*), and EP4 inhibitor-treated *Gan* (EP4i-*Gan*) mouse stomachs. Arrows indicate lymphocyte infiltration. Scale bars, 100  $\mu$ m. (B) Relative mRNA levels of inflammatory cytokines, chemokines and COX-2 in wild-type mouse stomach (*WT*), and SPF-*Gan* (*SPF*), GF-*Gan* (*GF*), *H. felis*-infected GF-*Gan* (*HF*), and EP4i-*Gan* (*EP4i*) mouse gastric tumors (mean  $\pm$  s.d.). Asterisks,  $P < 0.05$  versus wild-type level. (C) The mRNA level of inflammatory cytokines in the microdissected tumor epithelial cells (*Ep*) relative to the level in the tumor stroma (*St*). Asterisks,  $P < 0.05$ . (D) Immunostaining of F4/80 (*green*) with DAPI staining (*blue*) in gastric tumors of the indicated groups. Arrowheads indicate F4/80-positive macrophages. Scale bars, 100  $\mu$ m. (E) The mean number of F4/80-positive macrophages per microscopic field (mean  $\pm$  s.d.). Asterisks,  $P < 0.05$  versus SPF level. (F) Immunostaining of F4/80 (*red*) and  $\beta$ -catenin (*green*) in gastric tumors of a control *Gan* mouse (*left*) and clodronate liposome-treated *Gan* mouse (*center*). H&E staining of serial section of a clodronate liposome-treated *Gan* mouse tumor (*right*). Scale bars, 100  $\mu$ m. Arrowheads (*left*) indicate macrophages. Arrowheads and arrows in (*center* and *right*) indicate the mucosal surface of the tumor and tumor cells, respectively, in the macrophage-depleted area.

**Figure 4.** Chemokine induction by bacterial colonization and PGE<sub>2</sub> signaling. (A, B) The mRNA levels of the indicated chemokines in SPF-*K19-C2mE* (*red*) and GF-*K19-C2mE* (*blue*) gastric mucosa (A) and SPF-*Gan* (pink) and GF-*Gan* (purple) gastric tumors (B) relative to

that in the control SPF-wild-type mouse stomach (*gray*) (mean  $\pm$  s.d.). The indigenous bacterial colonization (*B.I.*) and PGE<sub>2</sub> transgenic status (*PGE<sub>2</sub>*) are indicated at the bottom. Asterisks,  $P < 0.05$  versus wild-type level. (C) Immunostaining of CCL2 (*red*, arrowheads) and  $\beta$ -catenin (*green*) with DAPI staining (*blue*) in SPF-*Gan* gastric tumor. Scale bar, 100 $\mu$ m. (D) The mRNA levels of CCL2 in the LPS-stimulated RAW264 cells relative to that in the unstimulated control cells (mean  $\pm$  s.d.). Asterisks,  $P < 0.05$  versus control level. (E) The mRNA levels of CCL2 in RAW264 cells with indicated treatment relative to the level in the unstimulated RAW264 cells (mean  $\pm$  s.d.). Asterisks,  $P < 0.05$ . (F) Immunostaining for F4/80 (*red*, arrowheads) and  $\beta$ -catenin (*green*) with DAPI staining (*blue*) in tumors of control SPF-*Gan* (*left*) and CCL2 antibody-treated SPF-*Gan* mice (*right*). Scale bars, 100 $\mu$ m. Arrows in (*right*) indicate regressed tumors in the macrophage-depleted area.

**Figure 5.** EP4 signaling on macrophages and epithelial cells. (A) Immunostaining of EP4 in the normal gastric mucosa of wild-type mouse and gastric tumors of SPF- and GF-*Gan* mice. Arrowheads and arrows indicate stromal EP4 expression in the tumor epithelia and tumor stromal cells, respectively. Scale bars, 100 $\mu$ m. (B) The mRNA levels of EP4 in gastric tumors of SPF-*Gan* and GF-*Gan* mice relative to the wild-type mouse level (mean  $\pm$  s.d.). Asterisks,  $P < 0.05$ . (C) The mRNA levels of inflammatory cytokines in the control or drug-treated RAW264 cells relative to the level of LPS-stimulated RAW264 cells (mean  $\pm$  s.d.). Asterisks,  $P < 0.05$  to the LPS-stimulated level. (D) Representative photographs of organoid structures formed by the primary cultured gastric epithelial cells in matrigel with EP4 inhibitor treatment (*top right*) and no-treatment control (*top left*). The mean number of organoids larger than 0.2

mm in diameter in the microscopic field on day 6 of culture (*bottom*) (mean  $\pm$  s.d.). Asterisks,  $P < 0.05$ .

**Figure 6.** Wnt promotion by bacterial infection and TNF- $\alpha$ . (A) Relative mRNA levels of Wnt-target genes, CD44 and EphB3, in the wild-type mouse stomach (*WT*), and SPF-*Gan* (*SPF*), GF-*Gan* (*GF*), *H. felis*-infected GF-*Gan* (*GF->HF*), and EP4 inhibitor-treated-*Gan* (*EP4i*) mouse gastric tumors (mean  $\pm$  s.d). Asterisks,  $P < 0.05$  versus wild-type level. (B) Immunostaining of CD44 (*top*) and EphB3 (*bottom*) in the wild-type mouse stomach (*left*) and SPF-*Gan* (*center*) and GF-*Gan* (*right*) mouse tumors. Arrowheads indicate immunostained epithelial cells in the wild-type (*WT*) and GF-*Gan* mice. Scale bars, 100 $\mu$ m. (C) Immunoblotting of active  $\beta$ -catenin in the SPF-*K19-Wnt1* and GF-*K19-Wnt1* mouse stomach (*top*), and SPF-*Gan* and GF-*Gan* mouse gastric tumors (*bottom*).  $\beta$ -Actin was used as an internal control. (D) The number of preneoplastic lesions developed in SPF-*K19-Wnt1* (*SPF*) and GF-*K19-Wnt1* (*GF*) mice. The mean numbers are indicated. (E) The mRNA levels of TNF- $\alpha$  and EphB3 in the gastric tumors of the anti-TNF- $\alpha$  neutralizing antibody-treated *Gan* ( $\alpha$ TNF- $\alpha$ ) relative to those of untreated control *Gan* mice (control) (mean  $\pm$  s.d.). Asterisks,  $P < 0.05$  versus control. (F) Immunoblotting of active  $\beta$ -catenin in control (cont) and anti-TNF- $\alpha$  antibody-treated ( $\alpha$ TNF- $\alpha$ ) *Gan* mouse tumors.