

Geochemical Characteristics of Mantle Peridotite Beneath a Mature Arc

著者	Ishimaru Satoko, Arai Shoji
journal or publication title	Promotion Environmental Research in Pan-Japan Sea Area -Young Researchers' Network- : Abottract
page range	114-115
year	2006-03-08
URL	http://hdl.handle.net/2297/6555

Geochemical Characteristics of Mantle Peridotite Beneath a Mature Arc

Satoko ISHIMARU^a and Shoji Arai^b

(a) Division of Environmental Science and Engineering, Graduate School of Natural Science and Technology, Kanazawa University, Kakuma, Kanazawa, 920-1192, JAPAN

(b) Graduate School of Natural Science and Technology, Kanazawa University, Kakuma, Kanazawa, 920-1192, JAPAN

Kamchatka arc is a mature arc, and the Pacific plate is subducting at relatively rapid rates (60-80 mm/yr) beneath the southern part of Kamchatka Peninsula from the Kuril-Kamchatka trench. There are three volcanic chains with the distance from 200 km, 320 km, and 400 km (Tatsumi *et al.* 1994). Avacha volcano is located at the southern Kamchatka Peninsula and on the frontal chain that forms the volcanic front, and the depth to the subducting slab is about 120 km there (Figure 1) (Gorbatov *et al.* 1997). Peridotite xenoliths from Avacha volcano are useful to understand the mantle materials, processes, and magma genesis beneath the mature arc.

The peridotite xenoliths from Avacha volcano, are mainly harzburgite with highly depleted character; high Fo (90-92) of olivine, high Cr# (0.5-0.7) of spinel and low modal clinopyroxene (< 1 vol.%). This harzburgite is the one of the most refractory peridotite xenoliths from arc setting (Figure 2; Arai *et al.*, submitted). Avacha peridotites are metasomatized by SiO₂-rich component in various extent and secondary orthopyroxenes replacing olivine are observed (Arai *et al.*, 2003). Geochemical characteristics of residual spinel lherzolite have been discussed, although those of highly depleted spinel harzburgite have not been described in detail. Trace-element compositions of





Figure 2 Modal amounts of olivine, orthopyroxene and clinopyroxene in spinel peridotite xenoliths from the Western Pacific. Note that the peridotite on average gradually changes from depleted harzburgite in the frontal side of arc (Avacha and Iraya) to fertile lherzolite in continent (Korea, Sikhote-Alin and China) through Japan arcs. Date source is Arai *et al.* (submitted) and references there in.




Figure 3 Relationships between the Fo content of olivine and Cr# of chromian spinel in Avacha xenoliths. OSMA, olivine-spinel-mantle array, is a residual trend of spinel peridotite (Arai, 1994). Shaded area represents abyssal peridotites (Arai, 1994). Bold-dashed line shows the area of the composition of coarse-grained peridotite from Iraya volcano, the Luzon Arc, Philippines (Arai & Kida, 2000).

bulk-rock peridotites are controlled by grain-boundary component, but are determined by composition of glasses, related with metasomatism, if any. The grain-boundary component is quite important to determine the geochemical characteristics of highly depleted spinel harzburgite without clinopyroxene and glass.

The Fo content of olivine is lower at a given Cr# of coexisting spinel in the Avacha peridotite than in ordinary peridotite (Figure 3). This characteristic is not related with the formation of secondary orthopyroxene at the expense of olivine, because there are no obvious relationships between the modal amounts of secondary orthopyroxene and the mineral and bulk-rock compositions. There is discrepancy between relatively low Fo content of olivine and other petrological features in the Avacha peridotite, for example, high Cr# of spinel and low abundance of HREE. The most refractory peridotite, which has high Cr# and low HREE, has relatively low Fo content of olivine and higher LREE. Then, the Avacha peridotites are residue after influx melting of upper mantle peridotite.

Metasomatism has occurred after influx melting, and crystallized secondary orthopyroxene at the expense of olivine. There are two types of secondary

orthopyroxene; one has complicated grain boundary, and the other is euhedral to subhedral and sometimes accompanying interstitial amphibole and silicic glass. The texture and chemistry of the secondary phases suggest SiO₂-oversaturated character of the metasomatic agent. The metasomatism by SiO₂-rich material has been reported from other arc, for example, Iraya volcano, Luzon arc (e.g., Schiano *et al.*, 1995, Arai & Kida, 2000), Papua New Guinea (Grégoire *et al.*, 2001) and Patagonia (Kilian & Stern, 2002). The silica enrichment is an important phenomenon in the sub-arc mantle.

References

- Arai, S. (1994) Characterization of spinel peridotites by olivine-spinel compositional relationships: review and interpretation. *Chemical Geology* **113**, 191-20
- Arai, S., Abe, N. & Ishimaru, S.. Mantle peridotites from the Western Pacific. *Gondwana Research*, submitted
- Arai, S., Ishimaru, S. & Okrugin, V. M. (2003) Metasomatized harzburgite xenoliths from Avacha volcano as fragments of mantle wedge of the Kamchatka arc: Implication for the metasomatic agent. *The Island Arc* **12**, 233-246.
- Arai, S. & Kida, M. (2000) Origin of fine-grained peridotite xenoliths from Iraya volcano of Batan Island, Philippines: deserpentinization or metasomatism at the wedge mantle beneath an incipient arc? *The Island Arc* **9**, 458-471.
- Gorbatov, A., Kostoglodov, V., Suárez, G. & Gordeev, E. (1997). Seismicity and structure of the Kamchatka subduction zone, *Journal of Geophysical Research* **102**, 17883-17898
- Grégoire, M., McInnes, B. I. A. & O'Reilly S. Y. (2001). Hydrous metasomatism of oceanic sub-arc mantle, Lihir, Papua New Guinea Part 2. Trace element characteristics of slab-derived fluids. *Lithos* **59**, 91-108.
- Kilian, R., Stern, C. R. (2002). Constraints on the interaction between slab melts and the mantle wedge from adakitic glass in peridotite xenoliths. *European Journal of Mineralogy* **14**, 25-36.
- Schiano, P., Clocchiatti, R., Shimizu, N., Maury, R. C., Jochum, K. P. & Hofmann, A. W. (1995). Hydrous, silica-rich melts in the sub-arc mantle and their relationship with erupted arc lavas. *Nature* **377**, 595-600
- Tatsumi, Y., Furukawa, Y., Kogiso T., Yamanaka, Y., Yokoyama, T. & Fedetov, S. A. (1994) A third volcanic chain in Kamchatka: thermal anomaly at transform/convergence plate boundary, *Geophysical Research Letters* **21**, 7, 537-540

^a Electronic Address: jaja@earth.s.kanazawa-u.ac.jp

^b Electronic Address: ultrasa@kenroku.kanazawa-u.ac.jp, URL: http://earth.s.kanazawa-u.ac.jp/Arai_Lab/