

Reply to: Over time, differences in survival and favorable neurologic outcomes between conventional and compression-only cardiopulmonary resuscitation have been gradually reduced in pediatric out-of-hospital cardiac arrest

著者	Goto Yoshikazu, Funada Akira, Goto Yumiko
著者別表示	後藤 由和, 舟田 晃
journal or publication title	Resuscitation
volume	124
page range	e9-e10
year	2018-03-01
URL	http://doi.org/10.24517/00050464

doi: 10.1016/j.resuscitation.2017.12.023



**Reply to: Over Time, Differences in Survival and Favorable Neurologic Outcomes
between Conventional and Compression-Only Cardiopulmonary Resuscitation Have
Been Gradually Reduced in Pediatric Out-of-Hospital Cardiac Arrest**

Yoshikazu Goto, MD, PhD¹, Akira Funada, MD, PhD¹, Yumiko Goto, MD, PhD²

¹Department of Emergency and Critical Care Medicine, Kanazawa University Hospital,
Kanazawa, Japan

²Department of Cardiology, Yawata Medical Centre, Komatsu, Japan

Corresponding Author:

Yoshikazu Goto, MD, PhD

Kanazawa University Hospital, Department of Emergency and Critical Care Medicine,
Takaramachi 13-1, Kanazawa 920-8640, Japan

Tel: +81-76-265-2020

Fax: +81-76-234-4287

E-mail: gotoyosh@med.kanazawa-u.ac.jp

Word count: 491

We thank Dr. Fukuda for his comments regarding our recent publication¹. Recently, the difference in 30-day neurologically intact survival rate between bystander conventional cardiopulmonary resuscitation (CPR) and bystander chest-compression-only CPR (CC-CPR) has been decreasing¹. In matched patients aged 1-17 years (n=2682, 67.6% of patients before propensity score matching) in our study¹, the 30-day neurological intact survival rates in overall patients receiving conventional CPR were equivalent to those in patients receiving CC-CPR in 2011-2012 (9.8% vs. 6.5%, $P=0.15$) and 2013-2014 (11.2% vs. 8.8%, $P=0.38$). Table 1 shows the results of the secondary analyses of 30-day outcomes after out-of-hospital cardiac arrest (OHCA) in the matched patients aged 1-17 years from 2011 to 2014 (n=1148), when we use same matched dataset in the manuscript¹. Risk difference, risk ratio, and odds ratio of overall patients receiving conventional CPR for 30-day neurologically intact survival was 0.03 (95% confidence interval [CI], -0.003 to 0.062), 1.40 (95% CI, 0.96 to 2.03), and 1.44 (95% CI, 0.96 to 1.04), respectively, compared to overall patients receiving CC-CPR. In subgroup analyses, only the cohort of patients aged 1-7 years had a significant difference in 30-day neurologically intact survival between two bystander CPR cohorts ($P=0.04$) with risk difference of 0.04 (95% CI, 0.002 to 0.075), risk ratio of 2.03 (95% CI, 1.01 to 4.09), and odds ratio of 2.11 (95% CI, 1.01 to 4.42). Moreover, the difference in 30-day survival rate between two bystander CPR cohorts is crucial in overall patients and in the three subgroups. These results show that conventional CPR is superior to CC-CPR in 30-day survival even in the recent years.

However, we must pay attention in interpreting these results. The number of matched patients for analyses is considered inappropriate for secondary analyses. The number of matched patients who received CC-CPR (n=574) accounted for only 39.7% of overall patients who received CC-CPR during the study period (n=1447, 2011-2014), eliminating 873 patients (60.3% of overall patients), while the number of patients who received conventional CPR (n=574) accounted for 98.1% of overall patients who received conventional CPR (n=585, 2011-2014). To focus on patients who were treated in recent years (from 2011 onward), we must analyse another matched patient cohort using not only propensity score matching method but also stratified analysis, inverse probability weighting methods, or doubly

robust estimator method². Accordingly, in our manuscript¹, we did not include any further data analyses for patients treated in recent years. The glass half-empty analysis may have introduced significant bias to the results³.

Considering the aforementioned circumstances, our study¹ strongly supports the 2017 International Liaison Committee on Resuscitation summary statement⁴ and the European Resuscitation Council 2017 guidelines update⁵: we suggest that bystanders provide CPR with ventilation for infants and children <18 years of age with OHCA; if bystanders cannot provide rescue breaths as part of CPR, they should at least provide chest compressions.

Funding Sources

This work was supported by the Japan Society for the Promotion of Science (KAKENHI Grant No.15K08543).

Conflicts of Interest

None

References:

1. Goto Y, Funada A, Goto Y. Conventional versus chest-compression-only cardiopulmonary resuscitation by bystanders for children with out-of-hospital cardiac arrest. *Resuscitation* 2017, <https://dx.doi.org/10.1016/j.resuscitation.2017.10.015>.
2. Bang H, Robins JM. Doubly robust estimation in missing data and causal inference models, *Biometrics* 2005;61:962-972.
3. Maconochie I, de Caen A. When should ADULT CPR be delivered to children? *Resuscitation* 2017. <https://dx.doi.org/10.1016/j.resuscitation.2017.11.050>.
4. Olasveengen TM, de Caen AR, Mancini ME, Maconochie IK, Aickin R, Atkins DL, et al. 2017 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations Summary. *Resuscitation* 2017;121:201-214.
5. Perkins GD, Olasveengen TM, Maconochie I, Soar J, Wyllie J, Greif R, et al. ERC 2017 Guidelines Update. *Resuscitation* 2017. <https://doi.org/10.1016/j.resuscitation.2017.12.007>.

Table 1. Comparison of 30-day Outcomes in Matched Patients Aged 1-17 Years (N=1148).

		Conventional CPR		Compression-only CPR		P-value
		N	% (95% CI)	N	% (95% CI)	
30-day survival						
Overall (n=1148)		109/574	19.0 (16.0-22.4)	81/574	14.1 (11.5-17.2)	0.03
Subgroup						
Aetiology	Cardiac, n=405 (35.3%)	48/200	24.0 (18.6-30.4)	42/205	20.5 (15.5-26.5)	0.40
	Non-cardiac, n=743 (64.7%)	61/374	16.3 (12.9-20.4)	39/369	10.6 (7.8-14.1)	0.02
Initial rhythm	Shockable, n=116 (10.1%)	29/57	50.9 (38.3-63.4)	25/59	42.4 (30.6-55.1)	0.36
	Non-shockable, n=1032 (89.9%)	80/517	15.5 (12.6-18.8)	56/515	10.9 (8.5-13.9)	0.03
Witnessed status	Witnessed, n=415 (36.1%)	64/210	30.5 (24.6-37.0)	46/205	22.4 (17.3-28.6)	0.06
	Unwitnessed, n=733 (63.9%)	45/364	12.4 (9.4-16.1)	35/369	9.5 (6.9-12.9)	0.21
Age	1-7 years, n=599 (52.2%)	54/304	17.8 (13.9-22.5)	30/295	10.2 (7.2-14.1)	0.008
	8-17 years, n=549 (47.8%)	55/270	20.4 (16.0-25.6)	51/279	18.3 (14.2-23.2)	0.53
30-day CPC 1-2						
Overall (n=1148)		60/574	10.5 (8.2-13.2)	43/574	7.5 (5.6-9.9)	0.08
Subgroup						
Aetiology	Cardiac, n=405 (35.3%)	34/200	17.0 (12.4-22.8)	28/205	13.7 (9.6-19.0)	0.35
	Non-cardiac, n=743 (64.7%)	26/374	6.9 (4.8-10.0)	15/369	4.1 (2.5-6.6)	0.08
Initial rhythm	Shockable, n=116 (10.1%)	26/57	45.6 (33.4-58.4)	20/59	33.9 (23.1-46.6)	0.20
	Non-shockable, n=1032 (89.9%)	34/517	6.6 (4.7-9.0)	23/515	4.5 (3.0-6.6)	0.14
Witnessed status	Witnessed, n=415 (36.1%)	44/210	20.9 (16.0-27.0)	29/205	14.2 (10.0-19.6)	0.07
	Unwitnessed, n=733 (63.9%)	16/364	4.4 (2.7-7.0)	14/369	3.8 (2.3-6.3)	0.68
Age	1-7 years, n=599 (52.2%)	23/304	7.6 (5.1-11.1)	11/295	3.7 (2.1-6.6)	0.04

8-17 years, n=549 (47.8%)

37/270 13.7 (10.1-18.3)

32/279 11.5 (8.2-15.7)

0.43

CI, confidence interval; CPC, cerebral performance category; CPR, cardiopulmonary resuscitation.