

# The sex-specific effect of dioxin exposure on the growth of children: A Vietnamese cohort study

Oanh Thi Phuong Nguyen, Teruhiko Kido<sup>1)</sup>

## Abstract

The effects of dioxins on children's growth have been investigated extensively. This follow-up study was conducted in 89 mother-child pairs in Vietnam (43 pairs from a dioxin hotspot and 46 pairs from an unsprayed area) to assess the effects of lactational dioxin exposure on postnatal growth. Maternal breast milk was collected in 2008 from women who had given birth 4–16 weeks previously. The children's growth parameters were measured at 3, 5, and 7 years of age. Dioxin levels in breast milk were determined by gas chromatography/high-resolution mass spectrometry. Toxic equivalency values of polychlorinated dibenzodioxins (PCDDs), polychlorinated dibenzofurans (PCDFs), and PCDDs + PCDFs in maternal breast milk were 2–4 times higher in women residing in the hotspot. Among the children from the dioxin hotspot, lower weight was observed for girls at 3 years of age, reduced body mass index (BMI) and head and chest circumference were observed in girls at 5 years of age, and reduced height and head and chest circumference were observed in girls at 7 years of age. There were no significant differences in body parameters in boys between the dioxin hotspot area and those from the unsprayed area. In girls, maternal breast milk dioxin levels were negatively correlated with BMI and head and chest circumference at 5 years of age, and with height and head circumference at 7 years of age. Multiple regression analysis indicated that BMI at 5 years of age and chest circumference at 5 and 7 years of age showed stronger inverse associations with maternal age than with dioxin exposure. In summary, we found sex-specific effects on postnatal growth in Vietnamese children exposed to dioxin.

## KEY WORDS

Dioxin, breast milk, mother-child pairs, growth retardation, sex-specific

## Introduction

Dioxin levels are still elevated in the environment, food, and body fluids of humans residing in contaminated areas in Southern Vietnam, nearly five decades after dioxin-contaminated herbicides were sprayed during the Vietnam War<sup>1-4)</sup>. The contaminated areas, such as Bien Hoa, Phu Cat, and Da Nang, are termed dioxin hotspots, and long-term exposure may cause adverse health effects in residents of these areas<sup>5)</sup>. Various epidemiological studies have found that dioxin levels in breast milk were related to steroid hormone disruptions in mothers and children in these hotspots, and may decrease expressive

communication scores and increase autistic traits and developmental coordination disorders in infants<sup>6-10)</sup>. High serum dioxin levels have also been related to increases in prostate cancer rates and hormone disruption in men<sup>3,11)</sup>.

Because it is highly lipophilic, dioxin can cross the placental barrier to fetus during pregnancy and be transmitted to infant through breastmilk<sup>12)</sup>. The cumulative dioxin dose of exclusively breastfed infants has been reported to be 3.6 and 4.6 times higher than the dose in formula-fed infants at 6 months and 1 year, respectively, and 10–18 times higher than levels at birth<sup>13)</sup>. Breastfeeding for 6 months can contribute

---

Division of Health Science, Graduate School of Medical Science, Kanazawa University

1) Faculty of Health Sciences, Institute of Medical, Pharmaceutical and Health Sciences, Kanazawa University

12% and 14% to the body burdens of dioxin in men and women, respectively, measured at the age of 25<sup>14</sup>). Thus, breast milk is a high-risk source of dioxin exposure in infants. Moreover, dioxin levels in breast milk are related to their levels in the placenta, maternal serum, and umbilical cord blood, suggesting that dioxin levels in breast milk could be used as a perinatal exposure marker in breastfed infants<sup>15</sup>).

Perinatal exposure to dioxin can cause growth retardation in infants and children<sup>16</sup>). Most published epidemiological studies have focused on perinatal exposure and its effects on infancy growth. High dioxin levels in maternal serum, breast milk, or umbilical cord blood have been correlated with reduced birth weight, head circumference, and weight for gestational age, shorter gestational age, and lower growth rate, weight, and body mass index (BMI) in infants<sup>17-21</sup>). Few studies, however, have evaluated the effect of dioxin exposure on older children.

Our previous studies of mother-infant pairs, conducted in a dioxin hotspot (Phu Cat) and a control (non-sprayed) area, found significant differences in the levels of dioxin congeners and hormones in mothers residing in the two areas<sup>9, 22</sup>). Furthermore, a high percentage of low birth weight was observed in the hotspot; birth weight was negatively correlated with toxic equivalency (TEQ) values for 2,3,7,8-tetrachlorodibenzo-p-dioxin and 2,3,4,7,8-pentachlorodibenzofuran<sup>22</sup>). To investigate whether early-life exposure to dioxin continues to affect childhood growth beyond infancy, we followed up with participants from that study and measured children's body parameters at 3, 5, and 7 years.

## Materials and methods

### Study areas

The hotspot area was the Phu Cat district of Binh Dinh province, the site of a US Army airbase in which dioxin-contaminated herbicides were stored and used during the Vietnam War (1965–1971)<sup>5</sup>). The non-sprayed area was the Kim Bang district of Ha Nam province, which was not exposed to dioxin during the war. Both areas are agricultural and have not been contaminated by industrial pollution.

### Study participants and measurements

Women (60 from Phu Cat and 63 from Kim Bang) who were breastfeeding their infants (from 4 to 16 weeks) and had resided in the study areas for more

than 5 years were invited to participate in 2008. Each woman donated 10–20 mL of breast milk and was measured for anthropometric indicators (height, weight) and interviewed regarding age, family income, length of residence, occupation, educational level, disease status, smoking habit, or alcohol consumption.

Body parameters in the children (height, weight, and head and chest circumference) were measured by medical staff at the ages of 3, 5, and 7 in 2011, 2013, and 2015, respectively. BMI was calculated from height and weight. The final number of participating mother-child pairs was 89 (43 from Phu Cat and 46 from Kim Bang).

The Medical Ethics Committee of Kanazawa University approved this study (No. 455). The mothers or fathers gave consent for medical examination of the children.

### Analysis of dioxin in breast milk

Polychlorinated dibenzodioxins (PCDDs) and polychlorinated dibenzofurans (PCDFs) levels in breast milk were determined by gas chromatography/high-resolution mass spectrometry (JMS-700 MS station mass spectrometer, JEOL Ltd., Tokyo, Japan) in accordance with a previously reported method<sup>23</sup>). Dioxin detection limits were determined at a signal-to-noise ratio of 3 on a lipid. Congener levels below the limit of detection were assigned a value equal to half the limit of detection. The determined values are shown as picograms per gram of lipid and were converted to TEQs using the World Health Organization toxic equivalency factor<sup>24</sup>).

### Statistical analysis

Data were analyzed using the JMP 12 software package (SAS Institute, Cary, NC, USA) and Microsoft Excel 2010 (Microsoft Corp., Redmond, WA, USA). The significance level was set at  $p \leq 0.05$ .

Dioxin levels were  $\log_{10}$  transformed to improve normality. Data are shown as mean and standard deviation or median (interquartile range) for original data and as geometric mean with geometric standard deviation for  $\log_{10}$ -transformed data. The mean difference in each indicator was calculated using Student's *t*-test or the Wilcoxon signed rank test for normal or non-normal distribution, as determined by the Shapiro-Wilk test. Pearson's correlation coefficient and Spearman's correlation coefficient were also calculated depending on distribution. To determine the influence of confounding

factors, we used multiple regression analysis with body parameters at 3, 5, and 7 years as dependent variables. Maternal age, BMI, and educational level, as well as the length of the breastfeeding period and TEQ-dioxin congener levels in breast milk were selected as independent variables.

## Results

### Demographic characteristics

Table 1 displays the demographic characteristics of the mothers and children, stratified by sex. In girls, TEQ of PCDDs, PCDFs, and PCDDs + PCDFs in the hotspot were more than twice the values in girls residing in the non-sprayed area. In boys, TEQ of PCDDs, PCDFs, and PCDDs + PCDFs in the hotspot were nearly four

times higher than the values in boys residing in the non-sprayed area. There was no significant difference in the lengths of breastfeeding periods or maternal age, BMI, educational level, smoking status, or alcohol consumption between the two groups.

### Comparison of growth parameters in children

Table 2 displays the growth parameters measured in children and stratified by sex. Weight at 3 years was significantly lower in girls from the hotspot area. BMI and head and chest circumference differed significantly in girls from the two areas at 5 years, and height and head and chest circumference differed in girls at 7 years. Growth parameters in boys did not differ significantly between the two areas.

Table 1. Demographic characteristic of mothers and children

Parameters	Girls			Boys		
	Hotspot	Non-sprayed	p-value	Hotspot	Non-sprayed	p-value
<b>Children</b>						
Subjects	n = 18 (41.9%)	n = 27 (58.7%)		n = 25 (58.1%)	n = 19 (41.3%)	
Full breastfeeding (months)	5.5 (4.0-6.0)	4.0 (3.0-6.0)	0.417 <sup>b</sup>	6.0 (4.0-6.0)	6.0 (3.0-6.0)	0.832 <sup>b</sup>
<b>Mothers</b>						
Age (years)	26.6 (4.1)	25.0 (2.8)	0.160 <sup>a</sup>	27.0 (3.4)	26.6 (2.3)	0.667 <sup>a</sup>
BMI	21.1 (1.9)	20.9 (2.0)	0.771 <sup>a</sup>	20.2 (19.0-21.7)	20.2 (19.3-22.0)	0.868 <sup>b</sup>
Education (%)						
Elementary (%)	n = 6 (33.3%)	n = 6 (22.2%)	0.702 <sup>c</sup>	n = 6 (24.0%)	n = 3 (15.8%)	0.660 <sup>c</sup>
Junior	n = 9 (50.0%)	n = 14 (51.9%)		n = 9 (36.0%)	n = 10 (52.6%)	
High school	n = 3 (16.7%)	n = 6 (22.2%)		n = 7 (28.0%)	n = 5 (26.3%)	
University	n = 0 (0%)	n = 1 (3.7%)		n = 3 (12.0%)	n = 1 (5.3%)	
Alcohol habit (% no)	n = 16 (88.9%)	n = 27 (100%)	0.154 <sup>d</sup>	n = 24 (96.0%)	n = 19 (100%)	1.000 <sup>d</sup>
Smoking habit (% no)	n = 18 (100%)	n = 25 (92.6%)	1.000 <sup>d</sup>	n = 25 (100%)	n = 19 (100%)	
TEQ PCDDs (pg/g lipid)	5.1 (1.3)	2.0 (1.6)	< 0.001 <sup>b</sup>	7.3 (1.4)	1.8 (1.4)	< 0.001 <sup>a</sup>
TEQ PCDFs (pg/g lipid)	3.2 (1.4)	1.4 (1.5)	< 0.001 <sup>a</sup>	5.1 (1.4)	1.3 (1.3)	< 0.001 <sup>a</sup>
TEQ PCDDs+PCDFs (pg/g lipid)	8.4 (1.3)	3.5 (1.5)	< 0.001 <sup>a</sup>	12.5 (1.4)	3.2 (1.5)	< 0.001 <sup>a</sup>

Data are shown as geometric mean ± geometric standard deviation for TEQ of dioxin levels and mean ± standard deviation or median (interquartile range) for other variables.

<sup>a</sup> t-test, <sup>b</sup>Wilcoxon signed rank's test, <sup>c</sup> Chi-square test, <sup>d</sup> Fisher exact test

Table 2. Comparison of growth parameters in children

Parameters	Girls			Boys		
	Hotspot (n = 18)	Non-sprayed (n = 27)	p-value	Hotspot (n = 25)	Non-sprayed (n = 19)	p-value
<b>3 years</b>						
Height (cm)	90.0 (3.1)	90.0 (2.2)	0.296*	93.3 (3.4)	92.1 (2.5)	0.203*
Weight (kg)	11.9 (1.2)	12.7 (1.0)	0.042*	13.1 (12.0-14.6)	13.0 (11.5-14.0)	0.403
BMI	14.8 (1.5)	15.4 (1.0)	0.158*	15.0 (14.2-15.8)	14.7 (14.2-15.8)	0.981
Head Cir. (cm)	48.0 (1.4)	48.5 (1.1)	0.226*	49.4 (1.6)	48.8 (1.0)	0.131*
Chest Cir.(cm)	48.0 (2.0)	48.8 (1.6)	0.156*	50.5 (2.6)	50.0 (2.0)	0.445*
<b>5 years</b>						
Height (cm)	104.9 (4.5)	105.9 (2.4)	0.379*	107.3 (4.4)	107.7 (3.2)	0.709*
Weight (kg)	15.7 (14.1-16.8)	16.2 (15.4-17.9)	0.056	16.9 (15.9-21.2)	17.4 (15.5-18.9)	0.619
BMI	14.1 (13.6-14.7)	14.8 (14.0-15.4)	0.046	15.4 (13.8-18.2)	14.6 (14.0-16.1)	0.441
Head Cir. (cm)	48.7 (48.5-49.7)	50.0 (40.9-50.9)	0.002	50.6 (1.8)	50.3 (1.6)	0.524*
Chest Cir.(cm)	50.3 (49.5-52.3)	53.9 (52.0-55.2)	<0.001	53.0 (51.0-58.3)	53.2 (50.2-54.4)	0.394
<b>7 years</b>						
Height (cm)	115.3 (4.2)	118.4 (3.2)	0.013*	120.5 (4.9)	120.6 (4.3)	0.940*
Weight (kg)	19.2 (17.2-21.5)	20.2 (19.3-23.7)	0.082	21.5 (19.7-27.6)	22.2 (19.9-23.7)	0.943
BMI	14.1 (13.2-16.5)	14.8 (14.0-16.0)	0.336	15.3 (13.8-17.7)	14.6 (13.7-16.6)	0.896
Head Cir. (cm)	50.5 (1.4)	51.4 (1.2)	0.037*	51.9 (1.9)	51.5 (1.5)	0.345*
Chest Cir.(cm)	55.0 (2.7)	57.3 (3.8)	0.002*	57.3 (54.9-64.3)	59.3 (56.3-60.8)	0.320

Data are shown as mean (standard deviation) or median (interquartile range). Cir. Circumference

\* t-test, others: Wilcoxon sign rank's test

**Correlations of dioxin levels in breast milk and body parameters in children**

Table 3 displays correlation data between TEQ values of dioxin in maternal breast milk and body parameters in children. In girls at 5 years, inverse correlations were found between BMI and head circumference and TEQ of PCDDs ( $r = -0.32, p = 0.031$  and  $r = -0.29, p = 0.050$ , respectively), and chest circumference and TEQ of PCDDs, PCDFs, and PCDDs + PCDFs ( $r = -0.38, p = 0.011$ ;  $r = -0.31, p = 0.039$ ;  $r = -0.35, p = 0.017$ , respectively). Similarly, negative correlations were observed between height and TEQ of PCDDs + PCDFs ( $r = -0.33, p = 0.027$ ) as well as between chest

circumference and TEQ of PCDFs ( $r = -0.30, p = 0.042$ ) in girls at 7 years. There was no significant correlation between body parameters and dioxin levels in girls aged 3 years.

In boys, no significant correlations were found between body parameters and dioxin levels, with the exception of height and TEQ of PCDF values at 3 years, chest circumference and TEQ of PCDD at 5 years, and head circumference and TEQ of PCDD at 7 years.

**Multiple regression analysis of dioxin congener levels in breast milk and body parameters in girls**

Table 4 displays data from multiple regression analysis of body parameters in girls as dependent variables and

Table 3. Correlations of dioxin levels in breastmilk and body parameters in children

Parameters	Girls						Boys					
	TEQ PCDDs		TEQ PCDFs		TEQ PCDDs+PCDFs		TEQ PCDDs		TEQ PCDFs		TEQ PCDDs+PCDFs	
	r	p	r	p	r	p	r	p	r	p	r	p
<b>3 years</b>												
Height	-0.06	0.714	-0.17	0.278*	-0.12	0.450*	0.30	0.051	0.30	0.046	0.280	0.069
Weight	-0.27	0.077	-0.12	0.449*	-0.20	0.196*	0.23	0.131	0.19	0.220	0.210	0.181
BMI	-0.29	0.053	-0.01	0.964*	-0.13	0.391*	0.13	0.412	0.05	0.759	0.100	0.504
Head Cir.	-0.06	0.682	0.05	0.721*	-0.01	0.959*	0.29	0.057	0.20	0.201	0.240	0.113
Chest Cir.	-0.17	0.278	-0.01	0.961*	-0.10	0.521*	0.17	0.262	0.10	0.537	0.140	0.353
<b>5 years</b>												
Height	-0.08	0.592	-0.01	0.925*	-0.03	0.824*	0.15	0.334	0.22	0.156	0.180	0.245
Weight	-0.24	0.118	-0.21	0.176	-0.22	0.146	0.23	0.125	0.16	0.296	0.200	0.192
BMI	-0.32	0.031	-0.27	0.072	-0.29	0.057	0.24	0.112	0.12	0.449	0.180	0.236
Head Cir.	-0.29	0.050	-0.24	0.106	-0.29	0.053	0.28	0.061	0.15	0.342	0.230	0.127
Chest Cir.	-0.38	0.011	-0.31	0.039*	-0.35	0.017*	0.30	0.046	0.22	0.151	0.260	0.086
<b>7 years</b>												
Height	-0.27	0.070	-0.29	0.058*	-0.33	0.027*	0.15	0.316	0.20	0.193	0.180	0.250
Weight	-0.25	0.099	-0.19	0.209	-0.20	0.181	0.18	0.252	0.15	0.328	0.170	0.276
BMI	-0.18	0.249	-0.17	0.277	-0.14	0.362	0.18	0.235	0.10	0.502	0.150	0.341
Head Cir.	-0.23	0.125	-0.12	0.448*	-0.20	0.197*	0.34	0.024	0.23	0.140	0.290	0.053
Chest Cir.	-0.24	0.120	-0.30	0.042	-0.27	0.070	0.05	0.767	0.00	0.990	0.030	0.835

p, p-value; r, correlation coefficient; Cir. Circumference  
 \*, Pearson correlation coefficient; others, Spearman correlation coefficient

Table 4. Multiple regression analysis of dioxin congener levels in breastmilk and body parameters in girls

Parameters	Height			Weight			BMI			Head Cir.			Chest Cir.		
	β	p	R <sup>2</sup>	β	p	R <sup>2</sup>	β	p	R <sup>2</sup>	β	p	R <sup>2</sup>	β	p	R <sup>2</sup>
<b>3 years</b>															
Mother Age	-0.03	0.869	0.06	-0.36	0.027	0.22	-0.39	0.020	0.19	-0.44	0.006	0.26	-0.48	0.003	0.20
TEQ PCDDs	-0.08	0.619		-0.24	0.122		-0.20	0.197		-0.03	0.831		-0.11	0.442	
Mother Age	-0.03	0.856	0.08	-0.38	0.024	0.18	-0.40	0.018	0.16	-0.45	0.006	0.27	-0.49	0.003	0.25
TEQ PCDFs	-0.16	0.316		-0.12	0.446		-0.01	0.953		0.07	0.626		0.00	0.978	
Mother Age	-0.03	0.874	0.07	-0.37	0.026	0.20	-0.39	0.020	0.17	-0.45	0.006	0.26	-0.48	0.003	0.26
TEQ PCDDs+PCDFs	-0.12	0.480		-0.19	0.218		-0.12	0.432		0.01	0.956		-0.06	0.681	
<b>5 years</b>															
Mother Age	-0.11	0.516	0.09	-0.32	0.050	0.21	-0.35	0.025	0.30	-0.51	0.001	0.31	-0.09	0.588	0.23
TEQ PCDDs	-0.02	0.904		-0.27	0.084		-0.35	0.019		-0.18	0.223		-0.33	0.036	
Mother Age	-0.11	0.509	0.09	-0.34	0.044	0.16	-0.37	0.023	0.23	-0.52	0.001	0.30	-0.10	0.514	0.20
TEQ PCDFs	0.01	0.968		-0.16	0.317		-0.21	0.157		-0.14	0.333		-0.28	0.076	
Mother Age	-0.11	0.513	0.09	-0.33	0.048	0.19	-0.36	0.025	0.26	-0.51	0.001	0.31	-0.09	0.567	0.22
TEQ PCDDs+PCDFs	-0.01	0.941		-0.23	0.148		-0.30	0.049		-0.17	0.247		-0.32	0.039	
<b>7 years</b>															
Mother Age	-0.09	0.581	0.14	-0.28	0.092	0.20	-0.26	0.132	0.12	-0.53	0.001	0.36	-0.42	0.007	0.31
TEQ PCDDs	-0.32	0.053		-0.31	0.049		-0.18	0.274		-0.23	0.101		-0.30	0.040	
Mother Age	-0.11	0.513	0.12	-0.30	0.082	0.12	-0.27	0.116	0.10	-0.55	0.001	0.32	-0.44	0.006	0.30
TEQ PCDFs	-0.27	0.092		-0.11	0.492		0.03	0.829		-0.10	0.478		-0.27	0.064	
Mother Age	-0.10	0.561	0.14	-0.28	0.089	0.16	-0.27	0.124	0.10	-0.54	0.001	0.35	-0.42	0.007	0.30
TEQ PCDDs+PCDFs	-0.31	0.056		-0.23	0.156		-0.08	0.619		-0.18	0.190		-0.29	0.049	

β, standardized coefficient; p, p-value; r<sup>2</sup>, coefficient of determination, Cir. Circumference  
 Dependent variables are height, weight, BMI, head and chest circumference at 3, 5 and 7 years old.  
 Independent variables are mother age, mother BMI, education, full breastfeeding period and TEQ of dioxin.

adjusted maternal dioxin levels as independent variables. Negative associations were observed between maternal age and children's weight and BMI at 3 and 5 years, as well as between head and chest circumference at all ages surveyed, except for chest circumference at 5 years. Furthermore, TEQ of PCDD and PCDDs + PCDFs in breast milk were inversely associated with BMI at 5 years ( $\beta = -0.35$ ,  $p = 0.019$  and  $\beta = -0.30$ ,  $p = 0.049$ , respectively) and chest circumference at 5 ( $\beta = -0.33$ ,  $p = 0.036$  and  $\beta = -0.32$ ,  $p = 0.039$ , respectively) and 7 years ( $\beta = -0.30$ ,  $p = 0.040$  and  $\beta = -0.29$ ,  $p = 0.049$ , respectively). No significant association was found between body parameters and other independent variables, such as maternal BMI, maternal educational level, or length of breastfeeding period (data not shown).

### Discussion

The findings of this study suggest that lactational exposure to PCDDs and PCDFs results in sex-specific postnatal growth effects. Differences in body parameters between children from the hotspot area and the non-sprayed area and correlations of these parameters with dioxin levels in maternal breast milk were found only in girls. In particular, a lower BMI at 5 years and chest circumference at 7 years were negatively correlated with breast milk levels of PCDDs and PCDDs + PCDFs, although no significant correlations between body parameters and breast milk dioxin levels were observed in children aged 3 years.

These results are partially consistent with those of epidemiological studies of the effects of perinatal exposure to dioxin or dioxin-like compounds on children's growth that found differences only in girls. Maternal polychlorinated biphenyl (PCB) levels have been correlated with reduced growth in Japanese girls whose mothers consumed contaminated oil and reduced weight in girls in an American cohort<sup>25, 26</sup>. One European study reported an association between maternal PCB levels and height and weight of children<sup>27</sup>. Lamb et al. reported that maternal levels of mono-ortho-substituted PCBs and ortho-substituted PCBs were associated with reduced weight in girls through 17 years of age<sup>28</sup>. In a cohort of German children, exposure to background levels of dichlorodiphenyldichloroethylene (DDE) was reported to correlate with an overall height reduction of 1.8 cm in girls, although the differences disappeared after the age of eight<sup>29</sup>. In contrast with our findings,

other studies reported a statistically significant BMI increase in 7-year-old girls perinatally exposed to dioxin and dioxin-like compounds<sup>27</sup>. Caucasian girls with high transplacental PCB exposures in an American cohort were reported to be 5.4 kg heavier for their height than other girls included in the analysis<sup>30</sup>. Prenatal DDE exposure was positively associated with waist circumference and the waist/height ratio in Belgian girls aged 7–9 years<sup>31</sup>.

Other studies have reported the effects of dioxin exposure on growth in 4-month-old Vietnamese boys (weight and BMI), in 8–12-year-old Russian boys (BMI and height), and in 4-year-old American children of both sexes (weight)<sup>18, 32, 33</sup>. Some animal studies reported that maternal PCB exposure resulted in growth retardation in offspring in guinea pigs and rats<sup>34, 35</sup>.

In the present study, girls appeared to be more sensitive to the effects of PCDDs and PCDFs. One potential reason is a reduction in thyroid hormones, responsible for regulating growth. A number of studies in humans have found that exposure to dioxin and PCBs was negatively correlated with levels of triiodothyronine and thyroxine in infants or with thyroid-stimulating hormone in females<sup>36–38</sup>. In juvenile fish, one PCB congener was shown to inhibit levels of triiodothyronine and thyroxine, leading to reduced total body weight and length and retarded metamorphic progress<sup>39</sup>. As mentioned above, our results have been consistent with some animal experiment and human studies about the effect of dioxin on growth in female<sup>25–29, 34, 35</sup>. However, the main reason for the sex-specific differences in effects is still unclear and we need more evidence in the further researches in the future.

The limitations of the study should be noted. First, PCB congener levels in maternal breast milk were not estimated, precluding analysis of perinatal PCB exposure and developmental delays in the study cohort. Second, the sample was small, and a larger sample may assist in elucidating the effects of dioxin on children's growth. Third, the study lacks data on postnatal exposure after breastfeeding. Dioxin exposure from contaminated sources could be contributed to the growth retardation of children in this study.

To our knowledge, this is the first longitudinal/cohort study to investigate the effects of lactational dioxin exposure on postnatal growth in Vietnamese children



living in a dioxin hotspot area. We found a negative association between TEQ of PCDDs and PCDFs and BMI at 5 years and head circumference at 7 years in girls. No significant effects were found in boys. Periodic follow-up is necessary for assessing growth retardation in children exposed to persistent organic pollutants.

## Reference

- 1) Hue N, Nam V, Thuong N, et al: Determination of PCDD/Fs in breast milk of women living in the vicinities of Da Nang Agent Orange hot spot (Vietnam) and estimation of the infant's daily intake. *Science of The Total Environment* 491: 212-218, 2014
- 2) Manh HD, Kido T, Tai PT, et al: Levels of polychlorinated dibenzodioxins and polychlorinated dibenzofurans in breast milk samples from three dioxin-contaminated hotspots of Vietnam. *The Science of the total environment* 511, 416-422, 2015
- 3) Schecter A, Quynh HT, Pavuk M, et al: Food as a source of dioxin exposure in the residents of Bien Hoa City, Vietnam. *Journal of Occupational and Environmental Medicine* 45: 781-788, 2003
- 4) Van Luong H, Nishijo M, Thao PN, et al: Association of dioxin exposure and reproductive hormone levels in men living near the Bien Hoa airbase, Vietnam. *Science of The Total Environment* 628: 484-489, 2018
- 5) Dwernychuk LW: Dioxin hot spots in Vietnam. *Chemosphere* 60: 998-999, 2005
- 6) Tran NN, Pham TT, Ozawa K, et al: Impacts of Perinatal Dioxin Exposure on Motor Coordination and Higher Cognitive Development in Vietnamese Preschool Children: A Five-Year Follow-Up. *PLoS ONE* 11: e0147655, 2016
- 7) Tai PT, Nishijo M, Nghi TN, et al: Effects of Perinatal Dioxin Exposure on Development of Children during the First 3 Years of Life. *The Journal of Pediatrics* 175: 159-166, 2016
- 8) Anh LT, Kido T, Honma S, et al: A relationship in adrenal androgen levels between mothers and their children from a dioxin-exposed region in Vietnam. *Science of the Total Environment* 607-608: 32-41, 2017
- 9) Kido T, Dao TV, Ho MD, et al: High cortisol and cortisone levels are associated with breast milk dioxin concentrations in Vietnamese women. *Eur J Endocrinol* 170: 131-139, 2014
- 10) Kido T, Honma S, Nhu DD, et al: Inverse association of highly chlorinated dioxin congeners in maternal breast milk with dehydroepiandrosterone levels in three-year-old Vietnamese children. *Sci Total Environ* 550: 248-55, 2016
- 11) Sun XL, Kido T, Honma S, et al: The relationship between dioxins exposure and risk of prostate cancer with steroid hormone and age in Vietnamese men. *Science of The Total Environment* 595: 842-848, 2017
- 12) Schecter A, Pöpke O, Ball M: Evidence for transplacental transfer of dioxins from mother to fetus: Chlorinated dioxin and dibenzofuran levels in the livers of stillborn infants. *Chemosphere* 21: 1017-1022, 1990
- 13) Chao H-R, Wang S-L, Lee C-C, et al: Level of polychlorinated dibenzo-p-dioxins, dibenzofurans and biphenyls (PCDD/Fs, PCBs) in human milk and the input to infant body burden. *Food and chemical toxicology* 42: 1299-1308, 2004
- 14) Patandin S, Dagnelie PC, Mulder P, et al: Dietary exposure to polychlorinated biphenyls and dioxins from infancy until adulthood: A comparison between breast-feeding, toddler, and long-term exposure. *Environmental Health Perspectives* 107: 45-51, 1999
- 15) Wang S-L, Lin C-Y, Guo YL, et al: Infant exposure to polychlorinated dibenzo-p-dioxins, dibenzofurans and biphenyls (PCDD/Fs, PCBs)—correlation between prenatal and postnatal exposure. *Chemosphere* 54: 1459-1473, 2004
- 16) Schecter A, Gasiewicz TA: *Dioxin and Health*. A John Wiley & Sons, INC, New Jersey pp 329-431, 2003
- 17) Tsukimori K, Uchi H, Tokunaga S, et al: Blood levels of PCDDs, PCDFs, and coplanar PCBs in Yusho mothers and their descendants: Association with fetal Yusho disease. *Chemosphere* pp 90: 1581-1588, 2013
- 18) Nishijo M, Nakagawa H, Maruzeni S, et al: Impact of perinatal dioxin exposure on infant growth: A cross-sectional and longitudinal studies in dioxin-contaminated areas in Vietnam. *PloS One* 7: e40273, 2012
- 19) Hertz-Picciotto I, Charles MJ, James RA, et al: In utero polychlorinated biphenyl exposures in relation to fetal and early childhood growth. *Epidemiology* 16: 648-656, 2005
- 20) Veiga-Lopez A, Kannan K, Liao C, et al: Gender-

## Acknowledgment

We would like to thank the participants as well as the organizations that supported the performance of this study, including 10-80 Division, Hanoi Medical University, and Kim Bang and Phu Cat medical centers in Vietnam. The authors declare that they have no conflict of interest.

- specific effects on gestational length and birth weight by early pregnancy BPA exposure. *The Journal of Clinical Endocrinology & Metabolism* 100: 1394-1403, 2015
- 21) Patandin S, Koopman-Esseboom C, De Ridder MA, et al: Effects of environmental exposure to polychlorinated biphenyls and dioxins on birth size and growth in Dutch children. *Pediatric Research* 44: 538-545, 1998
  - 22) Van Tung D, Kido T, Honma S, et al: Low birth weight of Vietnamese infants is related to their mother's dioxin and glucocorticoid levels. *Environ Sci Pollut Res Int* 23: 10922-10929, 2016
  - 23) The Tai P, Nishijo M, Kido T, et al: Dioxin concentrations in breast milk of Vietnamese nursing mothers: a survey four decades after the herbicide spraying. *Environmental science & technology* 45: 6625-6632, 2011
  - 24) Yamashita F, Hayashi M: Fetal PCB syndrome: clinical features, intrauterine growth retardation and possible alteration in calcium metabolism. *Environmental Health Perspectives* 59: 41-45, 1985
  - 25) Van den Berg M, Birnbaum LS, Denison M, et al: The 2005 World Health Organization reevaluation of human and mammalian toxic equivalency factors for dioxins and dioxin-like compounds. *Toxicological sciences* 93: 223-241, 2006
  - 26) Blanck HM, Marcus M, Rubin C, et al: Growth in girls exposed in utero and postnatally to polybrominated biphenyls and polychlorinated biphenyls. *Epidemiology* 13: 205-210, 2002
  - 27) Iszatt N, Stigum H, Govarts E, et al: Perinatal exposure to dioxins and dioxin-like compounds and infant growth and body mass index at seven years: A pooled analysis of three European birth cohorts. *Environment international* 94: 399-407, 2016
  - 28) Lamb MR, Taylor S, Liu X, et al: Prenatal exposure to polychlorinated biphenyls and postnatal growth: a structural analysis. *Environmental health perspectives* 114: 779-785, 2006
  - 29) Karmaus W, Asakevich S, Indurkha A, et al: Childhood growth and exposure to dichlorodiphenyl dichloroethene and polychlorinated biphenyls. *The Journal of pediatrics* 140: 33-39, 2002
  - 30) Jacobson JL, Jacobson SW, Humphrey HE: Effects of exposure to PCBs and related compounds on growth and activity in children. *Neurotoxicology and teratology* 12: 319-326, 1990
  - 31) Delvaux I, Van Cauwenberghe J, Den Hond E, et al: Prenatal exposure to environmental contaminants and body composition at age 7–9 years. *Environmental research* 132: 24-32, 2014
  - 32) Gladen BC, Ragan NB, Rogan WJ: Pubertal growth and development and prenatal and lactational exposure to polychlorinated biphenyls and dichlorodiphenyl dichloroethene. *The Journal of pediatrics* 136: 490-496, 2000
  - 33) Burns JS, Williams PL, Sergeev O, et al: Serum dioxins and polychlorinated biphenyls are associated with growth among Russian boys. *Pediatrics* 127: 59-68, 2011
  - 34) Lundkvist U: Clinical and reproductive effects of Clophen A50 (PCB) administered during gestation on pregnant guinea pigs and their offspring. *Toxicology* 61: 249-257, 1990
  - 35) Brezner E, Terkel J, Perry AS: The effect of Aroclor 1254 (PCB) on the physiology of reproduction in the female rat--I. *Comparative Biochemistry and Physiology Part C: Comparative Pharmacology* 77: 65-70, 1984
  - 36) Eguchi A, Nomiyama K, Minh Tue N, et al: Residue profiles of organohalogen compounds in human serum from e-waste recycling sites in North Vietnam: Association with thyroid hormone levels. *Environmental Research* 137: 440-449, 2015
  - 37) Nagayama J, Okamura K, Iida T, et al: Postnatal exposure to chlorinated dioxins and related chemicals on thyroid hormone status in Japanese breast-fed infants. *Chemosphere* 37: 1789-1793, 1998
  - 38) Koopman-Esseboom C, Morse DC, Weisglas-Kuperus N, et al: Effects of dioxins and polychlorinated biphenyls on thyroid hormone status of pregnant women and their infants. *Pediatric research* 36: 468-473, 1994
  - 39) Dong Y, Zhang X, Tian H, et al: Effects of polychlorinated biphenyls on metamorphosis of a marine fish Japanese flounder (*Paralichthys olivaceus*) in relation to thyroid disruption. *Marine pollution bulletin* 119: 325-331, 2017

## ダイオキシン暴露の子供の成長に関する性特異的な影響：ベトナムでのコホート研究

ニュエン・ティ・フオン・オアン，城戸照彦<sup>1)</sup>

### 要 旨

ダイオキシン類は残留性有機汚染物質であり、子供の成長に影響することが報告されている。本追跡研究は授乳中のダイオキシン暴露が生後の子どもの発達に影響するかを明らかにするため、ベトナムの枯葉剤撒布地区 (Phu Cat) と非撒布地区 (Kim Bang) で実施された。母乳は2008年に生後4-16週の児を持つ母親より収集した。その子供の身体計測が3才、5才、7才の時に実施された。本研究では、すべての調査に参加した89組の母子 (Phu Cat 43組、Kim Bang 46組) を対象とした。母乳中のダイオキシンの測定にはGC-HRMSを用いた。母乳中の polychlorinated dibenzo-dioxins (PCDDs)、-furans (PCDFs) と PCDDs/Fs の毒性当量は男女とも枯葉剤撒布地区の方が非撒布地区より2-4倍有意に高かった。3才児の体重、5才児のBMI、頭囲と胸囲、7才児の身長、頭囲と胸囲は女兒でのみ枯葉剤撒布地区の方が非撒布地区より有意に小さかった。男児では両地区間で身体計測値に有意差は無かった。ダイオキシンと5才児のBMI、頭囲と胸囲の間に、さらに7才児の頭囲と胸囲との間に有意な負の相関が女兒にのみ見られた。重回帰分析では5才児のBMIと胸囲、7才児の胸囲はダイオキシンよりも母親の年齢と強い負の相関を示した。結論として、ダイオキシンはベトナムの子供の生後の発達遅延に、性により異なる影響を与えていることが判明した。