

J. Math. Tokushima Univ.
Vol. 32 (1998), 15-18

On Some Numerical Relations of tetragonal Linear Systems

By

Akira Ohbuchi

*Department of Mathematical Sciences,
Faculty of Integrated Arts and Sciences,
The University of Tokushima,
1-1, Minamijosanjima-cho, Tokushima 770, JAPAN
email address: ohbuchi@ias.tokushima-u.ac.jp
(Received September 14, 1998)*

Abstract

Let \mathcal{L} be a pencil of degree 4 on a curve C and let e_1, e_2, e_3 be scollar invariants. We prove that $e_1 \leq e_2 + e_3 + 2$ if and only if e_1, e_2, e_3 are scollar invariants of some tetragonal curve.

1991 Mathematics Subject Classification. Primary 14H45

Introduction

Let C be a complete non-singular curve defined over an algebraically closed field k with $\text{char}(k) \neq 2$. Assume that C is a tetragonal curve. We assume that C is non-hyperelliptic of genus g .

Let $F_i = \Gamma(C, \omega \otimes \mathcal{O}(-ig_4^1))$. The modules F_i ($i = 1, 2, \dots$) give a filtration,

$$F_0 \supset F_1 \supset \dots \supset F_n \supset \dots$$

and by the definition of $\{F_i\}_{i=0}^{\infty}$ we have injective maps

$$F_0/F_1 \hookrightarrow F_1/F_2 \hookrightarrow \dots \hookrightarrow F_n/F_{n+1} \hookrightarrow \dots$$

By Riemann-Roch's Theorem, $\dim F_0/F_1 = 3$. We define the scollar invariants $e_i = e_i(g_4^1)$ ($i = 1, 2, 3$) by

$$e_i = e_i(g_4^1) = \#\{j \in \mathbb{N}; \dim(F_{j-1}/F_j) \geq i\} - 1 \quad (i = 1, 2, 3).$$

It is clear that $e_1 \geq e_2 \geq e_3 \geq 0$ and $e_1 + e_2 + e_3 = g - 3$. We now consider another description of scollar invariants. For any $i \in \mathbb{N} \cup \{0\}$, we define $\alpha_i =$

$\dim\Gamma(C, \mathcal{O}((i+1)g_4^1)) - \dim\Gamma(C, \mathcal{O}(ig_4^1))$. Then we have

$$e_i = \min\{j; \alpha_j \geq 4 - i + 1\} - 1$$

where $1 \leq i \leq 3$. This is proved by Riemann-Roch Theorem. Therefore we have that

$$\begin{aligned} \dim\Gamma(C, \mathcal{O}(g_4^1)) &= 2, \\ \dim\Gamma(C, \mathcal{O}(2g_4^1)) &= 3, \\ &\dots, \\ \dim\Gamma(C, \mathcal{O}((e_3 + 1)g_4^1)) &= e_3 + 2, \\ \dim\Gamma(C, \mathcal{O}((e_3 + 2)g_4^1)) &= (e_3 + 3) + 1, \\ &\dots, \\ \dim\Gamma(C, \mathcal{O}((e_2 + 1)g_4^1)) &= (e_2 + 2) + (e_2 - e_3), \\ \dim\Gamma(C, \mathcal{O}((e_2 + 2)g_4^1)) &= (e_2 + 3) + (e_2 - e_3 + 1) + 1, \\ &\dots, \\ \dim\Gamma(C, \mathcal{O}((e_1 + 1)g_4^1)) &= (e_1 + 2) + (e_1 - e_3) + (e_1 - e_2), \\ \dim\Gamma(C, \mathcal{O}((e_1 + 2)g_4^1)) &= (e_1 + 3) + (e_1 - e_3 + 1) + (e_1 - e_2 + 1) + 1, \\ &\dots. \end{aligned}$$

We know the following result (see [4] p.4588 Theorem 1).

Theorem 1 *Let $e_1 \geq e_2 \geq e_3 \geq 0$ be integers such that $e_1 + e_2 + e_3 = g - 3$. Then there is a 4-gonal curve $C = (C, g_4^1)$ with the scollar invariants (e_1, e_2, e_3) such that $|(e_3 + 2)g_4^1|$ is birationally very ample, if and only if $e_2 \leq 2e_3 + 2$ and $e_1 \leq e_2 + e_3 + 2$.*

In this paper we shall consider a generalization of this result. The Main result is the following.

Theorem 2 (Main Theorem) *Let $e_1 \geq e_2 \geq e_3 \geq 0$ be integers such that $e_1 + e_2 + e_3 = g - 3$. Then there is a tetragonal curve $C = (C, g_4^1)$ with the scollar invariants (e_1, e_2, e_3) , if and only if $e_1 \leq e_2 + e_3 + 2$.*

1 The Proof of Main Theorem

We use the following result.

Theorem 3 *Let C be a tetragonal curve with the scollar invariants (e_1, e_2, e_3) . Let g_4^1 be a base point free complete linear system on C . Let $\phi : C \rightarrow \mathbb{P}(\Gamma(C, \mathcal{O}((e_3 + 2)g_4^1)))$ be the morphism defined by $|(e_3 + 2)g_4^1|$. If ϕ is not birational onto its image, then $e_2 \neq e_3$, $\deg(\phi) = 2$ and $\phi(C)$ is a complete non-singular curve of genus $e_3 + 1$ admitting a g_2^1 with $\phi^*g_2^1 = g_4^1$.*

From Theorem 1 and Theorem 3, we may assume C is a two-sheeted cover of a hyperelliptic curve D of genus $e_3 + 1$ such that $e_3 \neq e_2$. And put $\pi : C \rightarrow D$ be a double-covering and we put $g_4^1 = \pi^*g_2^1$. Under these assumptions, we prove that Main Theorem.

We know that $\pi_*\mathcal{O}(ng_4^1) \cong \mathcal{O}(ng_2^1) \oplus \mathcal{O}(ng_2^1 - E)$ for some divisor D such that $2E$ is linearly equivalent to some effective divisor (see [6] p.326-p.328). We now consider (C, g_4^1) such that $g_4^1 = \pi^*g_2^1$ and let e_1, e_2, e_3 be scollar invariants. Then

$$\dim\Gamma(C, \mathcal{O}((e_3 + 2)g_4^1)) = (e_3 + 3) + 1$$

implies $\dim\Gamma(D, \mathcal{O}((e_3 + 2)g_2^1 - E)) = 0$ because $\pi_*\mathcal{O}(ng_4^1) \cong \mathcal{O}(ng_2^1) \oplus \mathcal{O}(ng_2^1 - E)$. And

$$\dim\Gamma(C, \mathcal{O}((e_2 + 2)g_4^1)) = (e_2 + 3) + (e_2 - e_3 + 1) + 1$$

implies $\dim\Gamma(D, \mathcal{O}((e_2 + 2)g_2^1 - E)) = 1$. Therefore we have an effective divisor $T = P_1 + \dots + P_t$ such that $\iota(P_i) \notin \{P_1, \dots, P_t\}$, for every $i = 1, \dots, t$ and $(e_2 + 2)g_2^1 - T \sim E$ where ι is a hyperelliptic involution on D . As $2E$ is linearly equivalent to a ramification divisor of ϕ (see [6] p.326-p.328), so we have

$$2g - 2 = 2(2(e_3 + 1) - 2) + 2(2(e_2 + 2) - t).$$

Hence we have

$$t = -e_1 + e_2 + e_3 + 2$$

because $e_1 + e_2 + e_3 = g - 3$. As $t \geq 0$, we have $e_1 \leq e_2 + e_3 + 2$. Now let $e_1 \geq e_2 \geq e_3 \geq 0$ be integers such that $e_1 + e_2 + e_3 = g - 3$ and assume that $e_1 \leq e_2 + e_3 + 2$. Let D be a hyperelliptic curve of genus $e_3 + 1$. Let $t = -e_1 + e_2 + e_3 + 2$. Take points $P_1, \dots, P_t \in D$ such that $\iota(P_i) \notin \{P_1, \dots, P_t\}$, for every $i = 1, \dots, t$. Let $T = P_1 + \dots + P_t$ and $E = (e_2 + 2)g_2^1 - T$. Let take an effective divisor R such that $R \sim 2E$. Because $2E \sim 2(e_2 + 2)g_2^1 - 2T$, so $e_2 + 2 \geq 2t = 2(-e_1 + e_2 + e_3 + 2)$ implies $2E$ is linearly equivalent to some effective divisor R . Therefore an isomorphism $\mathcal{O}(-2E) \cong \mathcal{O}(-R) \hookrightarrow \mathcal{O}$ induces an algebra structure on $\mathcal{O} \oplus \mathcal{O}(-E)$. Let $C = \text{Spec}(\mathcal{O} \oplus \mathcal{O}(-E))$. Then R is a ramification divisor of $\pi : C \rightarrow D$ (see [6] p.326-p.328), therefore C is of genus g . Let $g_4 = \pi^*(g_2^1)$. Then it is clear that scollar invariants of g_4^1 are e_1, e_2, e_3 .

Q.E.D.

References

- [1] M. Coppens, The Weierstrass gap sequence of the total ramification points of trigonal curve of \mathbb{P}^1 , *Indag. Math.*, **47** (1985) 245–270.
- [2] M. Coppens, The Weierstrass gap sequence of the ordinary ramification points of trigonal curve of \mathbb{P}^1 ; Existence of a kind of Weierstrass gap sequence, *J. Pure Appl. Algebra*, **43** (1986) 11–25.
- [3] R. C. Gunning, On the gonality ring of Riemann surfaces. preprint.
- [4] T. Kato, A. Ohbuchi Very ampleness of multiple of tetragonal linear systems, *Communications in Algebra* **21(12)** (1993), 4587–4597.
- [5] J. Komeda, The Weierstrass gap sequences of certain ramification points of tetragonal coverings of \mathbb{P}^1 , *Research Rep. of Ikutiku Tech. Univ.* **B-12** (1988), 185–191.
- [6] D. Mumford, Prym varieties I, *Contribution to Analysis*. Acad. Press, (1947) 325–355.
- [7] F.-O. Schreyer, Syzygies of Canonical Curves and Special Linear Series, *Math. Ann.*, **275** (1986) 105–137.