

## ORIGINAL

# Toll-like receptor 4 regulates gastric pit cell responses to *Helicobacter pylori* infection

Tsukasa Kawahara, Yuki Kuwano, Shigetada Teshima-Kondo, Tomoko Kawai, Takeshi Nikawa, Kyoichi Kishi, and Kazuhito Rokutan

*Department of Nutritional Physiology, The University of Tokushima School of Medicine, Tokushima, Japan*

**Abstract:** Gastric pit cells express mitogen oxidase1 (Mox1) and essential components for the phagocyte NADPH oxidase (p67-, p47-, p40-, and p22-phoxes). *Helicobacter pylori* (Hp) lipopolysaccharide (LPS) is a potent up-regulator of the Mox 1 oxidase. In this study, we examined the expression levels of several key members of the Toll-like receptor (TLR) family in primary cultures of guinea pig gastric pit cells. These cells expressed the TLR4 mRNA. Immunoblot analysis and immunofluorescence histochemistry with an anti-TLR4 antibody showed that gastric pit cells possessed significant amounts of TLR4 protein preferentially on the plasma membrane. In contrast, the cells did not express the TLR2 and TLR9 transcripts and did not contain detectable amounts of TLR2 protein. Neither peptidoglycan from *Staphylococcus aureus* nor Hp DNA with the CpG motif up-regulated Mox1 oxidase activity. Hp LPS activated nuclear factor- $\kappa$ B in association with the expression of cyclooxygenase II and tumor necrosis factor  $\alpha$  transcripts. These findings suggest that TLR 4 may play a crucial role in the initiation of inflammatory responses of gastric pit cells against Hp infection. *J. Med. Invest.* 48 : 190-197, 2001

**Keywords :** toll-like receptor, gastric pit cell, *Helicobacter pylori*, mitogen oxidase 1

## INTRODUCTION

*Helicobacter pylori* (Hp) is a Gram-negative, microaerophilic bacterial rod, associated with gastritis, peptic ulcer, and gastric cancer. (1-3). Hp infection elicits inflammatory and immunological responses in the host, although the magnitude of the responses varies from strain to strain and from host to host. Both cellular immune and antibody responses contribute to Hp-associated gastritis. Once the host has been infected, gastric epithelial cells secrete interleukin (IL)-6, IL-8 and other mediators that recruit neutrophils to gastric mucosa (4). Surface mucous cells

(pit cells) serve a primary protective role against irritants by providing a mucous coat. Hp preferentially colonizes in the mucous coat and selectively attaches to pit cells. There is increasing evidence that pit cells actively regulate the initiation of Hp-associated inflammation in gastric mucosa. In addition to inflammatory cytokines, such as IL-6, IL-8, and tumor necrosis factor (TNF- $\alpha$ ), major histocompatibility complex II antigen was shown to be strongly expressed on these cells (5). Thus, to understand the pathophysiology of Hp-associated gastritis, it is particularly important to elucidate the molecular basis of Hp-epithelial cell interactions.

Among the Hp genes, an insertion of approximately 40 kb of foreign DNA, named *cag* pathogenicity island (PAI), is now known as a major determinant of virulence (6). The *cag* PAI genes contains 31 genes, and six of the genes code for the core subunits of the type IV export machinery that can transfer the

Received for publication May 18, 2001 ; accepted May 25, 2001.

Address correspondence and reprint requests to Kazuhito Rokutan, Department of Nutritional Physiology, The University of Tokushima School of Medicine, Kuramoto-cho, Tokushima770-8503, Japan and Fax : +81-88-633-7086.

cytotoxin-associated protein A (CagA) into host epithelial cells, and translocated CagA has been shown to be tyrosine-phosphorylated by host cells (6-9). This route has been implicated as one of the major pathways to trigger gastric epithelial responses. Recently, we reported that gastric pit cells expressed a novel superoxide anion ( $O_2^-$ )-producing NADPH oxidase (10). This oxidase was then identified as a mitogen oxidase 1 (Mox1) (11). *Hp* lipopolysaccharide (LPS) up-regulated Mox1 oxidase activity, and enhanced production of oxygen radicals from Mox1 oxidase subsequently activated a crucial transcription activator of many inflammatory response genes, nuclear factor- $\kappa$ B (NF- $\kappa$ B) (10, 11). These findings suggested that gastric pit cells may have an ability to initiate an innate immune response against *Hp*. Based on these findings, in the present study, we examined the expression of several key members of the Toll-like receptor (TLR) family in primary cultures of guinea pig gastric pit cells. We also tested whether bacterial components actually stimulated distinct TLR signals.

## MATERIALS AND METHODS

### *Preparation and culture of gastric mucosal cells under LPS-free conditions*

Male guinea pigs weighing approximately 250 g were purchased from Shizuoka Laboratory Animal Center Inc. (Shizuoka, Japan). Gastric mucosal cells were isolated aseptically from guinea pig fundic glands as described previously (10). In the present experiments, all reagents used for culture were free from detectable amounts of LPS by the *Limulus* amoebocyte lysate assay (Endospecy; Seikagaku Kogyo Co. Tokyo, Japan). The isolated cells were cultured for 2 days in RPMI 1640 (GIBCO, Grand Island, NY), containing 50  $\mu$ g/ml gentamicin, 100 U/ml penicillin G, and 10% (vol/vol) fetal bovine serum (FBS; ICN Biomedicals, Aurora, OH). This complete medium contained less than 0.01 endotoxin unit (EU)/ml of LPS. Cell populations were determined by histochemical and immunohistochemical analyses, as described previously (10). After culturing for 48 h, growing cells consisted of pit cells (about 90%), pre-pit cells (about 5%), parietal cells (4-5%), mucous neck cells (less than 1%), and fibroblasts (less than 1%).

### *Measurement of $O_2^-$ production*

Mature pit cells were confirmed to be  $O_2^-$ -producing

cells by the nitroblue tetrazolium staining (10). The amount of  $O_2^-$  release was spectrophotometrically measured by the superoxide dismutase (SOD)-inhibitable reduction of cytochrome *c* and expressed as nmol/mg protein/h (10).

### *Preparation of *Hp* LPS and CpG DNA*

LPS was prepared from NCTC 11637 by the hot-phenol-water method (12) and subsequently treated with DNase 1, RNase A, and proteinase K, as described by Moran *et al.* (13). The treated LPS was boiled for 1 h to inactivate the enzymes. The LPS was dialyzed against LPS-free water (Otsuka Pharmaceutical Co., Tokushima, Japan) and ultracentrifuged at 100,000g for 18 h. Precipitated LPS was dissolved in LPS-free saline (Otsuka Pharmaceutical Co.). Phosphothioate-stabilized oligonucleotides containing CpG motif (CpG DNA) in *cagM* (5-TCCAATAACGTTTCTAATTTATTGAGCAGA-3) and *cagH* (5-TTCAATAGCGCTAGGATTTTTAGGGTGGT-3) were purchased from Nisshinbo Co. (Tokyo, Japan). These CpG DNAs were dissolved in LPS-free saline.

### *Detection of TLR, cyclooxygenase II (COX II), TNF- $\alpha$ , and actin transcripts*

Total RNA was isolated from guinea pig gastric mucosal cells (GMC), guinea pig or human peripheral blood lymphocytes (PBL) with an acid guanidium thiocyanate-phenol-chloroform mixture (14). Reverse transcriptase (RT)-PCR was done to detect the TLR2, 4, 9, COX II, TNF- $\alpha$  and actin transcripts using the following PCR primer sets:

TLR 2, 5'-GTCCAGGAGCTGGAGA-3' and 5'-GGAACCTAGGACTTTATCGCA-3'; TLR 4, 5'-TCACCTGATGCTTCTTGCTG-3' and 5'-AGTCGTCTCCAGAAGATGTG-3', TLR 9, 5'-ATGCCCTGCGCTTCCATTCATGGA-3' and 5'-GAACTGTCTTCAACACCAGGCCTT-3', COX II, 5'-CCAGTTTGTGAATCATTACC-3' and 5'-AAAGTACTCGGCTTCCAGTAG-3', TNF- $\alpha$ , 5'-AAAGTAGACCTGCCCGGACT-3' and 5'-GTACCTCATCTACTCCCAGG-3', and  $\beta$ -actin, 5'-TCACCGAGGCCCTCTGAACCCTA-3' and 5'-GGCAGTAATCTCCTTCTGCATCCT-3'. The resultant PCR products separated on an agarose gel were purified, ligated into a pCR 2.1-TOPO vector (Invitrogen, Carlsbad, CA.), and transformed into JM109 cells. Transformed plasmids containing the appropriate insert DNA were selected and sequenced with a DNA sequencer (model ABI 377; PE Biosystems Japan, Tokyo, Japan).

*Detection of TLR proteins*

An anti-TLR 2 or anti-TLR 4 antibody was generated by immunization of a rabbit with synthetic peptide of the amino acid residues 295-310 of human TLR2 or 183-199 of human TLR4 (15). The resultant serum was further purified by affinity chromatography with the synthetic peptide-conjugated agarose. A membrane fraction was prepared as previously described (10). Each sample of 20 µg protein per lane was separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) using a 7.5% polyacrylamide gel and transferred to a polyvinylidene difluoride filter. After nonspecific binding sites were blocked with 4% purified milk casein, the filter was incubated for 1 h at room temperature with the anti-TLR2 or anti-TLR4 antibody. Bound antibodies were detected by an enhanced chemiluminescence system (Amersham Pharmacia, Piscataway, NJ).

*Immunofluorescence histochemistry*

Cells growing on a LPS-free glass coverslip were washed with phosphate-buffered saline (PBS) and fixed with 4% paraformaldehyde in PBS for 20 min. After washing with PBS, the cells were blocked with 4% purified milk casein. They were incubated with a 1 : 100 dilution of rabbit anti-TLR4 antibody for 1 h at room temperature and then treated with a 1:500 dilution of Rhodamine-linked goat antibody against rabbit IgG (Amersham Pharmacia) for 1 h at room temperature. The samples were mounted with Vectashield Mounting Medium (Vector, Burlingame, CA). Subcellular localization of TLR4 was viewed using a confocal laser scanning microscopy (model TCSNT ; Leica, Heidelberg, Germany)

*Gel mobility shift assay.*

Nuclear proteins were prepared from cultured guinea pig gastric mucosal cells, and the activation of NF-κB was examined by gel mobility shift assay, as described previously (10, 11).

*Statistical analysis*

ANOVA and Scheffé's test were used to determine statistically significant differences. Differences were considered significant if  $P < 0.05$ .

**RESULTS***Effects of Hp components on O<sub>2</sub><sup>-</sup> production*

Guinea pig gastric pit cells, cultured under the

Table 1. O<sub>2</sub><sup>-</sup> release from guinea pig gastric mucosal cells

Treatment <sup>a</sup>	O <sub>2</sub> <sup>-</sup> release (nmol/ mg protein/h) <sup>b</sup>
none	11 ± 2
culture supernatant	19 ± 3*
boiled culture supernatant	19 ± 4*
LPS	105 ± 4*
DNase 1-treated LPS	102 ± 5*
RNase A-treated LPS	103 ± 4*
proteinase K-treated LPS	102 ± 3*
peptidoglycan	12 ± 3
CpG DNA ( <i>cagM</i> )	13 ± 3
CpG DNA ( <i>cagH</i> )	10 ± 3

<sup>a</sup>Culture supernatant and boiled supernatant of *Hp* were prepared as previously described (10). LPS was prepared by the hot-phenol-water method and treated with DNase 1, RNase A, or proteinase K, as described in MATERIALS AND METHODS. Cells were treated with the culture supernatants, these LPSs (1 EU/ml), *S. aureus* peptidoglycan (10 µg/ml), or CpG DNAs (50 µg/ml) at 37°C for 24 h in RPMI 1640 containing 10% FBS. <sup>b</sup>O<sub>2</sub><sup>-</sup> release was measured as described in MATERIALS AND METHODS. \*Significantly increased vs untreated control cells ( $p < 0.05$  by ANOVA and Scheffé's test).

conditions used in our previous studies, spontaneously released about 50 nmol O<sub>2</sub><sup>-</sup>/mg protein/h (10, 11). The basal rate of O<sub>2</sub><sup>-</sup> production decreased to 11 ± 2 nmol/mg protein/h (mean ± SD,  $n=12$ ) in the LPS-free system used in the present study (Table 1). Using this system, we re-evaluated the effects of bacterial components or products on Mox1 oxidase.

Culture supernatant was prepared after cultivation of *Hp* alone ( $1 \times 10^7$  cells/ml) in RPMI 1640 containing 10% FBS for 24 h was then added to gastric mucosal cells ( $1 \times 10^5$  cells). After been cultured for 24 h, O<sub>2</sub><sup>-</sup> production increased 1.7-fold (Table 1). The stimulatory effects of supernatants could not be eliminated by boiling (Table 1). When LPS was extracted by the hot-phenol method and added to the cells, it increased O<sub>2</sub><sup>-</sup> production 10-fold. This LPS possibly contained DNA, RNA, and denatured protein, while treatment of *Hp* LPS with DNase 1, RNase, or proteinase K did not eliminate the activity (Table 1).

Three different bacterial components have been identified as specific ligands for distinct TLRs; LPS for TLR4, peptidoglycan for TLR2, and CpG DNA for TLR9 (16, 17). We also tested whether stimulation of TLR9 by bacterial DNA with CpG motif resulted in the up-regulation of O<sub>2</sub><sup>-</sup> production. For this purpose, CpG DNAs that were coded in the *cagM* and *cagH* genes of *Hp* NCTC11638 (GenBank




Fig. 1. Detection of TLR 2, 4, and 9 transcripts in gastric mucosal cells. Total RNA was isolated from cultured guinea pig gastric mucosal cells (GMC), human or guinea pig peripheral blood lymphocytes (PBL), and RT-PCR was performed, as described in MATERIALS AND METHODS. The RT-PCR products (5 µg) were subjected to electrophoresis in a 6% polyacrylamide gel. Lanes 1 in A-C show molecular weight standard markers (MWSTD). Similar results were obtained in three separate experiments.




Fig. 2. Detection of TLR2 and 4 proteins in gastric mucosal cells. Membrane proteins were prepared from guinea pig GMC, human or guinea pig PBL, as described in MATERIALS AND METHODS. Immunoblot analysis with an antibody against TLR4 (A) or 2 (B) was performed. Guinea pig GMC growing on an LPS-free glass coverslip were stained with an antibody against TLR4 as described in MATERIALS AND METHODS (C). Cells were examined using a confocal laser scanning microscopy. The scale bar indicates 10 µm.

accession number AE000511) were synthesized, and gastric mucosal cells were stimulated by these CpG DNAs. However, these CpG DNAs did not increase O<sub>2</sub><sup>-</sup> production (Table 1). We also confirmed that stimulation of TLR2 by *Staphylococcus aureus* peptidoglycan (Fluka Chemie AG, Buchs, Switzerland) did not affect the O<sub>2</sub><sup>-</sup>-producing activity (Table 1). Thus, LPS was identified as a bioactive component of *Hp* that stimulated Mox1 oxidase expressed in gastric pit cells.

Cell viability was maintained throughout the experiments. This was based on continued trypan blue

exclusion, adherence to the culture plates, and unchanged morphological features (data not shown). The concentrations of LPS used in this study did not cause apoptosis of the cells, which was confirmed by DNA ladder formation and nuclear morphology (data not shown). However, *Hp* LPS at 42 EU/ml (71-560 ng/ml) or higher induced DNA fragmentation and apoptotic chromosomal condensation under low-serum conditions (submitted to JMI).

#### Expression of TLRs on gastric mucosal cells

Human and guinea pig peripheral blood lymphocytes




Fig. 3. Activation of NF- $\kappa$ B and expression of COX II or TNF- $\alpha$ . After been cultured for 48 h in RPMI 1640 containing 10% FBS, cells were treated with 1 EU/ml *Hp* LPS in RPMI 1640 containing 10% FBS, 200 U/ml SOD, and 350U/ml catalase for the indicated times. Nuclear extracts were prepared from the cells and analyzed by gel mobility shift assay as described in MATERIALS AND METHODS (A). Total RNA was isolated and analyzed by RT-PCR as described in MATERIALS AND METHODS (B). Cells were treated with 1 EU/ml *Hp* LPS in RPMI 1640 containing 10% FBS for the indicated times (C). The amounts of O<sub>2</sub><sup>-</sup> release were measured as described in MATERIALS AND METHODS, and they are expressed as means  $\pm$  SD ( $n=12$ ). \*Significantly increased vs untreated control cells ( $p<0.05$  by ANOVA and Scheffé's test).

(PBL) were isolated as described previously (10). Total RNA was extracted from these cells and used as positive controls to detect the transcripts of TLRs 2, 4, and 9 in gastric mucosal cells. RT-PCR analysis showed that gastric mucosal cells expressed neither the TLR2 (Fig. 1A) nor the TLR9 transcript (Fig. 1C). These findings also demonstrated that our cultures did not contain significant numbers of immune cells. RT-PCR amplified the TLR4 mRNA in gastric mucosa cells (Fig. 1B). The nucleotide sequence showed that the amplified product had 90% nucleotide sequence identity to the human TLR4 cDNA (GenBank accession number U88880).

#### Expression of TLR2 and 4 proteins in gastric mucosal cells

We also examined whether TLR2 and TLR4 proteins were expressed in gastric mucosal cells, since TLR2 was reported to be another candidate for an LPS-receptor (18). Immunoblot analysis with an anti-TLR4 antibody showed that guinea pig gastric mucosal cells expressed TLR4 protein, as was detected in human and guinea pig PBL (Fig. 2A). Immunocytochemical analysis showed that TLR4 was preferentially distributed on plasma membrane (Fig. 2C). In contrast, human and guinea pig PBL contained significant amounts of TLR2 protein, while gastric mucosal cells did not possess detectable amounts of TLR2 protein (Fig. 2B).

#### LPS-induced expression of COX II and TNF- $\alpha$ mRNAs

In previous studies (10, 11), we showed that *Hp* LPS could activate NF- $\kappa$ B in cultured gastric mucosal cells. After stimulation by *Hp* LPS, the activation of NF- $\kappa$ B occurred within 30 min and continued for longer than 12 h (10). Enhanced production of oxygen radicals by up-regulated Mox1 oxidase was suggested to mainly participate in the activation (10). However, the increase in O<sub>2</sub><sup>-</sup> production from the cells was first detected at 8 h after the treatment with *Hp* LPS (Fig. 3C). Furthermore, *Hp* LPS could activate NF- $\kappa$ B with a peak at 4 h even in the presence of SOD and catalase (Fig. 3A), showing that NF- $\kappa$ B was activated directly by *Hp* LPS, in addition to indirectly by Mox1-derived oxygen radicals up-regulated by *Hp*-LPS. In association with this activation of NF- $\kappa$ B, *Hp* LPS stimulated the expression of COX II (an upper panel in Fig. 3B) and TNF- $\alpha$  (a middle panel in Fig. 3B) transcripts in the presence of SOD plus catalase. These findings suggest that LPS directly initiates NF- $\kappa$ B-dependent inflammatory and immune responses.

## DISCUSSION

Using an LPS-free culture system, we reconfirmed that guinea pig gastric mucosal cells in culture were sensitive to *Hp* LPS. *Hp* does not attach to

guinea pig gastric mucosal cells ; therefore, our system is an excellent model for studying the contact-independent interactions between gastric epithelial cells and the Gram-negative bacterium. Mox1 oxidase-derived  $O_2^-$  and related oxygen intermediates were suggested to play crucial roles in the initiation of inflammatory and immune responses as well as in the regulation of cell growth (10, 11). *Hp* LPS that activates Mox1 oxidase appears to be one of the important virulent factors of *Hp*.

Recently, it was been shown that bacterial components, such as LPS, peptidoglycan, and DNA stimulate innate immune response through distinct TLR receptors. TLR2 has been suggested to be another possible candidate for an LPS receptor (18). We confirmed that gastric mucosal cells did not express the TLR2 mRNA and detectable amounts of TLR2 protein. In contrast, intestinal epithelial cells express TLR2 and TLR4 (19). The sensitivity to LPS is strikingly different between gastric mucosal cells and intestinal epithelial cells. Intestinal epithelial cells are insensitive to LPS, making them tolerant to constant exposure of resident microflora. In contrast, any bacteria, except for *Hp*, cannot colonize persistently under acidic conditions in the stomach. The high sensitivity to *Hp* LPS may be an important feature characteristic of gastric mucosal cells, when considering *Hp* infection. The TLR9 transcript was not expressed in gastric mucosal cells (Fig.1C) and intestinal epithelial cells (data not shown). DNA molecules are usually stable in acidic conditions in the stomach, and the intestine is persistently exposed to bacterial flora. Therefore, the absence of TLR9 may make gastrointestinal epithelial cells quiescent to an excessive response to bacterial DNA.

In phagocytes, activation of TLR4 signaling is reported to induce expression of other TLRs (20). However, *Hp* LPS did not induce expression of the TLR2 and 9 transcripts in gastric mucosal cells (data not shown). Thus, among the TLRs examined, TLR4 appeared to play a crucial role in the gastric mucosal cell response to *Hp* infection. More recently, it was reported that TLR5 is a receptor specific for bacterial flagellin and mediated an innate immune response to this bacterial component (21). Further studies are necessary to elucidate the role of TLR family in the innate immune response of gastric mucosa.

Recently, several novel isozymes of gp91-*phox* expressed in non-phagocytic cells, including Mox1 (22), Renox (23), and Thoxes (24), were molecularly identified. Among these non-phagocytic oxidases, the pit cell Mox1 oxidase, whose  $O_2^-$ -producing capacity

is equivalent to that of macrophages (10, 11), is the most potent. We previously studied the effects of growth factors (epidermal growth factor and transforming growth factor- $\beta$ ), cytokines (interferon- $\gamma$ , TNF- $\alpha$ , IL-1, IL-3, and IL-6), histamine, carbacol, and phorbol 12-myristate13-acetate (PMA) on the oxidase activity ; however, none were able to up-regulate the  $O_2^-$  production (10 and data not shown). Finally, we observed that LPS from *Hp* and *E. coli* similarly up-regulate Mox1 oxidase activity (10). These findings also support the important role of *Hp* LPS and TLR4 in the initiation of an inflammatory response against *Hp* infection.

In phagocytes, LPS stimulates TLR4 signal-transmittable molecules, such as myeloid differentiation factor 88 (MyD 88), IL-1R-associated kinase, TNF receptor-associated factor 6, TAK1, and NIK, leading to activation of NF- $\kappa$ B (25). *Hp* infection triggers the activation of NF- $\kappa$ B to produce inflammatory response genes, such as TNF- $\alpha$ , interleukin 6, and 8 (26, 27). In a previous study (10), it was suggested that enhanced production of oxygen intermediates by Mox1 oxidase might play a crucial role in the activation of NF- $\kappa$ B, leading to expression of proinflammatory mediators. *Hp* LPS stimulated distinct TLR4 signaling, such as phosphorylation of TAK1 and TAB1 (submitted to JMI). In the present study, we demonstrated that *Hp* LPS itself directly activated NF- $\kappa$ B and stimulated the COX II and TNF- $\alpha$  mRNA expression even when  $O_2^-$  and hydrogen peroxide were removed by SOD and catalase. Oxygen radicals derived from Mox1 oxidase appeared to significantly contribute to the activation of NF- $\kappa$ B particularly later than 8 h after treatment with *Hp* LPS, since, in this later phase, *Hp* LPS markedly enhanced the production of  $O_2^-$  from Mox1 oxidase, and SOD and catalase significantly inhibited NF- $\kappa$ B activation (10). Thus, NF- $\kappa$ B was rapidly activated directly by *Hp* LPS, and Mox1-derived oxygen intermediates prolonged this activation, leading to an enhanced inflammatory response of gastric mucosal cells.

The findings of the present study suggested that TLR4-mediated intracellular signals and Mox1 oxidase may play an important role in the *Hp*-epithelial interactions, leading to the initiation of an inflammatory response of gastric mucosa against *Hp* infection.

## REFERENCES

1. Blaser MJ: *Helicobacter pylori* and the pathogenesis of gastroduodenal inflammation. J Infect Dis

- 161 : 626-633, 1990
2. Parsonnet J, Friedman GD, Vandersteen DP, Chang Y, Vogelman JH, Orentreich N, Sibley RK : *Helicobacter pylori* infection and the risk of gastric carcinoma. N Engl J Med 325 : 1127-1131, 1991
  3. Parsonnet J, Hansen S, Rodriguez L., Gelb AB, Warnke RA, Jellum E, Orentreich N, Vogelman JH, Friedman GD : *Helicobacter pylori* and gastric lymphoma. N Engl J Med 330 : 1267-1271, 1994
  4. Sharma SA, Tummuru MKR, Blaser MJ, Kerr LD : Activation of IL-8 gene expression by *Helicobacter pylori* is regulated by transcription factor nuclear factor- $\kappa$ B in gastric epithelial cells. J Immunol 160 : 2401-2407, 1998
  5. Fan X, Crowe SE, Behar S, Gunasena H, Ye G, Haeberle H, Houten NV, Gourley WK, Ernst PB, Reyes VE. The effect of class II major histocompatibility complex expression on adherence of *Helicobacter pylori* and induction of apoptosis in gastric epithelial cells : a mechanism for T helper cell type1-mediated damage. J Exp Med 187 : 1659-1669, 1998
  6. Covacci A., Telford JL, Giudice GD, Parsonnet J, Rappuoli R : *Helicobacter pylori* virulence and genetic geography. Science 284 : 1328-1333, 1999
  7. Asahi M, Azuma T, Ito S, Ito Y, Suto H, Nagai Y, Tsubokawa M, Maeda S, Omata M, Suzuki T, Sasakawa C : *Helicobacter pylori* CagA protein can be tyrosine phosphorylated in gastric epithelial cells. J Exp Med 191 : 593-602, 2000
  8. Odenbreit S., Püls J, Sedlmaier B, Gerland E, Fisher W, Hass R: Translocation of *Helicobacter pylori* CagA into gastric epithelial cells by type IV secretion. Science 287 : 1497-1500, 2000
  9. Stein M., Rappuoli R, and Covacci A : Tyrosine phosphorylation of the *Helicobacter pylori* CagA antigen after *cag*-driven host cell translocation. Proc. Natl Acad Sci USA 97 : 1263-1268, 2000
  10. Teshima S, K Rokutan, Nikawa T, Kishi K : Guinea pig gastric mucosal cells produce abundant superoxide anion through an NADPH oxidase-like system. Gastroenterology 115 : 1186-1196, 1998
  11. Teshima S., Kutsumi H, Kawahara T, Kishi K, Rokutan K : Regulation of growth and apoptosis of cultured guinea pig gastric mucosal cells by mitogen oxidase 1. Am J Physiol 279 : G1169-G 1176, 2000
  12. Westphal O, Jann K : Bacterial lipopolysaccharides : extraction with phenol-water and further application of the procedure. Methods Carbohydr. Chem 5 : 83-91, 1965
  13. Moran AP, Helander IM, Kosunen TU : Compositional analysis of *Helicobacter pylori* rough-form lipopolysaccharides. J Bacteriol 174 : 1370-1377, 1992
  14. Chomczynski P, Sacchi N : Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. Anal Biochem 162 : 156-159, 1987
  15. Zhang FX, Kirschning CJ, Mancinelli R, Xu XP, Jin Y, Faure E, Montovani A, Roche M, Muzio M, Arditi M : Bacterial lipopolysaccharide activates nuclear factor- $\kappa$ B through interleukin-1 signaling mediators in cultured human dermal endothelial cells and mononuclear phagocytes. J Biol Chem 274 : 7611-7614, 1999
  16. Takeuchi O, Hoshino K, Kawai T, Sanjo H, Takeda H, Ogawa T, Takeda K, Akira S : Differential roles of TLR2 and TLR4 in recognition of gram-negative and gram-positive bacterial cell wall components. Immunity 11 : 443-451, 1999
  17. Hemmi H, Takeuchi O, Kawai T, Kaisho T, Sato S, Sanjo H, Matsumoto M, Hoshino K, Wagner H, Takeda K, Akira S : A Toll-like receptor recognizes bacterial DNA. Nature 408 : 740-745, 2000
  18. Yang RB, Mark MR, Gray A, Huang A, Xie MH, Zhang M, Goddard A, Wood WI, Gurney AL, Godowski PJ : Toll-like receptor-2 mediates lipopolysaccharide-induced cellular signaling. Nature 395 : 284-288, 1998
  19. Cario E, Rosenberg IM, Brandwein SL, Beck PL, Reinecker HC, Podolsky DK: Lipopolysaccharide activates distinct signaling pathways in intestinal epithelial cell lines expressing Toll-like receptors. J Immunol 164 : 966-972, 2000
  20. Liu Y, Wang Y, Yamakuchi M, Isowaki S, Nagata E, Kanmura Y, Kitajima I, Maruyama I : Upregulation of toll-like receptor 2 gene expression in macrophage response to peptidoglycan and high concentration of lipopolysaccharide is involved in NF-kappa b activation. Infect Immun 69 : 2788-2796, 2001
  21. Hayashi F, Smith KD, Ozinsky A, Hawn TR, Yi EC, Goodlett DR, Eng JK, Akira S, Underhill DM, Aderem A : The innate immune response to bacterial flagellin is mediated by Toll-like receptor 5. Nature 410 : 1099-1103, 2001
  22. Suh YA, Arnold RS, Lassegue B, Shi J, Xu X, Sorescu D, Chung AB, Griendling KK, Lambeth JD : Cell transformation by the superoxide-generating oxidase Mox1. Nature 401 : 79-82, 1999
  23. Geiszt M, Kopp JB, Varnai P, Leto TL : Identi-

- fication of renox, an NAD(P)H oxidase in kidney. Proc Natl Acad Sci USA 97 : 8010-8014, 2000
24. De Deken X, Wang D, Many MC, Costagliola S, Libert F, Vassart G, Dumont JE, Miot F: Cloning of two human thyroid cDNAs encoding new members of the NADPH oxidase family. J Biol Chem 275 : 23227-23233, 2000
  25. Irie T, Muta T, Takeshige K : TAK1 mediates an activation signal from toll-like receptor(s) to nuclear factor-kappaB in lipopolysaccharide-stimulated macrophages. FEBS Lett 467 : 160-164, 2000
  26. Sharma SA, Tummuru MK, Blaser MJ, Kerr LD : Activation of IL-8 gene expression by *Helicobacter pylori* is regulated by transcription factor nuclear factor-kappa B in gastric epithelial cells. J Immunol 160 : 2401-2407, 1998
  27. Crabtree JE, Shallcross TM, Heatley RV, Wyatt JI : Mucosal tumour necrosis factor alpha and interleukin-6 in patients with *Helicobacter pylori* associated gastritis. Gut 32 : 1473-1477, 1991