

**Epigenetic regulation of Hox gene expression by PcG genes  
in a primitive mode of insect embryogenesis in the cricket  
*Gryllus bimaculatus***

Yuji Matsuoka

Department of Biological Science and Technology  
Life and Materials Systems Engineering  
Graduate School of Advanced Technology and Science  
Tokushima University

September 2015

# Contents

<b>1. Abstract</b>	3
<b>2. General Introduction</b>	
2.1. Epigenetic regulation by Polycomb group genes	4
2.2. The cricket, <i>Gryllus bimaculatus</i> , is an emerging model organism	5
<b>3. Introduction</b>	6
<b>4. Results</b>	
4.1. Suppression of <i>Gryllus</i> PcG genes causes a homeotic phenotype	9
4.2. <i>Gryllus E(z)</i> has histone methylation activity	15
4.3. <i>E(z)</i> <sup>RNAi</sup> embryos show anterior misexpression of Hox genes	17
4.4. PcG genes and gap genes regulate Hox genes independently	22
4.5. Regional autonomy in the PcG silencing of Hox genes	24
<b>5. Discussion</b>	26
<b>6. Experimental procedures</b>	31
<b>7. References</b>	33
<b>8. Acknowledgement</b>	38
<b>9. Publications</b>	39

## 1. Abstract

In insect species that undergo long germ segmentation, such as *Drosophila*, all segments are specified simultaneously at the early blastoderm stage. As embryogenesis progresses, the expression boundaries of Hox genes are established by repression of gap genes, which is subsequently replaced by Polycomb group (PcG) silencing. At present, however, it is not known whether patterning occurs this way in a more ancestral (short germ) mode of embryogenesis, where segments are added gradually during posterior elongation. In this study, two members of the PcG family, *Enhancer of zeste (E(z))* and *Suppressor of zeste 12 (Su(z)12)*, were analyzed in the short germ cricket, *Gryllus bimaculatus*. Results suggest that although stepwise negative regulation by gap and PcG genes is present in anterior members of the Hox cluster, it does not account for regulation of two posterior Hox genes, *abdominal-A (abd-A)* and *Abdominal-B (Abd-B)*. Instead, *abd-A* and *Abd-B* are predominantly regulated by PcG genes, which is the mode present in vertebrates. These findings suggest that PcG-mediated silencing of Hox genes may have occurred during animal evolution. The ancestral bilaterian state may have resembled the current vertebrate mode of regulation, where PcG-mediated silencing of Hox genes occurs before their expression is initiated and is responsible for the establishment of individual expression domains. Then, during insect evolution, the repression by transcription factors may have been acquired in anterior Hox genes of short germ insects, while PcG silencing was maintained in posterior Hox genes.

## 2. General Introduction

### 2.1. Epigenetic regulation by Polycomb group genes

Epigenetic regulation is a way of gene-expression regulation without altering the underlying DNA sequence. That includes activation and silencing via chemical modification on DNA sequences or histone residues. These epigenetic changes play a role in cell memory by maintaining a gene expression profile throughout cell division. In this study, I focus on function of Polycomb group (PcG) genes, encoding epigenetic factor components.

PcG genes are genetically identified as a *trans* regulator that contributes to maintain the expression pattern of Hox genes in *Drosophila melanogaster* (Lewis, 1978). PcG proteins form three different classes of complexes termed Polycomb Repressive Complex (PRC) 1 and 2, and Pleiohomeotic Repressive complex (PhoRC). The stepwise regulation by these complexes is described as below. First, PhoRC contains the sequence specific DNA binding protein Pleiohomeotic (Pho), which binds specifically to Polycomb Response Elements (PREs), and recruits PRC2 to PREs with reaction of Pho with E(z), Polyhomeotic and Polycomb (Mohd-Sarip et al., 2002; Wang et al., 2004). Subsequently, PRC2 methylates histone 3 on lysine 27 (H3K27) residues within neighboring nucleotides. PRC2 contains core components: Enhancer of zeste (E(z)), Suppressor of zeste 12 (Su(z)12), Extra sex combs and Nurf-55. The SET domain containing E(z) subunit trimethylates H3K27, and Su(z)12 is necessary for the function in methylation supplementary. Then, PRC1 recognizes to be recruited to the trimethylated H3K27 via chromo-domain within Pc subunit. PRC1 contains Pc, Polyhomeotic, dRING, Posterior sex combs, Sex combs on midleg and TBP-associated factors. dRING possessing E3 ligase activity catalyzes ubiquitination in histone 2A on lysine 119, which might results in formation of heterochromatin. The molecular mechanisms of the formation of heterochromatin are not elucidated completely.

PcG genes are evolutionarily conserved from *Drosophila* to vertebrate. In *Drosophila*, PcG genes mutants show all body segments transform to most posterior abdominal segment and these embryos cause lethal (Soto et al., 1995). Multiple loci, whose mutations show similar phenotype with *Pc* mutant, are isolated, and are called as PcG genes. Genetic and biochemical analyses indicate that PcG proteins are necessary to maintain the repression of Hox genes, but are not necessary to establish segment specific expression domain of Hox genes (Schuettengruber et al., 2007). Knockout of

these genes in mouse indicate homeotic transformation in vertebra, suggesting that PcG proteins contribute to maintain the repression of Hox genes also in vertebrate (van der Lugt et al., 1994; Akasaka et al., 1996). In addition to the regulation of Hox genes expressions, PcG proteins repress another target genes expressions for control of cell growth, maintenance of stem cell, tumorigenesis, genome imprinting and X-chromosome inactivation (Gieni and Hendzel, 2009).

## **2.2. The cricket, *Gryllus bimaculatus*, is an emerging model organism**

In the present thesis, I studied function of PcG component genes in a two-spotted cricket, *Gryllus bimaculatus*, which is an emerging model animal to analyze morphogenesis and leg regeneration. In this insect, strong genetic tools for gene functional analysis, such as RNA interference, transgenesis, and gene-knockout, are available.

The one of such tools is RNA interference (RNAi). We can use four methods of RNAi in accordance with targeted genes; embryonic, parental, nymphal and regenerative RNAi. Embryonic RNAi is used for analyzing gene functions during embryogenesis and performed by injecting double-stranded RNA (dsRNA) into eggs (Shinmyo et al., 2005). Parental RNAi is also used for the same purpose, but in this case, dsRNA is injected in the body cavity of adult female crickets (Mito et al., 2005). Compared to embryonic RNAi, We can obtain plenty of RNAi-ed eggs by parental RNAi. Nymphal and regenerative RNAi is used for analysis of gene function during postembryonic development. In these methods, dsRNAs are injected into the body cavity of nymphal crickets. For studying regeneration mechanisms, we cut a leg of injected nymphs after injecting dsRNA.

The other technique is transgenesis by using *piggyback* transposase (Shinmyo et al., 2004; Nakamura et al., 2010). By using this technique, we generated transgenic cricket, of which cells or nuclei are fluorescently labeled. We succeeded live imaging to analyze a dynamics of cell movement during embryogenesis. Generation of transgenic crickets would be also useful for analyzing functions of genes and *cis*-regulatory elements. In this study, I used the parental RNAi method for analyzing PcG gene functions and also used eGFP expressing transgenic crickets to visualize RNAi effects on movements of developing embryos.

### 3. Introduction

Though all insects possess highly conserved adult body plans, there are two different ways developmental patterning can be accomplished. In long germ insects, all segments are specified simultaneously at the early blastoderm stage. In contrast, in short or intermediate germ (hereafter collectively called short germ) insects, only the anterior regions are specified at the blastoderm stage. The remaining posterior segments are gradually formed from the most posterior region, termed the growth zone, during posterior elongation. While the short germ type is believed to be the ancestral mode of segmentation in arthropods (reviewed by Davis and Patel, 2002), the actual molecular mechanisms that regulate it have not been identified.

Despite operational differences, previous studies had revealed that canonical functions of segmentation genes are fundamentally conserved between long germ and short germ segmentation (Mito et al., 2005; Mito et al., 2006; Mito et al., 2007). As embryogenesis progresses, a noticeable difference is observed in the expression of Hox genes. In long germ species, such as *Drosophila*, all Hox genes are expressed simultaneously at the blastoderm stage (Akam, 1987; Castelli-Gair, 1998; Bae et al., 2002). In short germ insects, on the other hand, the middle Hox genes, *Sex combs reduced* (*Scr*), *Antennapedia* (*Antp*), and *Ultrabithorax* (*Ubx*), are expressed in the anterior regions preceding initiation of posterior elongation. This is followed by expression of their posterior counterparts, *abdominal-A* (*abd-A*) and *Abdominal-B* (*Abd-B*), which are associated with the formation of posterior segments (Tear et al., 1990; Kelsh et al., 1993; Shippy et al., 1998; Peterson et al., 1999; Zhang et al., 2005). These observed differences in the temporal activation of posterior Hox genes might reflect different regulatory mechanisms of Hox genes.

In *Drosophila*, gap genes function as repressors and provide positional information, determining the anterior border of the Hox gene expression domain. This repressed state of each Hox gene is maintained by Polycomb group (PcG) genes after decay of gap gene activity (Simon et al., 1992; Struhl and Akam, 1985; Jones and Gelbart, 1990). Previous studies reveal that the functions of gap genes as Hox gene repressors are conserved in short germ insects (reviewed by Jaeger, 2011), while the functions of PcG genes have not yet been analyzed in short germ insect development.

Functionally, PcG genes are identified as trans regulators that contribute to maintaining the expression patterns of Hox genes in *Drosophila* (Lewis, 1978). PcG

gene products comprise 3 different types of complexes termed Polycomb repressive complex (PRC) 1 and 2 and Pleiohomeothic repressive complex (PhoRC). PhoRC, which binds specifically to Polycomb response elements (PREs), recruits PRC2 to PREs. PRC2 then trimethylates histone H3 on lysine 27 (H3K27) residues. This, in turn, provides a platform for recruiting PRC1. PRC1 catalyzes ubiquitination of lysine 119 on histone H2A, leading to silencing of target genes (Grimaud et al., 2006; Müller and Kassis, 2006; Simon and Kingston, 2009; Simon and Kingston, 2013). *Drosophila* PcG mutants exhibit homeotic phenotypes, in which multiple Hox genes are activated in body regions where they should be silent (Simon et al., 1992; Soto et al., 1995). In those mutants, transformation of all segments to posterior segments occurs as a result of misexpressed *Abd-B* (Jones and Gelbart, 1990; Birve et al., 2001). Misexpression of Hox genes starts after establishment of normal expression domains, indicating that PcG genes are involved in maintenance of Hox gene repression but not in initial repression (Simon et al., 1992; Struhl and Akam, 1985; Jones and Gelbart, 1990). Thus, during *Drosophila* embryogenesis, PcG-mediated silencing maintains Hox expression boundaries after they are set in early embryos by gap gene activity (White and Lehmann, 1986; Harding and Levine, 1988; Irish et al., 1989; Reinitz and Levine, 1990; Qian et al., 1991). Concordantly, trimethylation of histone H3 on lysine 27 (H3K27me3) is first detected after germband formation, 4–7 hours after egg laying (AEL) (Tie et al., 2009) and after the stage of assumed gap gene activity.

The PcG gene function to silence Hox genes is conserved in vertebrates. However, the establishment of Hox gene expression domains is different from that in *Drosophila*. In vertebrates, segments are progressively formed from anterior to posterior, along with posterior embryo growth. Hox genes are activated in time-dependent manner, during and after posterior growth, reflecting their positions in the gene cluster. This temporal and spatial collinear activation of Hox genes is accomplished through progressive demethylation of H3K27 in the gene cluster, which is silenced by PcG gene action prior to Hox gene activation (Soshnikova and Duboule, 2009). This sequential histone demethylation process is regulated by a gradient of retinoic acid signaling (Lee et al., 2007; Agger et al., 2007). Thus, in the regulatory machinery of vertebrate Hox genes, PcG-mediated silencing occurs before Hox gene expression is initiated, and it is responsible for the establishment of individual expression domains. This patterning mechanism differs substantially from *Drosophila*. Because the process of posterior

growth in vertebrates might be homologous to that in short germ insects with canonical Wnt signaling and the transcription factor caudal playing crucial roles (Martin and Kimelman, 2009), the regulatory machinery of the posterior Hox genes in short germ insects could be more similar to vertebrates than to *Drosophila*.

To examine this possibility, two members of the PcG family, the *Enhancer of zeste* (*E(z)*) and *Suppressor of zeste 12* (*Su(z)12*) genes, which are essential for histone methyltransferase activity of the PRC2 complex, were investigated in the short germ cricket *Gryllus bimaculatus*. To elucidate their functions in short germ insect development, RNA interference (RNAi)-based functional analyses of these genes were performed in the cricket. Here I discuss how the data provide novel insight into the key transitions and mechanisms governing regulation of Hox genes during animal evolution.




## 4. Results

### 4.1 Suppression of *Gryllus* PcG genes causes a homeotic phenotype

To investigate the functions of PcG genes during embryogenesis, I cloned *E(z)* and *Su(z)12* from *Gryllus* (Fig. 1). *E(z)* was first detected by *in situ* hybridization at stage 8 in a ubiquitous pattern (Fig. 2A,C). A parental RNAi approach was used to knock down these genes. *E(z)*<sup>RNAi</sup> embryos exhibited a membrane-enclosed, crescent shaped body that lacked cuticle formation (Fig. 3B). In these embryos, dorsal closure occurred at the ventral side due to failure in katanepsis, resulting in an “inside-out” morphology and 100% lethality (Fig. 3B; Fig. 4). As illustrated in Fig. 3D, the phenotype was characterized by a greatly contracted body and by transformation of antennae and mouthparts to leg-like appendages (Fig. 5). Two different dsRNAs, *Gb'E(z)\_N* and *Gb'E(z)\_C*, produced the same morphological phenotypes, excluding the possibility of off-targeting effects by dsRNAs. *Su(z)12*<sup>RNAi</sup> embryos exhibited a similar but less severe phenotype (Fig. 6C). This observation was consistent with real-time quantitative PCR (RT-qPCR) results showing the lesser reduction of *Su(z)12* mRNA levels in *Su(z)12*<sup>RNAi</sup> embryos compared to *E(z)* mRNA levels in *E(z)*<sup>RNAi</sup> embryos (Fig. 2D,E). For this reason, *E(z)*<sup>RNAi</sup> embryos were used for further detailed analyses.


The patterns of leg markers, *aristaless (al)* (Miyawaki et al., 2002) and *Distal-less (Dll)* (Niwa et al., 1997), were determined in *E(z)*<sup>RNAi</sup> embryos. With the exception of mandibles, the head appendages assumed leg-like expression patterns of *al* compared to wild type (Fig. 5A,B). In addition, in the affected head appendages, *Dll* expression patterns resembled those found in thoracic legs with distal and proximal domains (Fig. 5C,D). The number of segments was not affected, suggesting that the observed phenotypes were not caused by changes in segment specification mechanisms.

Expression of *wingless (wg)*, a ventral side marker of legs (Niwa et al., 2000), was restricted to the ventral margin of the elongating limb buds (Fig. 5E). However, in *E(z)*<sup>RNAi</sup> embryos, the *wg* pattern expanded to the dorsal limb margin (Fig. 5F). These observations indicated that, in depleted embryos, head appendages (antennae and mouthparts) assume leg-like identities, which also displayed altered dorsoventral polarity.




**Figure 1. Domain structure and conservation of *E(z)* and *Su(z)12* in *Gryllus*.**

The conserved regions (SET domain and VEFS-Box domain) in *E(z)* and *Su(z)12*, are indicated by the orange box. Identical and similarly charged amino acid residues are marked with asterisk and dots, respectively. Arrows show the corresponding regions used to generate dsRNA templates. The red bar shows corresponding regions used for the quantitative RT-PCR.




**Figure 2. Expression pattern and level of PcG genes in *Gryllus* embryos**

(A,B) Expression patterns of *E(z)* in wild type and *E(z)*<sup>RNAi</sup> embryos at stage (st.) 8. (C) Relative amount of *E(z)* mRNA present in st. 4 and st. 8 embryos. At st. 8, *E(z)* expression levels increased by 89% compared to st. 4. Error bars indicate standard deviation. (D,E) Relative amounts of *E(z)* and *Su(z)12* mRNA in wild type (white bar) and RNAi embryos (gray bar). Scale bar: 200  $\mu$ m.




**Figure 3. Effects of *E(z)* RNAi during *Gryllus* embryogenesis.**

(A) Wild type and (B) *E(z)*<sup>RNAi</sup> embryos ten days after egg laying (AEL). (C) SEM pictures of wild type and (D) *E(z)*<sup>RNAi</sup> dissected embryos. To enhance visibility, the appendages are artificially colored: antenna (red); maxilla (orange); labium (yellow); T1 leg (green); T2 leg (blue); and T3 leg (purple). Scale bars: 200  $\mu$ m.




**Figure 4. Effects of *E(z)* RNAi on embryonic movement in transgenic (pBGact-eGFP; Nakamura et al., 2010) embryos.**

(A–C) Time-lapse recording of embryonic movement in control embryos undergoing katatrepsis from 4 days after egg laying (AEL) to 6 days AEL. (D–F) In *E(z)*<sup>RNAi</sup> embryos, katatrepsis did not occur. Scale bars: 200  $\mu$ m. Abbreviation: H, Head.



**Figure 5. Effects of *E(z)* RNAi on limb specific genes**

(A,B) Expression pattern of *aristaless* (*al*). In wild type (wt) embryos, *al* was expressed in the most distal regions of developing appendages. In developing legs, *al* was expressed in the femur and tibia (A). In *E(z)*<sup>RNAi</sup> embryos, all appendages excluding the mandible showed an *al* expression pattern similar to the limbs of wt embryos (B). (C,D) Expression pattern of *Distal-less* (*Dll*). In wt embryos, *Dll* was expressed in appendages, excluding the mandible (C). The expression pattern was distinct for each appendage. Antenna showed a broad pattern. The maxilla and mandible showed expression in the distal tip. Limbs showed intense expression in distal regions and weak expression in proximal regions. In *E(z)*<sup>RNAi</sup> embryos, all appendages, excluding the mandible, showed homogeneous expression patterns of *Dll* with two distal and one proximal domain (shown by arrowheads) (D). (E,F) Expression patterns of *wingless* (*wg*). In wt embryos, *wg* was expressed along the ventral side of each appendage (E). In *E(z)*<sup>RNAi</sup> embryos, *wg* was expressed on both the ventral and dorsal sides of each limb (F). Abbreviations: An, antenna; Mn, mandible; Mx, maxilla; Lb, labrum; T1–3, thoracic segments 1–3. Scale bar: 200  $\mu$ m.



**Figure 6. *Su(z)12<sup>RNAi</sup>* embryos showed mild misexpression pattern of Hox genes**  
 (A–C) Morphology of late stage embryos halved at the middle line, along the A–P axis. Asterisks indicate the transformed leg-like appendages. (D–F') Expression patterns of *Scr* in wild type (wt), *E(z)<sup>RNAi</sup>* and *Su(z)12<sup>RNAi</sup>* embryos. In wt, *Scr* was expressed in the labium and T1 segment (D,D'). In *E(z)<sup>RNAi</sup>* embryos, *Scr* expression was greatly diminished (E,E'). In *Su(z)12<sup>RNAi</sup>* embryos, *Scr* was expressed in the labium and ectopically expressed in the maxilla (F,F'). (G–I') Expression patterns of *Ubx* in wt, *E(z)<sup>RNAi</sup>*, and *Su(z)12<sup>RNAi</sup>* embryos. In wt, there was no expression of *Ubx* in regions anterior to the T3 segment (G,G'). In *E(z)<sup>RNAi</sup>* embryos, *Ubx* was ectopically expressed in all appendages (H,H'). In *Su(z)12<sup>RNAi</sup>* embryos, *Ubx* expression spread to the anterior regions, except in the maxillae and labium (I,I'). Abbreviations: An, antenna; Mn, mandible; Mx, maxilla; Lb, labrum; T1–3, thoracic segments 1–3. Scale bars: 100  $\mu$ m.

#### 4.2. *Gryllus E(z)* is involved in histone methylation

To elucidate whether the observed  $E(z)^{\text{RNAi}}$  phenotype was caused by changes in histone methylation activity, the spatiotemporal distribution of methylated histone 3 on lysine 27 (H3K27me3) was examined. In wild type *Gryllus*, H3K27me3 signals were observed ubiquitously throughout development (Fig. 7, two left-most columns), starting at the early blastoderm stage and continuing through stage 8. In  $E(z)^{\text{RNAi}}$  embryos, H3K27me3 levels were greatly reduced at every stage, although a slight signal was observed at stages 6–8 (Fig. 7, right side), consistent with RT-qPCR results (supplementary material Fig. 2D). These results indicated that the function of  $E(z)$  in *Gryllus* is to activate and maintain histone methylation, and suggest that the phenotypes observed in Fig. 3B,D may be attributed to changes at the epigenetic level.

#### **Figure 7. Distribution of H3K27me3 in wild type and $E(z)^{\text{RNAi}}$ of eggs and embryos.**

Twelve hours AEL, H3K27me3 was detected only in cells on the egg surface. In embryos stage (st.) 3 to st. 8, the H3K27me3 signal was detected throughout the whole embryo. In  $E(z)^{\text{RNAi}}$  eggs and embryos, the H3K27me3 level was reduced, with a slight recovery at stage 6. White boxes indicate regions of higher magnification. Scale bars: 5000  $\mu\text{m}$  in low magnification, and 500  $\mu\text{m}$  in high magnification.




Figure 7.



### 4.3 $E(z)^{\text{RNAi}}$ embryos show anterior misexpression of Hox genes

To clarify the identity of transformed leg-like structures and determine whether the functions of PcG genes in regulation of Hox genes are conserved in *Gryllus*, the three middle and two posterior Hox genes, *Scr*, *Antp*, *Ubx*, *abd-A* (Zhang et al., 2005), and *Abd-B*, were investigated. Specifically, expression was determined in wild type and  $E(z)^{\text{RNAi}}$  embryos at stage 4 (germband formation stage), stage 5 (just after the initiation of posterior elongation and before A1 and/or A2 are formed), stage 6 (posterior elongation stage and beginning of limb bud formation), and stage 8 (posterior elongation completed). Consistent with the transformation of appendage morphology, all Hox gene patterns were also altered in  $E(z)^{\text{RNAi}}$  embryos (Figs 8 and 9). Similar phenotypes were observed in  $Su(z)12^{\text{RNAi}}$  embryos, although the effects were much milder (Fig. 6).

Initial expression domains of the middle genes, *Scr*, *Antp*, and *Ubx*, were established at stage 4 (Fig. 8A,I,Q), and throughout stage 4 and stage 5 expression domains of these three genes were identical in wild-type and  $E(z)^{\text{RNAi}}$  embryos (Fig. 8). Differences in expression were observed at stage 6 (Fig. 8G,O,W). At stage 6,  $E(z)^{\text{RNAi}}$  embryos misexpressed *Scr* in the prospective mandible segment, while *Scr* expression in the prospective labial segment was significantly reduced (Fig. 8G). *Scr* expression was abolished by stage 8 (Fig. 8H). *Antp* expression was reduced in  $E(z)^{\text{RNAi}}$  embryos at stage 5 (Fig. 8N), but it increased in intensity at stage 6 encompassing the whole embryo (Fig. 8O). Neurogenic expression began at stage 8 with anterior ectopic expression (Fig. 8P). Finally,  $E(z)^{\text{RNAi}}$  embryos expressed *Ubx* in the T1 and T2 limb buds at stage 6 (Fig. 8W), which was followed by expansion into the anterior region by stage 8 (Fig. 8X).

Wild-type expression of the two posterior genes, *abd-A* and *Abd-B*, first appeared at stage 5 and stage 8, respectively (Fig. 9B,K).  $E(z)^{\text{RNAi}}$  embryos exhibited anterior expansion of expression, similar to the middle Hox genes (Fig. 8), although ectopic expression of *abd-A* and *Abd-B* appeared almost simultaneously with wild-type expression. For example, in RNAi embryos, the normal (in A1 and A2, in Fig. 9B) and ectopic expression (in the lateral regions of the gnathal and thoracic segments; Fig. 9F) of *abd-A* appeared simultaneously. Ectopic expression continued to expand and encompassed the anterior head segments during stage 6 and stage 8 (Fig. 9G,H). This trend was even more striking in *Abd-B*; ectopic expression reached its full extent at the

same stage as wild type expression (Fig. 9N).

In summary, knocking down *E(z)* using RNAi results in anterior misexpression of all examined Hox genes, suggesting that Hox genes were epigenetically silenced through H3K27me3 in the anterior region. In addition, there is a temporal difference in the establishment of ectopic expression domains between the middle and posterior Hox genes. The process is stepwise in the former, but it is simultaneous in the latter.

**Figure 8. Effects of *E(z)* RNAi on the expression patterns of middle Hox genes.**

(A–D) Expression patterns of *Scr* in wild type (wt) embryos. Initial expression was restricted to the labrum from stage (st.) 4 to st. 6 (A–C), after which it spread to the T1 segment at st. 8 (D). (E–H) Expression patterns of *Scr* in *E(z)*<sup>RNAi</sup> embryos. Expression was similar to wt at st. 4 (E), then it became weaker as development progressed until it disappeared completely at st. 8 (F–H). At st. 6, anterior misexpression appeared (red bracket in G). (I–L) Expression patterns of *Antp* in wt. *Antp* was strongly expressed in T1–2 at st. 4 (I), then it expanded into the abdominal segments at st. 5 and st. 6 (J,K). At st. 8, the signal was observed in both the epidermis and central nervous system (CNS) of thoracic and abdominal segments (L). (M–P) Expression patterns of *Antp* in *E(z)*<sup>RNAi</sup> embryos. (M) Expression was normal at st. 4, then it was reduced at st. 5 (N), followed by ectopic expression encompassing the whole embryo at st. 6. At st. 8, the signal in the epidermis disappeared completely, but it was maintained in the CNS throughout the A–P axis (P). Anterior ectopic expression is shown by the red brackets in O,P. (Q–T) Expression patterns of *Ubx* in wt. (Q) Expression started in the growth zone at st. 4, then it expanded into T1 and abdominal segments at st. 5–8 (R–T). (U–X) Expression of *Ubx* in *E(z)*<sup>RNAi</sup> embryos. (U,V) The pattern appeared normal until st. 5, after which it expanded into the anterior regions (red brackets in W,X). Abbreviations: Mx: maxilla; Lb: labium; T1–3: thoracic segments 1 to 3; A1: abdominal segment 1; GZ: growth zone. Scale bars: 200  $\mu$ m.




Figure 8.




Figure 9.

**Figure 9. Effects of *E(z)* RNAi on the expression patterns of posterior Hox genes.**


(A–D) Expression patterns of *abd-A* in wild type (wt). Expression started in A1–2 at stage (st.) 5 (B), then it expanded posteriorly to encompass the whole abdomen during st. 6–8 (C,D). (E–H) Expression of *abd-A* in *E(z)*<sup>RNAi</sup> embryos. *abd-A* expression was not be detected at st. 4 (E). At st. 5 and 6, *abd-A* expanded into anterior regions, as depicted with red brackets (F,G). At st. 8, the signal was observed throughout the ventral regions, the head, and thorax, but was absent from the gnathal appendages and legs (H). (I–K) Expression patterns of *Abd-B* in wt. *Abd-B* was not present until posterior elongation was completed (I,J). Expression started at st. 8 and localized to A8–10 (K). (L–N) Expression of *Abd-B* in *E(z)*<sup>RNAi</sup> embryos. *Abd-B* expression was absent at st. 5 and 6, similar to wt (L,M). At st. 8, the signal expanded ventrally into the head region and thorax (N). Abbreviations: A1–10, abdominal segments 1 to 10. Scale bars: 200µm.

#### 4.4 PcG genes and gap genes regulate Hox genes independently

$E(z)^{\text{RNAi}}$  embryos exhibited ectopic expression of *abd-A* in the anterior regions at stage 5 (Fig. 9F). Coincidentally, embryos treated with RNAi directed to the gap gene *hunchback* (*hb*) also exhibit ectopic *abd-A* expression in the prospective gnathal and thoracic segments at stage 5 (Mito et al., 2005). Since PcG genes and *hb* knockdowns altered the regulation of *abd-A* in anterior regions, the genetic relationship between these genes was investigated.

The expression patterns of two gap genes, *hb* and *Krüppel* (*Kr*), were investigated in  $E(z)^{\text{RNAi}}$  embryos (Fig. 10A). Results showed that *hb* was not affected. Expression was confined to the prospective mandibular to labial segments (Mito et al., 2005). *Kr* spatial regulation was also preserved, appearing in the labial to T3 segments (Mito et al., 2006). However, overall expression levels were reduced, possibly due to secondary effects of derepression of other genes (Fig. 10A). In summary, spatial expression of *hb* and *Kr* were not altered due to  $E(z)$  depletion.

Finally, both  $E(z)$  and *hb* were knocked down simultaneously, and the Hox genes were examined (Fig. 10B). When *hb* was targeted with RNAi, the gnathal and thoracic regions were transformed to abdominal identities due to the expansion of *Ubx* and *abd-A* expression into the gnathal regions (Fig. 10B, left column). Interestingly, *Abd-B* remained unaffected, displaying a wild type pattern. In  $hb/E(z)^{\text{RNAi}}$  embryos, however, ectopic expression of *Ubx* and *abd-A* was observed in the anterior-most head region (antennal and ocular segments) as well. Furthermore, *Abd-B* expression expanded into the gnathal segments (compare Fig. 10B, bottom left and right panels). These observations revealed that the combined  $hb/E(z)^{\text{RNAi}}$  embryos exhibited a combined phenotype that resembled the sum of individual *hb* and  $E(z)$  knockdowns, indicating that there is no genetic interaction between *hb* and  $E(z)$ .



**Figure 10. *E(z)* and *hb* regulate posterior Hox genes expression independently.**


(A) Expression patterns of *hb* and *Kr* were not altered in *E(z)*<sup>RNAi</sup> embryos at stage (st.) 5. (B) In *hb*<sup>RNAi</sup> embryos, *Ubx* and *abd-A* were ectopically expressed in the gnathal and thoracic segments, while the *Abd-B* pattern was not altered. In *E(z)/hb*<sup>RNAi</sup> embryos, *Ubx* and *abd-A* were aberrantly expressed in the head regions. The same trend was observed for *Abd-B*, which was ectopically expressed in the gnathal and thoracic segments. Abbreviations: An, antenna; Lb, labium; T3, thoracic segment 3; Scale bars: 100  $\mu$ m.

#### 4.5 Regional autonomy in the PcG silencing of Hox genes

To elucidate whether temporal PcG-mediated silencing of Hox genes was related to posterior elongation of the embryo, the *caudal* (*cad*) gene was analyzed. Previous studies have shown that *cad*<sup>RNAi</sup> embryos lacked posterior segments (Shinmyo et al., 2005), revealing that *cad* is essential for posterior patterning in *Gryllus*. Consistent with previous experiments, the expression of *Ubx*, *abd-A*, and *Abd-B* in *cad*<sup>RNAi</sup> embryos was determined (Fig. 11). Both mild and severe phenotypes were observed. In mild phenotypes, anterior regions were formed normally, whereas the regions posterior from T2 were missing. In those embryos, *Ubx*, *abd-A*, and *Abd-B* expression was reduced in the remaining posterior regions (Fig. 11A–C). In severe phenotypes, embryos showed only anterior head morphology. Furthermore, the expression of all three genes (*Ubx*, *abd-A*, and *Abd-B*) could not be detected (Fig. 11G–I), suggesting that the growth zone activity was completely abolished in *cad*<sup>RNAi</sup> embryos.

Treatment with RNAi directed against both *cad* and *E(z)*, produced *cad/E(z)*<sup>RNAi</sup> embryos, which displayed a phenotype that is a combination of individual *cad* and *E(z)* depleted embryos. In mild phenotypes, *Ubx*, *abd-A*, and *Abd-B* exhibited anterior expansion compared to single *E(z)* RNAi embryos (Fig. 11D–F). Remarkably, this trend became even stronger in the severe phenotype. Indeed, while none of Hox genes were expressed in insects treated with the single *cad* RNAi, they all become overexpressed in the double RNAi mutant (Fig. 11J–L). These observations indicated that PcG-mediated silencing of Hox genes was region-specific and stage-specific, with silencing in the anterior regions occurring independently of posterior region development.





**Figure 11. Effects of *E(z)* and *cad* RNAi on the expression pattern of posterior Hox genes.**

Both mild and severe phenotypes were observed. (A–C) Mild phenotypes of *cad*<sup>RNAi</sup> embryos. Expression of *Ubx* (A), *abd-A* (B) and *Abd-B* (C) was detected in regions posterior to the T2 segment. (D–F) Mild phenotypes of *cad/E(z)*<sup>RNAi</sup> embryos. *Ubx* (D), *abd-A* (E), and *Abd-B* (F) were ubiquitously expressed in the whole embryo, consistent with patterns observed in *E(z)*<sup>RNAi</sup> embryos (Fig. 4). (G–I) Severe phenotypes of *cad*<sup>RNAi</sup> embryos. (J–L) Severe phenotypes of *cad/E(z)*<sup>RNAi</sup> embryos. Expression of *Ubx* (J), *abd-A* (K), and *Abd-B* (L) showed anterior expansion. Abbreviations: An, antenna; Mx, maxilla; Lb, labium; T1 and 2, thoracic segments 1 and 2. Scale bar: 100  $\mu$ m.

## 5. Discussion

### Functional conservation of the PRC2 complex in *Gryllus*

In the present study, *Gryllus* embryos treated with RNAi directed against *E(z)* displayed reduced H3K27me3 signals during embryogenesis (Fig. 7), suggesting that the *E(z)* protein plays an important role in histone methylation activity of the PRC2 complex. These observations are consistent with previous work in *Drosophila* (Diptera) and *Bombyx* (Lepidoptera) (Tie et al., 2009; Li et al., 2012). The phenotypes of insects treated with RNAi directed against *Su(z)12*, another component of the PRC2 complex, were similar to *E(z)* RNAi phenotypes, supporting the model of structural and functional conservation of the entire PRC2 complex in *Gryllus* (Orthoptera). Thus, my results strengthen the view that epigenetic regulation machinery involving PcG complexes may be shared among all insect lineages.

There was also a noticeable difference in the developmental dynamics of H3K27me3 in *Gryllus*, which was first detectable at the syncytial blastoderm stage before the germ anlage was formed (12 h AEL; corresponding to stage 4 in *Drosophila*) (Fig. 7). By comparison, H3K27me3 was first detected in *Drosophila* after germband formation at stages 9–11 (Tie et al., 2009). At present, I do not know how *E(z)*-mediated temporal differences in H3K27me3 could affect early embryo development. As indicated by data (Fig. 4), one possibility is that H3K27me3 in the early blastoderm may be involved in proper development of extraembryonic tissues and dictate katatrepsis defects observed in *E(z)*<sup>RNAi</sup> embryos.

### Mode of establishing expression domains varies among *Grylls* Hox genes

As demonstrated in Figs 8 and 9, treatment of *Gryllus* with *E(z)* RNAi resulted in anterior misexpression of *Scr*, *Antp*, *Ubx*, *abd-A*, and *Abd-B*, suggesting that these Hox genes are epigenetically silenced by PcG in the region anterior to each of their individual expression domains. This anterior silencing may be essential for providing proper segment identities in embryos, as the *E(z)*<sup>RNAi</sup> phenotype exhibited a homeotic transformation of head appendages, with the exception of mandibles, into leg-like structures (Fig. 3). However, while anterior misexpression of those same Hox genes was reported in *E(z)* loss-of-function mutants in *Drosophila*, mutant embryos displayed very different phenotypes; all trunk segments acquired the A8 identity (Jones and Gelbart, 1990). This discrepancy may be due to the fact that ectopic expression of

*Abd-B* in *Gryllus* occurs at a germband stage with well-developed leg buds (stage 8), later than the extended-germband stage in *Drosophila* (Simon et al., 1992). This may prevent the inhibition of leg development in *Gryllus*. In addition, ectopic expression of *Scr*, *Antp*, and *Ubx* in early and late appendage buds may activate the leg development program, leading to leg-like transformation of head appendages.

In addition to the different morphological phenotypes of *E(z)* loss-of-function mutants, significant differences in the patterns of Hox gene expression were observed. In *Gryllus E(z)<sup>RNAi</sup>* embryos, *Scr* expression was lost at stage 8 (Fig. 8H). On the other hand, *Drosophila E(z)* mutant embryos exhibited ectopic expression of *Scr* in the whole embryo at late stages (Soto et al., 1995). It is possible that *Scr* expression in *Gryllus* is repressed by *Ubx*, because *Ubx* is misexpressed in the whole body of late stage *E(z)<sup>RNAi</sup>* embryos, overlapping potential regions of *Scr* expression. This regulatory relationship may not be conserved in *Drosophila*, suggesting that there may be differences in transcriptional regulation of Hox genes between *Gryllus* and *Drosophila*.

This study revealed another critical difference in PcG silencing of posterior Hox genes (*abd-A* and *Abd-B*) between *Drosophila* and *Gryllus*. In *Drosophila*, the repression by gap genes determines the anterior expression boundary of each Hox gene (in both Antennapedia and Bithorax complexes). Subsequently, the role of a gap gene is replaced by epigenetic silencing from PcG genes, which, in turn, maintains anterior borders (Simon et al., 1992; Struhl and Akam, 1985; Jones and Gelbart, 1990). Such a stepwise change in the Hox repression system may also apply to regulation of *Scr*, *Antp*, and *Ubx* in *Gryllus*, as indicated by anterior expansion of expression in *E(z)<sup>RNAi</sup>* embryos (Fig. 8). On the other hand, *Gryllus abd-A* and *Abd-B* were misexpressed at the stage when normal expression appears (Fig. 9), suggesting that PcG silencing is involved in establishing expression domains of these genes. Indeed, *hb* represses *abd-A* expression in anterior (gnathal and thoracic) regions (Mito et al., 2005), as expected if a gap gene is involved in establishing the initial Hox expression domain. However, as shown by embryos treated with RNAi against *hb* and *E(z)*, both genes seemed to act in parallel, not in a stepwise manner (Fig. 10). In addition, *Abd-B* is normally activated after completion of segmentation, when gap gene activity cannot be assumed.

The transcription of *Gryllus* posterior Hox genes seems to be activated during or after posterior embryo elongation and is epigenetically silenced by PcG in the regions outside of their normal expression as a way of establishing their anterior expression

boundaries. This mechanism is reminiscent of Hox gene regulation in vertebrates. The initial state of the vertebrate Hox gene cluster is “closed (H3K27me3 positive)” via PcG silencing and subsequently “opened (H3K27me3 negative)” in a temporal manner to induce the expression of a particular Hox gene. Thus, PcG silencing is also required for establishment of the initial expression domains (Soshnikova and Duboule, 2009). This similarity between vertebrates and *Gryllus* with regard to dependence on PcG silencing in establishment of Hox expression domains suggests that this mode may represent the ancestral state in insects.

Activation of *Abd-B* in *Gryllus* may also be regulated in the manner similar to vertebrates. Indeed, this locus may also be silenced by PcG prior to transcription and subsequently turned into an active state when temporal and spatial patterns dictate. It should be noted, however, that inhibition of posterior elongation in *Gryllus* did not affect ectopic expression in the remaining embryonic regions (Fig. 11), implying the existence of autonomous mechanisms for Hox gene silencing and induction of transcriptional activators. Such a transcriptional activator may be upregulated throughout the wild type embryo at a specific developmental stage where Hox genes are expressed. This differs from vertebrates, in which intercellular signals, such as retinoic acid gradients, control where a specific Hox gene is activated or repressed (Kiecker and Lumsden, 2012). Alternatively, the *Gryllus Abd-B* locus might be kept “open” in prospective *Abd-B*-expressing cells throughout posterior embryo elongation until its activation stage, while in more anterior regions the locus might be “closed” prior to *Abd-B* activation. In this case, signals from the posterior growth zone, such as Wnt (Miyawaki et al., 2004), might be involved in inhibiting PcG silencing according to an activity gradient during posterior elongation.

### **Evolutionary transition of the PcG silencing mode of Hox genes**

Insights from the present and previous studies in vertebrates, *Drosophila*, and *Gryllus*, suggest that an intriguing transition of PcG-mediated silencing of Hox genes occurred during animal evolution (Montavon and Duboule, 2013; Bantignies and Cavalli, 2006). As illustrated in Fig. 12, the ancestral bilaterian state may have resembled the current vertebrate mode, where PcG-mediated silencing of Hox genes occurs before Hox gene expression is initiated, establishing individual expression domains. Then, during insect evolution, repression by transcription factors may have been acquired in anterior Hox

genes of short germ insects, while PcG silencing was maintained in posterior Hox genes. During long germ insect evolution, the involvement of transcription factors may have spread to encompass the posterior Hox genes, resulting in the stepwise repression governed by gap and PcG genes that is observed in present day dipterans.

**Figure 12. Model of Hox gene regulation machinery during bilaterian evolution.**

A diagram illustrating the regulation of Hox genes during the development of vertebrates, short germ insects (*Gryllus*), and long germ insects (*Drosophila*). In vertebrates, regulation of all Hox genes is controlled by the PcG complex. During protostome evolution, the introduction of transcription factors (TFs; e.g. gap genes) in short germ insects allowed for the regulation of anterior Hox loci, prior to PcG activity. During the evolution of holometabolous lineages, such as dipterans, all Hox gene loci became regulated by TFs, resulting in apparent stepwise regulation by gap genes and PcG. Different shades of green boxes indicate Hox gene loci. Gray ovals represent PcG repressed chromatin. Orange arrows indicate the start of gene transcription.




Figure 12.

## 6. Experimental procedures

### Animals

*Gryllus bimaculatus* nymphs and adults were reared at 28–30 °C with 70% humidity under a 10 light, 14 dark photoperiod, as previously described (Niwa et al., 1997). Fertilized eggs were collected with wet kitchen towels and incubated at 28 °C in a plastic dish. Embryos were staged as described previously (Zhang et al., 2005).

### Cloning of the *Gryllus bimaculatus* *E(z)* and *Su(z)12*

Partial nucleotide sequences of *Gryllus bimaculatus* *E(z)* and *Su(z)12* were identified from cDNA obtained from adult ovaries (Zeng et al., 2013). Partial fragments of *E(z)* and *Su(z)12* were PCR-amplified with gene specific primers and used as double-stranded RNA (dsRNA) templates. The primer sequences for *E(z)*<sub>N</sub> and *E(z)*<sub>C</sub> (see Fig. 1) were 5'-ATACTTGGGCACCAATCCAA-3' (forward) and 5'-TTCTTCTGGCCTCCCCTTAT-3' (reverse) and 5'-CTTGGAGTGGA-CTGCACTGA-3' (forward) and 5'-CTCGCACAGCAAGATAGCAG-3' (reverse), respectively. Primer sequences for *Su(z)12* are 5'-ATTGAAACGCACCAAGAACC-3' (forward) and 5'-ATGGGCCACATTCAAGGTAA-3' (reverse). The cDNA sequences of *Gryllus* *E(z)* and *Su(z)12* were deposited in the DNA Data Bank of Japan (DDBJ) (accession numbers: AB378079; LC005751).

### Parental RNAi

Cloned partial cDNAs of *E(z)* and *Su(z)12* were used as templates to prepare dsRNAs. The MEGAScript Kit (Ambion) was used for in vitro synthesis of dsRNAs, which were adjusted to a concentration of 20 µM. Parental RNAi treatment was performed by injecting dsRNAs into the body cavity of adult female crickets as described previously (Mito et al., 2005).

### Embryo fixation, whole mount *in situ* hybridization, and immunohistochemistry

Embryo fixation and whole-mount *in situ* hybridization with digoxigenin (DIG)-labeled

antisense RNA probes were performed as previously described (Niwa et al., 2000; Zhang et al., 2005). For immunohistochemistry, embryos were fixed with the same methods as *in situ* hybridization (Niwa et al., 2000). Fixed embryos were rehydrated stepwise in 75%, 50%, and 25% solutions of methanol/phosphate-buffered saline + 0.1% Tween (PBT) and PBT for 5 minutes each. Next, embryos were incubated for 1 hour in 1% bovine serum albumin (BSA) (Invitrogen)/PBT at room temperature and incubated with a rabbit polyclonal anti-trimethylated H3K27 antibody (Millipore 07-449) diluted 1 to 500 in 1% BSA/PBT overnight at 4 °C. After washing with PBT three times, embryos were incubated in 1% BSA/PBT for 1 hour at room temperature and then with Alexa Fluor 488 Goat Anti- mouse IgG(H+L) (Invitrogen) diluted 1 to 500 dilution in 1% BSA/PBT for 1 hour. After washing with PBT three times, embryos were counter stained with DAPI (Sigma) diluted 1 to 1000 in PBT for 10 minutes and then washed with PBT three times. PBT was then substituted by 25% and 50% glycerol/PBT to make embryos transparent for microscopy.

### **Real Time quantitative PCR**

Total RNA was extracted from embryos using ISOGEN (Nippon-Gene). After treatment with DNaseI (Invitrogen), RNA was reverse-transcribed to cDNA using SuperScriptIII reverse transcriptase (Invitrogen). Real-time quantitative PCR was performed using the power SYBR Green PCR Master Kit (Applied Biosystems) and an ABI 7900 Real Time PCR System (Applied Biosystems), as described previously (Nakamura et al., 2008). Primer pairs were: E(z)Fw; 5'-GGCAGATG- GCAAAGCAGTGT-3' and E(z)Rv; 5'-CTTCATGCAGGCAGCATGA- 3'; Su(z)12Fw; 5'-ACCGGTGGTGGTGTGTA CTT-3' and Su(z)12Rv; 5'-GCGATAAATCTGCGTTGGTT-3'.



## 7. References

- Akasaka, T., Kanno, M., Balling, R., Mieza, M.A., Taniguchi, M. and Koseki, H.** A role for *mel-18*, a Polycomb group-related vertebrate gene, during theanteroposterior specification of the axial skeleton. *Development* **122**, 1513-1522.
- Agger, K., Cloos, P. A. C., Christensen, J., Pasini, D., Rose, S., Rappsilber, J., Issaeva, I., Canaani, E., Salcini, A. E. and Helin, K.** (2007). UTX and JMJD3 are histone H3K27 demethylases involved in *HOX* gene regulation and development. *Nature* **449**, 731-735.
- Akam, M.** (1987). The molecular basis for metameric pattern in the *Drosophila* embryo. *Development* **101**, 1-22.
- Bae, E., Calhoun, V. C., Levine, M., Lewis, E. B. and Drewell, R. A.** (2002). Characterization of the intergenic RNA profile at *abdominal-A* and *Abdominal-B* in the *Drosophila* bithorax complex. *Proc. Natl. Acad. Sci. USA*. **99**, 16837-16852.
- Bantignies, F. and Cavalli, G.** (2006). Cellular memory and dynamic regulation of polycomb group proteins. *Curr. Opin. Cell. Biol.* **18**, 275-283.
- Birve, A., Sengupta, A. K., Beuchle, D., Larsson, J., Kennison, J. A., Rasmuson-Lestander, Å. and Müller, J.** (2001). *Su(z)12*, a novel *Drosophila* Polycomb group gene that is conserved in vertebrates and plants. *Development* **128**, 3371-3379.
- Castelli-Gair, J.** (1998). Implications of the spatial and temporal regulation of *Hox* genes on development and evolution. *Int. J. Dev. Biol.* **42**, 437-444.
- Davis, G. K. and Patel, N. H.** (2002). Short, long, and beyond: molecular and embryological approaches to insect segmentation. *Annu. Rev. Entomol.* **47**, 669-699.
- Grimaud, C., Nègre, N. and Cavalli, G.** (2006). From genetics to epigenetics: the tale of Polycomb group and trithorax group genes. *Chromosome Res.* **14**, 363-375.
- Gieni, R.S. and Hendzel, M.J.** (2009). Polycomb group protein gene silencing, non-coding RNA, stem cells, and cancer. *Biochem. Cell Biol.* **87**, 711-746.
- Harding, K. and Levine, M.** (1988). Gap genes define the limits of Antennapedia and bithorax gene expression during early development in *Drosophila*. *EMBO J.* **7**, 205-214.
- Irish, V. F., Martinez-Arias, A. and Akam, M.** (1989). Spatial regulation of the *Antennapedia* and *Ultrabithorax* homeotic genes during *Drosophila* early development. *EMBO J.* **8**, 1527-1537.
- Jaeger, J.** (2011). The gap gene network. *Cell. Mol. Life Sci.* **68**, 243–274.

- Jones, R. S. and Gelbart, W. M.** (1990). Genetic analysis of the *Enhancer of zeste* locus and its role in gene regulation in *Drosophila melanogaster*. *Genetics* **126**, 185-199.
- Kelsh, R., Dawson, I. and Akam, M.** (1993). An analysis of Abdominal-B expression in the locust *Schistocerca gregaria*. *Development* **117**, 293-305.
- Kiecker, C. and Lumsden, A.** (2012). The role of organizers in patterning the nervous system. *Annu. Rev. Neurosci.* **35**, 347-367.
- Lee, M. G., Villa, R., Trojer, P., Norman, J., Yan, K., Reinberg, D., Croce, L. D. and Shiekhatar, R.** (2007). Demethylation of H3K27 Regulates Polycomb Recruitment and H2A Ubiquitination. *Science* **318**, 447-450.
- Lewis, E. B.** (1978). A gene complex controlling segmentation in *Drosophila*. *Nature* **276**, 565-570.
- Li, Z., Tatsuke, T., Sakashita, K., Zhu, L., Xu, J., Mon, H., Lee, J. M. and Kusakabe, T.** (2011). Identification and characterization of Polycomb group genes in the silkworm, *Bombyx mori*. *Mol. Biol. Rep.* **39**, 5575-5588.
- Martin, B. L. and Kimelman, D.** (2009). Wnt signaling and the evolution of embryonic posterior development. *Curr. Biol.* **19**, 215-219.
- Mito, T., Sarashina, I., Zhang, H., Iwahashi, A., Okamoto, H., Miyawaki, K., Shinmyo, Y., Ohuchi, H. and Noji, S.** (2005). Non-canonical functions of *hunchback* in segment patterning of the intermediate germ cricket *Gryllus bimaculatus*. *Development* **132**, 2069-2079.
- Mito, T., Okamoto, H., Shinahara, W., Shinmyo, Y., Miyawaki, K., Ohuchi, H. and Noji, S.** (2006). *Krüppel* acts as a gap gene regulating expression of *hunchback* and *even-skipped* in the intermediate germ cricket *Gryllus bimaculatus*. *Dev. Biol.* **294**, 471-481.
- Mito, T., Kobayashi, C., Sarashina, I., Zhang, H., Shinahara, W., Miyawaki, K., Shinmyo, Y., Ohuchi, H. and Noji, S.** (2007). *even-skipped* has gap-like, pair-rule-like, and segmental functions in the cricket *Gryllus bimaculatus*, a basal, intermediate germ insect (Orthoptera). *Dev. Biol.* **303**, 202-213.
- Miyawaki, K., Inoue, Y., Taro, M., Fujimoto, T., Matsushima, K., Shinmyo, Y., Ohuchi, H. and Noji, S.** (2002). Expression patterns of *aristaless* in developing appendages of *Gryllus bimaculatus* (cricket). *Mech. Dev.* **113**, 181-184.

- Miyawaki, M., Mito, T., Sarashina, I., Zhang, H., Shinmyo, Y., Ohuchi, H. and Noji, S.** (2004). Involvement of Wingless/Armadillo signaling in the posterior sequential segmentation in the cricket, *Gryllus bimaculatus* (Orthoptera), as revealed by RNAi analysis. *Mech. Dev.* **121**, 119-130.
- Mohd-Sarip, A., Venturini, F., Chalkley, G.E. and Verrijzer, C.P.** (2002). Pleiohomeotic can link polycomb to DNA and mediate transcriptional repression. *Mol. Cell Biol.* **22**, 7473-83.
- Montavon, T. and Duboule, D.** (2013). Chromatin organization and global regulation of Hox gene clusters. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **368**, 20120367.
- Müller, J. and Kassis, J. A.** (2006). Polycomb response elements and targeting of Polycomb group proteins in *Drosophila*. *Curr. Opin. Genet. Dev.* **16**, 476-484.
- Nakamura, T., Mito, T., Miyawaki, K., Ohuchi, H. and Noji, S.** (2008). EGFR signaling is required for re-establishing the proximodistal axis during distal leg regeneration in the cricket *Gryllus bimaculatus* nymph. *Dev. Biol.* **319**, 46-55.
- Nakamura, T., Yoshizaki, M., Ogawa, S., Okamoto, H., Shinmyo, Y., Bando, T., Ohuchi, H., Noji, S. and Mito, T.** (2010). Imaging of transgenic cricket embryos reveals cell movements consistent with a syncytial patterning mechanism. *Curr. Biol.* **20**, 1641-1647.
- Niwa, N., Saitoh, M., Ohuchi, H., Yoshioka, H. and Noji, S.** (1997). Correlation between *Distal-less* expression patterns and structures of appendages in development of the two-spotted cricket, *Gryllus bimaculatus*. *Zool. Sci.* **14**, 115-125.
- Niwa, N., Inoue, Y., Nozawa, A., Saitoh, M., Misumi, Y., Ohuchi, H., Yoshioka, H. and Noji, S.** (2000). Correlation of diversity of leg morphology in *Gryllus bimaculatus* (cricket) with divergence in *dpp* expression pattern during leg development. *Development* **127**, 4373-4381.
- Pelegri, F. and Lehmann, R.** (1994). A Role of Polycomb Group Genes in the Regulation of Gap Gene Expression in *Drosophila*. *Genetics* **136**, 1341-1353.
- Peterson, M. D., Rogers, B. T., Popadi, A. and Kaufman, T. C.** (1999). The embryonic expression pattern of *labial*, posterior homeotic complex genes and the *teashirt* homologue in an apterygote insect. *Dev. Genes. Evol.* **209**, 77-90.
- Qian, S., CapoviHa, M. and Pirotta, V.** (1991). The *bx* region enhancer, a distal *cis*-control element of the *Drosophila Ubx* gene and its regulation by *hunchback* and other segmentation genes. *EMBO J.* **10**, 1415-1425.

- Reinltz, J. and Levine, M.** (1990). Control of the initiation of homeotic gene expression by the gap genes *giant* and *tailless* in *Drosophila*. *Dev. Biol.* **140**, 57-72.
- Schuettengruber, B., Chourrout, D., Vervoort, M., Leblanc, B. and Cavalli, G.** (2007). Genome regulation by polycomb and trithorax proteins. *Cell* **128**, 735-745.
- Shinmyo, Y., Mito, T., Matsushita, T., Sarashina, I., Miyawaki, K., Ohuchi, H. and Noji, S.** (2005). *caudal* is required for gnathal and thoracic patterning and for posterior elongation in the intermediate-germband cricket *Gryllus bimaculatus*. *Mech. Dev.* **122**, 231-239.
- Shippy, T. D., Brown, S. J. and Denell, R. E.** (1998). Molecular characterization of the *Tribolium abdominal-A* ortholog and implications for the products of the *Drosophila* gene. *Dev. Genes. Evol.* **207**, 446-452.
- Simon, J., Chiang, A. and Bender, W.** (1992). Ten different *Polycomb* group genes are required for spatial control of the *abdB* and *AbdB* homeotic products. *Development* **114**, 493-505.
- Simon, J. A. and Kingston, R. E.** (2009). Mechanisms of Polycomb gene silencing: knowns and unknowns. *Nat. Rev. Mol. Cell Biol.* **10**, 697-708.
- Simon, J. A. and Kingston, R. E.** (2013). Occupying chromatin: Polycomb mechanisms for getting to genomic targets, stopping transcriptional traffic, and staying put. *Mol. Cell.* **49**, 808-824.
- Soshnikova, N. and Duboule, D.** (2009). Epigenetic regulation of vertebrate *Hox* genes. *Epigenetics* **16**, 537-540.
- Soto, M., Chou, T. and Bender, W.** (1995). Comparison of Germline Mosaics of Genes in the Polycomb Group of *Drosophila melanogaster*. *Genetics* **140**, 231-243.
- Struhl, G. and Akam, M. E.** (1985). Altered distributions of *Ultrabithorax* transcripts in *extra sex combs* mutant embryos of *Drosophila*. *EMBO J.* **4**, 3259-3264.
- Tear, G., Akam, M. and Martinez-Arias, A.** (1990). Isolation of an *abdominal-A* gene from the locust *Schistocerca gregaria* and its expression during early embryogenesis. *Development* **110**, 915-925.
- Tie, F., Banerjee, R., Stratton, C. A., Prasad-Sinha, J., Stepanik, V., Zlobin, A., Diaz, M. O., Scacheri, P. C. and Harte, P. J.** (2009) CBP-mediated acetylation of histone H3 lysine 27 antagonizes *Drosophila* Polycomb silencing. *Development* **136**, 3131-3141.
- van der Lugt, N.M., Domen, L., Linders, K., van Roon, M., Robanus-Maandag, E.,**

- te Riele, H., van der Valk, M., Deschamps, J., Sofroniew, M., van Lohuizen, M. and Berns, A.** (1994). Posterior transformation, neurological abnormalities, and severe hematopoietic defects in mice with a targeted deletion of the *bmi-1* proto-oncogene. *Genes Dev.* **8**, 757-769.
- White, R. A. H. and Lehmann, R.** (1986). A gap gene, *hunchback*, regulates the spatial expression of *Ultrabithorax*. *Cell* **47**, 311-321.
- Zeng, V., Ewen-Campen, B., Horch, H. W., Roth, S., Mito, T. and Extavour, C. G.** (2013). Developmental Gene Discovery in a Hemimetabolous Insect: *De Novo* Assembly and Annotation of a Transcriptome for the Cricket *Gryllus bimaculatus*. *PLoS One* **8**, e61479.
- Zhang, H., Shinmyo, Y., Mito, T., Miyawaki, K., Sarashina, I., Ohuchi, H. and Noji, S.** (2005). Expression patterns of the homeotic genes *Scr*, *Antp*, *Ubx*, and *abd-A* during embryogenesis of the cricket *Gryllus bimaculatus*. *Gene Expr. Patterns* **5**, 491–502.

## **8. Acknowledgement**

I greatly appreciate Prof. Sumihare Noji for helpful advices and great condition for research. And I also thank Assistant Prof. Taro Mito, Associate Prof. Aleksandar Popadić , Dr. Tetsuya Bando, Dr. Taro Nakamura, Dr. Takahito Watanabe and Dr. Yoshiyasu Ishimaru for helpful discussion and comments on the manuscript.

## 9. Publications

1. **Matsuoka, Y., Bando, T., Watanabe, T., Ishimaru, Y., Noji, S., Popadic, A., and Mito, T.** (2015). Short germ insects utilize both the ancestral and derived mode of Polycomb group-mediated epigenetic silencing of Hox genes. *Biol. Open* **5**, 702-709.
2. **Ishimaru, Y., Nakamura, T., Bando, T., Matsuoka, Y., Ohuchi, H., Noji, S., and Mito, T.** (2015). Involvement of *dachshund* and *Distal-less* in distal pattern formation of the cricket leg during regeneration. *Sci. Rep.* **5** : 8387.
3. **Bando, T., Ishimaru, Y., Kida, T., Hamada, Y., Matsuoka, Y., Nakamura, T., Ohuchi, H., Noji, S., and Mito, T.** (2013). Analysis of RNA-Seq data reveals involvement of JAK/STAT signalling during leg regeneration in the cricket *Gryllus bimaculatus*. *Development*, **140**, 959-964.