

## ORIGINAL

# Simultaneous detection of *Bacteroides fragilis* group species by *leuB* -directed PCR

Tsuyoshi Miki, Tomomi Kuwahara\*, Haruyuki Nakayama, Natsumi Okada, Keiko Kataoka, Hideki Arimochi, and Yoshinari Ohnishi

*Department of Molecular Bacteriology, Institute of Health Biosciences, The University of Tokushima Graduate School, Tokushima, Japan*

**Abstract :** *Bacteroides* species, saccharolytic Gram-negative obligate anaerobes, are frequently isolated from human infections such as peritonitis, abscesses and bacteremia. Among the species in the genus *Bacteroides*, the species called “*B. fragilis* group” are particularly involved in human infections and are medically important because they account for a major part of anaerobic isolates from clinical specimens. The purpose of this study was to develop PCR primers that specifically and simultaneously amplify the *leuB* gene, which encodes L-isopropylmalate dehydrogenase in *B. fragilis* group species. We determined partial nucleotide sequences of *leuB* genes and compared them in seventeen strains of nine *B. fragilis* group species, and the regions that are conserved among *Bacteroides* strains but different from other species were used as a *B. fragilis* group-specific PCR primer set, BacLBF-BacLBR. Specificity tests of the primer set using 52 phenotypically characterized strains and 75 isolates from rat feces showed only one case each of false-positive and false-negative. The detection limit of the *leuB*-directed PCR using BacLBF and BacLBR was  $3.9 \times 10^3$  colony-forming units. These results indicate that *leuB* amplification using BacLBF and BacLBR is a useful tool for rapid diagnosis of *Bacteroides* infection and for rapid differential diagnosis of anaerobic infections. *J. Med. Invest.* 52 : 101-108, February, 2005

**Keywords :** *Bacteroides*, *leuB*, PCR, rapid detection

## INTRODUCTION

*Bacteroides* are saccharolytic Gram-negative obligate anaerobes that are most predominant in the human colon (1). The representative species of the genus *Bacteroides* is *B. fragilis*, and ten species that are phylogenetically close to *B. fragilis* have been defined as “*B. fragilis* group” (2). The *B. fragilis* group species are frequently isolated from mixed aerobic-anaerobic infections such as intra-abdominal, diabetic foot and soft tissue infections, though their virulence potentials are

different. *B. fragilis* is regarded as the most virulent species among *Bacteroides* species and often causes severe bacteremia (3, 4). Although the growth rate of *Bacteroides* species is faster than that of other obligate anaerobes, identification of *Bacteroides* species by the traditional anaerobic culture method based on carbohydrate fermentation patterns is laborious, and use of this procedure sometimes results in misidentification. These conditions often delay the useful application of species identification and susceptibility data for patients. Therefore, PCR-based rapid and accurate identification methods in which specific genes such as glutamine synthetase gene (5) and neuraminidase gene (6, 7) are amplified have been developed for *B. fragilis*. Results obtained by using these methods showed that these methods have sufficient levels of specificity and sensitivity to be used for clinical speci-

Received for publication November 30, 2004 ; accepted January 7, 2005.

Address correspondence and reprint requests to Tomomi Kuwahara\*, Department of Molecular Bacteriology, Institute of Health Biosciences, The University of Tokushima Graduate School, Kuramotocho, Tokushima 770-8503, Japan and Fax : +81-88-633-7069.

mens but that they provide limited information when species other than *B. fragilis* are contained in the specimens. A recent study on isolation rates of *B. fragilis* group species and antimicrobial susceptibility patterns demonstrated that the isolation rate has changed in the past ten years and that the increase in non-*B. fragilis* species has influenced the overall antimicrobial susceptibility of *Bacteroides* isolates (8). It is therefore important to develop a rapid and accurate method for identifying non-*B. fragilis* species for empirical antimicrobial therapy used by clinicians for treatment of mixed aerobic-anaerobic infections.

The aim of this study was to develop a PCR-based method for the simultaneous detection of all *B. fragilis* group species. For this purpose, we chose  $\beta$ -isopropylmalate dehydrogenase gene (*leuB*), which is involved in leucine biosynthesis, as a target because *B. fragilis* group species are leucine prototrophs and thought to have a complete set of genes for the leucine biosynthetic pathway (9).

## MATERIALS AND METHODS

### Bacterial strains

Bacterial strains used in this study are as follows. The anaerobes used were 9 strains of *B. fragilis* (2 laboratory strains, ATCC25285 and ATCC43859, and 7 clinical isolates), 6 strains each of *B. distasonis* (ATCC8503 and 5 clinical isolates) and *B. vulgatus* (ATCC8482 and 5 clinical isolates), 5 strains of *B. thetaiotaomicron* (ATCC 29148 and 4 clinical isolates), 4 strains of *B. uniformis* (2 laboratory strains, ATCC8492 and BU1001, and 2 clinical isolates), 3 strains of *B. ovatus* (ATCC8483 and 2 clinical isolates), and one strain each of *B. caccae* (JCM 9498), *B. eggerthii* (ATCC27754), *B. merdae* (JCM9497), *B. stercoris* (JCM9496), *Porphyromonas gingivalis* (381), *Por. asaccharolytica* (NCTC9337), *Por. endodontalis* (ATCC 35406), *Prevotella corporis* (JCM8480), *Prevotella melaninogenica* (ATCC25845), *Clostridium perfringens* (GAI 0668), *Eubacterium limosum* (ATCC8480), and *Fusobacterium nucleatum* (F-1). The aerobes used in this

study were one strain each of *Bacillus subtilis* (T), *Staphylococcus aureus* (209P), *Staphylococcus epidermidis* (IID 866), *Streptococcus pyogenes* (UTB111), *Klebsiella pneumoniae* (KUB13), *Enterobacter aerogenes* (UTB7), *Proteus vulgaris* (KUB22), *Serratia marcescens* (UTB66), *Pseudomonas aeruginosa* (UTC55), and *Escherichia coli* (W3110). All anaerobic bacteria were cultured on GAM agar plates (Nissui Pharmaceutical Co., Tokyo, Japan) under anaerobic conditions. Aerobes except for *Streptococcus pyogenes* were cultured on Luria-Bertani (LB) agar plates (10). *Streptococcus pyogenes* was cultured on a 5% sheep blood agar plate.

### Preparation of bacterial cells for PCR

Several colonies of each strain grown on GAM, LB, or 5% sheep blood agar plates were suspended in 1.0 ml of TE buffer (10 mM Tris-HCl [pH 8.0], 1 mM EDTA) and heated at 100 °C for 10 min. After centrifugation at 12,000 g for 1 min, 10  $\mu$ l of the supernatant was collected and used as a template for the PCR amplification of *leuB*. The mid-logarithmic phase culture (1.0 ml) of *B. fragilis* strain ATCC25285 in GAM broth was centrifuged, washed in 1.0 ml of TE buffer, and resuspended in 1.0 ml of TE. Serial 10-fold dilutions were prepared to give appropriate colony forming units (CFU), and they were lysed by heating at 100 °C for 10 min. The supernatant (10  $\mu$ l) of each dilution after centrifugation at 12,000 g for 1 min was used as a template to determine the sensitivity of *leuB*-PCR

### PCR amplification and partial sequencing of *leuB* genes from *B. fragilis* group species

Oligonucleotide primers used in this study are listed in Table 1. Three kinds of primer sets (LBF22-LBR44, LBF22-LBR55, and BacLBF-BacLBR) were used for amplification of *leuB*. PCR amplification of *leuB* was performed using 10  $\mu$ l of heated cell supernatant of each strain, 0.5  $\mu$ M of each oligonucleotide primer, and 2.5 units of *AmpliTaq* Gold (Perkin-Elmer Co., CT, U.S.A.) in the following setting: pre-heating (96 °C for 10 min), 40 cycles of DNA denaturation (96 °C for

Table 1. Oligonucleotide primers used for PCR amplification of *leuB*.

Primer	Sequence (5' to 3')*	Location **
LBF22	TGYGARAARTTYGGICA	85-101
LBR44	CCYTTCIACYTGIATYTC	982-998
LBR55	ATRTARTCIACIATCCA	1033-1049
BacLBF	TGGCIATGCGYAARAA	281-296
BacLBR	CCCATRGARCCRCTGAT	754-770

\* Y=T or C, R=A or G, and I=inosine.

\*\* The location in the nucleotide sequence of *B. fragilis* strain YCH 46 *leuB* gene (13) is shown.

1 min)/primer annealing (50 for 1 min)/DNA extension (72 for 1 min). PCR products were separated by 2.0% agarose-gel electrophoresis and visualized by ethidium bromide staining. The LBF22-LBR 55-generated *leuB* products were excised from agarose gel and purified using a QIA quick gel extraction kit (QIAGEN, Inc., CA, U.S.A.) according to the manufacturer's instructions. DNA concentration was determined spectrophotometrically, and 200 ng of each purified product was used for sequencing analysis. Nucleotide sequences of the internal portions of *leuB* genes of *B. fragilis* group species were determined on both strands with an automated sequencer (Genetic Analyzer 310, Applied Biosystems Inc., Foster City, Calif., U.S.A.) using BigDye terminator chemistry (Applied Biosystems). Nucleotide sequences obtained were aligned by using the multiple sequence alignment program Clustal W (11). The *B. fragilis* group-specific regions within *leuB* were determined by homology searches of the sequences obtained against public databases by BLASTN (12).

Isolation of bacteria from rat feces

One piece of fresh rat feces (ca. 0.1 g) was suspended in anaerobic solution (33 mM KH<sub>2</sub>PO<sub>4</sub>, 42 mM Na<sub>2</sub>HPO<sub>4</sub>, 0.05% L-cystein-HCl, 0.05% Tween 80, and 0.1% agar), and serial 10-fold dilutions were prepared with the anaerobic solution. A part of each dilution (0.1ml) was plated on GAM agar plates and incubated anaerobically for 48 hrs. Seventy-five colonies were randomly

selected and isolated on fresh GAM agar plates. Gram-staining was performed on all of the isolates, and biochemical tests using an API 20 A system (Bio Mérieux S.A., France) were performed on obligate anaerobes for species identification.

Nucleotide sequence accession numbers

The partial *leuB* sequences that we determined in this study have been assigned the following GenBank accession numbers : AB196153 (*B. caccae* strain JCM 9498), AB196154 (*B. distasonis* strain ATCC8503), AB196155 (*B. eggerthii* strain ATCC27754), AB196149 (*B. fragilis* strain YCH1), AB196148 (*B. fragilis* strain ATCC25285), AB196150 (*B. fragilis* strain YCH46), AB196152 (*B. ovatus* strain ATCC8483), AB196151 (*B. ovatus* strain KYU7), AB196156 (*B. stercoris* strain JCM9496), AB196157 (*B. thetaiotaomicron* strain ATCC29148), AB 196158 (*B. thetaiotaomicron* strain KYU12), AB196159 (*B. thetaiotaomicron* KYU16), AB 196160 (*B. uniformis* strain ATCC8492), AB196164 (*B. uniformis* strain BU1001), AB196163 (*B. uniformis* strain OMC37), AB196161(*B. vulgatus* strain ATCC 8482), and AB196162 (*B. vulgatus* strain KYU10).

RESULTS

Partial sequencing of the leuB genes of B. fragilis group species

Among *Bacteroides* species, the nucleotide sequences




Fig. 1. The nucleotide sequence and the deduced amino acid sequence of the *leuB* gene from *B. fragilis* strain YCH46(13). The variable amino acids when compared with *B. thetaiotaomicron* strain VPI-5482 are circled. The annealing sites of the PCR primers (LBF22, LBR44, and LBR55) used for amplification of *leuB* from *B. fragilis* group species are double-underlined.

of *leuB* have been determined in *B. fragilis* strain YCH 46(13, 14) and *B. thetaiotaomicron* strain VPI-5482 (15). The nucleotide sequences of *leuB* in other *B. fragilis* group species must be obtained to design universal primers for all species in the *B. fragilis* group. Firstly, we designed PCR primers to amplify the *leuB* gene from *B. fragilis* group species other than *B. fragilis* and *B. thetaiotaomicron*. The predicted amino acid sequences of *leuB* in *B. fragilis* and *B. thetaiotaomicron* were highly conserved and showed 94.9% identity. The nucleotide sequence identity was 79.9% and a total of 213 mismatches were found within a 1,062-bp *leuB* sequence. Of those, 172 mismatches (80.8%) were present in the third position of the codon, and only 15 of them were responsible for amino acid substitutions. Therefore, we designed degenerate primers (LBF22, LBR44, and LBR55) from the conserved regions between *B. fragilis* and *B. thetaiotaomicron*, in which amino acids with low degeneracy are abundant (Fig. 1).

The results of amplification of *leuB* using the primer set LBF22-LBR44 or LBF22-LBR55 from type strains of 10 *B. fragilis* group species are shown in Table 2. The primer set LBF22-LBR55 generated the band of expected size (965 bp) from all of the type strains tested except for *B. merdae* strain JCM9497. The other primer set, LBF22-LBR44, also failed to yield a product from *B. merdae* strain JCM9497 despite the fact that both primer sets were able to generate an expected product from *B. distasonis*, which is a species close phylogenetically to *B. merdae*. We then amplified *leuB* genes from seventeen strains of nine species in the *B. fragilis* group except for *B. merdae* using the primer set LBF22-LBR55 and determined the nucleotide sequences of the products. Overall nucleotide sequence identities among the strains tested ranged from 77 to 100%.

#### Design of PCR primers specific for species of the *B. fragilis* group

We aligned the partial *leuB* sequences from seventeen strains obtained as described above and searched for the conserved region among the species to design specific and universal primers for *B. fragilis* group species. Degenerate PCR primers, BacLBF and BacLBR, were synthesized on the basis of nucleotide sequences at positions 281-296 and 754-770, respectively, to be used for specific and simultaneous PCR detection of *B. fragilis* group species (Table 1 and Fig. 2).

#### Specificity test of the PCR primer set

To evaluate the specificity of the PCR primer set, BacLBF-BacLBR, we amplified *leuB* genes from 52 strains (37 strains of the *B. fragilis* group, 5 strains of

species closely related to the *B. fragilis* group, and 10 strains of unrelated aerobes) that had been stocked in our laboratory. As shown in Fig. 3 and Table 3, the PCR primer set, BacLBF-BacLBR, produced a 490-bp band from all strains of the *B. fragilis* group species tested except for one strain of *B. distasonis*. In contrast, none of the other species generated a 490-bp band.

The specificity of BacLBF-BacLBR was further examined in a blind fashion using 75 isolates from rat feces (43 aerobes and 32 anaerobes) (Table 4). Although only two *B. fragilis* group species (*B. caccae* and *B. vulgatus*) were identified within the isolates, a *leuB* amplification product was obtained from all strains of the two species. On the other hand, none of the other isolates generated any product except for one Gram-positive rod. This fastidious Gram-positive isolate was lost during cultivation and could not be identified at a species level. These results indicated that the *leuB*-directed PCR using the primer set, BacLBF-BacLBR, is a useful diagnostic tool for rapid detection of medically important *B. fragilis* group species.

#### Sensitivity of the PCR

As shown in Fig. 4, we prepared serial 10-fold dilutions of the *B. fragilis* strain ATCC25285, which contained cells from  $3.9 \times 10^1$  to  $3.9 \times 10^6$  CFU and were used as templates for amplification of *leuB* with the primer set BacLBF-BacLBR. The detection limit of *leuB*-directed PCR using BacLBF and BacLBR was  $3.9 \times 10^3$  CFU.

## DISCUSSION

*B. fragilis* group species are medically important Gram-negative obligate anaerobes and are often associated with a variety of human infections. Since local *Bacteroides* infection often causes severe bacteremia in immunocompromised patients and since *B. fragilis* group species are the most resistant to antimicrobial agents among all anaerobes, rapid and accurate differentiation of anaerobic isolates is needed for effective management of infected patients. However, phenotypic tests for anaerobes by traditional culture methods used in clinical laboratories are time-consuming. In addition, comparisons of the results from 16S rDNA sequence based-identification and phenotypic tests revealed that only 45.5% of *B. caccae*, 53.3% of *B. ovatus*, 66.7% of *B. merdae*, and 78.6% of *B. thetaiotaomicron* were correctly identified by phenotypic tests (16). PCR-based rapid diagnosis has been targeted to *B. fragilis* because this species is the most virulent among *Bac-*

Table 2. Results of PCR amplification of *leuB* from *Bacteroides fragilis* group species.

Bacterial strain	Primer set used	
	LBF22-LBR44	LBF22-LBR55
<i>B. caccae</i> JCM9498	NT	+
<i>B. distasonis</i> ATCC8503	+	+
<i>B. eggerthii</i> ATCC27754	NT	+
<i>B. fragilis</i> ATCC25285	NT	+
<i>B. merdae</i> JCM9497		
<i>B. ovatus</i> ATCC8483	NT	+
<i>B. stercoris</i> JCM9496	NT	+
<i>B. thetaiotaomicron</i> ATCC29148	NT	+
<i>B. uniformis</i> ATCC8492	NT	+
<i>B. vulgatus</i> ATCC8482	NT	+

NT, not tested.

Plus and minus indicate whether visible and invisible bands produced by the PCR, respectively.




Fig. 2. Alignment of partial *leuB* sequences of *B. fragilis* group species. Numbers on both sides indicate the nucleotide positions in *leuB* of *B. fragilis* strain YCH46. The annealing sites of the primers BacLBF and BacLBR are indicated and double-underlined. Asterisks indicate the conserved nucleotides among all of the strains tested.




Fig. 3. Specificity of the *leuB*-directed PCR using BacLBF and BacLBR. Lanes : 1, *B. caccae* JCM9498 ; 2, *B. distasonis* ATCC8503 ; 3, *B. eggerthii* ATCC27754 ; 4, *B. fragilis* ATCC25285 ; 5, *B. merdae* JCM9497 ; 6, *B. ovatus* ATCC8483 ; 7, *B. stercoris* JCM9496 ; 8, *B. thetaiotaomicron* ATCC29148 ; 9, *B. uniformis* ATCC8492 ; 10, *B. vulgatus* ATCC8482 ; 11, *Por. asaccharolytica* ATCC25260 ; 12, *Por. gingivalis* 381 ; 13, *Prevotella corporis* JCM8529 ; 14, *Prevotella melaninogenica* ATCC25845 ; 15, *Escherichia coli* W3110 ; M, molecular size markers.

Table 3. Specificity tests of the oligonucleotide primers, BacLBF and BacLBR, using phenotypically characterized bacterial strains.

Species classification	No. of strains	<i>leuB</i> -directed PCR	
		No. of positive	No. of negative
<b><i>Bacteroides fragilis</i> group</b>			
<i>B. caccae</i>	1	1	0
<i>B. distasonis</i>	6	5	1
<i>B. eggerthii</i>	1	1	0
<i>B. fragilis</i>	9	9	0
<i>B. merdae</i>	1	1	0
<i>B. ovatus</i>	3	3	0
<i>B. stercoris</i>	1	1	0
<i>B. thetaiotaomicron</i>	5	5	0
<i>B. uniformis</i>	4	4	0
<i>B. vulgatus</i>	6	6	0
<b>Species closely related to <i>B. fragilis</i> group</b>			
<i>Porphyromonas asaccharolytica</i>	1	0	1
<i>Por. gingivalis</i>	1	0	1
<i>Por. endodontalis</i>	1	0	1
<i>Prevotella corporis</i>	1	0	1
<i>Prevotella melaninogenica</i>	1	0	1
<b>Unrelated aerobic species</b>	<b>10</b>	<b>0</b>	<b>10</b>

*teroides* and accounts for 63% of all the *B. fragilis* group isolates (17). However, a recent survey has revealed that isolation rates of *Bacteroides* species other than *B. fragilis* have been increasing in the past ten years. It is therefore important to develop a rapid and accurate technique for identification of all *B. fragilis* group species for appropriate treatment of anaerobic infections.

Liu *et al.* have recently reported *B. fragilis* group-specific PCR primers designed on the basis of the 16S rDNA sequence (16). However, due to the high sequence similarities of 16S rDNA among closely related species, this primer set also reacts to several strains of *Porphyromonas* and *Prevotella*. In this study, we synthesized *B. fragilis* group-specific PCR primers

(BacLBF-BacLBR) on the basis of  $\beta$ -isopropylmalate dehydrogenase gene. This primer set showed high specificity for *B. fragilis* group species and did not cross-react with closely related species in *Porphyromonas* and *Prevotella*.

The minimum level of *leuB*-directed PCR using BacLBF and BacLBR was  $3.9 \times 10^3$  cells. It seems to be difficult to apply *leuB*-directed PCR for direct detection of *Bacteroides* in specimens that contain relatively small numbers of bacterial cells, such as in a blood sample from a patient with bacteremia. However, the PCR will be a useful diagnostic tool when it is applied to specimens that contain relatively large numbers of bacterial cells such as samples of pus or drainage fluids. It will also be




Fig. 4. Sensitivity of the *leuB*-directed PCR using BacLBF and BacLBR on the basis of viable cell number of *B. fragilis* strain ATCC 25285. Lanes: 1,  $3.9 \times 10^6$ ; 2,  $3.9 \times 10^5$ ; 3,  $3.9 \times 10^4$ ; 4,  $3.9 \times 10^3$ ; 5,  $3.9 \times 10^2$ ; 6,  $3.9 \times 10^1$ ; 7,  $3.9 \times 10^0$ ; 8,  $3.9 \times 10^{-1}$ ; M, molecular size markers.

Table 4. Specificity tests of the oligonucleotide primers, BacLBF and BacLBR, using fecal isolates from a rat.

Isolate classification *	No. of isolates tested	<i>leuB</i> -directed PCR	
		No. of positive	No. of negative
<b>Aerobes</b>			
Gram-positive cocci	8	0	8
Gram-positive rods	24	1	23
Gram-negative rods	11	0	11
<b>Anaerobes</b>			
Gram-positive rods			
<i>Bifidobacterium</i> spp	3	0	3
<i>Lactobacillus fermentum</i>	1	0	1
Unidentified	2	0	2
Gram-negative rods			
<i>Bacteroides caccae</i>	15	15	0
<i>Bacteroides vulgatus</i>	9	9	0
<i>Porphyromonas gingivalis</i>	2	0	2

\*Anaerobic isolates were identified at the species level by biochemical tests using an API20 A system.

useful to confirm the results obtained by traditional biochemical tests. The diagnostic potentials of the PCR developed in this study are expected to be increased by simultaneous use with species-specific primer sets such as *B. fragilis*-specific primers designed from *glnA* (5) or *nanH* (6, 7).

## ACKNOWLEDGEMENTS

The authors thank Miss Y. Tsutishashi, Mr. K. Nitta, and Mr. K. Honjo for their technical assistance. This work was supported in part by a Grant-in-Aid for Developmental Scientific Researches (B) (No.06557020) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

## REFERENCES

1. Salyers AA : *Bacteroides* of the human lower intestinal tract. Ann Rev Microbiol 38 : 293-313, 1984
2. Collins MD, Shah, HN : Recent advances in the taxonomy of the genus *Bacteroides*. In : Borriello SP, Hardie JM, Drasar BS, Duerden BI, Hudson MJ, Lyson RJ, eds. Recent Advances in Anaerobic Bacteriology. Martinus Nijhoff Publishers, Dordrecht, The Netherlands, 1987, pp.249-258
3. Frank BP, Dennis LK : *Bacteroides fragilis* subspecies in clinical isolates. Ann Intern Med 86 : 569-571, 1977
4. Goldstein EJ : Anaerobic bacteremia. Clin Infect Dis 23 : 97-101, 1996
5. Yamashita Y, Kohno S, Koga H, Tomono K, Kaku M : Detection of *Bacteroides fragilis* in clinical

- specimens by PCR. J Clin Microbiol 32 : 679-683, 1994
6. Jotwani R, Kato N, Kato H, Watanabe K, Ueno K : Detection of *Bacteroides fragilis* in clinical specimens by polymerase chain reaction amplification of the neuraminidase gene. Curr Microbiol 31: 215-219, 1995
  7. Kuwahara T, Akimoto S, Ugai H, Kamogashira T, Kinouchi T, Ohnishi Y : Detection of *Bacteroides fragilis* by PCR assay targeting the neuraminidase-encoding gene. Lett Appl Microbiol 22 : 361-365, 1996
  8. Aldridge KE, O'Brien M : *In vitro* susceptibilities of the *Bacteroides fragilis* group species : change in isolation rates significantly affects overall susceptibility data. J Clin Microbiol 40 : 4349-4352, 2002
  9. Varel VH, Bryant MP : Nutritional features of *Bacteroides fragilis* subsp. *fragilis*. Appl Microbiol 28 : 251-257, 1974
  10. Sambrook J, Fritsch EF, Maniatis T : Molecular Cloning. A Laboratory Manual, Cold Spring Harbor Laboratory Press, New York, 1989
  11. Thompson JD, Higgins DG, Gibson TJ : CLUSTAL W : improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. Nucleic Acids Res 22 : 4673-4680, 1994
  12. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ : Gapped BLAST and PSI-BLAST : a new generation of protein database search programs. Nucleic Acids Res 25 : 3389-3402, 1997
  13. Sarker MR, Akimoto S, Ugai H, Kuwahara T, Ohnishi Y : Nucleotide sequence of the gene encoding  $\beta$ -isopropylmalate dehydrogenase (*leuB*) from *Bacteroides fragilis*. Microbiol Immunol 39 : 525-529, 1995
  14. Kuwahara T, Yamashita A, Hirakawa H, Nakayama H, Toh H, Okada N, Kuhara S, Hottori M, Hayashi T, Ohnishi Y : Genomic analysis of *Bacteroides fragilis* reveals extensive DNA inversions regulating cell surface adaptation. Proc Natl Acad Sci USA 101 : 14919-14924, 2004
  15. Xu J, Bjursell MK, Himrod J, Deng S, Carmichael LK, Chiang HC, Hooper LV, Gordon JI : A genomic view of the human-*Bacteroides thetaiotaomicron* symbiosis. Science 299 : 2074-2076, 2003
  16. Liu C, Song Y, McTeague M, Vu AW, Wexler H, Finegold SM : Rapid identification of the species of the *Bacteroides fragilis* group by multiplex PCR assays using group- and species-specific primers. FEMS Microbiol Lett 222 : 9-16, 2003
  17. Brook I : Pathogenicity of the *Bacteroides fragilis* group. Ann Clin Lab Sci 19 : 360-376, 1989